

ORIGINAL ARTICLE

**Comparison of Antioxidant and Anti-Tyrosinase Activity  
between Black Soybean (*Glycine max* (L.) Merr.) and Daidzein**

**Delken Kuswanto<sup>1</sup>, I Nyoman Ehrich Lister<sup>2</sup>, ErmiGirsang<sup>3</sup>, Ali NapihNasution<sup>4</sup>,  
Wahyu Widowati<sup>5</sup>**

<sup>1</sup>Master Program of Biomedical Sciences, Universitas Prima Indonesia, Medan, Indonesia,

<sup>2</sup>Faculty of Medicine, Universitas Prima Indonesia, Medan, Indonesia,

<sup>3</sup>Faculty of Public Health, Universitas Prima Indonesia, Medan, Indonesia,

<sup>4</sup>Department of Tropical Medicine, Faculty of Medicine, Universitas Prima Indonesia, Medan, Indonesia,

<sup>5</sup>Faculty of Medicine, Maranatha Christian University, Bandung, Indonesia

Corresponding E-mail: [ermigirsang@unprimdn.ac.id](mailto:ermigirsang@unprimdn.ac.id)

**Abstract:** Free radicals in the body will increase with excessive ultraviolet (UV) light exposure, induce oxidative stress with the formation of Reactive Oxygen Species (ROS). A visible effect on skin tissue known as photoaging, including the process of melanogenesis catalyzed by the tyrosinase enzyme, risked for pigmentation or melanoma disorder.

The use of natural ingredients has been widely used by the community to prevent the aging process. Natural compounds from a plant can be a source of antioxidants and have anti-aging abilities through inhibition of the tyrosinase enzyme. Black soybean (*Glycine max* (L.) Merr.) are high in the isoflavone compound, one of which is daidzein.

This study evaluates the antioxidant and anti-aging potential of black soybean extract with daidzein. Antioxidants using the DPPH method and anti-aging tests carried out, namely the inhibition of the tyrosinase enzyme, are very important in the aging process.

Daidzein has an IC<sub>50</sub> value of DPPH scavenging activity around 109.34±2.80 µg/mL lower than black soybean extract with 116.52±2.50 µg/mL. Results on tyrosinase enzyme inhibition activity, black soybean extract had an IC<sub>50</sub> value of 70.71±1.83 µg/mL lower compared to daidzein with 72.65±2.81 µg/mL. In contrast to the better antioxidant activity of daidzein, the black soybean extract is more potential to inhibit the enzyme tyrosinase.

**Keywords:** black soybean, daidzein, antioxidant, anti-aging, anti-tyrosinase

## INTRODUCTION

The aging process is a physiological process that will occur in every living thing and the skin is a body tissue that gives a visual image of aging. Skin aging is a complex process influenced by genetic,

environmental and nutritional properties.<sup>(1)</sup>

Excessive ultraviolet (UV) light exposure increases the contribution of free radicals known as Reactive Oxygen Species (ROS). This certainly influences the process of melanogenesis on the skin catalyzed by the

enzyme tyrosinase. This enzyme regulates skin pigmentation through the synthesis of melanin. Increased UV radiation will increase the synthesis of melanin which causes a risk of pigmentation or melanoma disorders.<sup>(2)</sup>

Various types of modern medicine have been applied to prevent premature aging, one of which is cosmetic. Most cosmetics have side effects and are not safe, so natural ingredients are needed.<sup>(3)</sup> Currently, natural herbal plants are getting attentions to solve problems. The use of traditional material has long been used by the community as an alternative treatment, for disease prevention, healing, health recovery and improving health status. Based on WHO data, mentioning Africa, Asia and Latin America using traditional materials as a complement to their primary treatment.<sup>(4)</sup>

The use of natural materials as traditional ingredients has also been carried out by Indonesia since the days of our ancestors. The tendency of the use of traditional ingredients in the world is caused by plants containing compounds that are efficacious in medicine known as phytochemical compounds, namely a group of natural compounds that can be used to maintain health and treat diseases due to their antioxidant activity.<sup>(5)</sup>

In Indonesia, black soybeans are one of the plants which are the main food commodities after rice and corn.<sup>(6)</sup> Black soybeans contain high carbohydrates, vitamins, minerals, and proteins.<sup>(7)</sup> Besides, the advantages of black soybean plants have the most content of isoflavones (one of them

is *daidzein*) which acts as an antioxidant, only produced from plants and found in legumes.<sup>(8)</sup> Lately, antioxidants have become somewhat interesting in the medical world, known to have the effect of preventing premature aging (anti-aging) in fighting free radicals.<sup>(9)</sup>

It is necessary to conduct a study of the activity of black soybean extract as an antioxidant and anti-tyrosinase with the *daidzein* comparison compound. This needs to be done to prove the effectiveness of the black soybeans so that black soybeans can be used as natural food to slow down the aging process.

## METHODS

### Preparation of Black Soybean (*Glycine max* (L.) Merr.) Extract

Black soybean seeds (*Glycine max* (L.) Merr.) are washed and dried. *Glycine max* (L.) Merr.) were milled and extracted using 70% ethanol by maceration method for 3 days at room temperature. The filtrate was evaporated with a rotary evaporator so that a thick extract was obtained in paste form.<sup>(10)</sup>

### Qualitative Phytochemical Screening Assay

#### Flavonoid Identification

The test tube containing HCl 2N was dissolved by mixing 10 mg *Glycine max* (L.) Merr.) into it. Mg/Zn is added enough and heated for 10-15 minutes. Amyl alcohol 1 ml added to the test tube has been cooled and filtered. The presence of a red/orange

color indicates the sample contains a flavonoid compound.<sup>(11)</sup>

### Saponin Identification

*Glycine max* (L.) Merr.) (10 mg) was dissolved using ddH<sub>2</sub>O in a test tube, simmer for 5 minutes. HCl 1N is added to the solution which has been filtered and shaken strongly. The presence of foam that is still present and remained stable shows a sample containing a saponin compound.<sup>(11)</sup>

### Phenol Identification

*Glycine max* (L.) Merr.) 10 mg dissolved in 5 ml ddH<sub>2</sub>O, added 500 µl of FeCl<sub>3</sub> 1% solution. The presence of green/red/purple/blue/black color indicated *Glycine max* (L.) Merr.) contains phenol compounds.<sup>(11)</sup>

### Tannin Identification

HCl 2N around 2 mg was dissolved with 10 mg of *Glycine max* (L.) Merr.) in a test tube, heated in a water bath for 30 minutes. Amyl alcohol 500 µl is added after the solution cooled down. Orange/red color in the amyl alcohol layer indicated a tannin compound.<sup>(11)</sup>

### Alkaloid Identification

*Glycine max* (L.) Merr.) 10 mg dissolved in 5 ml ddH<sub>2</sub>O was evaporated in the water bath. The resulting residue is added 5 ml HCl 2N. The solution obtained is divided into 2 test tubes. The first tube is added 3 drops of HCl 2N which functions as a blank. The second tube solution was transferred as much as one drop to the drip

plate, then added 3 drops of Dragendorff reagent. The orange deposits formed indicate the presence of alkaloids.<sup>(11)</sup>

### Steroid/Triterpenoid Identification

The drip plate contains 10 mg *Glycine max* (L.) Merr.), added glacial acetic acid until submerged, left for 10-15 minutes and added with a drop of concentrated H<sub>2</sub>SO<sub>4</sub>. The presence of a blue-green color indicated the sample contains a steroid compound, whereas if it shows purple/red/orange color, the sample contains the triterpenoid compound.<sup>(11)</sup>

### Terpenoid Identification

Vanillin is added enough to the drip plate containing 10 mg of *Glycine max* (L.) Merr.) Addition of one drop H<sub>2</sub>SO<sub>4</sub> into it and homogenize. The presence of purple color contains the terpenoid compound.<sup>(11)</sup>

### DPPH (2,2-diphenyl-1-picryl-hydrazyl-hydrate) Assay

DPPH 0.077mmol 200 µl in methanol was added to each 50 µl of *Glycine max* (L.) Merr.) extract and daidzein respectively on the microplate. The mixture was incubated at room temperature for 30 minutes. DPPH 250 µl for negative control. The absorbance value is measured at a wavelength of 517 nm using a microplate reader.<sup>(12)</sup> DPPH method antioxidant activity (%):

$$DPPH \text{ scavenging activity (\%)} = 1 - \frac{\text{sample}}{\text{control}} \times 100$$

### Tyrosinase Inhibitor Assay

The mixture consisted of 20 µl *Glycine max* (L.) Merr.), 20 µl Tyrosinase from Mushroom (125 U/ml) enzyme, and 140 µl potassium phosphate buffer (20 mM, pH 6.8) were incubated at room temperature for 15 minutes. The control contained 20 µl enzyme and 160 µl phosphate buffer, then added 20 µl of L-DOPA substrate (1.5 mM) and incubated again at 10 minutes. The absorbance is measured using a wavelength of 470 nm.<sup>(13)</sup> The percentage of tyrosinase inhibitor activity is calculated using the formula:

$$\text{scavenging activity (\%)} = \frac{\text{control} - \text{sample}}{\text{control}} \times 100$$

## RESULTS

### Phytochemical Screening of *Glycine Max* (L.) Merr.)

Phytochemical screening of black soybean seed extract (*Glycine max* (L.) Merr.) includes alkaloid, tannin, steroid/triterpenoid, phenol, saponin, flavonoid, and terpenoid. Some of this substrate or component can affect the ability of antioxidants, such as alkaloid and terpenoid. The phytochemical test results provide an overview of the classes of compounds contained in *Glycine max* (L.) Merr.) which is saponin, tannin<sup>8</sup>, triterpenoid, terpenoid, alkaloid, and can be seen in Table 1.

### DPPH Scavenging Activity

The result of the antioxidant activity test using the DPPH<sup>15</sup> method in various concentrations can be seen in Table 2 and

Figure 1 with Tukey HSD statistical analysis. In the process of analyzing whether *Glycine max* (L.) Merr.) and the daidzein compound has antioxidant activity, it can be seen as a linear regression equation to find IC<sub>50</sub> values in Table 3.

**Table 1. Phytochemical Screening of *Glycine max* (L.) Merr.)**


Contain	Result
Flavonoid	-
Saponin	+
Phenol	-
Tanin	+
Steroid/triterpenoid	-/+
Terpenoid	+
Alkaloid	+

**Table 2. DPPH Scavenging Activity**

Concentration (µg/ml)	Average DPPH Scavenging (%)	
	<i>Glycine max</i> (L.) Merr.	Daidzein
200	65.01 ± 2.02 <sup>c</sup>	67.87 ± 1.02 <sup>e</sup>
100	49.85 ± 0.84 <sup>b</sup>	50.88 ± 0.98 <sup>d</sup>
50	34.52 ± 1.25 <sup>a</sup>	37.88 ± 0.46 <sup>c</sup>
25	31.59 ± 1.10 <sup>a</sup>	30.96 ± 0.44 <sup>b</sup>
12.5	31.97 ± 1.99 <sup>a</sup>	29.24 ± 0.51 <sup>a,b</sup>
6.25	29.90 ± 3.07 <sup>a</sup>	27.57 ± 1.23 <sup>a</sup>

Notes:

3) Data were presented as mean ± standard deviation. Different small letters in the same column are significant at P < .05 (Tukey HSD Post Hoc test).



**Figure 1. Various concentrations in DPPH method**



The antioxidant activity test used DPPH scavenging activity parameters as seen in Table 2, where *Glycine max* (L.) Merr). and *daidzein* had the highest antioxidant activity at a concentration of 200 µg/ml, each at 65.01±2.02 % and 67.87±1.02 %. The lower the concentration shows the small the percentage of antioxidant activity of *Glycine max* (L.) Merr). and *daidzein*.

Tabel 3 shows that *Glycine max* (L.) Merr). and *daidzein* with DPPH scavenging activity

obtained IC<sub>50</sub> values averaging 116.52±2.50 µg/ml and 109.34±2.80 µg/ml.

### Tyrosinase Inhibitor Activity

The test results of tyrosinase enzyme inhibition activity in various concentrations can be seen in Table 4 and Figure 2. The test results of tyrosinase inhibition activity in both samples were linear regression equations to find the IC<sub>50</sub> value in Table 5.

Table 3. The IC<sub>50</sub> Value of DPPH Scavenging- Activity

Sample	Equation	R <sub>2</sub>	IC <sub>50</sub> (µg/ml)	Average IC <sub>50</sub> (µg/ml)
Glycine max	1 Y = 0.1735x + 30.066	0.97	114.89	116.52 ± 2.50
	2 Y = 0.1766x + 28.914	0.97	119.40	
	3 Y = 0.2123x + 25.526	0.99	115.28	
	Mean Y = 0.1875x + 28.168	0.98	116.44	
Daidzein	1 Y = 0.2062x + 27.342	0.99	109.88	109.34 ± 2.80
	2 Y = 0.2190x + 26.718	0.99	106.31	
	3 Y = 0.2112x + 26.380	0.99	111.84	
	Mean Y = 0.2121x + 26.813	0.99	109.32	
Glycine max	1 Y = 0.1735x + 30.066	0.97	114.89	116.52 ± 2.50
	2 Y = 0.1766x + 28.914	0.97	119.40	
	3 Y = 0.2123x + 25.526	0.99	115.28	
	Mean Y = 0.1875x + 28.168	0.98	116.44	
Daidzein	1 Y = 0.2062x + 27.342	0.99	109.88	109.34 ± 2.80
	2 Y = 0.2190x + 26.718	0.99	106.31	
	3 Y = 0.2112x + 26.380	0.99	111.84	
	Mean Y = 0.2121x + 26.813	0.99	109.32	

Table 4. Tyrosinase Inhibition Assay

Concentration ( $\mu\text{g/ml}$ )	Tyrosinase Inhibition Activity (%)	
	Glycine max (L.) Merr.	Daidzein
100	$60.77 \pm 1.12^d$	$57.47 \pm 1.16^e$
50	$42.66 \pm 0.66^c$	$44.36 \pm 0.97^d$
25	$31.68 \pm 0.57^b$	$36.94 \pm 1.43^c$
12.5	$30.57 \pm 0.80^b$	$32.02 \pm 0.72^b$
6.25	$27.10 \pm 0.73^a$	$26.20 \pm 0.58^a$
3.13	$27.48 \pm 0.72^a$	$24.73 \pm 0.42^a$

Notes:

Data were presented as mean  $\pm$  standard deviation. Different small letters in the same column are significant at  $P < 0.05$  (Tukey HSD Post Hoc test).




Figure 2. Various concentrations in tyrosinase inhibition activity

12

Table 5. The  $\text{IC}_{50}$  Value of Tyrosinase Inhibitor Activity

Sample	Equation	$R^2$	$\text{IC}_{50}$ ( $\mu\text{g/ml}$ )	Average $\text{IC}_{50}$ ( $\mu\text{g/ml}$ )
Glycine max	1 $Y = 0.3676x + 24.629$	0.99	69.02	$70.71 \pm 1.83$
	2 $Y = 0.3374x + 25.489$	0.99	72.65	
	3 $Y = 0.3483x + 25.454$	0.99	70.47	
	Mean $Y = 0.3511x + 25.191$	0.99	70.66	
Daidzein	1 $Y = 0.3385x + 25.966$	0.97	71.00	$72.65 \pm 2.81$
	2 $Y = 0.3131x + 26.236$	0.97	75.90	
	3 $Y = 0.3327x + 26.360$	0.97	71.06	
	Mean $Y = 0.3281x + 26.187$	0.97	72.58	

Based on Tabel 4 and Figure 2, the tyrosinase inhibition activity of Glycine max (L.) Merr. was highest at a concentration of 100  $\mu\text{g/ml}$  of  $60.77 \pm 1.12\%$ . It was different at a concentration of 50  $\mu\text{g/ml}$ , 25  $\mu\text{g/ml}$ , and 12.5  $\mu\text{g/ml}$  where tyrosinase inhibition activity by Glycine max (L.) Merr. decreased with a percentage of  $42.66 \pm 0.66\%$ ,  $31.68 \pm 0.57\%$ , and  $30.57 \pm 0.80\%$ . Daidzein compound inhibition activity was higher at the same concentration of  $44.36 \pm 0.97\%$ ,  $36.94 \pm 1.43\%$  and  $32.02 \pm 0.72\%$  and higher than the daidzein compound at the same concentration of  $57.47 \pm 1.16\%$ .

Table 5 shows the Glycine max (L.) Merr. and daidzein compound showed that

both could inhibit the tyrosinase enzyme with  $\text{IC}_{50}$  at  $70.71 \pm 1.83 \mu\text{g/ml}$  and  $72.65 \pm 2.81 \mu\text{g/ml}$ .

## DISCUSSION

The phytochemical screening is carried out as a preliminary stage which aims to determine the class of bioactive compounds in a plant studied. The selection of the solvent and extraction method is the most important factor in conducting a phytochemical screening assay. The phytochemical screening is done qualitatively using a color reagent and see the color changes that occur. Black soybean seeds (Glycine max (L.) Merr.) is known as

nuts with nutritional value and is rich in protein. Not identified in our research, daidzein is one of the highest bioactive compounds in black soybean seeds and acts as an antioxidant<sup>(14)</sup> Therefore, *daidzein* was used as a comparison.

DPPH method is a simple method used to determine the antioxidant activity of plant material by capturing hydrogen atoms from the component of the extract of the plant. DPPH is sensitive to light, oxygen, and pH. However, it is stable in a radical form so that it may be quite an accurate measurement of antioxidant activity.<sup>(15)</sup>

The lower the concentration shows the smaller the percentage of antioxidant activity of *Glycine max* (L.) Merr). and *daidzein*. Antioxidant activity with DPPH scavenging on daidzein compound was higher than *Glycine max* (L.) Merr). at concentration of 200 µg/ml, 100 µg/ml, and 50 µg/ml.

*Glycine max* (L.) Merr). and *daidzein* with DPPH scavenging activity obtained IC<sub>50</sub> values averaging 116.52±2.50 µg/ml and 109.34±2.80 µg/ml. The result was different from the previous study conducted by Fidrianny et al., the IC<sub>50</sub> result by *Glycine max* (L.) Merr. was 3.13 µg/ml. (16) Kladna et al. reported the antioxidant test for several phytoestrogen compounds, one of which is daidzein, showed an IC<sub>50</sub> result of 2.81±0.03 mM through the DPPH scavenging method. (17) Both of these can be caused by different procedures in the extracted method, the solvent used and the

different wavelengths in the DPPH method Tyrosinase or polyphenol oxidase is an oxidoreductase that participates in the biosynthesis of melanin, the main pigment found in hair, eyes, and skin. The tyrosinase enzyme reaction with the L-DOPA substrate produces an orange color. Inhibition activity of tyrosinase enzyme is characterized by a reduction in the orange color formed or the result of a clearer colored reaction. This also marks the presence of antioxidant activity in the reaction.<sup>(13)</sup>

*Glycine max* (L.) Merr). and *daidzein* compound showed that both could inhibit the tyrosinase enzyme with IC<sub>50</sub> at 70.71±1.83 µg/ml and 72.65±2.81 µg/ml. The tyrosinase enzyme inhibition study showed anti-tyrosinase activity produced by *Glycine max* (L.) Merr). and *daidzein* compound, according to Lai et.al (2012) using black soybean sprouts extract which has a strong activity of 98% in inhibiting the tyrosinase enzyme and can be used as a lightening agent in skin cosmetic product.<sup>(18)</sup> Another study by Sitanggang et al. also showed a positive correlation in inhibiting >50% of the tyrosinase enzyme activity by yellow soybean.<sup>(19)</sup>

The effectiveness of a compound in biological or biochemical functions that can inhibit the oxidation process by 50% (IC<sub>50</sub>) is classified in several groups including <50 µg/ml (very strong); 50-100 µg/ml (strong); 101-150 µg/ml (medium); >150 µg/ml (weak).<sup>(20)</sup> In this study, the IC<sub>50</sub> value in each test is grouped in Table 6.

Table 6. The IC<sub>50</sub> Value of *Glycine max* (L.) Merr.) and *Daidzein*

Sample	Assay	IC <sub>50</sub> (µg/ml)	Activity
Glycine max (L.) Merr.	DPPH assay	116.52 ± 2.50	Medium
Daidzein		109.34 ± 2.80	Medium
Glycine max (L.) Merr.	Tyrosinase inhibition assay	70.71 ± 1.83	Strong
Daidzein		72.65 ± 2.81	Strong

## CONCLUSIONS

<sup>13</sup> Bioactive compounds that presence in black soybean (*Glycine max* (L.) Merr.) are saponin, t<sub>4</sub>nin, triterpenoid, terpenoid, an alkaloid. *Glycine max* (L.) Merr.) and *daidzein*, both have antioxidant and tyrosinase inhibition activity as anti-aging. From the result, it shows that *Glycine max* (L.) Merr.) has tyrosinase inhibitor activity better than *daidzein*. Therefore *Glycine max* (L.) Merr.) can be developed into anti-aging ingredients in the product.

## ACKNOWLEDGEMENTS

<sup>3</sup> The author would like to thank DR. Ermi Girsang, SKM, M.Kes and dr. Ali Napih Nasution, M.K.M as a mentor, DR. dr. I Nyoman Ehrich Lister, M.Kes., AIFM as head of the study program, dr. Linda Chiuman, M.K.M as dean, DR. Chrismis Nova linda Ginting, M.Kes as chancellor, and Aretha Medika Utama for supporting this research.

## REFERENCES

1. Tobin DJ. Introduction to skin aging. Journal of tissue. 2016; 26(1):1-5.
2. Lai X, Wichers HJ, Soler-Lopez M, Dijkstra BW. Structure and Function of Human Tyrosinase and Tyrosinase-Related Proteins. Chem. Eur. J. 2018; 24:50-53.
3. Siti ZR, Norkhadijah SIS, Praveena SM. Hazardous Ingredients in Cosmetics and Personal Care Products and Health Concern: A Review. Public Health Research. 2015; 5(1):1.
4. World Health Organization. Traditional Medicine Strategy. 2013.
5. Iris FFB & Sissi WG. Herbal medicine :biomolecular and clinical aspects, 2nd Ed. New York: Taylor & Francis Group, (Chapter 1). 2011.
6. Arifin AS. Kajian Morfologi Anatomi dan Agronomi antara Kedelai Sehat dengan Kedelai Terserang Cowpea Mild Mottle Virus serta Pemanfaatannya sebagai Bahan Ajar Sekolah Menengah Kejuruan. Jurnal Pendidikan Sains. 2013; 1(2):115.
7. Alghamdi et al. Comparative phytochemical profiling of different soybean (*Glycine max* (L.) Merr) genotypes using GC-MS. Saudi Journal of Biological Sciences. 2018; 25:15.
8. Sumardi et al. Potential of local black soybean as a source of the isoflavones *daidzein* and genistein. International Food Research Journal. 2017; 24(5):2140-2141.
9. Garg C, Khurana P, Garg M. Molecular Mechanisms of Skin Photoaging and



- Plant Inhibitors. *International Journal of Green Pharmacy*. 2017; 11(2):219-220.
10. Farnakope Herbal Indonesia. Jakarta: National Health Ministry; 2013.
11. Widowati et al. Antioxidant and Anti Aging Assays of *Oryza Sativa* Extracts, Vanillin and Coumaric Acid. *Journal of Natural Remedies*. 2016; 16(3):89-90.
12. Widowati et al. Antioxidant and Antiaging Activities of *Jasminum Sambac* Extract, and its Compounds. *Journal of Reports in Pharmaceutical Sciences*. 2018; 23(3):272-273.
13. Tu PTB, Tawata S. Anti-Oxidant, Anti-Aging, and Anti-Melanogenic Properties of the Essential Oils from Two Varieties of *Alpinia zerumbet*. *Molecules*. 2015; 20:16727-16728.
14. Altemimi A, Lakhssassi N, Baharlouei A, Watson DG, Lightfoot DA. Phytochemicals: Extraction, Isolation, and Identification of Bioactive Compounds from Plant Extracts. *Plants*. 2017; 6:2-4.
15. Al-Kawaz HS, Al-Mashhady LAM. Evaluation of The Phytochemical Constituents and Oxidant-Amtioxidant Status for *Actinidiadeliciosa* Extracts. *International Journal of Pharmacy & Therapeutics*. 2016;7(1):35-36.
16. Fidrianny I, Elvianna D, Ruslan K. In Vitro Antioxidant Activities in Various Beans Extracts of Five Legumes from West of Java-Indonesia Using DPPH and ABTS Methods. *International Journal of Pharmacognosy and Phytochemical Research*. 2016; 8(3):471-474.
17. Kladna A, Berczynski P, Kruk I, Piechowska T, Aboul-Enein HY. Studies on the antioxidant properties of some phytoestrogens. *Luminescence*. 2016; 31(6).
18. Lai et al. Study of Active Ingredients in Black Soybean Sprouts and Their Safety in Cosmetic Use. *Molecules*. 2012; 17:11672.
19. Sitanggang SSDH, Yanti, Lay BW. Effect of Purified Protein Fractions from Soybean on Tyrosinase Inhibition. *Sch. Acad. J. Biosci*. 2017; 5(8):594.
20. Budaraga IK, Arnim, Marlinda Y, Bulanin U. Antioxidant Properties of Liquid Smoke Production Variation of Pyrolysis Temperature Raw and Different Concentration. *International Journal of PharmTech Research*. 2016; 9(6):370.

## ORIGINALITY REPORT

21%

SIMILARITY INDEX

13%

INTERNET SOURCES

10%

PUBLICATIONS

14%

STUDENT PAPERS

## PRIMARY SOURCES

1

Submitted to Universitas Muhammadiyah  
Sumatera Utara

Student Paper

4%

2

S Prahastuti, M Hidayat, S T Hasianna, W  
Widowati, A Amalia, D T Yusepany, R Rizal, H S  
W Kusuma. " Antioxidant potential ethanolic  
extract of (L.) Merr. Var. Detam and daidzein ",  
Journal of Physics: Conference Series, 2019

Publication

4%

3

asrjetsjournal.org

Internet Source

4%

4

repository.maranatha.edu

Internet Source

1%

5

Submitted to Universitas Prima Indonesia

Student Paper

1%

6

rd.springer.com

Internet Source

1%

7

jurnal.umsu.ac.id

Internet Source

1%

8	repository.unand.ac.id Internet Source	1 %
9	Submitted to iGroup Student Paper	1 %
10	www.mdpi.com Internet Source	1 %
11	Submitted to Universidad Autónoma de Nuevo León Student Paper	<1 %
12	Submitted to Universitas Brawijaya Student Paper	<1 %
13	Yongxiang Hong, Xixi Cai, Biao Shao, Jing Hong, Shaoyun Wang, Pingfan Rao. "Isolation of a thermostable trypsin inhibitor with exploitable potential", European Food Research and Technology, 2013 Publication	<1 %
14	Submitted to Taylor's Education Group Student Paper	<1 %
15	Fifi Afiati, Siti Fakhriyyah Firza, Kusmiati, Lisana S. Aliya. "The effectiveness $\beta$ -glucan of shiitake mushrooms and <i>Saccharomyces cerevisiae</i> as antidiabetic and antioxidant in mice Sprague Dawley induced alloxan", AIP Publishing, 2019 Publication	<1 %

16

Wanying He, Xiaoyan Li, Ying Peng, Xiaoyan He, Siyi Pan. "Anti-Oxidant and Anti-Melanogenic Properties of Essential Oil from Peel of Pomelo cv. Guan Xi", *Molecules*, 2019

Publication

&lt;1 %

17

[publications.waset.org](http://publications.waset.org)

Internet Source

&lt;1 %

18

Submitted to Kyungpook National University

Student Paper

&lt;1 %

19

Submitted to School of Business and Management ITB

Student Paper

&lt;1 %

20

Yongjin Hu. "Characterization of fermented black soybean natto inoculated with *Bacillus natto* during fermentation", *Journal of the Science of Food and Agriculture*, 2010

Publication

&lt;1 %

21

Pier-Giorgio Pietta. "Flavonoids as Antioxidants", *Journal of Natural Products*, 2000

Publication

&lt;1 %

22

Betty Nurhayati, Ira Gustira Rahayu, Sonny Feisal Rinaldi, Wawan Sofwan Zaini et al. "The Antioxidant and Cytotoxic Effects of *Cosmos caudatus* Ethanolic Extract on Cervical Cancer", *The Indonesian Biomedical Journal*, 2018

Publication

&lt;1 %



23

Florence Alesandrini, René Mathis, Ghislaine Van de Syte, Didier Héroutart, Alain Puppo.  
"Possible roles for a cysteine protease and hydrogen peroxide in soybean nodule development and senescence", New Phytologist, 2003

Publication

<1 %

24

Submitted to Yeditepe University

Student Paper

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On