

**Current Clinical Practice**  
*Series Editor: Neil S. Skolnik*

Paul Lyons

# Obstetrics in Family Medicine

A Practical Guide

*Second Edition*



Humana Press

# Current Clinical Practice

## **Series Editor**

Neil S. Skolnik

Temple University, School of Medicine, Abington Jefferson Health,  
Abington, PA, USA

More information about this series at <http://www.springer.com/series/7633>



Paul Lyons

# Obstetrics in Family Medicine

A Practical Guide

Second Edition

 Humana Press

Paul Lyons, MD  
Department of Family Medicine  
University of California, Riverside  
Riverside, CA, USA

Current Clinical Practice  
ISBN 978-3-319-20076-7      ISBN 978-3-319-20077-4 (eBook)  
DOI 10.1007/978-3-319-20077-4

Library of Congress Control Number: 2015941444

Springer Cham Heidelberg New York Dordrecht London  
1st edition: © 2006 Humana Press Inc.

© Springer International Publishing Switzerland 2015

This work is subject to copyright. All rights are reserved by the Publisher, whether the whole or part of the material is concerned, specifically the rights of translation, reprinting, reuse of illustrations, recitation, broadcasting, reproduction on microfilms or in any other physical way, and transmission or information storage and retrieval, electronic adaptation, computer software, or by similar or dissimilar methodology now known or hereafter developed.

The use of general descriptive names, registered names, trademarks, service marks, etc. in this publication does not imply, even in the absence of a specific statement, that such names are exempt from the relevant protective laws and regulations and therefore free for general use.

The publisher, the authors and the editors are safe to assume that the advice and information in this book are believed to be true and accurate at the date of publication. Neither the publisher nor the authors or the editors give a warranty, express or implied, with respect to the material contained herein or for any errors or omissions that may have been made.

Printed on acid-free paper

Humana Press is a brand of Springer  
Springer International Publishing AG Switzerland is part of Springer Science+Business Media  
([www.springer.com](http://www.springer.com))

# Series Editor's Introduction

*Obstetrics in Family Medicine: A Practical Guide, 2nd edition*, by Paul Lyons, MD, is an important book for the approximately 21,000 family physicians in the USA who regularly perform obstetrics, as well as the 7500 physicians currently in residency training for whom obstetrics is considered a core body of knowledge. This title should serve as an important addition to the literature in its scope and orientation.

Dr. Lyons' approach to the material is carefully organized and gives particular attention to the relevant aspects of prenatal and pregnancy care that family doctors need to know. He speaks with clarity of voice as an inner-city family physician who has done a great deal of obstetrics care and understands both the practical realities of the delivery rooms and the evidence based on which obstetrical decisions are made.

*Obstetrics in Family Medicine: A Practical Guide* will find its place in the practices of family doctors who are doing obstetrics, in the offices of family doctors who may not practice obstetrics but desire to know more so that they can better understand what is happening to their pregnant patients, and as a useful resource for family medicine residents in training.

Neil Skolnik, MD  
Family Medicine Residency Program  
Abington Jefferson Health  
Abington, PA, USA  
Family and Community Medicine  
Temple University School of Medicine  
Philadelphia, PA, USA



# Preface

When I began the first edition of this textbook, I was certain of the topic's importance but less certain of the response to such a text. With the introduction of the second edition, I am grateful for the response to the previous edition as a reflection of the broad interest in providing quality care during this critical period in women's lives.

Much has changed in medicine since that time and obstetrical care has advanced as well. New diagnostic options have been introduced, new medications have found their way into routine care, and HIV has evolved from an acute and often fatal disease to a chronic disease that can be well managed with careful collaborative partnerships. While no text can fully capture all the intricacies of a field, I have tried to reflect these advances in this text.

While not all practitioners will include obstetrical care in their practice, it remains my conviction that all of us who care for women may find it helpful to have a reference that addresses key clinical issues in this important element of women's health.

I would like to once again recognize that this work would not be possible without the guidance and assistance of many others. The insight, critique, feedback, and suggestions have been invaluable in the final form. While there are too many to name individually, I would like to acknowledge Dimitrios Mastrogianis, MD, whose wisdom, humor, insight, and collaboration have been essential. I would like to recognize once again the ongoing support of my wife, Cynthia, and the forbearance of my children, Devin and Dylan. They are not only why I would do this work they are responsible, in large part, for how I am able to do it.

Riverside, CA

Paul Lyons, MD



# Contents

## Part I Preconception and Prenatal Care

<b>1 Physiology</b> .....	3
Background .....	3
Physiology of Menstruation .....	4
Physiology of Fertility .....	5
Hypothalamic Function .....	5
Pituitary Function .....	6
Ovulation .....	6
Physiology of Pregnancy .....	7
Cardiovascular Changes .....	7
Renal/Urinary Changes .....	8
Gastrointestinal Changes .....	9
Hematological Changes .....	9
<b>2 Preconception Counseling</b> .....	11
Background .....	11
The Patient .....	12
The Environment .....	13
Getting Started .....	13
Screening .....	14
Education .....	14
Intervention .....	16
<b>3 Prenatal Care</b> .....	17
Background .....	17
Key Domains of Prenatal Information .....	18
Frequency of Prenatal Visits .....	19
The First Prenatal Visit (See Table 3.1) .....	20
Follow-Up Prenatal Visits .....	25

<b>4 Medications in Pregnancy .....</b>	27
Background .....	28
General Principles of Medication Use in Pregnancy .....	28
Chronic Medical Conditions .....	28
Acute Medical Conditions .....	29
Acute Obstetrical Conditions .....	29
Therapeutic Categories and Considerations .....	30
Proven Human Teratogens .....	30
Special Considerations .....	32
Tobacco .....	32
Alcohol .....	32
Illicit Drugs .....	33
Over-the-Counter Medications .....	33
OTC Pain Medications .....	33
OTC Cough/Cold/Allergy Medications .....	34
<b>Part II Complications of Pregnancy</b>	
<b>5 Dysmorphic Growth and Genetic Abnormalities .....</b>	39
Background .....	39
History .....	40
Physical Examination .....	41
Diagnosis .....	41
Obstetrical Screening Tests .....	41
What Is Measured .....	41
What Is Detected .....	42
Confirmation/Follow-Up .....	42
Amniocentesis .....	42
Chorionic Villus Sampling .....	43
<b>6 Intrauterine Growth Restriction .....</b>	45
Background .....	46
Risk Factors .....	47
Fetal-Genetic Factors .....	47
Uterine-Environmental Factors .....	47
Maternal Factors .....	47
Toxic Exposures .....	48
Constitutional Factors .....	48
Complications of Growth Restriction .....	49
Antenatal Tracking and Diagnosis .....	49
Screening for Risks .....	49
Diagnosis .....	51
Management .....	52
Risk Reduction .....	52

<b>7 Preterm Labor</b> .....	55
Background.....	56
Factors Associated with Preterm Labor.....	56
Preconception Factor .....	57
Postconception Factors .....	57
Diagnosis.....	58
Intake Assessment.....	58
History.....	58
Physical Examination.....	58
Laboratory Studies.....	59
Management.....	59
Management Prior to 34 Weeks' Gestation .....	60
Management at 34–37 Weeks .....	62
Assessment of Fetal Lung Maturity.....	62
Lecithin-to-Sphingomyelin Ratio .....	62
Phosphatidylglycerol.....	63
Additional Tests .....	63
<b>8 Premature Rupture of Membranes</b> .....	65
Background.....	65
Diagnosis.....	66
History.....	66
Physical Examination.....	67
Laboratory .....	68
Management.....	69
Management at Term .....	70
Preterm Management .....	70
<b>9 Early Pregnancy Bleeding</b> .....	71
Background.....	72
General Approach to Early Pregnancy Bleeding per Vagina.....	72
History.....	74
Physical Examination.....	74
Ultrasound.....	75
Laboratory Studies .....	75
Ectopic Pregnancy.....	75
Background .....	75
Diagnosis.....	76
Management.....	77
Spontaneous Abortion.....	78
Background .....	78
Diagnosis.....	79
Management.....	80

<b>10 Late-Pregnancy Bleeding .....</b>	81
Background .....	82
General Approach to Late-Pregnancy Bleeding Per Vagina .....	82
History .....	83
Ultrasound .....	84
Physical Examination .....	84
Laboratory Studies .....	85
Placenta Previa .....	85
Background .....	85
History .....	85
Ultrasound Examination .....	86
Physical Examination .....	86
Laboratory Studies .....	86
Management .....	86
Abruptio Placenta .....	87
Background .....	87
History .....	88
Ultrasound .....	88
Physical Examination .....	88
Laboratory Studies .....	88
Management .....	88
<b>11 Recurrent Pregnancy Loss .....</b>	91
Background .....	91
Diagnosis .....	93
Management .....	93
History .....	93
Physical Examination .....	94
Diagnostic Studies .....	94
<b>12 Rh Isoimmunization .....</b>	95
Background .....	95
Fetal Consequences of Isoimmunization .....	96
Newborn Consequences of Isoimmunization .....	96
Diagnosis .....	97
History .....	97
Physical Examination .....	97
Diagnostic Studies .....	97
Management Prior to Isoimmunization .....	97
Management of Pregnancies with Rh-Sensitized Mothers .....	99
Amniotic Fluid Assessment .....	99
Ultrasoundography .....	100
Percutaneous Umbilical Blood Sampling .....	100

<b>13 Infection in Pregnancy .....</b>	101
Background .....	102
Symptoms of Infection in Pregnancy .....	102
Maternal Infection .....	104
Fetal Infection .....	104
Antibiotic Use in Pregnancy .....	105
Sulfonamides .....	105
Fluoroquinolones .....	105
Aminoglycosides .....	106
Vaginitis/Vaginosis .....	106
Background .....	106
History .....	106
Physical Examination .....	107
Laboratory Studies .....	107
pH Testing .....	107
KOH “Whiff” .....	107
Microscopic Examination .....	107
Treatment .....	108
Bacterial Vaginosis .....	108
Trichomoniasis .....	108
Yeast Vaginitis .....	108
Gonorrhea/Chlamydia .....	109
Urinary Tract Infections .....	109
Background .....	109
Diagnosis .....	110
History .....	110
Physical Examination .....	110
Laboratory Examination .....	110
Treatment .....	111
Group B Strep .....	111
Background .....	111
Diagnosis .....	112
Universal Screening .....	112
Treatment .....	112
<b>14 Hypertension in Pregnancy .....</b>	113
Background .....	114
Diagnosis .....	114
Diagnostic Criteria .....	114
History .....	115
Physical Examination .....	116
Laboratory Studies .....	116
Management .....	116
Chronic Hypertension .....	116

Pre-Eclampsia .....	117
Prevention .....	117
Management.....	117
Seizure Prevention .....	119
<b>15 Diabetes in Pregnancy .....</b>	<b>121</b>
Background.....	121
Diagnosis.....	122
Pregestational Diabetes .....	123
Management.....	123
Gestational Diabetes .....	126
Preconception Management.....	126
<b>16 Multigestational Pregnancy .....</b>	<b>127</b>
Background.....	127
Diagnosis.....	129
History.....	129
Physical Examination.....	130
Laboratory and Diagnostic Studies.....	130
Management.....	130
Prenatal Care .....	130
Labor and Delivery .....	131
<b>17 Postdates Pregnancy .....</b>	<b>133</b>
Background.....	133
Diagnosis.....	134
History.....	134
Physical Examination.....	135
Diagnostic Studies .....	135
Management.....	135
Assessment of Fetal Well-Being .....	135
<b>Part III Labor and Delivery</b>	
<b>18 Normal Labor .....</b>	<b>141</b>
Background .....	141
Prelabor .....	141
Labor .....	142
Assessment.....	143
History.....	143
Physical Examination.....	143
Laboratory Studies .....	144
Management.....	145
<b>19 Induction and Augmentation .....</b>	<b>147</b>
Background .....	147
Preparation .....	148
Pharmacological Options .....	148

Mechanical Options .....	148
Induction .....	149
<b>20 Pain Management in Labor .....</b>	<b>151</b>
Background .....	151
Nonpharmacological Management .....	152
Pharmacological Management .....	152
Narcotic Analgesia .....	152
Local Anesthesia .....	153
Epidural–Spinal Anesthesia .....	153
<b>21 Operative Delivery .....</b>	<b>155</b>
Background .....	155
Forceps Delivery .....	156
Use of Forceps .....	157
Vacuum-Assist Delivery .....	157
Cesarean Section .....	158
<b>Part IV Complications of Labor and Delivery</b>	
<b>22 Prolonged Labor .....</b>	<b>161</b>
Background .....	161
Complications of Labor .....	162
Prolonged Latent-Phase Labor .....	162
Failure to Dilate/Efface .....	163
Failure to Descend .....	165
<b>23 Shoulder Dystocia .....</b>	<b>167</b>
Background .....	167
Diagnosis .....	168
Management .....	169
<b>24 Malpresentation .....</b>	<b>171</b>
Background .....	171
Occiput Positions .....	172
Diagnosis .....	172
Management .....	172
Nonoccipital Presentations .....	172
Breech Presentation .....	173
Compound Presentation .....	174
<b>25 Fetal Heart Rate Monitoring .....</b>	<b>175</b>
Background .....	175
Normal Fetal Heart Tracings .....	176
Evaluation of Fetal Heart Rate Baseline .....	176
Tachycardia .....	177
Bradycardia .....	177

Evaluation of Fetal Heart Rate Variability .....	177
Acceleration .....	177
Early Deceleration .....	178
Variable Deceleration.....	178
Late Decelerations .....	178
<b>26 Maternal Fever in Labor .....</b>	<b>179</b>
Background .....	179
Diagnosis.....	180
History.....	180
Physical Examination.....	180
Diagnostic Studies .....	180
Management.....	181
<b>27 Postpartum Hemorrhage.....</b>	<b>183</b>
Background .....	184
Complications Causing Hemorrhage .....	184
Uterine Atony.....	184
Lacerations .....	184
Retained Placenta.....	185
Coagulopathy .....	185
Uterine Inversion.....	185
Diagnosis.....	185
Management.....	185
Laceration .....	186
Persistent Bleeding.....	186
<b>28 Perineal Laceration and Episiotomy .....</b>	<b>189</b>
Episiotomy .....	189
Background .....	189
Perineal Laceration .....	190
Background .....	190
<b>Part V Postpartum Management</b>	
<b>29 Newborn Evaluation .....</b>	<b>195</b>
Background .....	195
The Examination.....	196
History.....	196
Physical Examination.....	196
Laboratory and Diagnostic Studies .....	198
Vaccination.....	198
<b>30 Routine Hospital Postpartum Management .....</b>	<b>199</b>
Background .....	199
Postpartum Day 1.....	200
History.....	200
Physical Examination.....	200

Laboratory Studies .....	201
Management.....	201
Postpartum Day 2.....	202
History.....	202
Physical Examination.....	202
Laboratory Studies .....	203
Management.....	203
<b>31 Complications of the Hospital Postpartum Period .....</b>	<b>205</b>
Background.....	205
Fever.....	206
Infection .....	206
Endometritis.....	206
Urinary Tract Infections.....	207
<b>Index.....</b>	<b>209</b>

# **Part I**

## **Preconception and Prenatal Care**

# Chapter 1

## Physiology

### Contents

Background .....	3
Physiology of Menstruation .....	4
Physiology of Fertility .....	5
Hypothalamic Function .....	5
Pituitary Function .....	6
Ovulation .....	6
Physiology of Pregnancy .....	7
Cardiovascular Changes .....	7
Renal/Urinary Changes .....	8
Gastrointestinal Changes .....	9
Hematological Changes .....	9

### Key Points

1. The menstrual cycle can be considered a comprehensive physiological adaptation for potential pregnancy.
2. Normal menstrual cycles last 21–45 days (average 28 days), counted from the first day of menstrual bleeding.
3. Physiological adaptations of pregnancy affect most major organ systems including cardiac, renal, gastrointestinal, and endocrine systems.

### Background

Although most patients will not present to their providers with questions concerning the specifics of reproductive physiology, the care and management of pregnant patients begins with an understanding of the physiological environment in which

pregnancy occurs (or in some instances, does not occur). Many women's health providers will face questions concerning menstrual function prior to caring for a patient's obstetrical needs. Conversely, routine gynecological care may provide an opportunity to begin discussions of pregnancy planning and preconception counseling. For many women, a "routine" gynecological examination is the primary point of contact with the health-care system early in life. For this reason, all providers who care for women should have some understanding of normal reproductive physiological function. A brief overview of menstruation, fertility, and pregnancy follows.

## Physiology of Menstruation

Menstruation represents the cyclical physiological preparation for potential pregnancy, followed by removal of endometrial contents if pregnancy does not occur. Most women of reproductive age are familiar with menstruation. The average age of menarche in the United States is approximately 11.5 years. Most menstrual cycles are anovulatory in the first year following menarche and may remain irregularly ovulatory for up to 3 years (although women and providers should be aware that ovulation and/or pregnancy may occur). For the next three to four decades, most women will menstruate every 21–35 days (average  $28 \pm 7$  days). Bleeding is variable but generally lasts 3–5 days (1–7 days may be considered normal) and is of variable intensity (but generally less than 3 oz or  $90 \text{ cm}^3$ ).

Although generally considered an ovarian and uterine phenomenon, the normal menstrual cycle may be considered as a comprehensive physiological adaptation in preparation for possible pregnancy. In addition to the uterine and ovarian changes described here, changes can be noted in the cervix, vagina, breast, and core body temperature. The cervical mucus becomes thinner with increased pH to facilitate entry of sperm. Vaginal epithelial cells also undergo change. Mammary ducts proliferate under estrogen and progesterone stimulation, which may lead to breast swelling and tenderness. A small spike in basal body temperature can be seen at the time of ovulation. This observation has contributed to the use of basal body monitoring in fertility management.

Physiologically, bleeding represents the end of one cycle. From the perspective of the patient and the provider, however, bleeding is the most easily identified aspect of the menstrual cycle and is, therefore, used to mark the beginning of each cycle. The first day of menstrual bleeding is day 1 with each day numbered sequentially through the last day prior to the reoccurrence of bleeding. Each menstrual cycle can be divided into two halves that differ in hormonal and physiological events. In a typical or average menstrual cycle, each half is approximately 14 days in duration.

The first half of each menstrual cycle is marked by endometrial proliferation and follicular development. In the first week of each menstrual cycle, multiple follicles enlarge. At approximately 1 week, a single follicle becomes dominant and the others involute, becoming atretic. The dominant follicle will, with appropriate hormonal regulation, continue to develop and will eventually rupture releasing an ovum

for possible fertilization. With release of the ovum on day 14, the follicle undergoes a series of stereotypic changes filling with blood, granulose, and thecal cell proliferation and displacement of blood by luteal cells (corpus luteum). The luteal cells produce progesterone, which serves to stabilize the thickened endometrium through the second half of the menstrual cycle. The period of follicle development is referred to as the follicular phase. The period of luteal production of progesterone is referred to as the luteal phase.

Follicular development in the first half of each menstrual cycle is marked by follicular production of estrogen and endometrial proliferation in anticipation of possible implantation of a fertilized ovum. This generally occurs late in the first week and throughout the second week of the menstrual cycle. The first half of the menstrual cycle is, for this reason, sometimes referred to as the proliferative phase. With ovulation and luteal production of estrogen and progesterone, uterine glands become active, secreting clear fluid. This phase is referred to as the secretory phase. The endometrium will remain stable and secretory for as long as the progesterone stimulation continues.

If fertilization fails to occur, the corpus luteum will lose function beginning in the second half of the fourth week (corpus albicans). With the loss of hormonal support, endometrial thinning and localized necrosis lead to sloughing of the proliferative portion of the endometrial lining and the onset of menses. Until menopause, this cycle will repeat more or less regularly each month.

## **Physiology of Fertility**

The hormonal changes just described relate to preparation for release of the ovum and subsequent fertilization by sperm. As noted, however, these menstrual changes may occur in the absence of ovulation. In addition, under normal physiological conditions, pregnancy requires the presence of functional sperm in sufficient quantity to ensure fertilization of the released ovum.

In women, the release of an ovum is under the control of the hypothalamic–pituitary–ovarian endocrinological axis. Each of these components must function normally to ensure ovum release. Two pituitary hormones, in particular, are critical to normal ovulatory cycles—follicle-stimulating hormone (FSH) and luteinizing hormone (LH).

### ***Hypothalamic Function***

Release of pituitary hormones depends on hypothalamic stimulation. The hypothalamus is responsible for stimulation of a variety of pituitary hormones and hypothalamic dysfunction may manifest with altered fertility or a variety of other endocrinological signs or symptoms. In addition to pituitary stimulation, the hypothalamus is responsible for direct release of oxytocin (of import at the time of labor).

In relation to fertility, hypothalamic release of gonadotropin-releasing hormone (GnRH) stimulates the anterior pituitary production of FSH and LH. GnRH is produced in the hypothalamus and released directly to the pituitary via local blood vessels. Release of GnRH is episodic in brief, timed bursts. Although GnRH cannot be measured directly, pulsatile GnRH release results in pulsatile release of LH which can be measured providing indirect evidence of hypothalamic function. Failure to maintain this episodic release will inhibit pituitary stimulation, probably secondary to downregulation of pituitary receptors. Disruption of the timing of the episodic release will also impair fertility by disrupting the appropriate timing of FSH and LH stimulation of the ovary. In addition, appropriately episodic and timed GnRH stimulates pituitary GnRH receptors enhancing sensitivity at mid-cycle and facilitating a surge in LH at the time of ovulation.

## ***Pituitary Function***

As with the hypothalamus, the pituitary is responsible for the release of several hormones regulating a variety of physiological functions. In relation to fertility, the two key hormones are the gonadotropins, FSH and LH. These two agents are released cyclically and in a pulsatile fashion in response to GnRH stimulation. Together they are responsible for regulation ovarian hormonal secretion. Pituitary release of FSH and LH are also regulated by ovarian hormone release. Ovarian release of estradiol results in negative feedback (inhibition) of FSH release and positive feedback (stimulation) of LH release.

FSH, as the name implies, is responsible for stimulating early follicle development within the ovary. LH fosters ovarian production of estrogen and progesterone from the corpus luteum. In conjunction with LH, FSH is also responsible for terminal maturation. At the point of maturation, a surge in LH levels precipitates follicular rupture and ovum release.

## ***Ovulation***

Early in the menstrual cycle, FSH levels are slightly elevated (stimulating follicular development) and LH levels are low. In this phase of the menstrual cycle, estrogen serves an inhibitory role on LH. GnRH stimulation of the pituitary continues and the sensitivity of the pituitary is enhanced. Approximately 2 days prior to ovulation, the estrogen inhibition is reversed, becoming stimulatory and a positive feedback loop is established. Approximately 8–10 h prior to ovulation, LH levels reach a peak (LH surge). Ovulation then occurs. Following ovulation, estrogen once again becomes inhibitory and, in conjunction with elevated progesterone levels, serves to inhibit LH and FSH levels in the second half of the menstrual cycle.

## Physiology of Pregnancy

The physiological changes associated with pregnancy are numerous and the full scope of such changes is beyond the scope of this text. Common physiological changes with pregnancy are summarized in Table 1.1. Recognition of normal physiological changes is necessary not only to understand normal function while pregnant but also to facilitate recognition of physiological abnormalities that lie outside the normal range.

### ***Cardiovascular Changes***

Pregnancy can be considered an adaptive high-volume, hyperdynamic cardiovascular state. Increased volume, a newly developed peripheral vascular bed, and anatomic changes associated with an enlarging uterus all serve to alter normal cardiovascular status. The heart, itself, enlarges and cardiac output increases by nearly 50 %.

**Table 1.1** Physiological changes of pregnancy

Cardiovascular
Cardiac enlargement
Increased cardiac output
Systolic flow murmur
Decreased venous return
Decreased peripheral vascular resistance
Decreased blood pressure
Increased blood flow to the uterus, kidneys, skin, breasts
Renal/urinary
Increased urinary stasis
Increased urinary system volume
Kidney enlargement
Renal pelvis dilation
Ureteral elongation
Increased bladder capacity
Increased glomerular filtration rate
Elevation of rennin, aldosterone, angiotensin
Glucosuria
Gastrointestinal
Early satiety
Nausea, vomiting
Constipation
Gingival hypertrophy
Progression of periodontal disease
Decreased gastric emptying
Relaxation of lower esophageal sphincter
Hematological
Increased red blood cell volume
Anemia
Leukocytosis

This increased output is initially facilitated by an increase in cardiac volume and subsequently by an increase in heart rate. The increase in output reaches a peak near the end of the second trimester and then remains stable until the end of pregnancy.

The increase in volume may lead to increased flow turbulence within the heart. This turbulence may be apparent clinically as a systolic ejection murmur. Such a murmur will manifest in 80–90 % of all pregnant women. This murmur is a normal physiological finding and does not warrant further cardiovascular investigation.

Vascular changes are also common in pregnancy. With an increase in uterine size, venous return via the inferior vena cava may be directly impaired. Placing the patient in the left lateral recumbent position may alleviate the direct pressure of the uterus on the vena cava and facilitate enhanced venous return. The direct compression of venous return from the lower extremities may lead to peripheral edema. Peripheral vascular resistance declines with pregnancy as maternal cardiac output increases. Compensatory venous response to rapid position changes may also be impaired in pregnancy causing light-headedness or dizziness with rapid positional changes. Blood pressure often declines slightly (approximately 10 mmHg diastolic) with a nadir in the second trimester and a slight rise (to near prepregnant levels) near the end of pregnancy.

Blood flow is altered in pregnancy as well. The most obvious change is the increase in uterine blood flow with the development of the uteroplacental vascular bed. Blood flow through this vascular bed is facilitated by vascular resistance that is low relative to the overall peripheral vascular resistance. In addition to increased blood flow to the uterus, maternal blood flow is increased to the kidneys, breast, and extremities (including increased flow to the skin). Although concern has been raised that exercise may divert blood flow from these key areas to muscles, this has generally not been found to be clinically significant except for women who significantly increase their activity level from their prepregnancy baseline. A reasonable recommendation would be that women may continue exercise through pregnancy at a level not to exceed their usual degree of exertion.

## ***Renal/Urinary Changes***

Pregnancy is marked by an increase in urinary stasis. The direct impingement of the uterus and fetus on the bladder contributes to this effect as do anatomic changes within the urinary tract. Kidneys enlarge, the renal pelvis dilates, and the course of the ureter elongates contributing to increased volume within the urinary tract. This increased volume in turn contributes to an increase in post-void residual urine within the tract and subsequent stasis.

Renal function is also changed in pregnancy. A combination of hormonal modulation with increased plasma volume leads to an increase of nearly 50 % in the glomerular filtration rate. Renin, angiotensin, and aldosterone levels are all elevated

in pregnancy. Despite this increase in glomerular filtration, urinary output is not increased during pregnancy. Although many patients will report increased urinary frequency, the total output volume remains similar to the prepregnancy levels (the functional capacity of the bladder is, in fact, increased in pregnancy with a total capacity of approximately 1.5 L). Aldosterone-mediated sodium resorption in turn leads to fluid resorption and maintenance of intravascular homeostasis. Increased renal filtration does lead to increased creatinine clearance and a concomitant reduction in serum creatinine levels (along with decreased blood urea nitrogen levels). Sporadic glucosuria is a common finding in pregnancy and may be an artifact of increased glomerular filtration.

### ***Gastrointestinal Changes***

Gastrointestinal (GI) symptoms are among the most common complaints of pregnancy. Early in pregnancy nausea and vomiting are often reported. Later in pregnancy, early satiety and constipation are both commonly observed. Although these symptoms are generally not related to specific changes within the GI tract, some physiological alterations are noted with pregnancy. Gingival hypertrophy and worsening of gingival disease have both been reported. Some investigators have suggested a possible link between periodontal disease and an increased risk of preterm labor although the results are preliminary and inconclusive. GI motility is decreased, including a decrease in gastric emptying and increased transit time through the large intestine. Decreased gastric emptying combined with relaxation of the terminal portion of the esophagus may lead to an increase in reports of gastroesophageal reflux symptoms. This may be exacerbated late in pregnancy as the uterus increasingly displaces the stomach upward.

### ***Hematological Changes***

Pregnancy is associated with a variety of changes in hematological status. With an increase in intravascular volume, patients also experience an increase in red cell volume. This increase in red cell volume, in turn, increases the patient's need for iron. With inadequate dietary iron (either from food or supplementation), many pregnant patients will develop an iron deficiency anemia with the usual change in red cell indices (decreased mean corpuscular volume and decreased mean corpuscular hemoglobin content). An increase in blood leukocytes is common in pregnancy. Levels rise throughout pregnancy and peak during labor. Such a rise in white blood cells may make determination of infection more complicated.

# Chapter 2

## Preconception Counseling

### Contents


Background .....	11
The Patient.....	12
The Environment.....	13
Getting Started .....	13
Screening.....	14
Education .....	14
Intervention .....	16

### Key Points

1. Preconception counseling is medical evaluation and intervention performed prior to conception with the expectation that the course and outcome of subsequent pregnancies will be improved.
2. Preconception counseling and intervention may occur in the context of care for other medical conditions.
3. Preconception counseling consists of three primary activities: (a) risk identification/assessment, (b) patient education, and (c) risk intervention, when possible.
4. The postpartum period is often an ideal opportunity for preconception counseling for subsequent pregnancies.

### Background

Multiple independent factors impact the course and outcome of pregnancy. Figure 2.1 represents a schematic diagram of many of these factors and their interrelationship during the course of pregnancy. Prior to conception, a number of factors combine to provide the background environment in which subsequent pregnancies will develop.



**Fig. 2.1** Pregnancy outline

In particular, a complex interaction between the patient-related factors and environmentally related factors contribute to the likelihood of pregnancy, the prenatal course, pregnancy outcomes, and postpartum complications.

## ***The Patient***

Any woman who becomes pregnant brings with her a variety of genetic and acquired factors that have the potential to impact the course of pregnancy. Genetic predisposition plays a significant role in obvious ways (e.g., sickle cell trait, Tay–Sachs, cystic fibrosis) or in more subtle manners (e.g., polymorphic tumor necrosis factor- $\alpha$  which may contribute to a predisposition to preterm labor). Anatomic factors may also contribute (e.g., congenital cervical incompetence, anomalous uterus). Physiological considerations such as the function of the hypothalamic–pituitary–ovarian axis (e.g., oligomenorrhea, anovulatory cycles) also play a role in becoming pregnant and in maintaining pregnancy.

Pregnancy is one potential medical event in the life of a woman, but it is by no means the only one. In addition to the genetic, anatomic, and physiological factors, many women will also acquire medical conditions that may impact the course or outcome of pregnancy. Such conditions as diabetes mellitus, hypertension, and cardiac, renal, or thyroid disease all impact the course of subsequent pregnancies. Not only do underlying disease conditions have the potential to directly impact obstetrical outcomes, many of the medications and treatments for these conditions may also

have obstetrical implications. Surgical interventions involving cervical, pelvic, or intra-abdominal manipulation may also have consequences in pregnancy.

A variety of behavioral issues have direct bearing on pregnancy. Nutritional status is critical and has been the focus of recent recommended interventions such as folate supplementation prior to pregnancy. Much work has focused on the detrimental effects that tobacco, alcohol, and other drug use may have during pregnancy. Intrauterine growth rates, congenital anomalies, infant addiction, and preterm complications—among others—have all been shown to be impacted by the use of these substances. Patient exercise, fitness, and activity levels may also contribute to pregnancy outcomes.

## ***The Environment***

Although potentially less obvious, the environment in which the woman lives may contribute significantly to obstetrical outcomes. For example, lower socioeconomic status is associated with higher complication rates and/or less good outcomes in a variety of obstetrical conditions such as preterm labor, pregnancy-induced hypertension, and others. Family and partner support for the pregnancy may impact the degree to which women access care during pregnancy.

All of these factors are present prior to conception providing the biopsychosocial environment in which the pregnancy will occur. Because these factors are all present—potentially identifiable and potentially modifiable—prior to conception, it is helpful to consider pregnancy and prenatal care as beginning prior to conception as well.

Preconception counseling is medical evaluation and intervention performed prior to conception with the expectation that the course and outcome of subsequent pregnancies will be improved. Preconception counseling offers providers an opportunity to assess, document, and potentially alter many of the factors that influence pregnancy outcome. Many patients will present to their provider only after discovering they are pregnant. Under such circumstances, the opportunity to impact this potentially critical period is lost. For this reason, preconception counseling is of paramount importance for all providers who care for women of childbearing age. Prior to pregnancy, some women will only seek care for other medical problems. Preconception counseling and intervention, therefore, may need to occur only in the context of care for other medical conditions.

## **Getting Started**

Some women may raise the issue of planning for pregnancy providing the opportunity for the provider to begin the preconception assessment. For these women, providers can begin the process of risk identification/assessment, patient education, and

risk intervention/reduction (when possible). Other women, however, may benefit from prompting by their provider. There are a number of ways in which to approach the issue and each provider will determine for him or herself the approach that works best. Because preconception counseling will often be initiated during visits for nonobstetrical care, providers should be prepared to raise the issue in these contexts. A review of menstrual history as part of a routine exam might be followed by open-ended questioning such as “Tell me about any plans you might have to become pregnant.” Visits for chronic medical conditions might lead providers to raise the issue of “the impact of this condition should you choose to become pregnant.” Routine gynecological visits provide another opportunity for beginning pre pregnancy planning. Visits related to unprotected sex (sexually transmitted disease screening, late menses, pregnancy testing, etc.) provide an excellent and natural opportunity to discuss issues of importance prior to pregnancy. An often underutilized opportunity for preconception counseling is during routine postpartum care. Planning for subsequent pregnancies (or their prevention) can begin while the patient is still in the hospital and continue when she returns for routine outpatient postpartum care.

## Screening

As noted in Table 2.1, preconception counseling should include screening for issues related to the patient and to her environment (including occupational, financial, and family-related issues, among others). A variety of standardized prenatal care flow sheets exist that capture much of this data. Less formal screening may be appropriate under many circumstances. For patients who are seen regularly, screening may occur sequentially over time and should be updated periodically to ensure accuracy.

## Education

An appropriate preconception history will allow providers to develop a list of important pregnancy-related concerns prior to conception. This list can form the focus of ongoing education designed to allow patients to make the best possible decisions concerning their health. Modifiable risk factors can be identified and addressed as noted here. For other issues, intervention or modification may be less important than education and discussion of the identified risk factors. Providers of preconception counseling can facilitate their patients’ decisions concerning the desirability and timing of and preparation for pregnancy.

Genetics counseling may be beneficial for those patients at high risk for genetic complications during pregnancy. For patients with chronic medical conditions, the risks of pregnancy, optimal timing for pregnancy, contraception (if pregnancy is medically contraindicated), and prenatal management in the case of pregnancy should all be discussed.

**Table 2.1** Content of preconception counseling

<i>Patient-related factors</i>
Psychosocial issues
Tobacco, alcohol, illicit drug use
Psychiatric illness
Literacy/language barriers
Medications
Prescription
Over-the-counter
Herbal, natural, and complementary/alternative therapeutics
Medical
Hypertension
Diabetes
Thyroid disease
Systemic lupus erythematosus
Renal disease
Cardiac disease
Thromboembolic disease
Sickle cell disease (or trait)
Hepatitis
HIV/AIDS
Measles (including immune status)
Varicella (including immune status)
Intra-abdominal or pelvic surgery
Obstetrics/gynecology
Pelvic anomalies
Pelvic inflammatory disease
Prior obstetrical history (all outcomes including full-term, preterm, spontaneous and elective abortions, living children)
Macrosomic infants
Fetal/neonatal death
Genetic (patient, patient's family and partner)
Down syndrome
Neural tube defects
Cystic fibrosis
Congenital anomalies
Multiple gestation
<i>Environmental factors</i>
Psychosocial issues
Physical/sexual abuse
Partner/family support
Child care
Transportation
Financial support
Insurance status
Occupational issues
High-risk occupations
Occupational exposures

## Intervention

A number of important interventions can be offered for women who are not yet pregnant. Patients should be counseled concerning the role of prepregnancy well-being including nutrition, exercise, and fitness. Patients should be counseled concerning the benefits of smoking cessation both for overall health and specifically related to pregnancy. Patients should be made aware of the effects of alcohol and the need to eliminate alcohol consumption prior to becoming pregnant. Use of illicit drugs should also be discouraged.

For those patients who do not plan to become pregnant for at least 1 month, review of the vaccination history may reveal the need for administration of measles mumps rubella (MMR), varicella, and Hep B vaccines. For patients planning to become pregnant in the near future, folic acid supplementation is recommended. For patients with known risk factors (see Table 2.1) for neural tube defects, 4 mg per day is recommended. Because fewer than half of all cases of neural tube defects occur in patients with known risk factors starting prenatal, vitamins and/or folic acid supplementation (0.4 mg per day) should be recommended for all patients prior to conception.

Patients' medical conditions and medications should be reviewed for potential complications in pregnancy. For patients who take medications contraindicated or relatively contraindicated in pregnancy, providers should discuss the risks and benefits of such medications in pregnancy. Consultation with a maternal–fetal medicine specialist may be beneficial in these circumstances.

Other issues such as language, financial, or family support barriers can be discussed and problem-solving can begin prior to rather than after conception. Patients for whom insurance or financial issues may be an issue should be made aware of insurance programs—available in most states—to provide insurance for pregnant patients.

# Chapter 3

## Prenatal Care

### Contents

Background .....	17
Key Domains of Prenatal Information .....	18
Frequency of Prenatal Visits .....	19
The First Prenatal Visit (See Table 3.1) .....	20
Follow-Up Prenatal Visits .....	25

### Key Points

1. Prenatal care is a process not an event.
2. Excellent prenatal care represents a partnership between the provider, the patient, and her family.
3. Key domains of intake information in prenatal care include pregnancy dating, baseline maternal health status, family health history, medical conditions impacted by pregnancy, medical conditions impacting pregnancy, and infection.
4. Key domains of information during follow-up visits include normal growth and development, medical and/or obstetrical complications of pregnancy, and onset of labor.

### Background

Prenatal care is generally the most prolonged and sustained component of pregnancy care. Such care is often delivered by a single provider who will follow the course of the pregnancy with nearly as much interest as the pregnant patient. Although the prenatal period is often filled with anxiety, it is generally more relaxed and almost always less pressured than the labor and delivery setting. For these

reasons, prenatal care provides patients with an opportunity to educate themselves and participate in the process of preparing for a new infant.

Prenatal care can best be thought of as a process rather than a specific event. For patients who received preconception counseling, it is a continuation of the threefold process of education, risk identification, and risk reduction/intervention. For those patients who did not receive preconception counseling, it should be the beginning of such a process. Many prenatal care providers also participate in the delivery of their patients. In this regard, prenatal care provides an opportunity to establish or enhance a relationship that stretches from preconception through prenatal management to delivery. Excellent prenatal care represents a partnership between the provider and the pregnant patient (and her family) with ample opportunity for discussion, questions, and answers and active participation by each person involved.

Prenatal care is a cornerstone of modern obstetric care and has accompanied a reduction in the historic risk associated with pregnancy. Maternal mortality in 1935 was 582 per 100,000 pregnancies. By 1993, that figure had decreased to 7.5 per 100,000. Over approximately the same period, infant mortality decreased from 47 per 1000 to 8 per 100,000. Although many factors unrelated to prenatal care have contributed to these reductions, prenatal care has directly contributed to improved outcomes in a variety of areas: (a) fetal organogenesis (folic acid supplementation, glycemic control in diabetes), (b) infectious disease detection/treatment (e.g., chlamydia, bacteriuria), (c) infectious disease transmission prevention (e.g., syphilis, HIV), and (d) fetal growth (e.g., glycemic control in gestational diabetes).

Figure 2.1 in Chap. 2 provides a schematic outline for the interrelated factors involved in pregnancy. Factors present during the preconception period were reviewed in the prior chapter. With the advent of conception, additional factors come into play. Pregnancy itself introduces a variety of new conditions that can impact the health of both the mother and the infant. Although less well described, a variety of paternal factors can also affect the course and outcome of pregnancy. Although not specifically addressed in this chapter, there are a number of physician- and system-related factors that can, likewise, impact pregnancy. The process of prenatal care is cyclical and repetitive. Each visit allows providers to review what has already happened and to determine what has developed in the interval since the last visit. Each visit will focus on patient education, determination of normal growth and development, and compilation of critical information in a variety of important domains.

## Key Domains of Prenatal Information

The first step in any prenatal assessment is to confirm that the patient is, in fact, pregnant. Many patients will arrive for a first prenatal visit with a confirmed positive pregnancy test result. Other patients will arrive for their first “prenatal” visit with a variety of less clear presentations. They may have missed their period but not have

been tested. They may have self-tested but not trust the results (either positive or negative). Regardless of the nature of the uncertainty, all patients with uncertain pregnancy status should have their status confirmed via urine or serum human chorionic gonadotropin (hCG) testing. Following confirmation of pregnancy, all patients should have the opportunity to consider what the most appropriate next step should be. When uncertain regarding the desirability of a confirmed pregnancy, providers should be cautious in their choice of language. In particular, providers should avoid the use of language that implies a specific outcome such as “Congratulations, you’re pregnant!” or “We will need to get you started with prenatal care soon.” The use of open-ended questions such as “The test confirms that you are pregnant. What do you think you would like to do at this point?” will allow patients to more easily express their thoughts concerning subsequent management.

Initial evaluation during prenatal care is similar to the evaluation for preconception counseling. It is designed to focus on the patient’s baseline condition and any development in the time frame from the last menstrual period (LMP) through the first prenatal visit. (This is a time period during which patients may not recognize that they are pregnant and may include significant exposures to infectious or toxic agents.) At intake prenatal screening provides important information in six key domains:

1. Baseline maternal health including prior obstetrical history, if any
2. Family health (e.g., twins, Down syndrome, Tay–Sachs)
3. Medical conditions impacted by pregnancy (e.g., cardiac disease, renal disease)
4. Medical conditions impacting pregnancy (e.g., diabetes mellitus, hypertension, medications, toxic exposures)
5. Infection
6. Pregnancy dating

As discussed in the previous chapter, these domains of information can form the basis for patient education and intervention throughout the course of pregnancy (and beyond).

Following the initial prenatal screen, subsequent visits focus on interval developments and review of previously identified issues. It is often helpful to develop a prenatal problem list to organize ongoing issues under management. Such a list could include the problem, the date identified, interventions, if any, and the date resolved, if applicable. Follow-up prenatal care will focus on three key domains: (a) normal growth and development, (b) medical and/or obstetrical complications of pregnancy, and (c) onset of labor.

## Frequency of Prenatal Visits

The frequency of visits during prenatal care is dependent on the complexity of the care being provided. For those patients with complex presentations, individualized decisions must be made concerning the frequency of prenatal visits. For patients

with uncomplicated pregnancies, the currently recommended frequency for visits is as follows:

First prenatal visit: in the first trimester

Follow-up visits: every 4 weeks until 30 weeks gestation, then  
every 2 weeks until 36 weeks gestation, then  
every week until delivery at term, or  
twice weekly from 40 weeks for assessment of  
fetal well-being (see Chap. 18).

### ***The First Prenatal Visit (See Table 3.1)***

The first prenatal visit lays the groundwork for all subsequent visits and therefore includes the most comprehensive history, physical examination, and diagnostic testing. For patients who have had one or more preconception counseling visits, much

**Table 3.1** First prenatal visit

Last menstrual period (LMP)
Duration, abnormalities, fertility treatment, contraceptive use, and prior pregnancies
Prior obstetrical history: all prior pregnancies including elective and spontaneous abortions
All pregnancies: date, outcome, weeks of gestation
Deliveries: gender and weight of infant, duration of labor, delivery method, anesthesia, and complications, if any
General medical history: including all pertinent medical conditions but focused on
Anesthesia history
Medications
Medication allergies
Toxic exposures (including tobacco, alcohol, and illicit drugs)
History since LMP: events or exposures during early organogenesis
Drugs, medications, and radiation
Infectious diseases (cytomegalovirus, toxoplasmosis, tuberculosis)
Possible pregnancy-related symptoms (bleeding or discharge per vagina, abdominal pain, headache, visual complaints, emesis)
Family history: comprehensive but with an emphasis on
Fetal deaths or abnormalities
Multiple gestations
Genetic history (cystic fibrosis, Down syndrome, thalassemia)
Chronic medical conditions with strong familial link (e.g., hypertension, diabetes mellitus, renal disease, substance abuse)
Safety
Abuse, including sexual abuse
Domestic violence
Seat belt use
Environmental risks including work-related hazards

of the information can be obtained from the records of those visits. Information subject to change will need to be updated and a brief review of all material should be performed to ensure accuracy and completeness. Many offices utilize one of the many standardized obstetrical health history and prenatal flow sheet forms that exist in both paper and electronic form. In addition to providing a standardized method for recording necessary information, the flow sheets can serve as a prompt for providers throughout the course of pregnancy.

## History

*Menstrual History.* The preconception history begins with the LMP. Information should include the date of the first day of bleeding and the duration. The certainty of this date is of considerable importance as this forms the first basis for predicting the estimated date of delivery (EDD, or confinement in some older references). Depending on a variety of factors, including regularity of menses, duration since LMP, and whether patients track menses regularly, patient recall of their LMP may be of variable reliability. Some notation of the patient's subjective "certainty" concerning the date should be recorded. First-trimester bleeding is a relatively common complaint and may mimic menstruation. For this reason, patients should be questioned concerning the timing and quantity of bleeding as well. A "period" that is abnormally light or heavy or fell out of the normal cycle may represent an event of pregnancy rather than the true LMP.

*Prior Obstetrical History.* Prior obstetrical history can significantly impact the current pregnancy. For this reason all prior pregnancies should be summarized in the prenatal record. For all prior pregnancies, documentation should include the date of the pregnancy, the outcome (full-term delivery, preterm delivery, spontaneous abortion, elective abortion) and the weeks of gestation completed. For those pregnancies that resulted in live delivery, additional key information would include date of delivery, gender and weight of the infant, duration of labor, delivery method and anesthesia use, or complications, if any.

This information can be summarized in short-hand form as gravida/para figures. This notation takes the form of  $G_xP_{xxxx}$ .  $G$ (ravida) represents the total number of pregnancies regardless of outcome.  $P$ (ara) represents, in order from left to right, full-term deliveries, preterm deliveries, abortions (elective or spontaneous), and living children. Although it does not include all of the details of each pregnancy, this short-hand form provides a quick and convenient summary that is especially useful in oral or written presentations.

*General Medical History.* All pregnancies occur within the context of the patient's baseline health status. Many medical conditions may impact the course of pregnancy. At the same time, pregnancy may have a significant effect on a variety of pre-existing medical conditions. It should also be recognized that delivery is a surgical procedure in many instances with approximately one-fourth of all deliveries occurring via cesarean section. For this reason, a comprehensive review of the

patient's general medical history should be recorded. Special emphasis should be placed on chronic medical conditions, prior surgical and anesthesia history, medications, medication allergies, and substance use including tobacco, alcohol, and illicit drug use.

*History Since LMP.* Particular attention should be paid to medical events between the LMP and the first prenatal period. This represents the critical period of organogenesis during which a variety of medical conditions and exposures may have significant impact on the developing fetus. In particular, note should be made concerning management and/or control of chronic medical conditions that may impact fetal development such as hypertension and diabetes mellitus. Exposure to a variety of toxic and infectious agents is also important during this time period. Note should be made of exposure to drugs, medications, live vaccine preparations, or radiation. Known or suspected exposure to infectious agents should also be documented including sexually transmitted disease, cytomegalovirus, toxoplasmosis, tuberculosis, HIV, rubella, or varicella.

In addition to exposures, this period of early pregnancy is also marked by a variety of potential complications of pregnancy such as abnormal implantation or hyperemesis gravidarum. Patients should be screened for possible pregnancy-related symptoms such as bleeding or abnormal discharge per vagina, abdominal pain, headache, visual complaints, and severe nausea or vomiting.

*Family History.* A number of familial conditions and events are of potential consequence in pregnancy. A family history should be obtained with an emphasis on fetal deaths or abnormalities, multiple gestations, gestational diabetes, large- or small-for-gestational-age infants, significant genetic history (e.g., cystic fibrosis, Down syndrome, thalassemia), and chronic medical conditions with a strong familial link such as hypertension, diabetes, renal disease, and substance abuse.

## **Physical Examination**

All patients should receive a comprehensive physical examination. The purpose of the examination is threefold:

1. Diagnostic or supportive evidence of underlying medical conditions
2. Notation of normal physical changes often associated with pregnancy
3. Evaluation of the uterus and bony pelvis

A variety of physical changes can be noted during pregnancy. Although underlying medical conditions must be excluded, these findings generally represent benign changes and should be noted primarily for reference in case of change. Such findings include split S1 and/or a systolic ejection murmur on cardiac examination, mild thyroid enlargement, skin changes including malar rash and striae gravidarum, and accentuated lordosis noted on musculoskeletal examination.

Evaluation of uterine size is helpful in confirming the gestational age. At 6 weeks' gestation, the uterus has enlarged beyond its prepregnant dimensions and is described as the size of an orange. By 8–10 weeks, the uterus is described as grapefruit sized. At 10–12 weeks, the uterus is palpable on abdominal examination at the symphysis pubis. Once the uterus is palpable from the abdomen, measurement is made of the fundal height. This measurement represents the distance from the symphysis pubis to the top of the uterine fundus. By 20 weeks' gestation, the uterine fundus should normally be found at the level of the umbilicus. From 20 to 34 weeks' gestation, measurement of the fundal height (in cm) should be equivalent to the gestational age (in weeks). For example, at 28 weeks' gestation, the fundal height should be approximately 28 cm from symphysis to the top of the uterine fundus ( $\pm 2$  cm). Any significant deviation from these expected measurements should prompt the provider to reassess the gestational age and/or fetal development.

Assessment of the bony pelvic configuration (clinical pelvimetry) is often performed early in pregnancy to determine potential structural impediments to successful delivery. Abnormal findings do not rule out the possibility of a trial of labor and successful vaginal delivery but added caution may be warranted at the time of delivery. Notation should be made of the diagonal conjugate (distance from symphysis pubis to sacral promontory), ischial spines (blunt, prominent) sacrum (concave, straight), coccyx (fixed, mobile), and the pubic arch (normal, wide, narrow).

## **Laboratory and Diagnostic Testing**

As previously noted, all patients for whom pregnancy status is uncertain should have a confirmatory pregnancy test. This may take the form of either a urine or serum test for hCG. The urine hCG test is generally positive beginning at the time of the first missed period (approximately 4 weeks gestation). The serum test is generally positive at the time of implantation.

For all prenatal patients, routine obstetrical laboratory screening should include complete blood count with platelets, blood type (ABO and Rh), rapid plasma reagins, rubella titer, hepatitis B surface antigen (to detect active disease), HIV antibody, Papanicolaou smear (if not performed within the preceding 3 months), gonorrhea and chlamydia screen, and routine urine analysis with culture.

Selected patients may also benefit from screening for the following conditions: sickle cell disease (via routine screen or hemoglobin electrophoresis), Tay–Sachs, toxoplasmosis, cytomegalovirus, elevated serum lead, elevated glucose, substance use, and/or herpes simplex virus.

These screening tests are recommended at the first prenatal visit, which should routinely occur in the first trimester. For those patients who present for a first prenatal visit later in pregnancy, laboratory and diagnostic testing should include those tests just mentioned as well as all tests appropriate to their estimated gestational age at the time of presentation. These additional tests are described later.

## **Estimating Gestational Age**

At the conclusion of the first prenatal visit, a clinical assessment of gestational age and estimated date of delivery should be established. If the available information is incomplete or contradictory, a tentative date may be assigned with arrangements for acquisition of more definitive data.

The available data that may contribute to establishment of the gestational age includes the following:

1. LMP: if the menstrual history is certain, an accurate calculation can be made based on the first day of the LMP. Obstetrical calendars (often referred to as an “OB wheel”) are available (both physical and electronic, online or as an app for smartphones) that allow such calculation. In the absence of such a wheel, the EDD can be calculated by subtracting 3 months and adding 1 week to the LMP. For example, if the first day of the LMP was April 14, subtracting 3 months would yield January 14 and adding 1 week would then yield January 21 of the following year as the EDD.
2. Physical examination: as noted previously, uterine size can be used to give a clinical estimate of gestational age. When this data is congruent with the LMP, it is generally accurate to within 1 week.
3. Developmental milestones: the fetal heartbeat should be detectable with a hand-held Doppler at approximately 10 weeks of gestation and with a fetoscope at 18–20 weeks. Fetal quickening (fetal movement) should be reported by the mother at 18–20 weeks’ gestation.
4. Ultrasound: an obstetric ultrasound obtained early in pregnancy (first trimester or early second trimester) is accurate to within 1 week of gestation. The accuracy of ultrasound dating diminishes with advancing fetal age. In the third trimester, fetal ultrasound is accurate to within 2 weeks up to 36 weeks’ gestation and to within 3 weeks thereafter.

## **Laboratory Testing**

Levels of hCG obtained via quantitative testing are not considered accurate for dating purposes as significant variability is noted at any given gestational age.

As a general rule, the first gestational age/EDD assigned should remain unchanged throughout the pregnancy unless significant doubt exists concerning the data used to establish that date. Caution should be exercised when “correcting” an EDD based on later data.

## **Treatment and Follow-Up**

All patients not already taking prenatal vitamins should be given a prescription and encouraged to immediately begin taking one vitamin per day. In addition, iron supplementation should be considered for those patients with documented anemia.

At the first prenatal visit, patients should be given an overview of the course of prenatal care, a general caution concerning warning symptoms that should prompt early follow-up, and arrangements should be made for an appropriate follow-up visit. Patients should be given the opportunity to ask questions and to clarify any areas of uncertainty. It may be helpful to suggest that patients keep a list of written questions between visits to help ensure that all important concerns are addressed at each visit.

## ***Follow-Up Prenatal Visits***

Follow-up visits are designed to meet four purposes: to track the progress of the pregnancy; early identification of complications, if any; completion of testing/evaluation at specific gestational-age milestones; and patient education/anticipatory guidance.

Follow-up visits provide the opportunity to elicit patient questions and concerns and to provide education and counseling. Topics for review include nutritional status, activity and exercise in pregnancy, and symptoms consistent with preterm labor. Plans for delivery, such as when to contact the provider and when to go to the hospital, should be addressed. Also discussed should be issues related to delivery and the postpartum period, including use of anesthesia, breastfeeding, circumcision, and the duration of the postpartum stay.

Follow-up intervals noted here are for uncomplicated pregnancies. Follow-up must be arranged as indicated should abnormalities present at any point.

## **Interval History and Physical Exam**

The interval history should focus primarily on events and developments in the time period from the last visit to the current visit. Review of unresolved or ongoing concerns should also occur. Aspects of the history that should be addressed at every visit would include fetal movement (presence or absence and quantity), uterine contractions, pelvic pain or pressure, abdominal or back pain, and discharge or bleeding per vagina.

The physical examination contributes to assessment of fetal development and may also provide clues to the detection of pregnancy-related complications. The core elements of the physical examination include blood pressure, notation of edema, and weight. Assessments of fetal development include measurement of fundal height and documentation of the fetal heart rate.

Appropriate weight gain in pregnancy is a very common concern for pregnant patients. In population studies, optimal outcomes have been associated with maternal weight gain of 20–25 lb during pregnancy. Current recommendations for weight gain during pregnancy suggest 25–35 lb for a woman of average weight at the onset of pregnancy. Decreased weight gain can be associated with inadequate nutritional intake, inaccurate dating, intrauterine growth retardation, oligohydramnios, and

fetal demise. Increased weight gain can be associated with excessive caloric intake, inaccurate dating, macrosomia, multiple gestation, and polyhydramnios. Abnormal weight gain should prompt review of the prenatal course, eating patterns, and fetal well-being surveillance.

### **Interval Laboratory and Diagnostic Studies**

Each pregnancy will require individualization of appropriate interval laboratory and diagnostic studies. For most pregnancies, however, a variety of interval studies should be considered. These include the following:

1. At 10–13 weeks' gestation, screening for chromosomal abnormalities can be performed using ultrasound assessment of fetal nuchal translucency in combination with B-HCG and pregnancy associated.
2. At 15–20 weeks gestation, patients not screened earlier should be offered maternal triple (hCG, alpha fetal protein, and estriol) or quadruple (hCG, alpha fetal protein, estriol, inhibin) screening for developmental abnormalities. In addition, an obstetrical ultrasound is often ordered in this time period although it may be ordered earlier or later for specific indications. Indications for obstetrical ultrasound include uncertain gestational age, termination of pregnancy, induction, scheduled repeat cesarean section, fetal growth evaluation, suspected multiple gestation, size–date discrepancy, adjunctive to amniocentesis or chorionic villus sampling, suspected hydatidiform mole, suspected ectopic pregnancy, suspected fetal demise, suspected poly-/oligohydramnios, bleeding per vagina, maternal pelvic mass, uterine abnormality/uterine scar assessment, ovarian follicle surveillance, fetal biophysical assessment, placenta localization, past history of fetal congenital anomaly, or late initiation of prenatal care.
3. At 24–28 weeks' gestation, patients should undergo glucose challenge testing for gestational diabetes, repeat hemoglobin (28 weeks), and antibody screen (28 weeks).
4. At 35–37 weeks, RPR, Hep B sAg, HIV, GC/chlamydia, and Group B strep are all indicated.

# Chapter 4

## Medications in Pregnancy

### Contents

Background .....	28
General Principles of Medication Use in Pregnancy .....	28
Chronic Medical Conditions .....	28
Acute Medical Conditions.....	29
Acute Obstetrical Conditions .....	29
Therapeutic Categories and Considerations.....	30
Proven Human Teratogens .....	30
Special Considerations.....	32
Tobacco .....	32
Alcohol .....	32
Illicit Drugs .....	33
Over-the-Counter Medications .....	33
OTC Pain Medications .....	33
OTC Cough/Cold/Allergy Medications .....	34

### Key Points

1. All medications should be viewed with caution in pregnancy.
2. Management of medication use in pregnancy ideally begins with adequate pre-conception counseling and prepregnancy planning.
3. Medications used in pregnancy require clear identification of indication for use, duration of treatment, expected outcome, and signs or symptoms requiring early termination of their use.
4. When in doubt consultation with an expert in maternal–fetal medicine is strongly recommended.

## Background

With the exception of prenatal vitamins and possibly iron supplementation, all medications should be used with caution during pregnancy. Although clinical experience with many medications in pregnancy is quite extensive and the safety and efficacy is reasonably established, pregnancy represents a unique challenge in medication assessment. It would not be ethical, under most circumstances, to randomize pregnant patients to receive increasing doses of medications to assess safety and efficacy of a medication known to produce or suspected of producing harm in pregnancy. This limits the degree to which safety can be categorically stated for the use of any medication in pregnancy. Many medications once thought to be safe in pregnancy have subsequently been shown to be harmful. Other medications originally thought to be harmful have been shown to have beneficial effects when used for specific medically indicated purposes.

Many medications can and should be used in pregnancy for a variety of legitimate medical indications. Although it is beyond the scope of this chapter to discuss in detail the use of all medications in pregnancy, a few general guidelines can be offered. Texts exist that detail the risks and benefits of many available medications. Such a text should be a routine part of every obstetrical provider's library. When doubt exists concerning the indications for or the safety or efficacy of any medication, consultation with an expert in maternal-fetal medicine is strongly recommended.

## General Principles of Medication Use in Pregnancy

The three general principles are as follows:

1. Chronic medications should be reviewed to assess safety and efficacy.
2. The risk of not treating (or treating less effectively) an identified disease (acute or chronic) should be weighed against the risks of the proposed treatment.
3. All medications used in pregnancy require clear identification of indication for use, duration of treatment, expected outcomes, and signs or symptoms that require early termination of use.

## *Chronic Medical Conditions*

Some patients have chronic medical conditions that predate pregnancy. The management of many of these conditions will include the use of medication. Although it is always important to consider safety and efficacy when using medications, with the onset of pregnancy these considerations become considerably more complex. Ideally, a consideration of the impact of pregnancy on the medical condition as well as the impact of the medical condition on pregnancy would occur prior to

pregnancy. For some patients, this leads to a recommendation to delay pregnancy until the medical condition can be more adequately controlled. In other circumstances, it may lead to a recommendation to avoid pregnancy altogether.

Under many circumstances (both planned and unplanned), however, management of pregnancy will overlap with management of chronic medical conditions and their associated medications. The first consideration should be for the safety of the mother. Disease processes that are life threatening to the mother may require continuation of treatment even if pregnancy is continued. The provider should explore treatment alternatives with equal efficacy and better established safety profiles when possible. When safer alternatives are not available, providers should discuss with patients the potential risks of continuing the pregnancy while simultaneously continuing the use of the required medication versus the potential risks of terminating medication use for the duration of pregnancy. Patients must be given sufficient information to make an informed decision concerning their health and the health of their developing fetus, especially during the critical period of organogenesis early in pregnancy. In all circumstances providers and patients must make individualized treatment decisions based on the medical conditions of the specific patient.

### ***Acute Medical Conditions***

Pregnant patients are vulnerable to all the acute medical conditions of nonpregnant patients. Medical decisions concerning the treatment of acute medical conditions that arise during pregnancy must follow the same general guidelines as those for chronic conditions. Will the medical condition adversely affect the pregnancy? Will treatment of the condition ameliorate or eliminate these potential effects? Will the proposed treatment adversely affect the pregnancy? Are there safer or more well-established alternative treatment options? What are the *likely* consequences of not treating the medical condition? What are the *potential* consequences of not treating the medical condition?

As with the treatment of chronic medical conditions, it is critical that both providers and patients have sufficient information concerning the risks and benefits of treatment options to make informed, individualized decisions. When providers cannot adequately answer these questions, patients should be referred to a provider with sufficient expertise to provide more complete information.

### ***Acute Obstetrical Conditions***

Pregnancy may be accompanied by a variety of complications that require consideration of medication use. The same general principles apply and the same questions must be answered. When these complications are relatively common, much established data may exist to guide providers and patients in their decision-making process. When the complications are less common, consultation may be required.

## ***Therapeutic Categories and Considerations***

Any drug used during pregnancy should be checked for safety prior to use. Keeping in mind the general considerations just given, the following recommendations may be considered:

1. When antibiotics are indicated consider penicillin, cephalosporins (except cefotetan), clindamycin, and macrolides. Avoid sulfa drugs (contraindicated in the first and third trimester), quinolones, tetracyclines, and aminoglycosides (ototoxic, may be indicated for severe Gram-negative infections).
2. When analgesics are indicated consider acetaminophen and narcotic analgesia (consult reference for specific agents). Narcotic analgesics do cross the placenta and may affect the fetus transiently. Long-term narcotic analgesia use (or abuse) during pregnancy can be associated with withdrawal symptoms in the newborn. Narcotic analgesia at or near delivery has been associated with respiratory depression in newborns, which can be reversed, if necessary, with Narcan. Avoid aspirin (in analgesic doses) and nonsteroidal anti-inflammatory drugs (NSAIDs, contraindicated in late pregnancy).
3. For treatment of hypertension, consider labetalol (individual  $\beta$ -blockers should be reviewed prior to use as some  $\beta$ -blockers have been associated with adverse effects on uteroplacental and fetal hemodynamics and fetal growth), methyldopa, and hydralazine. Avoid angiotensin-converting enzymes (and angiotensin receptor blockade agents).
4. For patients with diabetes, consider insulin and regular- and intermediate-acting agents, but avoid oral hypoglycemics (recent data suggests that some oral hypoglycemics may be safely used in pregnancy but experience is limited and individual agents should be reviewed prior to use).
5. For patients suffering from nausea, consider using Diclectin (doxylamine/pyridoxine) or chlorpromazine.
6. In cases of gastritis/peptic ulcer disease, consider magnesium hydroxide, aluminum hydroxide, calcium carbonate, and bismuth subsalicylate.

## **Proven Human Teratogens**

Some agents have proven teratogenic potential. These agents are summarized in Table 4.1. Although the effects of such agents are potentially variable and predictable, their use should be very limited or avoided during pregnancy. Category D agents have proven teratogenic potential but may, under certain circumstances, be indicated. Prior to using any category D agent, providers should perform a careful review of indications, duration of therapy, all potential effects, and all potential alternatives to the proposed therapy. In addition, patients should be informed of these considerations, allowing for informed consent to the proposed therapy. Category X medications have proven teratogenic potential and use should be avoided in pregnancy.

**Table 4.1** Selected proven human teratogens

Category D
ACE inhibitors/ARB
Cyclophosphamide
Lithium
Paramethadione
Phenytoin
Barbiturates
Carbamazepine
Benzodiazepines
Systemic retinoids
Tetracycline
Tamoxifen
Trimethadione
Valproic acid
Warfarin
Category X
Thalidomide
Methotrexate
Danazol
Misoprostol
Diethylstilbestrol

**Table 4.2** Medication safety ratings in pregnancy

Category A: controlled human studies have demonstrated no increased risk of fetal harm
Category B: no controlled human studies suggest increased risk; probably safe in pregnancy when use is indicated
Category C: no evidence in human studies of increased risk but animal studies show possible increased risk
Category D: evidence in human studies suggest increased risk but benefits of treatment may outweigh risk
Category X: clear evidence of teratogenic effects exists; contraindicated in pregnancy

Teratogenicity is a multifactorial outcome that is related to several factors including (1) the teratogenic potential of the agent, (2) the susceptibility of the fetus, (3) the dose/duration of exposure, and (4) the timing of the exposure. As noted in Table 4.2, medications can be classified by their observed correlation with fetal harm in animal or human studies. Susceptibility may be based on the genetic factors as well as the gestational age of the fetus. As a general rule very early exposure (approximately first 2 weeks of gestation) results in an all-or-none phenomenon which may result in fetal demise. Exposure during the period of organogenesis (approximately weeks 3–9) are more likely to result in major morphologic abnormalities. Exposure thereafter may result in variable defects. Although this may serve as a guideline, careful assessment and possible consultation should be considered for any patient with a significant antenatal exposure.

## Special Considerations

In addition to the use of medications in pregnancy, a variety of other exposures may occur with possible effects on pregnancy. These might include occupational exposures, legal and illegal drugs or nonpharmacological items such as exercising, lifting, or other activities. As previously noted, most such exposures will be subject to limited data concerning possible pregnancy effects. For this reason, the same general principles should apply that apply to medication use:

1. Is there any data available to guide the decision?
2. Is there a specific and compelling reason for the exposure?
3. Do safer alternatives exist?
4. Can potential adverse effects be monitored?
5. Can exposure be limited or modified in such a way as to minimize potential risks?

Although a full discussion of all such exposures is the work of an entire text in its own right, three common and frequently encountered exposures deserve attention.

### ***Tobacco***

Tobacco is associated with a variety of adverse outcomes including low birth weight, increased risk of fetal demise, abruptio placenta, and placenta previa. Although the absolute risk associated with tobacco use is not clearly defined, the outcomes are potentially quite severe. As tobacco has no known benefits in pregnancy, every effort should be made to reduce or eliminate tobacco exposure during pregnancy.

### ***Alcohol***

Fetal alcohol syndrome (FAS) is a constellation of developmental and physical findings in neonates born to mothers who consumed large quantities of alcohol during pregnancy. FAS is associated with growth retardation, microcephaly, microphthalmia, and central nervous system deficiencies. The use of alcohol is quite prevalent and a significant number of pregnant mothers will have consumed alcohol prior to becoming pregnant. The question frequently arises whether alcohol can be safely used in any quantity during pregnancy. Although data is limited, there is no established safe level of alcohol use in pregnancy. For this reason, patients should be encouraged to eliminate or significantly limit alcohol use during pregnancy.

## ***Illicit Drugs***

A variety of illicit drugs are associated with adverse pregnancy outcomes. Each drug should be reviewed individually for specific concerns. In addition to the medical considerations, all such drugs are, by definition, illegal and carry with them significant social risk. All pregnant patients should be screened by history for illicit drug use and, when present, counseled concerning the desirability of reduction or elimination. This screening and counseling should be approached in a nonjudgmental and nonthreatening manner. A threatening or legalistic approach to patients is likely to reduce patient reporting and therefore limit providers' ability to effectively intervene.

## **Over-the-Counter Medications**

Questions concerning the use of over-the-counter (OTC) medications arise frequently during the course of pregnancy. The general considerations for use are the same as for prescription medications. The ease of access combined with the frequency with which OTC medications are used makes recommendations in pregnancy particularly challenging. It is estimated that more than half of all medication use in the United States is OTC. Approximately 75 % of all pregnant patients will use one or more OTC medication during the course of pregnancy.

Although providers can control access to prescription medications, OTC medications are, by definition, available to patients without the necessity of a prescription. In addition, the safety of OTC medications may vary with the time in pregnancy when they are used. For example, OTC NSAIDs that may be safely used early in pregnancy are generally contraindicated in the third trimester. For this reason, providers must be able to discuss in detail the appropriate uses and precautions that patients musts keep in mind when deciding whether to use OTC medications during pregnancy.

Food and Drug Administration (FDA) pregnancy safety ratings are available for all OTC medications and should be reviewed prior to their use. It should be noted that supplements such as herbal and/or natural preparations are not subject. FDA oversight and information concerning safety in pregnancy may be very limited. The breadth of OTC medications available make a comprehensive review beyond the scope of this text; however, two common categories of medications warrant consideration: pain medications and cough/cold/allergy medications.

## ***OTC Pain Medications***

Pain is among the most common of medical complaints, and although the frequency of pain in pregnancy may not be higher than in the nonpregnant state, it is certainly not any less common. For this reason, patients will often seek advice from their providers on OTC pain medication options.

## **Acetaminophen**

Acetaminophen is widely used in pregnancy and early childhood. Although randomized, controlled trials concerning the safety of acetaminophen in pregnancy are lacking, the extensive experience combined with few reports of complications makes the use of acetaminophen a safe choice in pregnancy. Acetaminophen is a category B drug in all stages of pregnancy.

## **Aspirin**

Use of aspirin has been associated with a variety of potential pregnancy complications, including prolonged gestation and decreased birth weight. In addition, aspirin has potent antiplatelet activity that may predispose patients to bleeding. The use of aspirin in pregnancy has been associated with neonatal hemorrhage. Aspirin is a category D drug in all stages of pregnancy and its use as an OTC medication should be discouraged during pregnancy.

## **Nonsteroidal Anti-inflammatory Drugs**

Although NSAIDs are often combined as a single class, significant differences may exist in their use during pregnancy. Indomethacin (a prescription NSAID) has been used in pregnancy for preterm labor (see Chap. 7) but has been associated with significant complications and should only be used with caution and for specific indications. OTC NSAIDs, including ibuprofen and naproxen, are category B drugs in early pregnancy but category D drugs in the third trimester. In the absence of a compelling indication for the use of NSAIDs in pregnancy, their use as an OTC medication should probably be limited.

## ***OTC Cough/Cold/Allergy Medications***

Given the duration of pregnancy, the likelihood of experiencing symptoms of allergies or viral respiratory infections is quite high. A wide variety of OTC medications are available to treat the various symptoms of viral respiratory infections and allergies and their use is quite common in pregnancy.

## **Antihistamines**

Chlorpheniramine is a commonly used antihistamine in a variety of allergy and cold formulations. Chlorpheniramine is a category B drug and is probably safe for use during pregnancy. Diphenhydramine is the second commonly used antihistamine

and is also a category B medication. It should, however, be used with caution as it has been shown to cross the placenta, may have oxytocin-like effects at high doses, and may interact with other drugs in pregnancy.

## **Decongestants**

Pseudoephedrine has been the subject of animal studies and has had widespread human use in pregnancy. Pseudoephedrine is a category B medication and its use in pregnancy is probably safe. Because it has been associated with a possible increase in the risk of gastroschisis, its use in the first trimester of pregnancy should probably be avoided when possible.

## **Cough Medications**

Cough medications fall into two broad categories: antitussive medications and expectorants. Both categories of medication are available in a wide variety of OTC formulations. Guaifenesin, a common expectorant, is a category C drug. Its use in the first trimester of pregnancy has been associated with a possible increased risk of neural tube defects but data is limited. It is probably safest to avoid the use of guaifenesin in the first trimester of pregnancy when possible. Dextromethorphan is also a category C drug. Animal studies have shown an association between dextromethorphan exposure and birth defects. Human data has not found a similar association.

## **Part II**

# **Complications of Pregnancy**

# Chapter 5

## Dysmorphic Growth and Genetic Abnormalities

### Contents

Background .....	39
History.....	40
Physical Examination.....	41
Diagnosis.....	41
Obstetrical Screening Tests .....	41
What Is Measured .....	41
What Is Detected.....	42
Confirmation/Follow-Up.....	42
Amniocentesis.....	42
Chorionic Villus Sampling.....	43

### Key Points

1. Growth and development represent a complex interaction between genetic predisposition and environmental exposure.
2. Screening for genetic risk factors ideally begins in the preconception period.

### Background

Most fetuses are free from identifiable genetic abnormalities. Their pattern of growth and development falls within the range of normal parameters. Under some circumstances physical growth is restricted (intrauterine growth restriction). This is discussed in Chap. 6. Under other circumstances, however, genetic abnormalities lead to abnormalities in growth, development, or both. These abnormalities may be relatively minor (e.g., color blindness) or they might be more significant (e.g., muscular dystrophy). Primary care providers must be familiar with common abnormalities and available screening options to identify these conditions when they arise.

Genetic inheritance is a complex interaction of maternal and paternal genotypic predisposition with a variety of environmental factors. The resulting phenotypic expression represents final outcome of these two factors. A review of both familial genetic predisposition and environmental risk factors will allow providers to discuss with expectant parents the developmental risks, if any, associated with a particular pregnancy.

Chromosomal abnormalities fall within two major groups, abnormalities of chromosomal number (e.g., trisomy 21, Down syndrome) and abnormalities of chromosomal structure (e.g., hemophilia). Chromosomal abnormalities (genotypes) may have variable phenotypes. This variability can represent variable expressivity (variation in expression of a disease within a population with a specified genetic abnormality) and/or variable penetrance (likelihood of expression within an individual with a genetic abnormality).

## History

Ideally genetic screening by history would begin in the preconception period. If that is not possible, it should begin as early in the prenatal period as possible. A variety of standardized screening tools exist to assist providers with the evaluation of maternal and paternal genetic risk factors. Table 5.1 provides a selected list of conditions commonly screened for in pregnancy. Common medical conditions with a strong genetic predisposition should be reviewed, including congenital abnormalities, cystic fibrosis, Down syndrome, hemophilia, sickle cell disease, neural tube defects, Tay–Sachs disease, and muscular dystrophy, among others. Although these standardized screening tools are helpful as a starting point, providers should be aware that additional detail may be required for some high-risk patients. If the primary care provider does not feel adequately prepared to complete such a genetic screening, referral should be made to a genetics counselor or a maternal–fetal medicine specialist.

**Table 5.1** Selected congenital conditions screened for in preconception/prenatal care

Endocrine disease
Autoimmune disease
Congenital abnormalities
Cleft lip/palate
Congenital heart disease
Cystic fibrosis
Down syndrome
Mental retardation
Neural tube defects
Hemophilia
Sickle cell disease
Thalassemia
Huntington's chorea
Tay–Sachs disease

In addition to a thorough family history, a history of recurrent early spontaneous abortions should be noted if present. Approximately 50 % of early spontaneous abortions are estimated to demonstrate genetic abnormalities. The age of the mother should be noted, as fetal genetic abnormalities increase with increasing maternal age. The prevalence of such disorders ranges from approximately 2 per 1000 live births at a maternal age of 25 years to approximately 150 per 1000 by maternal age 50.

## **Physical Examination**

Although a physical examination may, rarely, uncover a genetic abnormality not discovered through careful history taking, in general the physical examination contributes little to the assessment of risk for genetic abnormalities.

## **Diagnosis**

Although careful historical screening will identify many patients at risk for genetic and/or developmental abnormalities, adjunctive diagnostic testing is available and should be offered during the course of prenatal care. A wide variety of screening options are available and patients requiring specialized testing should be managed in conjunction with a genetics counselor and an experienced maternal-fetal medicine specialist. Three such tests (obstetrical triple screen, amniocentesis, chorionic villus sampling [CVS]), however, are offered with sufficient frequency that all primary care providers should be familiar with their basic function.

## **Obstetrical Screening Tests**

### First Trimester Combined Test

#### ***What Is Measured***

This test measures maternal serum levels of pregnancy-associated plasma protein-A, human chorionic gonadotropin, in conjunction with an ultrasound measure of nuchal translucency. Standardized values are established for these three components during weeks 9–13 of pregnancy. Screen positive results are followed by CVS (see below).

## ***What Is Detected***

### **Down Syndrome**

Down syndrome is a genetic abnormality of abnormal chromosomal distribution (trisomy) associated developmentally with mental retardation of variable degree, characteristic physical stigmata, and an association with a variety of other medical conditions including cardiac and hematological diseases. The prevalence of Down syndrome increases with increasing maternal age and with prior history of Down syndrome.

Down syndrome should be suspected with abnormally low triple screen results. Triple screen testing will identify approximately 60 % of all cases of Down syndrome. Although this is sufficiently sensitive for low-risk populations, higher-risk populations may require more accurate testing methods (see the next section). Abnormally low triple screen testing is falsely positive in approximately 5 % of cases. The most common cause of false-positive testing is inaccurate gestational dating.

### **Neural Tube Defects**

Although the serum tests of this screen can be correlated with the risk of neural tube defect, it is more accurately measured during the second trimester.

### ***Confirmation/Follow-Up***

Triple screen testing is, by definition, a screening test and all abnormal values require confirmation and diagnostic follow-up. For all abnormal values, confirmation of gestational age is critical. All data used to establish the gestational age should be reviewed and confirmed. Under some circumstances repeat triple screening may be indicated.

Confirmed abnormal elevations in triple screen results should be followed up with obstetrical ultrasound. This study will allow for careful examination of the anatomy of the developing fetus. In addition, ultrasonographic studies should allow for identification of multigestation and fetal demise.

Confirmed abnormally low triple screen results should be followed by genetics counseling with amniocentesis and fetal karyotyping.

### ***Amniocentesis***

Amniocentesis is a diagnostic procedure that consists of introduction of a sampling needle through the abdominal wall into the amniotic sac. A small sample of amniotic fluid is withdrawn, allowing for a variety of potential studies. The most common study performed with the fluid is determination of the fetal karyotype.

As noted earlier, an abnormal low triple screen is one indication for amniocentesis. Other indications include maternal age over 35 years, prior chromosomal abnormality, three or more prior spontaneous abortions, and known parental chromosomal abnormality.

Amniocentesis can be performed between 12 and 17 weeks' gestation. The risk of complications is higher earlier in pregnancy. For this reason, most are performed between 15 and 17 weeks' gestation. The most significant complication associated with amniocentesis is spontaneous abortion. The reported rate of post-amniocentesis spontaneous abortion is approximately 1 per 200 procedures, although the risk may be significantly lower in settings with highly experienced operators. Sensitivity and specificity are both >99 %.

### ***Chorionic Villus Sampling***

The need to wait until 15 weeks' gestation represents a significant limitation for amniocentesis in some cases. For this reason, alternative methods that may be performed earlier have been developed. One such method is CVS. The procedure is performed via transcervical or transabdominal approach with the target being at least 5 mg of chorionic villus tissue. CVS may be performed in earlier than amniocentesis (after 10 weeks' gestation) with similar sensitivity, specificity, and complication rates. Indications for CVS include:

1. Maternal age >35
2. Prior chromosomal abnormality
3. Positive serum screening test
4. Ultrasonographic abnormality
5. Parental chromosomal abnormality

# Chapter 6

## Intrauterine Growth Restriction

### Contents

Background .....	46
Risk Factors .....	47
Fetal-Genetic Factors .....	47
Uterine-Environmental Factors .....	47
Maternal Factors .....	47
Toxic Exposures .....	48
Constitutional Factors .....	48
Complications of Growth Restriction .....	49
Antenatal Tracking and Diagnosis .....	49
Screening for Risks .....	49
Diagnosis .....	51
Management .....	52
Risk Reduction .....	52

### Key Points

1. Intrauterine growth restriction (IUGR) is defined as fetal weight below the tenth percentile and an abdominal girth below the 2.5 percentile.
2. Factors associated with IUGR can be categorized as fetal genetic, uterine environmental, maternal, toxic exposures, and constitutional.
3. Complications associated with IUGR include early complications (increased mortality, pre-eclampsia, preterm labor, still birth) and late complications (learning, behavioral, and developmental abnormalities).

## Background

Fetal growth is among the most important of parameters monitored during the course of prenatal care. Although the majority of pregnancies proceed with no complications in fetal growth, a small number will show evidence of growth restriction. A variety of conditions are associated with or increase the risk for intrauterine growth restriction (IUGR). These factors are summarized in Table 6.1.

**Table 6.1** Risk factors for intrauterine growth restriction (IUGR)

Fetal-genetic factors
Chromosomal abnormalities (e.g., trisomy 21)
Neural tube defects
Achondroplasia
Osteogenesis imperfecta
Gastrointestinal (gastroschisis, duodenal atresia, pancreatic agenesis)
Renal disease (renal agenesis)
Neurofibromatosis
Uterine-environmental factors
Infection
Cytomegalovirus
Rubella
Herpes, varicella
Influenza
Toxoplasmosis
Oligohydramnios
Placental abnormalities
Placenta previa
Abruptio placentae
Placental malformation
Multigestation
Uterine anatomic abnormalities
Maternal factors
Prior IUGR infant
Hypertension
Diabetes (may also be associated with macrosomic infants)
Nutritional deficits
Gastrointestinal malabsorption
Constitutional short stature
Vascular disease
Toxic exposures
Smoking
Alcohol
Illicit drugs (heroin, cocaine)
Prescription medications (folic acid antagonists, warfarin)
Constitutional
Constitutional short stature
Female
Birth order

IUGR is defined as fetal weight below the tenth percentile and an abdominal girth below the 2.5 percentile for gestational age. As this definition implies, accurate fetal dating is critical to the diagnosis. At term, this corresponds to a birth weight of 2500 g (~5.5 lb). By this definition, approximately 5 % of all US infants demonstrate evidence of IUGR, accounting for approximately 175,000 infants annually in the United States.

## **Risk Factors**

Factors associated with IUGR can be categorized into five categories: fetal genetic, uterine environmental, maternal, toxic exposures, and constitutional factors.

### ***Fetal-Genetic Factors***

A variety of genetic conditions are associated with restricted fetal growth and account for approximately 10 % of all cases. Early fetal growth is predominantly driven by growth in the number of fetal cells, cellular hyperplasia. This process is strongly influenced by genetic and fetal factors. As such early growth restriction is predominantly associated with fetal-genetic factors. Such conditions as Down syndrome (trisomy 21) and Turner's syndrome are associated with an increased risk for IUGR. Structural growth defects such as anencephaly, achondroplasia, osteogenesis imperfecta, and renal and gastrointestinal (GI) defects are all associated with an increased risk of IUGR.

### ***Uterine-Environmental Factors***

Abnormalities of the uterine environment may also contribute to growth restriction. Congenital infections such as cytomegalovirus, toxoplasmosis, or rubella are associated with restricted fetal growth. Although these infections can cause significant fetal growth restriction, they are a relatively uncommon cause of IUGR, <5 %. Fetal urinary tract outflow obstruction with concomitant oligohydramnios can be associated with IUGR. Placental abnormalities (placenta previa, abruption, placental malformation) and multiple gestations can both alter the uterine environment and therefore lead to growth restriction. The structure of the uterus itself may also contribute to growth restriction. Uteroplacental insufficiency is most commonly associated with late-onset growth restriction.

### ***Maternal Factors***

Growth in pregnancy is fundamentally reliant on a balance between fetal needs (especially nutrition and oxygen) and maternal ability to meet those needs. Maternal health and nutrition contribute significantly to fetal growth and a number of maternal

factors can be associated with decreased fetal growth. Although specific maternal factors may not always be identified, a pregnant patient with a past history of IUGR is approximately twice as likely to have subsequent IUGR compared to those without such a history. Following the period of cellular hyperplasia, fetal growth is augmented by cellular hypertrophy. Growth restriction due to decreased cellular hypertrophy may be more responsive to correction of nutritional imbalances.

Underlying maternal diseases such as diabetes, hypertension, GI, and vascular disease are all associated with an increased risk for IUGR. Hypertension is associated with decreased placental perfusion resulting in decreased delivery of both oxygen and nutrients. GI disease with significant malabsorption will result in decreased delivery of nutrients to the fetus and may contribute to subsequent IUGR. It should be noted that less significant nutritional defects may not result in diminished fetal growth as nutrients are delivered preferentially to the fetus.

Fetal growth is dependent on adequate fetal oxygenation and delivery of necessary nutrients. Adequate delivery is dependent, in turn, on adequate vascular function. Maternal vascular disease associated with a number of diseases impairs maternal-fetal perfusion and therefore is associated with increased risk of IUGR. Such maternal conditions as diabetes mellitus, peripheral vascular disease associated with tobacco use, collagen vascular disease, and pregnancy complications such as pre-eclampsia can all decrease maternal-fetal perfusion, thus increasing the risk of IUGR.

## ***Toxic Exposures***

The use of toxic substances (tobacco, alcohol, and illicit drugs) has been associated with decreased fetal growth. Of these, tobacco use is by far the most common risk factor for diminished fetal growth. Pregnant patients who smoke approximately double the risk of IUGR, a risk that is proportional to the quantity of cigarettes smoked. Discontinuation of smoking before or during pregnancy is associated with reduction or elimination of this risk. Both alcohol and illicit drug use (especially heroin and cocaine) are associated with diminished fetal growth. Fetal alcohol syndrome infants are generally small for gestational age.

Prescription drug use may also be associated with decreased fetal growth. Folic acid is a critical component in fetal development. Supplementation with folate is recommended during the preconception and early prenatal course. Drugs that inhibit folic acid metabolism may result in decreased fetal growth among other complications. The use of warfarin for anticoagulation has also been associated with IUGR.

## ***Constitutional Factors***

Small mothers are more likely to have small infants. Although these infants are small for gestational age, they are otherwise healthy. If no maternal, fetal, or uterine factors can be identified, these infants are generally healthy and normal but small.

There is some variation in size based on birth order and gender. First infants are slightly smaller, on average, than subsequent infants; female infants are somewhat smaller, on average, than male infants.

## Complications of Growth Restriction

Although not all small babies have complications, IUGR represents the second leading cause of perinatal morbidity and mortality after prematurity. Infants with a birth weight below 2500 g have a mortality rate of 5–30 times that of infants weighing more than 2500 g. IUGR (or associated underlying disease processes) is associated with an increased risk for pre-eclampsia, preterm labor/delivery, stillbirth, fetal electrolyte abnormalities, hypoglycemia, and aspiration. Long-term consequences associated with IUGR include increased risk for learning, behavioral, and developmental abnormalities.


## Antenatal Tracking and Diagnosis

Every prenatal visit includes assessment of fetal growth either indirectly or directly. A general approach to tracking fetal growth is presented in Fig. 6.1. Indirect measures of fetal growth include maternal weight gain, fundal height, and age-appropriate landmarks (e.g., fetal heart tones heard with handheld Doppler at approximately 10–12 weeks' gestation). Direct measures of fetal growth can be obtained with obstetrical ultrasound. Interpretation of these markers depends on accurate pregnancy dating. Parameters for establishing gestational age and estimated date of delivery (EDD) are covered in Chap. 3. In addition to assessing markers of fetal growth, all patients should be screened for risk factors that may contribute to an increased risk for IUGR.

## Screening for Risks

The first prenatal visit will include a comprehensive historical review of the patient's, patient's family, and paternal risk factors. Particular note should be made of prior obstetrical history, including infant size at delivery, prior history of genetic abnormalities including neural tube defects and Down syndrome, and multiple gestations. The mother's history should be reviewed for chronic disease such as hypertension, cardiovascular disease, renal disease, diabetes mellitus (including prior macrosomic infants with or without a known diagnosis of diabetes mellitus), GI disease, and nutritional status. The note should be made of alcohol, tobacco, or illicit drug use.

Physical examination at the first prenatal visit should include assessment of uterine size (and comparison of this finding to estimated gestational age). If appropriate,



**Fig. 6.1** Tracking fetal growth

fetal heart tones should be documented at this visit. Initial laboratory studies that may contribute to risk screening would include protein or glucose on urinalysis and evidence of cytomegalovirus, herpes, rubella, and/or toxoplasmosis.

Each subsequent visit will include interval updates for each of these risk factors. Patients should be screened for changes in nutritional status, toxic substance use, medications, or new exposures. Each prenatal visit will include documentation of

blood pressure, maternal weight, fundal height, and fetal heart rate. Particular attention should be paid to fundal height measurement as this can be a significant clue to growth abnormalities. Fundal height (measured symphysis pubis to top of fundus) in centimeters should be equal to gestational age in weeks ( $\pm 2$  cm).

At 15–20 weeks' gestation, patients should be offered  $\alpha$ -fetoprotein (or triple test) screening. Also, in this time period many patients will undergo obstetrical ultrasound examination. Key parameters on ultrasound include biparietal diameter, head circumference, abdominal girth, estimated fetal weight, and amniotic fluid index (AFI). Routine screening for gestational diabetes is performed at 24–28 weeks' gestation but may be performed earlier in patients with increased risk.

Patients who show abnormalities in any of these parameters should be considered at increased risk for IUGR with appropriate follow-up and diagnostic testing when indicated.

## Diagnosis

Diagnosis of IUGR requires correlation of gestational age with ultrasonographic measurements of fetal growth. The accuracy of the diagnosis is ultimately dependent on the accuracy of each of these components. IUGR must also be understood to be a *growth* phenomenon; a single static measurement may be suggestive but is not diagnostic. Serial measurements are generally required to diagnose IUGR.

Estimation of gestational age is based on the last menstrual period, key developmental markers (such as heartbeat), and sonographic data. Sonographic data can be useful for three purposes: (a) estimation of gestational age, (b) identification of structural or developmental abnormalities, and (c) measurement of fetal growth parameters. The most accurate ultrasound parameters for estimating gestational age vary with the age of the fetus. Crown–rump length is most accurate in early pregnancy; biparietal diameter and head circumference are most accurate for second trimester evaluation. Third trimester estimates are decreasingly accurate but head circumference may be most accurate. The increased sophistication of obstetrical ultrasounds has yielded images capable of remarkable anatomic detail. Cardiac, renal, GI, and neurological anomalies may all be detected on ultrasound examination. Patients with suspected IUGR should undergo ultrasound examination with particular attention paid to detailed anatomic survey.

In addition to the parameters mentioned here, abdominal girth is particularly helpful in measuring fetal growth. Abdominal girth reflects subcutaneous fat that, in turn, is a marker for adequate fetal nutrition. IUGR is often associated with an oligohydramnios (decreased amniotic fluid) that can be measured via the AFI. The AFI is the sum of the largest fluid pocket in each of the four quadrants. IUGR with oligohydramnios is associated with increased morbidity compared with IUGR alone.

Although not diagnostic of IUGR, measurement of fetal vascular flow, Doppler velocimetry, can provide additional clinical information regarding fetal well-being. Blood flow is measured in the umbilical artery and the ratio of systolic to diastolic

flow is calculated. Peak systolic frequency shift/end-diastolic frequency shift (*S/D*) ratios of 1.8–2.0 are normal. *S/D* ratios  $>3.0$  indicate a pregnancy at high risk for adverse outcomes. Measurement of flow can also be measured in the middle cerebral artery and the ductus venosus.

## Management

### ***Risk Reduction***

Although early identification and management can reduce the morbidity associated with IUGR, risk reduction remains a critical component of management. Ideally, such risk evaluation and reduction would begin in the preconception period with identification of pre-existing risk factors. For patients with a strong history of genetic abnormalities, genetics counseling may be indicated prior to conception. Identification of pre-existing medical conditions and nutritional deficits may also contribute to a reduction in the risk for IUGR. Although the benefits of such management in IUGR have not been well established, control of blood pressure, euglycemia prior to conception, and nutritional augmentation, among others, are recommended. Patients who smoke should be counseled to discontinue smoking; alcohol use should also be discontinued. The use of illicit drugs should be identified and cessation should be counseled. For many patients, pregnancy may represent a compelling reason to reduce or discontinue the use of toxic substances with benefit to both the mother and fetus.

Infectious risks for IUGR should be reviewed with the patient and reduction of exposure counseled. Toxoplasmosis is associated with raw or undercooked meat and cat feces. Patients should be counseled concerning exposure to those with varicella and rubella. If patients are seen in the preconceptual period, immune status should be documented for varicella and rubella. Vaccinations for both are available and should be administered to nonimmune patients who do not plan to become pregnant in the subsequent 3 months.

With diagnosis of IUGR, management involves three basic tasks: (a) identification and modification (when possible) of underlying etiology, (b) monitoring for anticipated or possible complications, and (c) evaluation of risks and benefits of continued pregnancy versus early delivery. Identification of the potential etiology is similar to preconception evaluation. The benefits of intervention are not well documented but management of medical conditions such as hypertension and diabetes, augmentation of nutritional status, and reduction or elimination of tobacco, alcohol, and illicit drug use are recommended.

Complications associated with IUGR include prenatal and perinatal complications such as increased risk of hypoxia, metabolic acidosis, preterm labor, pre-eclampsia, and surgical delivery. Neonatal complications include aspiration, apnea, intubation, sepsis, hypoglycemia, electrolyte abnormalities (especially hypocalcemia), seizure,

and death. Careful intrapartum monitoring is critical to early identification and management of these potential complications.

In patients with documented IUGR, careful assessment of fetal growth and well-being is critical to determining the timing and manner of delivery. Weekly prenatal visits are recommended, with careful monitoring of fetal status with review of fetal movement and kick counts. As noted, serial measurements (every 2–6 weeks) of fetal growth are important for the diagnosis as well as for tracking. Antenatal fetal monitoring may also include nonstress testing, contraction stress testing, and/or bio-physical profile measurements at least weekly. Amniotic fluid should be measured via AFI once weekly.

Management of IUGR requires individualized assessment and decision-making. Timing of delivery should balance the benefits of further fetal maturity against the risks of continued exposure to an intrauterine environment that is less than optimal. Evidence of fetal maturity (well-documented gestational age of 38 weeks' gestation or documentation of fetal lung maturity) or significant fetal stress should prompt delivery. For infants with reassuring antenatal monitoring and without evidence of maturity, the benefits of intrauterine development may outweigh the risks of early delivery. Delivery should be planned for a facility that is experienced with and capable of managing the high-risk infants of a pregnancy complicated by IUGR. Management of perinatal complications in the infant is best managed by an experienced neonatologist.

# Chapter 7

## Preterm Labor

### Contents

Background .....	56
Factors Associated with Preterm Labor .....	56
Preconception Factor.....	57
Postconception Factors .....	57
Diagnosis.....	58
Intake Assessment.....	58
History .....	58
Physical Examination.....	58
Laboratory Studies .....	59
Management.....	59
Management Prior to 34 Weeks' Gestation .....	60
Management at 34–37 Weeks .....	62
Assessment of Fetal Lung Maturity .....	62
Lecithin-to-Sphingomyelin Ratio .....	62
Phosphatidylglycerol.....	63
Additional Tests .....	63

### Key Points

1. Preterm labor is uterine contractions resulting in progressive cervical change prior to 37 weeks' gestation. Preterm delivery is delivery prior to 37 weeks' gestation; low-birth weight infants are those that weigh less than 2500 g at delivery.
2. Prior to 34 weeks' gestation, most patients should be considered for tocolysis; from 34 to 37 weeks' gestation such decisions must be made on a case-by-case basis.
3. Complications associated with preterm delivery include increased perinatal mortality and complications of prematurity (including respiratory distress, gastrointestinal dysfunction, hemorrhage, and abnormalities of growth and development).

## Background

Preterm labor is among the most common and most serious of prenatal complications. Preterm labor and its potential sequelae of preterm delivery and low-birth weight (LBW) infants remain one of the most significant challenges of current obstetrical practice. Preterm labor is defined as uterine contractions resulting in progressive cervical change prior to 37 weeks' gestation. Preterm delivery is delivery prior to 37 weeks' gestation. LBW infants are defined as those infants weighing less than 2500 g at delivery regardless of gestational age. LBW infants should be distinguished from small-for-gestational-age (SGA) infants who are defined as those infants below the fifth percentile for weight based on gestational age.

Preterm labor affects approximately 10 % of all pregnancies. Preterm delivery affects approximately 13 % of all live births. Preterm delivery and LBW infants represent approximately 70 % of all perinatal mortality (~25,000 deaths annually) and 50 % of all neurological morbidity.

## Factors Associated with Preterm Labor

A number of factors have been associated with an increased risk of preterm labor. These are summarized in Table 7.1. These factors can be divided into pre- and postconception factors. Although the mechanisms that link these factors to the onset of preterm labor is, in most instances, poorly understood, a thorough review of the patient's history will allow providers to more carefully outline the risk of preterm labor, preterm delivery, and LBW infants.

**Table 7.1** Risk factors for preterm labor

Preconception factors
Lower socioeconomic status
Anatomic abnormalities (e.g., septate/bicornuate uterus, cervical incompetence)
Prior uterine surgery
Myomata
Diethylstilbestrol exposure
Past history of preterm labor
Under 18 years old, over 40 years old
Possible genetic predisposition
Postconception factors
Tobacco
Cocaine
Infection (e.g., group B streptococcus, <i>N. gonorrhoea</i> , <i>C. trachomatis</i> , trichomonas, <i>gardnerella</i> , <i>ureaplasma</i> , <i>mycoplasma</i> )

Although preterm labor alone is not associated with perinatal complications, concomitant conditions and outcomes are. Preterm labor may be complicated by preterm premature rupture of membranes (PPROM). PPROM is associated with a variety of complications discussed in Chap. 8. Preterm labor may also result in preterm delivery. Prematurity, in turn, is potentially associated with pulmonary dysfunction, gastrointestinal abnormalities, neurological complications, abnormalities of growth and development, and a significant risk of perinatal mortality. Complications of preterm delivery are the leading cause of perinatal mortality, responsible for approximately two-thirds of all deaths.

### ***Preconception Factor***

#### **Environmental Factors**

A number of environmental factors have been associated with an increased risk of preterm labor. The most significant environmental factor associated with preterm labor is lower socioeconomic status.

#### **Patient-Related Factors**

A number of pre-existing patient conditions also contribute to the risk of preterm labor. Patients may have a pre-existing genetic risk, a congenital anomaly (e.g., septate/bicornuate uterus or cervical incompetence), a pre-existing acquired obstetrical/gynecological risk (e.g., myomata, uterine surgery, diethylstilbestrol exposure), or a past history of preterm labor or second trimester spontaneous abortions. The recurrence rate of preterm labor is approximately 25 %. Additionally, the risk of preterm labor is highest among younger (<18 years old) and older (>40 years old) obstetrical patients. These conditions can be screened for early in pregnancy (or during preconception counseling). Although many of these factors are not modifiable, their presence can usefully contribute to a conversation between the provider and the patient concerning the risk for preterm labor during the current pregnancy.

#### ***Postconception Factors***

Once conception occurs, a number of additional factors contribute to the risk of preterm labor. An increased risk for preterm labor is associated with tobacco and cocaine use. Infections such as group B streptococcus, gonorrhea, chlamydia, trichomonas, gardnerella, ureaplasma, and mycoplasma have all been associated with increased preterm labor risk. Such exposures should be screened for at the first prenatal history (either directly through testing or via history) and as appropriate throughout the course of pregnancy.

## Diagnosis

As noted earlier, the diagnosis of preterm labor consists of three components: gestational age less than 37 weeks, presence of uterine contractions, and progressive cervical change.

The diagnosis of preterm labor begins with confirmation of the gestational age of the fetus. All data that contributed to the estimated date of delivery (EDD) should be reviewed for accuracy. The patient's last menstrual period should be reviewed for accuracy. Additional data such as prenatal ultrasound, sequential fundal height measurements, and gestational age at quickening should also be reviewed. If no such data is available, an obstetrical ultrasound may be indicated. It should be emphasized, however, that an ultrasound obtained late in pregnancy has significantly less accuracy for purposes of gestational dating.

The patient should be questioned concerning the presence of contractions (although the absence of patient-reported contractions does not exclude the possibility of clinically significant contractile activity). If preterm labor is suspected, patients should be placed on tocometric monitoring to confirm the presence of uterine contractions.

Documentation of progressive cervical change, under most circumstances, requires serial cervical examinations. After confirming the absence of bleeding per vagina, providers should document cervical dilation and effacement as well as fetal station. Although the patient may demonstrate unequivocal cervical evidence of labor on initial examination, generally the diagnosis will require comparison of initial findings to findings on a follow-up examination.

## Intake Assessment

### *History*

In addition to the history noted earlier, patients should be asked about bleeding per vagina, rupture of membranes or fluid leak, and/or symptoms of infection. Special caution should be exercised if the patient reports a history of bleeding per vagina. The management of third-trimester bleeding is covered in Chap. 10. A review of the past history should note the presence of cardiac, renal pulmonary, and/or endocrine abnormalities.

### *Physical Examination*

In addition to the pelvic examination for assessment of cervical change, the intake physical examination should document blood pressure, pulse, temperature, rupture of membranes (see Chap. 8), fetal heart rate, and uterine contractions.

## ***Laboratory Studies***

Patients admitted with preterm labor should have all prenatal laboratory values reviewed with lab values ordered or updated as necessary. Patients may require testing for infection, including gonorrhea, chlamydia, group B strep trichomonas, and bacterial vaginosis. Other studies that may contribute to evaluation of possible infection include increased interleukin-6 in amniotic or cervical samples, elevated ferritin in cervical or serum samples, and elevated granulocyte colony-stimulating factor in serum samples. Patients demonstrating clinical signs or symptoms of other obstetrical conditions (e.g., pregnancy-induced hypertension) should have laboratory evaluation as indicated for those conditions.


Controversy exists concerning the role of routine fibronectin testing in the management of suspected preterm labor. Specimens should be obtained via sterile speculum examination. A swap is placed in the posterior fornix for 10 s. Care should be taken to avoid manipulation of the cervix or use of lubricant. Known rupture of membranes is a contraindication to fibronectin testing. After 22 weeks' gestation, a result greater than 50 ng/mL is associated with an increased risk for preterm delivery with a sensitivity of 70–90 % and a specificity of 70–85 %. The negative predictive value is approximately 99 %. A negative test is a strong predictor of no preterm labor in the week following the test. Fetal breathing movements and measurement of cervical length can be used in conjunction with or as an alternative to fibronectin testing. Fibronectin has been shown to have the highest sensitivity and fetal breathing movements the highest specificity for delivery within 7 days of testing. In patients with a cervical length >30 mm, risk of delivery within 7 days is less than 5 % and fibronectin does not add to diagnostic accuracy.

## **Management**

The management of preterm labor is often limited in efficacy and duration and few modifiable factors have been identified. A general outline of management is shown in Fig. 7.1. Goals of management are focused on delay of delivery to allow four key outcomes:

1. Transfer, if necessary, to a facility with appropriate neonatal intensive care services
2. Administration of corticosteroids to decrease postpartum complications for the infant
3. Administration of magnesium sulfate (associated with a decreased risk of cerebral palsy)
4. Administration of appropriate prophylaxis for group B strep

Decisions concerning appropriate management should be tailored to the individual patient. Despite the challenge and variability involved in managing preterm labor, a few general guidelines can be given.



**Fig. 7.1** Management of preterm labor

### ***Management Prior to 34 Weeks' Gestation***

In general, fetal lung maturity cannot be assumed in infants prior to 34 weeks' gestation. For this reason, tocolysis is generally recommended. Although the efficacy and duration of such therapy is limited, a brief delay in delivery allows for administration of corticosteroids to enhance fetal lung maturity. All patients should be

**Table 7.2** Contraindications to tocolysis

Medical contraindications
Cardiac disease
Renal insufficiency
Pyelonephritis
Pulmonary hypertension
Untreated diabetes mellitus
Electrolyte abnormalities
Obstetrical contraindications
Fetal stress
Chorioamnionitis
Eclampsia
Fetal demise
Hemodynamic instability

screened for contraindications to tocolysis (see Table 7.2). Contraindications to tocolysis include underlying medical contraindications (cardiac disease, renal insufficiency, pyelonephritis, pulmonary hypertension, untreated diabetes mellitus, and electrolyte abnormalities) and obstetrical contraindications (fetal stress, chorioamnionitis, eclampsia, fetal demise, and hemodynamic instability). Tocolytic options include the following:

1. Terbutaline ( $\beta$ -2-sympathomimetic): 250  $\mu$ g subcutaneously every 3–4 h.
2. Ritodrine ( $\beta$ -2-sympathomimetic): 100  $\mu$ g per minute intravenous starting dose. Dose increased 50  $\mu$ g per minute every 20 min until contractions cease.
3. Magnesium sulfate ( $MgSO_4$ ): 6 g intravenous load over 15 min then 2 g per hour. It may be increased every hour until contractions cease, maximum dose of 5 g per hour is reached, or signs or symptoms of magnesium toxicity occur. Magnesium toxicity may be noted as neurological depression, cardiac depression or arrest, tetany, and hypotension. Monitoring of all patients on magnesium should include serum magnesium levels, maternal deep tendon reflexes, blood pressure, and strict recording of fluid input and output. Urinary retention is associated with magnesium use. Magnesium levels above 7 mEq/L are associated with diminished deep tendon reflexes, above 10 mEq/L with respiratory depression, and above 12 mEq/L with cardiac depression and arrest. Magnesium toxicity is treated with calcium gluconate 1 g intravenous.
4. Indomethacin (nonsteroidal anti-inflammatory, prostaglandin inhibitor): 100 mg per rectum or 50 mg orally loading dose and 50 mg per rectum or 25 mg orally every 4–6 h. Prior to 32 weeks, indomethacin has been shown to have equal efficacy to  $\beta$ -2-sympathomimetic agents probably by inhibiting prostaglandin synthesis. Use of indomethacin is associated with oligohydramnios and premature closure of the ductus arteriosus. For this reason, all patients on indomethacin should undergo frequent (every other day) ultrasound examinations to monitor for oligohydramnios. Because of the concern for premature ductus arteriosus closure, the use of indomethacin after 32 weeks gestation is controversial.

5. Nifedipine (calcium channel blockade): 20 mg orally loading dose and 10 mg orally every 6 h. Nifedipine's action in blocking calcium channel function in the smooth muscle is postulated to explain its efficacy in reducing uterine contractility. Studies have demonstrated efficacy similar to  $\beta$ -2-sympathomimetic agents.

Patients should be admitted and placed on bed rest. If significant cervical dilation has occurred, patients may be placed in a head-down position. Routine management includes monitoring of fluid status, hemodynamic status, and fetal well-being. If there is question concerning the gestational age or fetal lung status, fetal lung maturity testing may be considered. Mothers of infants at risk for fetal lung immaturity (24–35 weeks' gestation) should be treated with 12 mg of betamethasone, intramuscularly. Two doses should be given 24 h apart. Patients with evidence of contributory infection should be treated as appropriate for the infection.

### ***Management at 34–37 Weeks***

Fetal lung maturity in this range is highly variable and decisions to initiate tocolysis must be individualized. When time permits, assessment of fetal lung maturity may assist in decisions concerning tocolysis versus expectant management.

### ***Assessment of Fetal Lung Maturity***

Delivery of an infant prior to fetal lung maturation is associated with considerable neonatal morbidity and mortality. For this reason, assessment of fetal lung maturity is critical in all instances where gestational age cannot be firmly established or when prenatal complications require consideration of an early delivery.

Confirmation of the gestational age is critical. Gestational age can be confirmed by review of the last menstrual period, early obstetrical ultrasound results, and key developmental milestones such as quickening and fetal heart tones. Although these data may allow for accurate gestational dating when available, not all data will be available in all cases. Even with such data, a more accurate assessment of fetal lung maturity may be necessary to guide management decisions. A variety of options are available to assist in this assessment.

### ***Lecithin-to-Sphingomyelin Ratio***

As fetal lung maturity progresses, pulmonary secretions are accumulated in the amniotic fluid allowing for assessment of fetal lung maturity based on amniotic fluid sampling. Lecithin and sphingomyelin are present in approximately equal quantities until approximately 8 weeks prior to the EDD. Beginning at this point,

lecithin concentrations increase and sphingomyelin concentrations remain stable. As the fetus nears maturation, therefore, the ratio of lecithin to sphingomyelin will increase. Although the exact interpretation of the results may be site dependent, a lecithin-to-sphingomyelin ratio of 2:1 or greater is associated with generally favorable neonatal pulmonary outcomes.

### ***Phosphatidylglycerol***

The presence of blood or meconium in the amniotic fluid may alter the results of the lecithin-to-sphingomyelin ratio. For this reason, alternative tests have been developed that are not sensitive to the presence of these substances. One such test is phosphatidylglycerol, a component of surfactant that is present in amniotic fluid and increases in quantity as fetal lung maturity advances. Amniotic fluid samples may be tested for phosphatidylglycerol alone or in conjunction with lecithin–sphingomyelin testing. The presence of phosphatidylglycerol is associated with more advanced fetal lung maturity and therefore with generally improved neonatal pulmonary outcomes. The results may be reported either qualitatively or quantitatively.

### ***Additional Tests***

Because of the importance of documenting fetal lung maturity and the complexity/limitations of traditionally available tests, a variety of alternative tests are now available. FLM-TDx II is a simple and reliable test (although it can be affected by blood or meconium). A value of  $>55$  mg/g of albumin is considered indicative of fetal lung maturity. Lamellar body count (affected by blood but not meconium) with a value  $>50,000$  is also indicative of fetal lung maturity.

# Chapter 8

## Premature Rupture of Membranes

### Contents

Background .....	65
Diagnosis.....	66
History .....	66
Physical Examination.....	67
Laboratory.....	68
Management.....	69
Management at Term .....	70
Preterm Management.....	70

### Key Points

1. Premature rupture of membranes (PROM) is defined as rupture prior to the onset of labor.
2. Preterm premature rupture of membranes (PPROM) is defined as PROM occurring prior to 37 weeks' gestation.
3. Rupture of membranes is followed by onset of labor within 24 h in 90 % of term patients and 50 % of preterm patients.
4. PROM is associated with an increased risk of ascending infection. This risk increases with duration of rupture.

### Background

For most women, the pattern of labor is predictable. In general, women first note the onset of contractions that are relatively mild and irregular. As labor progresses, the contractions become stronger and more regular and of increased duration. Spontaneous rupture of membranes generally follows the development of a regular contraction pattern as cervical dilation progresses. In approximately 10 % of cases,

**Table 8.1** Risk factors for PROM

Infection
Hydramnios
Incompetent cervix
Placental abruption
Amniocentesis

however, spontaneous rupture of membranes occurs prior to the onset of labor. This is defined as premature rupture of membranes (PROM).

Premature, in this case, does not refer to gestational age but to labor. If rupture of membranes precedes labor *and* is prior to 37 weeks' gestation, the condition is referred to as preterm premature rupture of membranes (PPROM). Rupture of membranes is generally followed by onset of labor within 24 h. Ninety percent of patients at term and 50 % of preterm patients will begin labor within 24 h of spontaneous rupture of membranes. For term patients, PROM will generally mark impending labor and management can be expectant. PPROM often marks the onset of preterm labor and patients should be managed appropriately for their degree of prematurity. PPROM complicates approximately 30 % of all preterm deliveries and is associated with significant morbidity and mortality. The most significant complication of PPROM is infection.

A number of factors have been associated with PROM including infectious, anatomic, and pregnancy-related factors (see Table 8.1). A considerable number of cases, however, are idiopathic. Trichomonas, bacterial vaginosis, urinary tract infection (UTI), gonorrhea, chlamydia, and group B strep are among the infectious agents known or suspected to be associated with PROM. Women with documented cervical incompetence are also at increased risk for PROM. Amniocentesis is associated with an increased risk of PROM. This risk may be, in part, related to the experience of the provider performing the procedure. For this reason, patients requiring amniocentesis should be referred to providers with considerable experience. Placental abruption is occasionally associated with PROM and should be considered in the evaluation of patients with PROM.

## Diagnosis

### History

Patients with PROM will often report discharge or “leaking” per vagina. This fluid leak may be subtle (e.g., increased wetness noted on undergarments or pants) or may be substantial (e.g., a “gush” of fluid). A careful history should be obtained to distinguish the causes of discharge such as cervical infection, physiological mucus production (or loss of the mucus plug), urinary incontinence, or UTI. Although each of these requires evaluation and diagnosis, management varies considerably from that for PROM.

Patients with PROM have, by definition, ruptured the membranes that serve to protect the infant from ascending infection. Patients with PROM are, therefore, at increased risk for perinatal transmission of genital infection and/or vaginal flora such as group B strep. This risk increases with the duration of rupture. For this reason, the best possible estimate of the time of rupture is an important part of the history. Because PROM represents the loss of protective membranes (and the subsequent risk of infection), every patient presenting with suggestive symptoms must be fully evaluated before eliminating PROM as a diagnosis.

Patients may also present with reported “urinary” symptoms such as urinary incontinence or urinary frequency. Such symptoms are both common and challenging. Anatomic changes associated with pregnancy such as increased uterine and fetal size increase urinary incontinence. Physiological changes associated with pregnancy such as relative outflow obstruction and urinary stasis increase the likelihood of urinary tract infection. For this reason, such urinary symptoms should be carefully detailed to identify other symptoms consistent with UTI such as urgency, dysuria, hematuria, abdominal pain, fever chills, nausea, vomiting, or back/costovertebral angle pain. As a general rule, evaluation of PROM should include assessment for UTI and evaluation of UTI should include assessment for PROM.

In addition to assessment for symptoms of urinary pathology, history should include information concerning bleeding per vagina and symptoms consistent with early or impending labor such as contractions, abdominal pain/cramps, back pain, or mucus plug loss. When symptoms that might be suggestive of more serious pathology are discovered, such as fever, bleeding, or severe pain, the history should be further expanded to include symptoms consistent with abruptio placenta, placenta previa, or infection/sepsis.

In addition to obtaining the history associated with the leakage of fluid, a brief review of the prenatal course is important. In particular, attention should be paid to prior infections, previous episodes of contractions or preterm labor, and prior episodes of bleeding or multiple gestations. Review of the gestational dating, the estimated date of delivery, and the data used to calculate these is also critical.

## ***Physical Examination***

If the history is consistent with possible PROM, providers should assume that the membranes have ruptured until this is ruled out. The primary concern under such circumstances is to minimize the possibility of ascending infection. Manual examinations should be minimized. Sterile speculum examination should be performed and assessment should include testing for common infectious agents.

Vital signs should be documented and should include temperature, pulse, and blood pressure. A note should be made of patient discomfort or pain. Abdominal examination should include documentation of fundal height, abdominal tenderness, and fetal position (via Leopold’s maneuvers; see Table 8.2). The external genitalia should be examined for evidence of infection (such as herpes), discharge, and

**Table 8.2** Leopold's maneuvers

First maneuver	Identify fetal head, fetal body, and fundal height
Second maneuver	Palpate fetal body to determine position of the back (smooth and uninterrupted) and front (palpable fetal arms and legs)
Third maneuver	Move body side to side. Resistance to movement suggests fetal engagement
Fourth maneuver	Identify cephalic prominence

trauma. On sterile speculum examination, note should be made of blood, fluid, or discharge in the vaginal vault or at the cervical os. Samples should be obtained for gonorrhea, chlamydia, group B strep, herpes (if characteristic or suspicious lesions are noted), yeast, and trichomonas/bacterial vaginosis.

Fetal heart rate should be documented including rate and variability. Uterine contractions should be noted on tocometer.

## ***Laboratory***


As noted previously, laboratory studies may include evaluation for gonorrhea, chlamydia, group B strep, herpes, trichomonas, and bacterial vaginosis. Although gonorrhea, chlamydia, group B strep, and herpes will not be immediately available, evaluation for yeast, trichomonas, and bacterial vaginosis can be performed quickly and accurately in the office. A microscopic examination of discharge/fluid may reveal clue cells (bacterial vaginosis), flagellated organisms (trichomonas), or fungal elements (yeast). A more comprehensive review of these conditions can be found in Chap. 13. A urine sample should be obtained for urinalysis, culture, and microscopic evaluation.

PROM is most accurately diagnosed via laboratory evidence of amniotic fluid in the vaginal vault. The two most common tests for rupture of membranes are nitrazine testing and “ferning.” A sample of amniotic fluid placed on nitrazine paper will turn the paper blue. False positives may occur in the presence of blood, semen, or infection. A sample of the fluid should also be spread thinly on a microscope slide and allowed to air-dry. The dried sample should show characteristic fern-shaped crystalline pattern on microscopic examination.

Occasionally, suspected PROM may represent other more serious prenatal complications such as pyelonephritis, abruption, or pre-eclampsia. If these conditions are suspected on the basis of history or physical examination, additional laboratory studies will be necessary. Evaluation for these conditions should be reviewed in the appropriate chapters for each.

## Management

The initial step in managing PROM is to distinguish PROM from PPROM. For this reason accurate gestational dating is critical. Figure 8.1 outlines the general management of PROM.



**Fig. 8.1** Management of PROM

## ***Management at Term***

For patients determined to be at term (>37 weeks' gestation), management is similar to that for term labor. As the majority of these patients will begin labor within 24 h, no additional intervention will be necessary for many. Because the risk of infection increases as the duration of rupture increases, induction of labor may be necessary at 12 h or with signs/symptoms of fetal stress or infection (tachycardia, fetal tachycardia, fever).

## ***Preterm Management***

Management of PPROM is considerably more complicated than for PROM. In addition to the considerations associated with the management of ruptured membranes, providers must also manage the impending preterm labor. (A complete discussion of preterm labor management can be found in Chap. 7.)

Patients with PPROM should be admitted with appropriate monitoring of fetal heart rate and uterine contractions. Baseline evaluation should include a history, physical examination, and laboratory studies, as noted previously.

Because delay in delivery is often desirable in preterm patients, particular attention should be paid to evaluation and management of possible infection. In addition to the studies previously mentioned, additional studies might include amniocentesis for culture and Gram stain, complete blood count, and blood and urine cultures. Patients should be closely monitored for uterine tenderness, fever, or discharge or bleeding per vagina.

Antibiotics are indicated for patients with evidence of infection. Suspected chorioamnionitis should be treated with an appropriate broad-spectrum regimen such as ampicillin and gentamicin (e.g., 2 g intravenous ampicillin every 4 h and 2 mg/kg of gentamicin loading dose then 1.5 mg/kg every 8 h). For patients with evidence of group B strep, penicillin or ampicillin is indicated (e.g., 2 g intravenous ampicillin loading dose and then 1 g intravenously every 4 h).

Of patients with PPROM, 50 % will begin labor within 24 h. For this reason, providers should consider the use of steroids to decrease neonatal pulmonary complications of immaturity. One such regimen is two 12 mg doses of betamethasone 24 h apart.

# Chapter 9

## Early Pregnancy Bleeding

### Contents

Background .....	72
General Approach to Early Pregnancy Bleeding per Vagina .....	72
History .....	74
Physical Examination.....	74
Ultrasound.....	75
Laboratory Studies.....	75
Ectopic Pregnancy.....	75
Background .....	75
Diagnosis .....	76
Management.....	77
Spontaneous Abortion .....	78
Background .....	78
Diagnosis .....	79
Management.....	80

### Key Points

1. Evaluation of early pregnancy bleeding should focus initially on identification of ectopic pregnancy and spontaneous abortion.
2. The primary concern of initial management is assessment of hemodynamic stability and possible peritonitis.
3. Management of ectopic pregnancy is designed to (a) minimize maternal morbidity and mortality, (b) remove the ectopic pregnancy, and (c) maximize potential future fertility.
4. Management of possible spontaneous abortion begins with ruling out the possibility of ectopic pregnancy.

## Background

Bleeding per vagina may occur at any point during the course of pregnancy and always warrants careful attention to identification and management of the underlying etiology. Bleeding that occurs late in pregnancy (generally within the third trimester) is covered in Chap. 10. Bleeding that occurs early in pregnancy (generally within the first trimester) is of particular concern because of concerns with pregnancy viability and possible ectopic pregnancy with its associated morbidity and mortality. A careful review of common etiologies combined with a careful history, directed physical examination, and selected diagnostic studies will allow the provider to identify the underlying cause and initiate appropriate management in a timely manner.

Early pregnancy bleeding per vagina is a relatively common presentation. Most cases of early bleeding are mild, self-limited, and of indeterminate etiology. Among the cases in which an etiology can be determined most are caused by either spontaneous abortion or ectopic pregnancy. Approximately 5–15 % of all pregnancies will end in a clinically recognized spontaneous abortion (involuntary expulsion prior to 20 weeks' gestation). A considerably larger percentage (up to one-third) will end in an unrecognized abortion that is perceived to be menstrual bleeding. Although less common than spontaneous abortion, ectopic pregnancy is responsible for approximately 15 % of all maternal deaths. Although the differential diagnosis for early pregnancy bleeding includes several disparate conditions (see Table 9.1), the focus of initial evaluation is on identification of these two conditions. Less common causes of early pregnancy bleeding include trophoblastic disease and cervical pathology such as ulceration, infection, or cytopathology.

## General Approach to Early Pregnancy Bleeding per Vagina

As noted, most cases of early pregnancy bleeding per vagina are of indeterminate cause. A significant minority are related to spontaneous abortion. A smaller but significant number are associated with ectopic pregnancy. Initial evaluation will be directed toward identification of patients with these two conditions, documentation of the viability of the pregnancy, assessment of the medical stability of the mother, and reassurance for patients without evidence of either condition. Figure 9.1 outlines a general approach to the evaluation and management of early pregnancy bleeding.

**Table 9.1** Conditions associated with early pregnancy bleeding

Ectopic pregnancy
Threatened abortion
Incomplete abortion
Complete abortion
Trophoblastic disease
Cervical polyps
Cervical ulceration
Cervical cytological abnormalities




Fig. 9.1 Evaluation and management of early pregnancy bleeding

## ***History***

A careful history should be obtained of the bleeding itself as well as associated symptoms. In regard to the bleeding itself, information should be obtained concerning the following:

1. Characteristics of the bleeding (e.g., is it scant or copious, bright red or brown, and clot-like?).
2. Onset of the bleeding (e.g., did it start abruptly or gradually?).
3. Duration of the bleeding (how long has the bleeding lasted?).
4. Intensity of the bleeding (how much bleeding has been noticed? It may be helpful to use as comparison to normal menstrual flow, although providers should note that there is considerable physiological variability in the quantity of normal menstrual flow).
5. Exacerbating factors (did anything appear to precipitate the bleeding or make the bleeding worse once it had started?).
6. Relieving factors (has anything made the bleeding diminish or stop?).

Particular attention should be paid to trauma, activity (including sexual activity), and associated symptoms such as pain, dizziness, or weakness. The patient's past history should be reviewed for risk factors such as smoking, multiple sexual partners, past history of pelvic inflammatory disease, or intrauterine device use. Prior gynecological surgery should be reviewed, including tubal ligation and uterine surgery (including myomectomy). Any prior diagnosis of anatomic abnormalities of the genital tract should be noted. The patient's obstetrical history should be reviewed for prior ectopic pregnancies and/or spontaneous abortions. The prenatal course should be reviewed for any high-risk conditions that may have been previously noted, including a family history of significant genetic abnormalities or significant toxic or infectious exposures. A note should also be made of the patient's gestational age and the means used to determine the estimated delivery date.

## ***Physical Examination***

Physical examination begins with a rapid assessment of the hemodynamic stability of the mother. Depending on the gestational age of the patient, assessment of fetal heart rate may also be indicated. The mother's blood pressure (BP) and pulse should be recorded including orthostatic measurements. (To measure orthostatic—positional—changes in BP and pulse, patients should first be placed in a recumbent position. BP and pulse are measured in the usual manner. The patient is then raised to a sitting position and the pulse and BP are measured a second time. A note is made of a significant drop in BP or a significant rise in pulse.) The general assessment of the patient should include obvious signs of distress or discomfort. Abdominal examination should include assessment of uterine size. A pelvic examination should be performed. A note should be made of blood (location, at the cervical os, pooled

in the vaginal vault; quantity: from scant discoloration to copious active bleeding; character, bright red, clotted, brown). Cervical status should be noted including dilation. The presence of other findings including discharge, amniotic fluid, or cervical lesions such as polyps or erosions should be documented. A bimanual examination should be performed to assess for adnexal mass or tenderness. The uterus should be assessed for size and tenderness.

### ***Ultrasound***

Ultrasound examination may be critical in the assessment of early pregnancy bleeding. Confirmation of an intrauterine pregnancy is possible as early as 5 weeks' gestation. In general, the presence of an intrauterine pregnancy significantly reduces *but does not eliminate* the possibility of ectopic pregnancy. Ultrasound examination is also diagnostic of trophoblastic disease. The absence of fetal heart activity may be indicative of a threatened or incomplete abortion. Transvaginal ultrasound is study of choice for early pregnancy assessment. The absence of an intrauterine gestational sac after 6 weeks from the last menstrual period is strongly suggestive of nonviable pregnancy (ectopic or aborted). Transabdominal ultrasound may be used to assess for intra-abdominal fluid including blood.

### ***Laboratory Studies***

Early pregnancy bleeding per vagina may represent a significant obstetrical emergency in the case of an ectopic pregnancy that has ruptured. Blood loss may be rapid and difficult to control. For this reason, laboratory studies are ordered to establish baseline hematological status and in anticipation of a potential need for transfusion. Studies should include complete blood count with platelets, blood type, and crossmatch. If not previously documented Rh status should be confirmed. The second key function of laboratory surveillance is to document the viability of the pregnancy. For this purpose, quantitative  $\beta$ -human chorionic gonadotropin ( $\beta$ -hCG) should be obtained.

## **Ectopic Pregnancy**

### ***Background***

Ectopic pregnancy is diagnosed when pregnancy implantation occurs in a location outside the uterine cavity. The exact location can be variable and may include within the fallopian tubes (98 %) or in an intraperitoneal/extruterine location. Because blood supply and/or space is limited outside the uterine cavity, these pregnancies are

capable of growing for a brief period but will eventually outgrow their space or blood supply. If the pregnancy implants in a location (such as the fallopian tube) where space is limited, rupture may occur with associated intraperitoneal bleeding.

Risk factors associated with an increased risk for ectopic pregnancy include prior history of ectopic pregnancy, prior history of pelvic inflammatory disease, prior pelvic surgery (including tubal ligation and reversal of tubal ligation), multiple sexual partners (probably related to an increased risk for unrecognized pelvic inflammatory disease and subsequent scarring), and smoking.

## ***Diagnosis***

### **History**

Patients who present with early pregnancy bleeding per vagina should be carefully screened for risk factors that increase the probability of ectopic pregnancy as noted previously. Patients with ectopic pregnancy may report bleeding, abdominal pain, and symptoms of peritonitis, or it may be relatively asymptomatic. The bleeding may be variable depending on the location of the pregnancy. Although copious bleeding should prompt evaluation for ectopic pregnancy, providers should maintain a high index of suspicion as the absence of clinically apparent bleeding does not eliminate the possibility of ectopic pregnancy. The onset of symptoms may be acute but is often subacute with gradual worsening of symptoms that may be initially quite mild.

### **Physical Examination**

The physical examination should include those elements noted previously to assess maternal and fetal well-being, including vital signs, abdominal examination, and fetal heart rate monitoring when appropriate. The physical examination may be quite variable. For some patients, the findings on physical examination are all normal. For others with rupture and significant intraperitoneal bleeding, findings may be indicative of acute peritonitis including diminished bowel sounds and a rigid, tender abdomen with rebound and guarding. A pelvic examination should be performed to assess for the presence of blood and/or products of conception, cervical status, uterine tenderness, adnexal tenderness, and adnexal mass.

### **Ultrasound Examination**

Although history and physical examination may contribute to the diagnosis of ectopic pregnancy, all patients with early pregnancy bleeding per vagina should be assumed to have ectopic pregnancy until it is ruled out. For this reason, all patients

should have ultrasound confirmation of the presence, viability, and location of the developing fetus. Ultrasound may demonstrate a gestational sac in the extrauterine space. Alternately, the ultrasound may show no evidence of a gestational sac despite elevated  $\beta$ -hCG levels. Transvaginal ultrasound should demonstrate intrauterine pregnancy when hCG levels reach approximately 6000 mIU/mL. Ectopic pregnancy should be strongly suspected if the ultrasound does not demonstrate either characteristic findings of trophoblastic disease or intrauterine pregnancy when  $\beta$ -hCG levels are greater than 2000 mIU/mL.

## **Laboratory Studies**

Supportive laboratory studies include serum progesterone levels and serial quantitative  $\beta$ -hCG. Progesterone levels are rarely less than 5 ng/mL in viable pregnancies. Conversely, less than 1 % of ectopic pregnancies will demonstrate serum progesterone levels of more than 25 ng/mL. Serum  $\beta$ -hCG may also add useful data. Ectopic pregnancy is rarely associated with  $\beta$ -hCG levels greater than 50,000 mIU/mL. If initial findings are equivocal, serial  $\beta$ -hCG levels are very helpful. In normal pregnancies, serum  $\beta$ -hCG doubles every 2–3 days. An increase in  $\beta$ -hCG less than 50 % in 48 h is considered abnormal and should warrant further evaluation.

## **Management**

Management of ectopic pregnancy depends considerably on the stability of the patient. Following diagnosis of ectopic pregnancy, patients should be assessed for clinical stability. Patients with hemodynamic instability or evidence of peritonitis should be considered surgical emergencies. For patients who are stable, evaluation for surgical or medical management may be performed. The purpose of management is threefold: (a) to minimize morbidity and mortality for the mother, (b) to remove the ectopic pregnancy, and (c) to maximize future potential fertility options.

In addition to the conditions just listed, indications for surgical intervention include anemia, gestational sac greater than 4 cm on ultrasound evaluation, pain of more than 24 h duration, suspected heterotopic pregnancy, or uncertainty of diagnosis requiring laparoscopic confirmation.

Indications for medical management include gestational sac smaller than 4 cm on ultrasound examination and hemodynamic stability.

## **Surgical Management**

Patients with clinical evidence of hemodynamic instability or peritonitis should be treated as a surgical emergency. Providers should assume that blood products and fluid resuscitation will be necessary and should be prepared to provide both. Patients

should be crossmatched for at least four units of blood and should have at least two large bore intravenous access sites established as quickly as possible. A variety of surgical options exist depending on the clinical presentation of the patient, the desire to preserve future fertility, and the experience of the operator. These options include salpingostomy, segmental resection, fimbrial expression, salpingectomy (in cases with significant bleeding, significant tubal damage, or recurrent ectopic pregnancies at the same location), and laparotomy (in cases with severe hemodynamic stability).

## **Medical Management**

The use of methotrexate has emerged as a medical alternative to surgical management for appropriate patients. The success rate with single-dose methotrexate regimens is 80–90 %. One such regimen is 50 mg/m<sup>2</sup> of methotrexate via intramuscular injection. Serum  $\beta$ -hCG levels are obtained on days 1, 4, and 7. A second dose is administered on day 7 if serum  $\beta$ -hCG has not demonstrated a significant (~15 %) decrease. Serial weekly serum  $\beta$ -hCG levels should be obtained until levels are less than 15 mIU/mL.

## **Spontaneous Abortion**

### ***Background***

Among cases of early pregnancy bleeding with a known etiology, spontaneous abortion is by far the most common. As noted, 5–15 % of all pregnancies will end with a clinically apparent spontaneous abortion; up to one-third of all pregnancies end in spontaneous abortion. Although generally spontaneous abortion is not as medically concerning as ectopic pregnancy, the significance of spontaneous abortion for the patient (and her family) will often be quite high. For this reason, appropriate diagnosis and management of spontaneous abortion is critical.

Risk factors for spontaneous abortion include genetic abnormalities, toxic exposures, structural abnormalities (such as atypical uterine anatomy), smoking, and toxic exposure. Most cases of spontaneous abortion occur without a clinically apparent risk factor.

Spontaneous abortion may be divided into three general subtypes: threatened, incomplete, and complete.

Threatened abortion is defined as bleeding per vagina before 20 weeks without passage of tissue or premature rupture of membranes. Threatened abortion may present with or without cervical dilation. Bleeding per vagina prior to 20 weeks' gestation in conjunction with cervical dilation is referred to as inevitable abortion.

Incomplete abortion is defined as bleeding and incomplete passage of the products of conception. This condition may be associated with hemodynamic instability in which case it should be considered an obstetrical emergency and should include all the usual elements associated with management of such unstable patients.

Complete abortion is defined as bleeding with complete passage of all products of conception. The distinction between incomplete and complete abortion is not always clear, and care should be exercised to confirm the passage of all products prior to making the diagnosis of complete abortion.

## ***Diagnosis***

It should be recognized that patients will not present with a chief complaint of spontaneous abortion; they will present with a complaint of bleeding per vagina. For this reason, the general history for these patients will mirror that of patients with ectopic pregnancy.

## **History**

History should include the elements described earlier, including a history of the bleeding, associated symptoms, a review of historical risk factors, and symptoms of hemodynamic instability. Particular attention should be paid to reports of passage of tissue or “clots.” A past history of spontaneous abortion should raise providers’ suspicion of abortion.

## **Physical Examination**

Physical examination will begin with assessment of the hemodynamic stability of the patient. Pelvic examination will assess cervical dilation, bleeding, and/or products of conception. To assess for other potential causes of bleeding, providers should make note of adnexal tenderness or mass, cervical lesions, discharge, or other abnormalities.

## **Ultrasound Examination**

The use of ultrasound in suspected abortion may provide limited but useful information. Ultrasound examination may demonstrate the presence or absence of a gestational sac, the presence or absence of fetal cardiac activity, and the evidence of retained products of conception.

## **Laboratory Studies**

Laboratory studies are generally limited for patients who are clinically stable. Serial serum  $\beta$ -hCG levels should be obtained to document continued viability (for threatened abortion) or appropriate decline in levels (incomplete or complete abortion). For patients who are clinically unstable, laboratory studies are the same as noted for ectopic pregnancy.

## ***Management***

### **Threatened Abortion**

Effective management of threatened abortion is quite limited. Although few proven interventions exist, recommendations include bed rest, abstention from intercourse, and close monitoring for persistent bleeding. As noted earlier, documentation of serum  $\beta$ -hCG levels may be helpful.

### **Incomplete Abortion**

By definition, patients with incomplete abortion no longer have a viable pregnancy and have not yet passed all of the products of conception. Management requires (a) assessment of hemodynamic stability, (b) removal of retained products of conception, and (c) documentation of declining serum  $\beta$ -hCG levels to confirm adequacy of treatment. For hemodynamically unstable patients, management includes fluids, blood, and rapid surgical intervention. The use of oxytocin (30–40 U/L of fluid) may help control bleeding. For stable patients, surgical evacuation should be performed and serial serum  $\beta$ -hCG levels obtained. Patients who are Rh negative should receive rhogam at the time of dilation and evacuation to prevent isoimmunization.

### **Complete Abortion**

Management of patients with confirmed complete abortion is relatively straightforward. No intervention is required and serial serum  $\beta$ -hCG levels should be sufficient to document adequacy of management. As previously noted, however, providers should exercise caution as the distinction between complete and incomplete abortions is not always clinically apparent.

# Chapter 10

## Late-Pregnancy Bleeding

### Contents

Background .....	82
General Approach to Late-Pregnancy Bleeding Per Vagina .....	82
History .....	83
Ultrasound.....	84
Physical Examination.....	84
Laboratory Studies.....	85
Placenta Previa.....	85
Background .....	85
History .....	85
Ultrasound Examination .....	86
Physical Examination.....	86
Laboratory Studies .....	86
Management.....	86
Abruption Placenta.....	87
Background .....	87
History .....	88
Ultrasound.....	88
Physical Examination.....	88
Laboratory Studies .....	88
Management.....	88

### Key Points

1. Initial assessment of late-pregnancy bleeding is designed to identify potential placenta previa and abruptio placentae.
2. No manual or speculum examination should be performed until placenta previa has been ruled out.
3. Placenta previa is an absolute contraindication to vaginal delivery.

4. Placenta previa and abruptio placentae are obstetrical emergencies and require rapid assessment and management.
5. Initial management of placenta previa and abruptio placentae is directed toward ensuring hemodynamic stability and safe delivery.

## Background

Bleeding per vagina may occur at any point during the course of pregnancy and always warrants careful attention to identification and management of the underlying etiology. Bleeding that occurs early in pregnancy (generally within the first trimester) is covered in Chap. 9. Bleeding that occurs later in pregnancy (generally within the third trimester) is of particular concern because of the potentially serious underlying etiologies and the possibility of significant morbidity that exists for both the mother and the infant. A careful review of common etiologies combined with a careful history, directed physical examination, and selected diagnostic studies will allow the provider to identify the underlying cause and initiate appropriate management in a timely manner.


Late-pregnancy bleeding per vagina is a relatively common presentation. Approximately 5 % of all pregnancies is complicated by such bleeding. Although the differential diagnosis includes several disparate conditions, placental abnormalities comprise the majority of such bleeding (see Table 10.1). Half of all third-trimester bleeding is caused by either placental abruption or placenta previa. Other causes include cervical cytopathology, polyps, and “bloody show.” Bloody show refers to limited bleeding per vagina just prior to or at the onset of labor. Such bleeding is a variant of normal, although patients may require evaluation for other causes of bleeding.

## General Approach to Late-Pregnancy Bleeding Per Vagina

Most cases of significant late-pregnancy bleeding per vagina are caused by placental abnormalities: placental abruption and placenta previa. The diagnosis and management of each is detailed here. When faced with a patient who presents with bleeding per vagina, however, a general approach to evaluation (outlined in Fig. 10.1) will allow for rapid evaluation and triage of those patients with such significant obstetrical problems from those with other less common presentations.

**Table 10.1** Conditions associated with late-pregnancy bleeding

Placenta previa
Placental abruption
Cervical cytopathology
Polyps
“Bloody show”



**Fig. 10.1** Management of late-pregnancy bleeding

## History

A careful history should be obtained of the bleeding itself as well as associated symptoms. In regard to the bleeding itself, information should be obtained concerning the following:

1. Characteristics of the bleeding (e.g., is it scant or copious, bright red or brown, and clot-like?).
2. Onset of the bleeding (did it start abruptly or gradually?).
3. Duration of the bleeding (how long has the bleeding lasted?).
4. Intensity of the bleeding (how much bleeding has been noticed? It may be helpful to use as comparison to normal menstrual flow although providers should note

that there is considerable physiological variability in the quantity of normal menstrual flow).

5. Exacerbating factors (did anything appear to precipitate the bleeding or make the bleeding worse once it had started?).
6. Relieving factors (has anything made the bleeding diminish or stop?).

Particular attention should be paid to trauma, activity (including sexual activity), and associated symptoms. Of particular importance are pain with bleeding and symptoms consistent with labor such as cramping, contractions, or back pain. In addition, the patient should be questioned concerning fetal movement and/or fetal kick counts. The prenatal course should be reviewed for any high-risk conditions that may have been previously noted. Note should also be made of the patient's gestational age and the means used to determine the estimated date of delivery. The patient's past obstetrical history, if any, should be reviewed for prior episodes of late-pregnancy bleeding.

## ***Ultrasound***

Evaluation of late trimester bleeding will often require a pelvic examination and cervical assessment. *It is critical that no manual or speculum examination be performed prior to confirming the location of the placenta.* In the case of placenta previa, such an examination may cause placental damage and acute severe hemorrhage leading to an obstetrical emergency. Ultrasound examination should be performed before any vaginal examination. Ultrasound may be used to identify the location of the placenta and specifically to rule out placenta previa. It should be noted that although placental abruption *may* be seen on ultrasound, ultrasound is generally ineffective for detecting this condition with a sensitivity of approximately 15 %.

## ***Physical Examination***

Physical examination begins with a rapid assessment of the hemodynamic stability of the mother and of the fetal well-being. The mother's blood pressure and pulse should be recorded. The patient should be placed on electronic monitoring to assess fetal heart rate and contractions, if any. The general assessment of the patient should include obvious signs of distress or discomfort. Abdominal examination should include assessment of fetal size and position. Following ultrasound examination to determine the placental position, a pelvic examination should be performed. Note should be made of blood (location, at the cervical os, pooled in the vaginal vault; quantity, from scant discoloration to copious active bleeding; character, bright red, clotted, brown). Cervical status should be noted, including dilation and effacement.

The presence of other findings, including discharge, amniotic fluid, or cervical lesions such as polyps or erosions, should be documented. The position of the fetus should be confirmed, including presenting part and station. The uterus should be assessed for size and tenderness.

### ***Laboratory Studies***

Late-pregnancy bleeding per vagina may represent a significant obstetrical emergency. Blood loss may be rapid and difficult to control. For this reason, laboratory studies are ordered to establish baseline hematological status and in anticipation of a potential need for transfusion. Studies should include complete blood count with platelets, international normalized ratio, partial thromboplastin time, fibrinogen, fibrin split products, blood type, and crossmatch.

## **Placenta Previa**

### ***Background***

Placenta previa may be diagnosed when the placenta is located in such a position as to partially or completely occlude the internal cervical os (and therefore to occlude the outflow tract of the birth canal). Total or complete placenta previa, as the name implies, represents occlusion of the entire cervical os. Partial placenta previa represents less than complete occlusion. A marginal placenta signifies a location adjacent to but not occluding the cervical os. Although these categories are useful guides, it should be noted that the distinction may not be entirely clear on ultrasound examination. For this reason, particular caution should be exercised whenever the location of the placenta appears to be near or over the cervical os.

Under such circumstances, cervical dilation and/or direct trauma (such as fetal descent or pelvic examination) may cause laceration of the placenta and subsequent bleeding. Because of the highly vascular nature of the placenta, such bleeding may be rapid, copious, and life threatening to both the mother and the fetus. Placental previa is estimated to occur in approximately 1 out of every 100 pregnancies.

### ***History***

Patients who present with late-pregnancy bleeding per vagina should be carefully screened for risk factors that increase the probability of placenta previa. These risk factors include prior dilation and curettage, myomectomy, cesarean section,

maternal age (prevalence increases with increasing maternal age), grand multiparity, and multiple gestation. The primary symptom associated with placenta previa is bleeding per vagina. The onset is usually acute, may be associated with trauma or physical manipulation such as examination, is often continuous, and may be variable but can be significant. It is classically not associated with pain (although the presence of pain *does not* rule out the possibility of placenta previa).

### ***Ultrasound Examination***

Although these historical factors may contribute to the diagnosis of placenta previa, all patients with late-pregnancy bleeding per vagina should be assumed to have placenta previa until it is ruled out. For this reason, all patients should have ultrasound confirmation of the location of the placenta prior to manual or speculum examination. If prior ultrasound examinations have demonstrated a placental location away from the cervical os, this is sufficient to rule out placenta previa as the placenta may migrate away from the cervical os but does not migrate toward the cervical os.

### ***Physical Examination***

The physical examination should include those elements noted previously to assess maternal and fetal well-being, including vital signs, abdominal examination, fetal heart rate monitoring, and monitoring for contractions. For patients in whom the ultrasound confirms placenta previa, no manual or speculum examination should be performed.

### ***Laboratory Studies***

Laboratory studies are the same as those noted previously.

### ***Management***

Providers should assume that blood products and fluid resuscitation will be necessary and be prepared to provide both. Patients with confirmed placenta previa should be crossmatched for at least four units of blood and should have at least two large bore intravenous access sites established as quickly as possible. If the patient is actively contracting, tocolysis may be indicated to lessen the risk of further placental damage and bleeding.

## Management at Term

For patients who have reached at least 36 weeks' gestation, cesarean section is indicated. Placenta previa is an absolute contraindication to vaginal delivery. If the patient is actively contracting and the cesarean section cannot be performed immediately, tocolysis is indicated. Delivery of the infant and the placenta is the definitive treatment for placenta previa. Patients should be monitored for hemodynamic stability with blood and/or fluids administered to maintain adequate blood pressure and to replete circulating blood volume.

## Preterm Management

Placenta previa with active bleeding per vagina is an obstetrical emergency. If the mother or infant are unstable and/or the bleeding cannot be controlled, cesarean section is indicated. Management of maternal hemodynamic status is the same as at term. Neonatal specialists should be present at the delivery to manage the newborn.

For preterm patients who are stable and not bleeding, care should be individualized. Such care might include tocolysis, assessment of fetal lung maturity, and preparation for cesarean section.

## Abruptio Placenta

### *Background*

The maintenance of fetal perfusion, oxygenation, and nutrition requires the maintenance of the maternal–fetal placental unit. The interruption of placental function secondary to separation of the placental from the uterine wall is referred to as abruptio placenta or placental abruption. As with placenta previa, the severity of the condition may vary from minor separation with minimal signs or symptoms to complete abruption with significant compromise of maternal and/or fetal well-being. Although the variability in presentation makes exact prevalence difficult to calculate, it is estimated that 1 out of 100 pregnancies are affected by significant placental abruption.

The classic presentation for placental abruption is abrupt onset painful bleeding with associated uterine contractions. However, the variability in severity as well as location of the abruption makes such “classic” presentations rather uncommon. Up to 80 % of all cases of placental abruption present with no clinically apparent bleeding. The severity of the condition cannot be reliably predicted based on the quantity of clinically apparent bleeding. Patients with known or suspected placental abruption should be assumed to have bled significantly until proven otherwise. Because of the variability in presentation, providers should have a low threshold for suspecting placental abruption and manage such patients diligently.

## ***History***

As noted, the classic presentation for placental abruption is late-pregnancy bleeding per vagina. The bleeding is associated with abdominal or back pain, acute in onset, and variable in quantity (or perhaps absent). Patients will often report the acute onset (or an acute increase in severity) of uterine contractions.

## ***Ultrasound***

As with all patients with late-pregnancy bleeding per vagina, no pelvic examination should be performed until ultrasound confirmation of placental location has been obtained. Although ultrasound has a very low sensitivity for detection of placental abruption, approximately 15 % of cases are ultrasonographically apparent.

## ***Physical Examination***

In general, the physical examination does not contribute significantly to the diagnosis of placental abruption. Suggestive findings include hemodynamic instability, evidence of fetal stress on fetal heart rate monitoring, firm or tender uterus, and/or back tenderness. Although neither sensitive nor specific, the finding of a very firm uterus that does not relax (or hypertonic uterus on tocometer) should prompt immediate consideration of the possibility of placental abruption. All patients with suspected abruption should be placed on fetal heart rate monitoring, tocometer, and hemodynamic monitoring, including blood pressure and pulse.

## ***Laboratory Studies***

Laboratory studies are as noted in the previous sections.

## ***Management***

Appropriate management depends on a clinical assessment of the severity of the condition. As noted, however, the physical findings associated with abruption may be highly variable. When doubt exists concerning the severity of the condition, it is advisable to make preparations for management of severe abruption with significant maternal or fetal compromise.

## **Management of Mild Abruptio**

If the bleeding is mild or absent and both mother and fetus are clinically stable, management can be expectant. Patients should be placed on continuous hemodynamic, fetal heart rate and uterine contraction monitoring. Evidence of stability includes normal maternal blood pressure and pulse and minimal symptoms. Fetal heart rate monitoring should demonstrate normal heart rate with adequate variability. Uterine contraction pattern should be monitored for signs of hypertonicity or tetanus. If the patient is preterm, tocolysis may be indicated to facilitate fetal lung maturity. Maternal coagulation status should also be monitored. Because of the possibility of worsening abruption and hemodynamic instability, all patients should have at least two large bore intravenous access sites and should have adequate crossmatched blood available.

## **Management of Moderate or Severe Abruptio**

For patients with more significant clinical abruption, management must focus on two primary concerns: management of the hemodynamic status of the mother and rapid delivery of the infant. Maternal shock is a possible sequela of severe placental abruption. These patients will require emergent management with particular attention to the “ABCs” (airway, breathing, circulation) as dictated by the clinical presentation. Patients may require fluid infusion and/or blood product transfusion that may include fresh frozen plasma, cryoprecipitate, or platelets.

A clinical decision must be made concerning the instability of the mother and the likelihood of delivery. In less severe cases, induction of labor may be appropriate. In more severe cases, cesarean section may be indicated for more rapid delivery of the infant. Neonatology support should be present at the time of delivery for management of the newborn.

# Chapter 11

## Recurrent Pregnancy Loss

### Contents

Background .....	91
Diagnosis.....	93
Management.....	93
History .....	93
Physical Examination.....	94
Diagnostic Studies .....	94

### Key Points

1. Spontaneous abortion is defined as delivery prior to 20 weeks' gestation.
2. Recurrent pregnancy loss is defined as three or more spontaneous abortions prior to 20 weeks' gestation.

### Background

Spontaneous abortion is a common outcome of pregnancy, especially in the first trimester of pregnancy. Defined as delivery prior to 20 weeks of gestation, spontaneous abortion is often an unexpected and traumatic experience for the patient and her family. Spontaneous abortion is discussed in detail in Chap. 9. Briefly, abortion can be considered to fall into one of two categories: complete abortion, which is defined as delivery of all products of conception prior to 20 weeks' gestation, or incomplete abortion, defined as delivery of some but not all products of conception prior to 20 weeks' gestation.

In addition to these two categories, first-trimester bleeding can also indicate a possible impending abortion. Uterine bleeding prior to 20 weeks' gestation without delivery of any products of conception is considered a threatened abortion. Uterine bleeding prior to 20 weeks with cervical dilation but without delivery of products of conception is referred to as an inevitable abortion.

Although spontaneous abortion is common, recurrent pregnancy loss (three or more consecutive abortions prior to 20 weeks' delivery) is considerably less common, affecting approximately 1 % of all women of childbearing age. In addition to the evaluation and management considerations that pertain to a single episode of spontaneous abortion, recurrent spontaneous abortions require further evaluation and more significant management considerations (see Table 11.1).

Prior spontaneous abortion increases the risk for subsequent abortion, although the degree of risk is not well quantified. As noted in the previous chapter, spontaneous abortion, often not clinically diagnosed, occurs in up to one-third of all pregnancies. Among patients with a history of spontaneous abortion, the risk of spontaneous abortion rises to approximately 30 % after two prior spontaneous abortions and to approximately 50 % after three prior episodes. Although that makes recurrence a frequent event, it should be emphasized that it is by no means inevitable.

**Table 11.1** Risk factors for recurrent pregnancy loss

Exogenous exposures
Alcohol
Tobacco
Cocaine
Obstetrical/gynecological abnormalities
Uterine abnormalities
Leiomyomas
Genetic uterine anatomic abnormalities (bicornuate, septate)
Cervical incompetence
Diethylstilbestrol exposure
Asherman's syndrome
Infection (gonorrhea, chlamydia, syphilis, listeria, mycoplasma, ureaplasma, toxoplasma)
Progesterone deficiency
Genetic abnormalities
Chronic medical conditions
Thyroid disease (hypothyroid and hyperthyroid)
Renal disease
Uncontrolled diabetes mellitus
Collagen vascular disease
Uncontrolled hypertension
Antiphospholipid antibodies (anticardiolipin antibody, lupus anticoagulant)

## Diagnosis

The diagnosis of recurrent spontaneous abortion is made clinically in patients with an appropriate history. Although additional evaluation may be appropriate for evaluation and management purposes, no additional studies are required to make the diagnosis.

## Management

Management of recurrent spontaneous abortion begins with a careful history, a directed physical examination, and diagnostic studies to elucidate an etiology. Although not all cases of recurrent spontaneous abortion will have an identified cause, identification of such a cause when present may contribute to successful management. In general, identified causes will fall into the categories of exogenous exposures (such as medication, tobacco, alcohol, illicit drugs, occupational exposures), obstetrical or gynecological abnormalities, genetic abnormalities, or chronic medical conditions (such as renal disease, diabetes). In general, outcomes for patients with recurrent abortions are good, with at least half of such patients eventually carrying a pregnancy to term.

## History

A careful history should be obtained. The history should focus on likely underlying causes. As would be expected, the history begins with a focus on prior obstetrical and gynecological conditions. All previous pregnancies should be reviewed for complications and outcomes. Abnormal menstrual cycles may be indicative of hormonal dysregulation or previously unrecognized spontaneous abortions. Gynecological history should focus on fertility, especially hypofertility, and any known anatomic abnormalities. Gynecological surgeries, especially those involving the cervix, should be reviewed. A past history of sexually transmitted disease (STD) should be obtained as well as a history of any current symptoms consistent with STD.

The patient's exposure to exogenous agents should include medications, occupational exposures, and tobacco, alcohol, or illicit drug use. All medications, including over-the-counter, prescription, and complementary/alternative modalities, should be reviewed. If the provider is unfamiliar with any of the medications, referral to a maternal–fetal medicine specialist might be considered. The patient's work history should be reviewed for possible occupational exposures. Although less common as a source of exogenous toxins, the patient's hobbies and recreational activities should also be reviewed.

A review of the patient's family history may be helpful in at least two regards. A careful genogram for possible genetic abnormalities should be obtained. A genetics counselor may provide assistance with obtaining this history and the genogram.

Additionally, the family history may reveal a familial predisposition to a variety of medical conditions for which the patient is at risk. Hypertension, diabetes, renal disease, and coagulopathies, among other disease processes, may be familial and may contribute to an increased risk for spontaneous.

## ***Physical Examination***

The physical examination is often directed by the findings of the history and is more limited in scope. The patient's vital signs, particularly her blood pressure, should be measured. Examination of the head/neck should include palpation of the thyroid for enlargement and/or nodularity. Additionally, examination of the eyes may reveal evidence of hypertensive or diabetic retinal changes. Examination of the skin should be performed to detect the physical stigmata of endocrine disorders such as striae or changes in skin texture, consistency, or color. Cardiovascular examination should include documentation of murmurs, if any (especially renal bruits), and peripheral pulses. Neurological examination should include documentation of the peripheral sensory function.

Although unlikely to demonstrate any abnormalities, a pelvic examination should be performed to document normal anatomy and to obtain cervical samples for gonococcus and chlamydia.

## ***Diagnostic Studies***

Diagnostic studies will be directed toward the common underlying etiologies for recurrent spontaneous abortion. If genetic causes are suspected, these studies include karyotyping of both parents. Routine laboratory studies include cervical sampling for gonorrhea/chlamydia, progesterone, thyroid-stimulating hormone, thyroxine, lupus anticoagulant, and anticardiolipin antibody. If diabetes is suspected on the basis of family history or patient symptoms, a fasting blood sugar may help to clarify the diagnosis. Renal function may be assessed via routine blood serum electrolyte testing and urinalysis. Under limited circumstances, a serum drug screen may be indicated, although this diagnosis is generally made by history rather than by laboratory studies. If anatomic abnormalities are suspected, evaluation of the lower reproductive tract via radiological (hysterosalpingogram, pelvic ultrasound) or endoscopic (hysteroscopy) means is indicated.

Management of subsequent pregnancies will depend in large measure on the findings of this diagnostic evaluation. Although several of the identified conditions are not directly correctable, others are. Treatment of infection, management of hypertension, treatment of thyroid disease, and elimination or reduction of toxic exposures may all enhance the likelihood of a successful pregnancy. Many patients with recurrent pregnancy loss will be managed in conjunction with a provider who specializes in this area of obstetrics.

# Chapter 12

## Rh Isoimmunization

### Contents

Background .....	95
Fetal Consequences of Isoimmunization .....	96
Newborn Consequences of Isoimmunization .....	96
Diagnosis.....	97
History .....	97
Physical Examination.....	97
Diagnostic Studies .....	97
Management Prior to Isoimmunization .....	97
Management of Pregnancies with Rh-Sensitized Mothers .....	99
Amniotic Fluid Assessment .....	99
Ultrasonography.....	100
Percutaneous Umbilical Blood Sampling .....	100

### Key Points

1. Rh isoimmunization represents a maternal antibody response to immunologically incompatible fetal blood.
2. Rh isoimmunization is a preventable outcome of maternal–fetal Rh incompatibility.
3. Fetal blood should be considered Rh positive unless documented otherwise.

### Background

Transfer of nutrients, proteins, and antibodies between the mother and fetus generally occurs across the placenta without direct transfer of blood. Although the maternal and fetal blood supplies are separated from each other, they are, of necessity, in relative proximity and can, under certain circumstances, come into contact with each other. Under these circumstances, exposure to foreign proteins may elicit an immune response with potential health effects for either the mother or fetus.

Rh isoimmunization represents a serious maternal–fetal complication. All red blood cells are produced with a variety of surface antibodies that serve to identify one's own red cells from those of others. The presence of red blood cells from an outside source will elicit an immune response with subsequent hemolysis/removal of the foreign cells. This response may be relatively minor or may represent a significant medical complication.

The primary surface proteins serve to identify blood within the ABO blood type categories. Exposure to immunologically incompatible blood types within the ABO category will result in significant hematological complications. A second set of proteins (rhesus factor) also provide significant identification. Rh-negative ( $[Rh^-]$  those without Rh factor) patients exposed to blood containing Rh factor (Rh positive  $[Rh^+]$ ) will produce an antibody response to that blood. It is this incompatibility that results in the complications of Rh isoimmunization. An initial exposure will lead to development of antibodies. Subsequent exposures will yield a significant antibody response and an associated reaction.

Pregnant Rh-negative patients exposed to fetal Rh-positive blood may also develop such an antibody response (approximately 20 % will isoimmunize if not treated). Because such antibodies can cross the placenta and reach the fetus, all subsequent pregnancies that involve Rh-positive fetuses will potentially result in hemolysis and in complications for the fetus.

Rh factor is a genetically inherited trait with an autosomal-dominant inheritance pattern. An Rh-negative mother is, by definition, homozygous ( $Rh^-/Rh^-$ ). An Rh-positive father may be homozygous ( $Rh^+/Rh^+$ ) or heterozygous ( $Rh^+/Rh^-$ ). As all children will receive one Rh-negative gene from such mothers, the Rh status of the child is dependent on the Rh status of the father. All children of homozygous Rh-positive fathers will be Rh positive. One-half of all children of heterozygous fathers will be Rh positive.

## Fetal Consequences of Isoimmunization

Although maternal blood does not generally cross the placenta, maternal antibodies do. The placental transmission of maternal antibodies to Rh factor will result in fetal hemolysis and subsequent anemia. In addition, the red blood cell destruction results in the release of heme and bilirubin. These breakdown products are cleared by the maternal circulation after crossing the placenta. The results for the fetus are generally related to complications of severe anemia (erythroblastosis fetalis) and may include heart failure, acute pericardial effusion, hypoxia, acidosis, and death.

## Newborn Consequences of Isoimmunization

Following delivery, Rh-isoimmunized infants lose access to the maternal circulation but will continue to have circulating maternal antibodies. Under these circumstances, the neonate may be incapable of adequately clearing the bilirubin and heme that

result from continued red blood cell destruction. The potential complications for infants will include those anemia, cardiac complications, acidosis, and death, as well as deposition of heme within the basal ganglia of the developing brain (kernicterus).

## Diagnosis

### *History*

The Rh status of the infant is usually unknown; therefore, particular care should be taken to note the Rh status of the mother. When known, the Rh status of the father should also be noted. Prior pregnancies should be recorded, including the Rh status of the resulting infant. If the patient has had prior pregnancies, any administration of Rh immunoglobulin (discussed later) should be recorded. As noted previously, the initial exposure is less significant than subsequent exposures. For this reason, the history is particularly important in second and subsequent pregnancies.

### *Physical Examination*

In general, the physical examination will not contribute to the diagnosis of Rh isoimmunization, but careful monitoring of fetal growth and development is critical in the management of pregnancies complicated by Rh isoimmunization.


### *Diagnostic Studies*

Rh factor should be documented for all pregnant patients. For those patients found to be Rh negative, paternal testing may be indicated. If paternity is uncertain or paternal testing cannot be performed, the infants should be assumed to be Rh positive and management should proceed accordingly.

## Management Prior to Isoimmunization

Overall management of Rh- patients is outlined in Fig. 12.1. The management of pregnancies with the potential for isoimmunization consists of the following three key principles:

1. Minimize the potential for maternal-fetal transfusion (see Table 12.1).
2. Administer Rh immunoglobulin if maternal-fetal transfusion is known or suspected.



**Fig. 12.1** Evaluation and management of Rh- patient

**Table 12.1** Potential sources of maternal-fetal transfusion

Spontaneous or induced abortion
Amniocentesis
Chorionic villus sampling
Hemorrhage
External cephalic version
Trauma

3. Because all pregnancies are potentially complicated by minor maternal–fetal transfusions not recognizable clinically, all Rh-negative patients will receive immunoglobulin at about 28 weeks' gestation.

At 28 weeks' gestation, Rh-negative patients should be tested for evidence of Rh antibodies. If no antibodies are detected, patients should receive 300 µg of Rh immunoglobulin as a single intramuscular dose. Following delivery, the Rh status of the infant should be determined. If the infant is found to be Rh positive, a second dose of Rh immunoglobulin should be administered within the first 72 h postpartum.

Under all circumstances where maternal–fetal transfusion is possible or suspected, patients should receive Rh immunoglobulin. In general, a single dose is sufficient for most exposures. If a larger-than-usual transfusion is suspected, further testing should be performed to determine the extent of exposure and additional doses of immunoglobulin may be necessary. This occurs in less than 1 out of 200 cases.

## **Management of Pregnancies with Rh-Sensitized Mothers**

Management of the first pregnancy associated with isoimmunization is generally uncomplicated and will follow the guidelines for usual pregnancy management. Infants may require additional monitoring postpartum but can be expected to follow the usual newborn course. All Rh-sensitized mothers, however, should be informed of the development, the nature of the problem, and its potential effects in subsequent pregnancies. Critical discussions of contraception, preconception counseling, and future pregnancy planning should begin in the immediate postpartum period.

Patients known to be previously Rh sensitized require careful management and involvement of a skilled maternal–fetal medicine provider. Although a full discussion of the management of such patients is beyond the scope of this text, the general course of management is reviewed here.

For previously sensitized patients, antibody titers should be drawn at intake, at 20 weeks' gestation, and every 4 weeks thereafter. If antibody titers remain below 1:8, careful antenatal monitoring and continued routine prenatal care are recommended. When titers rise above a critical threshold (1:8–1:32), management will proceed as described below:

### ***Amniotic Fluid Assessment***

Beginning in the second trimester, bilirubin levels in the amniotic fluid decrease. In patients with significant isoimmunization, levels of bilirubin will rise as the pregnancy progresses. Serial measurement of bilirubin in the amniotic fluid can be used to determine the presence and severity of rising bilirubin levels. Fetal involvement is graded as mild, moderate, or severe based on the results of these studies. Mild involvement will require amniocentesis every 2–3 weeks with delivery at term in the

absence of a worsening of the fetal condition. Moderate involvement requires amniocentesis every 1–2 weeks with delivery when fetal lung maturity can be demonstrated (see Chap. 7). Severe disease requires weekly amniocentesis and may also require intrauterine transfusion. In all cases assessment of fetal well-being is necessary (see Chap. 17).

### ***Ultrasonography***

With advances in fetal ultrasonography, assessment of blood flow in the middle cerebral artery now allows for noninvasive assessment of fetal status. The decrease in blood viscosity (due to red blood cell lysis) results in higher velocity blood flow. Normalized values correlated with fetal hematocrit allow for identification affected fetuses with moderate or severe anemia.

### ***Percutaneous Umbilical Blood Sampling***

Under circumstances requiring direct measurement of fetal hematocrit, sampling of the umbilical vein allows for such measurement. Normal fetal hematocrit is approximately 40 % (36–44 %) but may be <30 % in severe anemia.

# Chapter 13

## Infection in Pregnancy

### Contents

Background .....	102
Symptoms of Infection in Pregnancy .....	102
Maternal Infection .....	104
Fetal Infection .....	104
Antibiotic Use in Pregnancy .....	105
Sulfonamides.....	105
Fluoroquinolones .....	105
Aminoglycosides.....	106
Vaginitis/Vaginosis.....	106
Background .....	106
History .....	106
Physical Examination.....	107
Laboratory Studies .....	107
pH Testing .....	107
KOH “Whiff” .....	107
Microscopic Examination .....	107
Treatment .....	108
Bacterial Vaginosis.....	108
Trichomoniasis.....	108
Yeast Vaginitis.....	108
Gonorrhea/Chlamydia.....	109
Urinary Tract Infections .....	109
Background .....	109
Diagnosis .....	110
History .....	110
Physical Examination.....	110
Laboratory Examination .....	110
Treatment .....	111
Group B Strep .....	111
Background .....	111
Diagnosis .....	112
Universal Screening .....	112
Treatment .....	112

## Key Points

1. Infectious complications in pregnancy include maternal or fetal morbidity, teratogenic or developmental abnormalities, and disruption of the course of pregnancy.
2. Management of infection during pregnancy begins with thorough preconception evaluation.
3. Routine screening of all prenatal patients for common asymptomatic infections is a cornerstone of prenatal care.
4. Women should be screened by history for signs or symptoms that are suggestive of common infectious complications of pregnancy at each prenatal visit.

## Background

Infection presents a particular challenge in pregnancy. Symptoms of infection may be subtle, unusual in presentation, or masked by other pregnancy-related symptoms. Infection may affect both the mother and fetus; antibiotic choices may be limited by concerns related to pregnancy and fetal development. As with most elements of obstetrical care, management of infection in pregnancy ideally begins in the preconception period. The preconception history can provide important data concerning infection risk such as recent exposures (toxoplasmosis, tuberculosis, cytomegalovirus), immunization status (influenza, measles, and varicella), and chronic infectious conditions (HIV, herpes, viral hepatitis). For some of these infections, appropriate preconception planning and risk reduction is the only effective intervention. A thorough physical examination may reveal evidence of such infections as active herpes, gonorrhea, or chlamydia. Preconception screening for HIV, tuberculosis, and syphilis, among other conditions, may allow for treatment prior to pregnancy, thereby decreasing the likelihood of adverse prenatal outcomes.

Infections occurring during the course of pregnancy may have two significant consequences. Direct maternal-fetal transmission may lead to infection not just in the mother but also in the fetus. Such infections may have very serious consequences for the fetus or newborn. Herpetic infection in newborns, for example, is associated with a 50 % mortality rate and significant morbidity among those infants who survive. In addition, some infectious agents are known to cause complications in the course of pregnancy, particularly preterm labor. Although most studies have shown conflicting data concerning the demonstrated benefit of diagnosis and treatment, most authorities recommend screening and treatment.

## Symptoms of Infection in Pregnancy

Although diagnosis of infection in pregnancy is critically important, symptoms of infection in prenatal patients may be absent or difficult to detect or interpret. Many infectious etiologies are asymptomatic. Syphilis, latent-phase herpes, HIV, and

others may be present without producing any symptoms in the pregnant patient. For these conditions, routine screening is recommended for all pregnant patients by history, physical examination, and/or diagnostic studies.

Some disease processes produce characteristic symptoms, which should be further evaluated whenever present. Such symptoms as fever, chills, bleeding per vagina, severe abdominal pain, or dysuria should lead to prompt evaluation. For other disease processes, the symptoms associated with infection may be difficult to interpret or may be masked by some of the common symptoms associated with pregnancy. Cervical mucus production may be increased during pregnancy and some women may interpret this increased mucus as infectious in cause. By contrast, some women may interpret an abnormal vaginal discharge as “normal” for pregnancy, delaying evaluation or failing to mention such symptoms during the course of prenatal care. Nonspecific abdominal pain is also a relatively common symptom occurring during pregnancy. Providers should be diligent in eliciting suggestive symptoms and should maintain a low threshold for following up such symptoms with appropriate diagnostic tests. As a general rule, women should be screened by history for symptoms that are suggestive of common infectious complications of pregnancy at each prenatal visit. Those patients who report such symptoms should be further evaluated for the possible presence of infection (Table 13.1).

**Table 13.1** Antibiotics in pregnancy

Infection	Antibiotics
Gram positive	Penicillins
	First-generation cephalosporins
	Clindamycin
Gram negative	Aminoglycosides ( <i>see text</i> )
	Third-generation cephalosporins
Anaerobic	Clindamycin
<i>Special conditions</i>	
Urinary tract infection	Cephalosporins
	Nitrofurantoin
	Ampicillin ( <i>see text</i> )
	Sulfisoxazole ( <i>see text</i> )
Chlamydia	Erythromycin
	Azithromycin
Gonorrhea	Ceftriaxone
Syphilis	Penicillin
Yeast	Azole antifungal agents after the first trimester
Bacterial vaginosis	Clindamycin
	Metronidazole after the first trimester
Trichomoniasis	Metronidazole after the first trimester
Herpes	Acyclovir; cesarean section indicated if present at time of labor
HIV	<i>See Chap. 16</i>
Group B strep	Ampicillin in labor

## Maternal Infection

The primary concern in pregnancy is with the health and welfare of the mother. Timely diagnosis and treatment of infection has direct health consequences for the mother and should be approached in the same manner as in nonpregnant patients. Routine prenatal care includes screening for gonorrhea, chlamydia, syphilis, HIV, hepatitis B, and urine culture for bacteriuria. These tests are performed in all patients regardless of symptoms. It is worth noting that most of these infections are sexually transmitted. Pregnancy can be taken as good evidence of unprotected sexual activity and therefore confers the infection risk associated with unprotected sex. Prenatal patients are also routinely screened for evidence of human papilloma virus through Pap smear screening. Although treatment is generally deferred until the patient is no longer pregnant, documentation of the presence and severity of dysplastic lesions is an important element of the comprehensive care plan. The timing and method of screening for these infectious conditions are reviewed in Chap. 3.

Patients with suggestive symptoms may also be screened for other infectious conditions. Patients with vaginal discharge should be evaluated for the presence of bacterial vaginosis and trichomoniasis in addition to gonorrhea and chlamydia. The presence of bacterial vaginosis has been associated with an increased risk of preterm contractions and preterm labor. Treatment of bacterial vaginosis or trichomoniasis is recommended after the first trimester. Patients with urinary symptoms such as dysuria or hematuria should be evaluated for possible urinary tract infection (UTI).

## Fetal Infection

In addition to maternal considerations, many infectious processes have health consequences for the developing fetus. Infection in pregnancy may result in three related fetal complications: direct infection of the fetus or newborn, alteration in the normal course of pregnancy, and alteration of fetal development. Many maternal infections can be transmitted to the fetus either during pregnancy or at the time of delivery. Syphilis, gonorrhea, group B strep (GBS), and HIV, among others, all have demonstrated maternal–fetal transmission. Syphilis is associated with neonatal long bone infection; gonorrhea may cause neonatal ophthalmological disease; GBS is a significant contributor to neonatal febrile illness including meningitis; maternal–fetal transmission of HIV remains a critical source of new HIV infections in many parts of the world.

Infectious agents may result in pregnancy complications even in the absence of direct fetal infection. Maternal infections, such as bacterial vaginosis or GBS, may affect the course of pregnancy resulting in such complications as preterm labor and delivery. Universal or directed screening of prenatal patients can identify a significant number of patients at risk. Several infectious complications have been associated with significant developmental abnormalities in fetuses. Cytomegalovirus, toxoplasmosis, and measles all have potentially serious effects on fetal development. Appropriate screening for exposure and/or timely vaccination can reduce the risk for these complications.

## Antibiotic Use in Pregnancy

Antibiotic choice in pregnancy is complicated by the fact that several commonly used classes of antibiotics are contraindicated or should be used with caution in pregnancy. Prior to the use of any antibiotic in pregnancy, providers should review its safety and efficacy in pregnancy. If questions concerning safety remain, consultation with a maternal–fetal expert is recommended. A more complete review of the use of medications in pregnancy can be found in Chap. 4.

Although microbial sensitivity to antibiotics is important in the management of any patient, it is of particular importance in pregnant patients. The potentially limited number of available choices makes culture and sensitivity studies critical in pregnancy. Allergies to such common antibiotics as penicillin may further limit antibiotic choices. In such circumstances, providers may be considering third- or fourth-line antibiotic choices. Under such circumstances, knowing the specific microbe and its sensitivity to available antibiotics is often very helpful.

Although a comprehensive review of all antibiotic classes is beyond the scope of this text, a few key classes should be reviewed (a review of the risk classification for drugs used in pregnancy can be found in Chap. 4).

### *Sulfonamides*

Sulfonamides are most commonly used as sulfamethoxazole in combination with trimethoprim. This antibiotic is routinely used for respiratory tract and UTIs in non-pregnant patients. Sulfonamides readily cross the placenta and levels persist in newborns for several days postpartum. Sulfonamides are a category B drug early in pregnancy but are category D near term. Complications include jaundice and hemolytic anemia. Currently available data do not suggest teratogenesis with the use of sulfonamides. Trimethoprim also crosses the placenta freely. Trimethoprim acts as a folate antagonist and may theoretically contribute to congenital abnormalities if used in the first trimester. There is no definitive data linking trimethoprim to congenital defect; however, because of its mechanism of action and some data to suggest a possible increased risk, trimethoprim is a category C drug.

### *Fluoroquinolones*

This class of drugs includes such agents as ciprofloxacin, levofloxacin, lomefloxacin, norfloxacin, and ofloxacin. These are used for a wide variety of infections including UTIs and respiratory tract infections. Ciprofloxacin has been associated with cartilaginous defects in some animal studies. Data from human studies does not demonstrate a consistent pattern of congenital abnormalities. In general, safer alternatives to fluoroquinolones exist for most conditions. Fluoroquinolones are generally a category C drug.

## ***Aminoglycosides***

This class of drugs includes such agents as gentamicin, used widely for serious Gram-negative infections. Although aminoglycosides are used when indicated in pregnancy, there are several issues that should be considered prior to their use. Aminoglycosides cross the placenta and are present at significant but reduced levels (compared with maternal serum levels). Gentamicin is associated with nephrotoxicity in laboratory animals, although only one possible case of possible prenatal gentamicin-associated renal disease has been reported in the literature. Ototoxicity has also been reported with gentamicin use, although no cases associated with in utero exposure have been reported. Aminoglycosides are generally considered class C drugs in pregnancy.

## ***Vaginitis/Vaginosis***

### ***Background***

A number of infectious agents may result in significant discharge per vagina during pregnancy. Many of the etiologic agents have significance in pregnancy; therefore, a search for the cause and appropriate treatment is particularly important in pregnant patients. In the presence of a significant host inflammatory response, this condition is referred to as vaginitis. In other circumstances, the presence of discharge in the absence of inflammation is referred to as vaginosis. Alteration in the normal vaginal environment including change in pH or elimination of normal flora with the use of some antibiotics allows for proliferation of such abnormal flora as *Gardnerella*, *Trichomonas*, or *Candida* species.

Bacterial vaginosis has been associated with a variety of complications in pregnancy including amniotic fluid infection, chorioamnionitis, postpartum endometritis, premature rupture of membranes (rupture of membranes prior to the onset of labor), preterm labor, preterm delivery, and low birth weight. Complications associated with yeast vaginitis are less clearly defined. In addition to the complications just noted, infection with gonorrhea or chlamydia has been associated with maternal-fetal transmission at the time of delivery.

### ***History***

A careful history will often assist in elucidation of the underlying cause for symptoms of vaginitis. The most common, and often most prominent symptom, is discharge per vagina. Patients should also be questioned concerning systemic symptoms such as fever, nausea, vomiting, change in bowel habits, or urinary symptoms. Recent sexual contact should be reviewed (in addition to prior laboratory studies for gonorrhea, chlamydia, urinalysis, urine culture, or wet mount studies). Symptoms in the partner should also be elicited. A review of recent medications and a past history of similar complaints may yield additional information.

## ***Physical Examination***

In addition to the routine elements of a prenatal physical examination, a focused exam should be performed to identify key findings consistent with vaginitis or, less commonly, significant peritoneal infections. On abdominal examination, providers should note pain rebound or guarding. On the pelvic examination, a note should be made of external or internal erythema or edema, discharge (including the quality and quantity of such discharge), blood, vaginal or cervical lesions, cervical motion tenderness, and adnexal masses or tenderness.

## ***Laboratory Studies***

Further testing is warranted when discharge is noted on exam. Patients should be screened for gonorrhea and chlamydia, especially in the presence of significant mucopurulent discharge. In addition, a sample of discharge should be obtained and tested for pH, KOH (potassium hydroxide) “whiff,” and microscopic examination with saline and KOH solutions.

### ***pH Testing***

pH paper placed directly in discharge may be useful in screening for abnormalities. A pH greater than 4.5 is consistent with bacterial vaginosis or trichomoniasis. A pH less than 4.5 is consistent with yeast.

### ***KOH “Whiff”***

KOH mixed with a sample of the discharge can be helpful in diagnosing bacterial vaginosis. A positive “whiff” test reveals a foul or fishy odor when the KOH is added.

### ***Microscopic Examination***

One slide prepared with saline and discharge; one slide prepared with KOH and discharge. The saline-prepared slide may show evidence of bacterial vaginosis (clue cells) or trichomoniasis (flagellated protozoa). KOH will lyse most cellular elements within the discharge, leaving only yeast on the slide. The presence of fungal elements on a KOH-prepared slide is evidence of candidal infection.

## ***Treatment***

When an infectious agent is identified, most authorities recommend treatment. The benefits of treatment may include reduction of pregnancy-related complications and/or symptomatic relief for the patient. Care should be exercised to select agents with a good safety profile and proven efficacy in pregnancy. If appropriate treatment is unclear, consultation with a maternal–fetal expert is recommended.

### ***Bacterial Vaginosis***

Patients with bacterial vaginosis present with mild or no pruritus, minimal pain, thin gray discharge, pH greater than 4.5, positive whiff test, and clue cells on microscopic examination. Bacterial vaginosis may be treated with clindamycin 2 % cream or oral clindamycin 300 mg by mouth twice a day for 7 days. After the first trimester, metronidazole is an alternative (0.75 % gel 5 g daily for 5 days or 250 mg by mouth three times daily or 500 mg by mouth twice daily for 7 days).

### ***Trichomoniasis***

Trichomoniasis presents as mild pruritus, minimal pain, copious yellow thin discharge, pH greater than 4.5, negative whiff test, and flagellated protozoa on microscopic examination. It may be treated after the first trimester with oral metronidazole. Some experts recommend against treating asymptomatic trichomoniasis in pregnancy as studies do not demonstrate improved outcomes. Topical metronidazole is associated with a higher rate of treatment failure and/or recurrence. Treatment is 500 mg of oral metronidazole twice a day for 7 days. An alternative regimen is 2 g of oral metronidazole as a single dose. This regimen is associated with a higher recurrence rate and increased gastrointestinal side effects. Tinidazole is effective in nonpregnant patients but safety data in pregnancy is limited and therefore it is not currently recommended.

### ***Yeast Vaginitis***

Patients with yeast vaginitis present with mild-severe pruritus, perineal irritation or pain, thick “curd-like” whitish discharge, pH less than 4.5, negative whiff test, and fungal elements on KOH-prepared microscopic examination. Yeast vaginitis may be treated with topical vaginal antifungal preparations, many of which are now available over the counter (OTC). Because of the desirability of obtaining a clear

diagnosis and of minimizing the use of unnecessary medications during pregnancy, most patients should be cautioned against self-treating with OTC preparations without consulting their provider.

### ***Gonorrhea/Chlamydia***

Gonorrhea/chlamydia should be treated in the usual manner to prevent maternal and/or fetal complications. In pregnancy, the use of quinolones and most variants of tetracyclines should be avoided. Gonorrhea should be treated with dual therapy including ceftriaxone and azithromycin. In the case of cephalosporin allergy, recommendations include either (1) desensitization followed by treatment with cephalosporin or (2) azithromycin 2 g orally. This treatment is also effective for chlamydia but has lower efficacy against gonorrhea and therefore should be followed by a test of cure. Treatment for chlamydia is either azithromycin 1 g orally as a single dose or amoxicillin 500 mg orally three times daily for 7 days. Because treatment efficacy is reduced in pregnancy, a test of cure is recommended no sooner than 3 weeks following treatment.

## **Urinary Tract Infections**

### ***Background***

Infections of both the lower and upper urinary tracts are more common in pregnancy. Anatomic and hormonal changes associated with pregnancy increase urinary stasis and may enhance ascending bacterial seeding. The increase in size of the uterus combined, later in pregnancy, with the presence of fetal body parts yields a functional partial obstruction of the outflow tract thereby increasing urinary stasis. Increased progesterone may yield ureteral smooth muscle relaxation, thus increasing the risk for ascending bacterial seeding and proliferation. Such changes appear to peak late in the second trimester or early in the third trimester. UTIs are associated with an increased risk of preterm delivery and low birth weight. In addition, the bacterial seeding can have serious consequences for the mother, including cystitis and pyelonephritis.

Women often report increased urinary frequency during the course of pregnancy. However, bacteriuria and its sequelae are also common during pregnancy. Approximately 5–10 % of all pregnant women develop asymptomatic bacteriuria (with higher prevalence among some at-risk populations). Of these, approximately one-third will develop clinically significant UTIs if not treated. Cystitis complicates about 1 % of all pregnancies with an additional 1–2 % of all pregnant women developing pyelonephritis. Commonly occurring pathogens include *E. coli* (80 %), *Klebsiella*, *Proteus*, and *Staphylococcus saprophyticus*. GBS is an uncommon cause of UTIs but is of particular significance in pregnancy because of its association with an increased risk for preterm labor and neonatal infection.

## ***Diagnosis***

The challenge, then, in pregnancy is to distinguish “normal” pregnancy-related urinary frequency from the urinary symptoms (including frequency) associated with UTIs.

## ***History***

Many patients may be asymptomatic. Others may report urinary frequency, urgency, dysuria, hematuria, fever, abdominal pain, cramping or contractions, and flank or back pain. Although frequency and urgency may be normal in pregnancy, the presence of the other symptoms should always prompt further evaluation. A change in baseline frequency or urgency should also prompt reevaluation. Because of the subtle or potentially confusing nature of the patients’ presenting complaints, providers should maintain a high index of suspicion for UTI. The presence of any symptoms should generally prompt evaluation and treatment if indicated.

## ***Physical Examination***

The physical examination should focus on physical signs of infection and/or signs of ascending infection. Key elements of the physical examination should include temperature, abdominal pain, rebound or guarding, and costovertebral angle tenderness. A pelvic examination is indicated for patients who present with abdominal or back symptoms or for those who report bleeding, contractions, or cramping.

## ***Laboratory Examination***

Urine culture remains the gold standard for diagnosis of UTIs. More than 100,000 colony-forming units per milliliter are considered a positive urine culture. In addition to assisting with the diagnosis, culture may also be useful in identifying the bacterial etiology and the antibiotic sensitivities of the isolated organism. More rapid assessment is often necessary, however, and a variety of additional tests are available that can provide results while the patient is still in the office (or within a few hours). Formal urinalysis performed in a laboratory may be helpful in assisting in diagnosis.

Other office-based tests are very specific but not sensitive. Dipstick leukocyte esterase testing is approximately 20 % sensitive and a 95 % specific. Dipstick nitrite is approximately 57 % sensitive and 95 % specific. Urine microscopy with more than ten white blood cells per high-powered field on a spun urine sample is about 25 % sensitive and 99 % specific. These tests can be very useful if positive but may need to be followed up with urine culture if negative.

## ***Treatment***

Because of the potentially significant complications of untreated UTIs, treatment is recommended regardless of patient symptoms. There are several treatment options but, as always, each treatment option should be reviewed for safety and efficacy in pregnancy. Ampicillin has traditionally been the treatment of choice for UTIs during pregnancy. Its safety profile is excellent and it has been used widely during pregnancy. However, emerging ampicillin resistance in *E. coli* (more than one-third resistant in some regions) has led many providers to use alternatives as first-line therapy. Among the options for treatment are the following:

- Cephalexin (category B, 250–500 mg orally, twice a day)
- Nitrofurantoin (category B, 50–100 mg orally, twice a day)
- Sulfa drugs (sulfisoxazole 1 g orally, four times a day, TMP/SMZ 160/180 mg orally, twice a day; they may be used in the second trimester but should be avoided in either first or third trimester)
- Fosfomycin (category B; 3 g single dose)

With the exception of fosfomycin, all treatment should be for 7–10 days for lower tract infections and up to 2 weeks for pyelonephritis.

## **Group B Strep**

### ***Background***

GBS (most commonly *Streptococcus agalactiae*) are normal colonic flora. Under appropriate conditions, GBS colonize in the vagina, complicating pregnancy and the early neonatal course. Of all pregnant women, 10–30 % are colonized with GBS in either the vaginal or rectal areas. GBS is a common and significant cause of prenatal and neonatal morbidity. GBS is associated with preterm labor, premature rupture of membranes, and preterm delivery. Less commonly, GBS is associated with UTIs. It is also associated with neonatal infection, including meningitis. Infants born to colonize women often develop early onset invasive disease, 80 % within the first 7 days of life. The prevalence of invasive disease is about 0.6 per 1000 live births. In the United States, GBS is responsible for about 2000 neonatal infections and 100 fatalities annually.

Treatment of GBS has been shown to decrease the incidence of such complications. The use of intrapartum antibiotic therapy to treat women at increased risk for transmission to their newborns can significantly reduce peripartum transmission and neonatal infection. Treatment of invasive GBS infection is associated with a reduction in rates of premature rupture of membranes and preterm delivery. The key to successful management is identification and treatment of those patients who are at risk for preterm complications or peripartum transmission.

## ***Diagnosis***

Diagnosis of GBS can be challenging. Most patients do not consistently demonstrate colonization throughout the course of pregnancy. Early GBS cultures do not correlate well with GBS status at the time of delivery. In addition, patients colonized with GBS are generally asymptomatic. For this reason, authorities recommend universal screening of prenatal patients regardless of risk or symptoms near term (35–37 weeks' gestation). For patients who present with suggestive signs or symptoms, GBS testing should be performed. High-risk conditions include preterm labor, premature rupture of membranes, or evidence of GBS UTI. Those patients with a past history of premature rupture of membranes, preterm delivery, or neonatal strep infection should also be considered high risk.

## ***Universal Screening***

All patients are screened during the course of pregnancy with treatment reserved for those who have positive cultures or who are at high risk.

- GBS testing is performed on all women at 35–37 weeks' gestation (the results of which more nearly correlate with intrapartum GBS status than cultures obtained earlier in pregnancy).
- In the absence of GBS culture results, all patients with fever, prolonged rupture of membranes (>18 h), or gestational age less than 37 weeks should be presumed GBS positive.
- In the absence of GBS cultures, all patients with a past history of GBS UTI or neonatal GBS infection should be presumed GBS positive.
- In patients with preterm premature rupture of membranes, GBS cultures should be obtained. Patients may be presumed GBS positive until culture results are available.

## ***Treatment***

Penicillin is the gold standard treatment for GBS in patients without penicillin allergy. Acceptable penicillin-based regimens include penicillin G intravenous, five million units loading dose followed by 2.5 million intravenous units every 4 h until delivery or 2 g of ampicillin intravenous loading dose followed by 1 g intravenously every 4 h.

For patients with a nonserious penicillin sensitivity, cefazolin 2 g IV once and then 1 g IV every 8 h until delivery is recommended. Alternatives for patients with more significant penicillin allergies include clindamycin (if sensitivity data is available and indicates clindamycin sensitivity) or vancomycin 20 mg/kg every 12 h (max 2 g dose).

# Chapter 14

## Hypertension in Pregnancy

### Contents

Background .....	114
Diagnosis.....	114
Diagnostic Criteria.....	114
History.....	115
Physical Examination.....	116
Laboratory Studies .....	116
Management.....	116
Chronic Hypertension .....	116
Pre-Eclampsia .....	117
Prevention .....	117
Management.....	117
Seizure Prevention .....	119

### Key Points

1. Hypertension in pregnancy is defined as blood pressure (BP) higher than 140 mmHg systolic or 90 mmHg diastolic on two occasions separated by at least 6 h.
2. Pregnancy-induced hypertension is defined as hypertension diagnosed at or after 20 weeks' gestation.
3. Pre-eclampsia is a multisystem disease characterized by hypertension and proteinuria.
4. Pre-eclampsia may lead to fetal complications including preterm delivery, intrauterine growth restriction (IUGR), fetal demise, and perinatal death, as well as maternal complications of seizure, stroke, and death.

## Background

Pregnancy may be complicated by hypertension either as a pre-existing condition or as a newly diagnosed condition during pregnancy. Each condition carries with it significant risks and important management considerations that may impact the well-being of both the mother and fetus. Pre-eclampsia, a multisystem disorder that is marked by pregnancy-induced hypertension and proteinuria, is a significant obstetrical risk that affects approximately 5 % of pregnancies.

The exact cause of pre-eclampsia is unknown, but its multisystem complications are well described. Physiologically, pre-eclampsia is marked by increased vascular resistance, platelet aggregation, and endothelial dysfunction. Clinically, pre-eclampsia may be identified with hypertension (occurring after 20 weeks' gestation), proteinuria, HELLP syndrome (**H**emolysis, **E**levated **L**iver enzymes, **L**ow **P**latelets), and seizures (eclampsia).

Chronic hypertension is defined as hypertension (repeated BP readings of systolic  $\geq 140$  or diastolic  $\geq 90$  mmHg) existing prior to pregnancy or first diagnosed prior to 20 weeks' gestation. Obstetrical complications associated with chronic hypertension include increased risk for pre-eclampsia (discussed later), abruptio placentae, premature delivery, IUGR, fetal demise, and fetal stress. Pregnancy itself may worsen hypertensive renal disease. The majority of such complications occur in women with diastolic BPs higher than 110 mmHg although such complications may occur in women with lower BP.

Pre-eclampsia occurs in approximately 5 % of all pregnancies and may be associated with many of the same obstetrical risks as chronic hypertension (Table 14.1): HELLP syndrome (10–20 %), abruptio placentae (1–4 %), and eclampsia (<1 %). Rarely, it may also be associated with maternal stroke or death. Complications for the neonate include IUGR (10–25 %), preterm delivery (15–67 %), and perinatal death (1–2 %). For both the mother and infant, the presence of pre-eclampsia may be associated with long-term cardiovascular morbidity.

## Diagnosis

### *Diagnostic Criteria*

The diagnostic criteria for hypertension in pregnant patients are similar to those for nonpregnant patients. Hypertension may be diagnosed in patients with at least two BP readings of more than 140 mmHg systolic or more than 90 mmHg diastolic separated by at least 6 h but not more than 7 days.

Pre-eclampsia is diagnosed by a combination of hypertension first diagnosed after 20 weeks' gestation and proteinuria ( $\geq 300$  mg per 24 h or 1+ dipstick protein in two random urine samples separated by at least 4 h). In addition, hypertension associated with neurological (headache, visual changes, altered mental status), gastrointestinal (right upper quadrant pain, nausea, vomiting, elevated liver enzymes),

**Table 14.1** Risk factors for pre-eclampsia

Maternal
Family history of pre-eclampsia
Early or late maternal age
Nulliparity
Prior history of pre-eclampsia
Assisted reproduction
Vascular disease
Diabetes
Obesity
Hypertension
Renal disease
Thrombophilia
Rheumatic disease
Infection
Paternal
Primipaternity
Prior pregnancy complicated by pre-eclampsia
Donated sperm
Fetal
Multifetal gestation
Hydrops fetalis
Chromosomal abnormalities
Congenital abnormalities

or hematological (decreased platelet count) findings should also suggest pre-eclampsia. Ten percent of patients who develop HELLP syndrome and up to 33 % of patients who develop eclampsia will not demonstrate the traditional findings of both hypertension and proteinuria.

Severe pre-eclampsia is diagnosed if any one of the three criteria (hypertension, proteinuria, multiorgan involvement) is severe. The criterion for severe hypertension in pregnancy is a systolic BP >160 mmHg and/or diastolic BP >110 mmHg. Proteinuria is considered severe at levels 5 g or more per day. Multiorgan involvement may be demonstrated by pulmonary edema, seizures, altered mental status, headaches, visual disturbance, persistent right upper quadrant pain with elevated liver enzymes, oliguria (<500 cc/day), or thrombocytopenia (<100,000).

## History

Although the diagnosis of pre-eclampsia is generally made on the basis of BP readings and proteinuria, a number of risk factors have been identified and should be reviewed in the history. These risk factors are shown in Table 14.2 and include family history of pre-eclampsia, past history of pre-eclampsia, nulliparity, primipaternity, age over 40, assisted fertility, multiple gestation, chronic hypertension, diabetes mellitus, renal disease, thrombophilia, obesity, maternal infection, and smoking.

**Table 14.2** Complications of pre-eclampsia

Preterm delivery
Intrauterine growth restriction
HELLP syndrome
Pulmonary edema
Acute renal failure
Abruptio placentae
Perinatal death
Eclampsia
Stroke
Death

In addition, all patients should be screened at each visit for symptoms suggestive of possible pre-eclampsia including headache, visual changes, altered mental status, abdominal pain, nausea, or vomiting.

## Physical Examination

Each prenatal visit should include documentation of the patient's BP. For patients who demonstrate elevated BP, a return visit should be scheduled within 1 week for a recheck of BP.

For patients with elevated BP, the physical examination should also include ophthalmological, neurological, and abdominal examinations as well as notation of peripheral edema (feet, hands, and face).

## Laboratory Studies

The most accurate test for proteinuria is a 24-h urine collection with measurement of protein excretion. In settings where such collections are not practical, two separate dipstick urinalysis tests demonstrating at least 1+ protein may substitute. Additional labs for patients with suspected pre-eclampsia should include complete blood count with platelets, liver enzymes, and a coagulation panel.

## Management

### *Chronic Hypertension*

Patients with a pre-existing diagnosis of hypertension should continue to receive treatment during the course of pregnancy. Providers should review the safety of their existing antihypertensive regimen and make adjustments as necessary.

Antihypertensive medications are reviewed in Chap. 4. Antihypertensive medications commonly used during pregnancy include methyldopa, hydralazine, and  $\beta$ -blockers. For patients diagnosed with hypertension after conception but prior to 20 weeks' gestation, the role of antihypertensives is less clear. Patients with severe hypertension (diastolic over 110 mmHg) should probably receive pharmacological treatment but treatment of women with mild essential hypertension in pregnancy has not been shown to improve outcomes. Patients with chronic hypertension should be carefully followed for signs or symptoms of pre-eclampsia as 15–25 % will develop superimposed pre-eclampsia.

## Pre-Eclampsia

### *Prevention*


As the risk factors associated with increased risk for pre-eclampsia have become increasingly well defined, interest has been focused on prevention of pre-eclampsia in patients at high risk. Proposed interventions have included dietary supplements, aspirin, and antihypertensive medications. Although the results have been mixed, there is little evidence to support the preventive benefits of diet and exercise, protein or salt restriction, magnesium, fish oil or antioxidant supplementation, heparin, or antihypertensive medications. Although some studies have shown potential benefit with the use of low-dose aspirin or calcium supplementation, insufficient evidence exists to make general recommendations concerning their use in the prevention of pre-eclampsia.

### *Management*

The management of pre-eclampsia involves balancing the maternal risks of prolonged pregnancy against the neonatal risks of premature delivery. The potential risks and benefits for each patient must be considered individually. Although no universally acceptable management protocol can be recommended, a general approach to the management of pre-eclampsia is outlined in Fig. 14.1.

Delivery is the definitive maternal management plan. For this reason, delivery should occur as soon as it can reasonably be achieved. For patients at term with mild disease, induction of labor is indicated. For patients with severe pre-eclampsia at or beyond 34 weeks, delivery is indicated with appropriate neonatal support. If the maternal condition appears stable but there is evidence of fetal compromise, management should follow a protocol similar to that of IUGR (see Chap. 6).

For patients with mild disease, no evidence of fetal compromise, and a gestational age of less than 34 weeks, ideal management is not well established. Under these circumstances, careful monitoring of maternal condition should be combined with close evaluation of fetal well-being. If both maternal and fetal conditions remain



**Fig. 14.1** Management of pre-eclampsia

stable, delivery can occur at 38 weeks. If either maternal or fetal condition worsens, delivery should occur as soon as possible.

As noted for chronic hypertension, the use of antihypertensives has shown mixed results. For patients with severe hypertension, the use of antihypertensive

medications has been shown to reduce maternal cerebrovascular complications. However, such treatment has not been shown to reduce neonatal complications and does not alter the maternal course of disease in relation to multiorgan complications. The use of antihypertensives has not been shown to improve maternal or neonatal outcomes in patients with mild disease. There is little evidence to support the use of any specific class of antihypertensive medications in the treatment of pre-eclampsia.

The use of corticosteroids has not been shown to improve maternal outcomes, although their use has been shown to improve neonatal outcomes for infants born prior to 34 weeks' gestation.

### ***Seizure Prevention***

Seizure (eclampsia) is one of the most significant complications of pre-eclampsia and may occur without preceding warning signs. For this reason, seizure prophylaxis with magnesium sulfate is indicated for patients with pre-eclampsia in the intrapartum period. Specific regimens for the use of magnesium may vary between institutions and the specific protocol should be reviewed by each provider. One possible protocol is as follows: loading dose of 6 g of intravenous magnesium sulfate followed by 2 g per hour of continuous intravenous infusion. Magnesium may be toxic in high doses and patients should be monitored closely while undergoing magnesium therapy. Maternal BP, deep tendon reflexes, mental status, and urinary output should be monitored. Serum magnesium levels should be measured. Magnesium levels above 7 mEq/L are associated with diminished deep tendon reflexes. Magnesium levels above 10 mEq/L are associated with respiratory depression. Magnesium levels above 12 mEq/L are associated with cardiac depression and arrest. Magnesium elevation is reversible with 10 % calcium gluconate 10 mL intravenous given over 10–15 min.

# Chapter 15

## Diabetes in Pregnancy

### Contents

Background .....	121
Diagnosis.....	122
Pregestational Diabetes .....	123
Management.....	123
Gestational Diabetes .....	126
Preconception Management.....	126

### Key Points

1. Diabetes mellitus (DM) is defined as carbohydrate intolerance resulting from either insulin deficiency or insulin insensitivity.
2. Exposure to elevated serum glucose is associated with increased risk for organogenic birth defects, macrosomic infants, shoulder dystocia, and birth trauma.
3. Gestational diabetes is defined as glucose intolerance first recognized during pregnancy.

### Background

DM (hyperglycemia secondary to either insulin deficiency or relative insulin insensitivity) is a significant medical condition with a potentially profound impact on pregnancy. Uncontrolled diabetes prior to or early in the course of pregnancy is associated with a variety of birth defects, including renal, gastrointestinal, cardiac, central nervous system, and skeletal abnormalities (see Table 15.1). The presence of diabetes may alter the interpretation of some prenatal tests including  $\alpha$ -fetoprotein and obstetric triple screen. Pregnancy may, in turn, affect the course of diabetes, worsening glucose control in patients with pre-existing diabetes.

**Table 15.1** Congenital abnormalities associated with diabetes

Duodenal and anorectal atresia
Hydronephrosis
Renal agenesis
Neural tube defects
Anencephaly
Ventricular septal defects
Aortic coarctation
Transposition of the great vessels

DM may be classified as type 1 diabetes, associated with pancreatic failure and insulin deficiency; type 2 diabetes, associated with ineffective insulin utilization (generally associated with hyperinsulinemia); or gestational diabetes mellitus (GDM), which is diabetes first diagnosed or recognized in pregnancy (generally associated with insulin resistance and similar to type 2 diabetes). GDM is associated with increased glucose intolerance during the course of pregnancy. In most cases this glucose intolerance will resolve following pregnancy. Hormonal changes associated with pregnancy increase maternal glucose intolerance. Human placental lactogen decreases cellular glucose uptake and increases lipolysis. Estrogen and progesterone are also known to affect glucose metabolism although to a lesser extent than human placental lactogen.

## Diagnosis

Diagnosis of types 1 and 2 DM is made prior to pregnancy. The diagnosis is made on the basis of documented hyperglycemia with either insulin deficiency (type 1) or insulin resistance (type 2). The diagnosis of GDM is generally made on the basis of oral glucose testing during the course of prenatal care, although, in some cases, the diagnosis may be made on the basis of fasting blood glucose values.

Screening for GDM is an area of some controversy. Most authorities recommend universal screening of all patients at 24–28 weeks' gestation. Earlier testing for those patients identified as high risk (*see* Table 15.2) is suggested by some authorities but has limited data demonstrating improved outcomes.

Initial screening is via 50 g glucose tolerance testing. Patients are not required to fast and do not need to alter their dietary intake. A standardized 50 g glucose load is administered orally and serum glucose level is tested at 1 h. The threshold for a positive screening test is also an area of some controversy and providers should be aware of the standard of care in their institution. Serum glucose values should be less than 130 or 140 mg/dL. The lower value has a higher sensitivity (will miss fewer true cases of diabetes) but is associated with a higher false-positive rate. The higher value reduces false-positive findings (has a higher specificity) but may have a lower sensitivity.

**Table 15.2** Risk factors for gestational diabetes

Family history of diabetes
Family history of macrosomic infants
Ethnicity
Obesity
Past history of gestational diabetes
Past history of macrosomic infant
Multigestation

**Table 15.3** Gestational diabetes diagnosis (100 g, 3-h oral glucose tolerance test)

Sample	Threshold
Fasting	105
1 h	190
2 h	165
3 h	145

*Note:* the diagnosis of gestational diabetes is made if any two values exceed the values listed in the table

Regardless of which threshold is used, all patients with a positive screening test should undergo diagnostic testing with a 3-h 100 g oral glucose challenge test. Patients should eat a carbohydrate-rich or unrestricted diet for 3 days prior to testing but should fast the night before the test. A standardized 100 g oral glucose load is administered, and serum glucose levels are drawn fasting 1, 2, and 3 h following administration. There is debate concerning the threshold values for this test as well. The National Diabetes Data Group's recommendation is presented in Table 15.3. According to this protocol, blood glucose levels should be below 105 mg/dL fasting and below 190 mg/dL, 165 mg/dL, and 145 mg/dL at 1 h, 2 h, and 3 h, respectively. The test is considered diagnostic if any two values exceed the threshold.

## Pregestational Diabetes

### *Management*

#### **Preconception Management**

The management of pregestational diabetes ideally should begin during the preconception period. Preconception management should include discussion of the effects of diabetes on pregnancy as well as discussion of the effects of pregnancy on diabetes. Patients should be counseled concerning the risks of maternal hyperglycemia on the developing infant and the importance of strict glycemic control prior to and throughout the course of pregnancy.

**Table 15.4** White's classification of diabetes in pregnancy

Class	Definition
A	Diet controlled
B	Age of onset $\geq 20$ years, duration $< 10$ years
C	Age of onset 10–19 years or duration 10–19 years
D	Age of onset $< 10$ or duration $> 20$ years
F	Nephropathy ( $\geq 500$ mg/day proteinuria)
H	Clinically evident arteriosclerotic heart disease
R	Retinopathy
T	Renal transplant

Preconception management should also include classification of the severity of diabetes. The modified White's classification system is shown in Table 15.4. This classification is based on four factors: (a) treatment regimen, (b) duration of disease or age of onset, (c) associated disease, and (d) diabetic complications.

Because of the association of poor glycemic control with an increased risk for congenital defects, all patients should be intensively monitored and tightly controlled prior to conception. Because of the complexity of achieving tight glycemic control, such efforts should begin well before anticipated conception when possible.

Oral hypoglycemic agents are not generally utilized during pregnancy. Patients managed on oral hypoglycemic agents prior to conception may require additional time (and education) to transition to the use of insulin. Patients may also require comprehensive diabetes education on diet, exercise, and blood glucose self-monitoring. Prenatal vitamins with folate should be prescribed as a part of preconception management as well.

## Management in Pregnancy

For patients who did not receive preconception evaluation and treatment, all of the management just discussed should begin with the first prenatal visit. Because the early weeks of pregnancy are the period of organogenesis, rapid assessment and intervention are critical to improve outcomes. Patients may require more frequent visits and will probably benefit from intensive educational interventions, beginning with a review of the patient's own knowledge of her disease and its management.

### Diet

All patients with diabetes should be counseled concerning the role of diet in the management of diabetes. Although specific recommendations may vary between patients, general recommendations include the following:

1. Caloric intake of 1800–2400 kcal per day generally as several small meals rather than fewer large meals.
2. Restriction of simple carbohydrates.
3. Inclusion of complex carbohydrates in an overall diet that includes approximately 40 % carbohydrates, 30 % fat, and 30 % protein.

4. Regular and predictable caloric intake will facilitate glycemic control and reduce the likelihood of hypoglycemia.
5. A small snack should be available at all times in case of hypoglycemia.

## **Insulin**

Pharmacological management of diabetes in pregnancy generally utilizes a combination of intermediate- and short-acting insulin. This combination allows for more controlled adjustment of glycemic control than either oral hypoglycemic agents or long-acting insulin alone. The exact dose of insulin will be based on measured blood glucose values with target levels of less than 105 mg/dL fasting (95 mg/dL according to some authorities) and less than 120 mg/dL 2 h postprandial. Insulin should be adjusted as necessary to achieve these target levels. Tight glycemic control has been demonstrated to improve obstetrical outcomes but have also been associated with increased risk of hypoglycemia. For this reason, patients should be educated concerning the symptoms of hypoglycemia and the appropriate steps to be taken should hypoglycemia occur.

## **Oral Hypoglycemics**

Although insulin has been the traditional pharmacologic treatment choice (and remains the most common choice) there is increasing acceptance of oral hypoglycemic medications in pregnancy. Glyburide, which does not cross the placenta, is the most common oral hypoglycemic agent used in pregnancy. Outcomes appear to be similar to insulin for patients who achieve equal level of glycemic control. Alternately, metformin can be used in the second and third trimester. Patients on metformin will often require the addition of insulin to achieve glycemic goals. Limited studies of acarbose have suggested a possible role although safety data is limited and abdominal cramping may limit efficacy. Older sulfonylureas and the thiazolidinedione medications are not currently recommended in pregnancy.

## **Blood Sugar Monitoring**

As noted, insulin dosing should be based on the results of blood glucose monitoring. For this reason, frequent finger-stick glucose monitoring should be a part of the management of patients treated with insulin. Although glycosylated hemoglobin values may provide adjunctive data concerning the level of glucose control, the long-term (3-month) retrospective nature of such testing makes it unsuitable for insulin management in pregnancy. Patients should be counseled to test blood sugar fasting and 2 h after every meal on a daily basis. If such intense monitoring is not feasible, patients should be counseled to vary daily (e.g., fasting and 2 h after lunch one day, 2 h after breakfast and 2 h after dinner the next) to allow for adequate assessment of insulin needs. Additional testing may be recommended at times when patients feel the symptoms of hypoglycemia to document the decrease in glucose levels.

## Gestational Diabetes

### *Preconception Management*

Preconception management of GDM begins with assessment of risk factors for the development of diabetes during pregnancy. The risk factors for GDM are listed in Table 15.1. All patients should be screened for risk factors for GDM. Patients found to be at increased risk may benefit from preconceptual diabetes testing as type 2 DM and GDM share common risk factors. Patients discovered to have diabetes prior to becoming pregnant should receive additional preconception counseling as noted earlier.

### **Management in Pregnancy**

Patients with GDM should be managed similarly to those patients with pre-existing diabetes. Although many patients with GDM can be appropriately managed with lifestyle modification alone, all patients should be counseled concerning the indications for further treatment with insulin if necessary. All patients with diabetes should test blood sugar regularly to assess the adequacy of their treatment regimen.

Because of the fetal risks associated with diabetes in pregnancy, all patients with diabetes should begin fetal surveillance in the third trimester of pregnancy. Antenatal assessment is discussed in Chap. 18. In addition, because of the risk of macrosomia, careful assessment of fetal growth should be included in prenatal assessment. Patients with evidence of macrosomic or small-for-gestational-age fetuses should receive ultrasonographic assessment of fetal growth.

### **Management in Labor**

During labor, efforts should be made to maintain the patient's serum glucose levels between 80 and 110 mg/dL. Care should be taken, however, not to induce maternal hypoglycemia in patients who are restricted from eating or drinking during the course of labor. This is especially true for patients with prolonged second-stage labor.

### **Postpartum Management**

All patients diagnosed with GDM should be screened for diabetes in the postpartum period to document resolution of the diabetes. Such patients are at increased risk for subsequent diagnosis of type 2 diabetes.

# Chapter 16

## Multigestational Pregnancy

### Contents

Background .....	127
Diagnosis.....	129
History .....	129
Physical Examination.....	130
Laboratory and Diagnostic Studies.....	130
Management.....	130
Prenatal Care .....	130
Labor and Delivery .....	131

### Key Points

1. Multigestational pregnancy presents unique management challenges beyond those encountered in singleton pregnancies.
2. Multigestational pregnancy may result from either a single or multiple fertilized ovum (ova).
3. Complications associated with multigestational pregnancies include spontaneous abortion, preterm delivery, pre-eclampsia, postpartum hemorrhage, and increased perinatal mortality.

### Background

Most pregnancies are the product of a single fertilized ovum and result in a single fetus. One to two percent of pregnancies, however, result in multiple fetuses, multigestational pregnancy. Such pregnancies present unique challenges for both prenatal management and delivery. Although uncommon, multigestational pregnancy occurs with sufficient frequency that primary care providers should be familiar with basic management considerations.

Multiple gestations may be the result of either a single fertilized ovum that divides early in development or multiple fertilized ova from the same cycle. Monozygotic (MZ) twins are genetically identical fetuses produced from a single fertilized ovum. MZ twins make up about 30 % of all twin pregnancies. Dizygotic (DZ) twins are the product of two separate fertilized ova. Although they are genetically similar, they are not genetically identical. DZ twins make up 70 % of all twin pregnancies with a frequency of approximately 1 out of 80 pregnancies. Although significantly less common, multigestational pregnancy may result in more than two developing fetuses. The expected natural frequency of multigestational pregnancy is approximately 1 in 80 twins, 1 in 6400 triplets, and 1 in 512,000 quadruplets. The increase in assisted fertility has significantly altered the frequency of multigestational pregnancies, however, and such figures may no longer be entirely applicable.

In addition to multiple fetuses, multigestational pregnancies may have several variations of multiple chorions and placentae. MZ twins may have a single placenta and a single chorion (~60 %) or may have two chorions with either a fused placenta or two separate placentae (~20 % each). DZ twins have two chorions and two placentae, which may be fused or separate (~50 % each).

Multigestational pregnancies are associated with increased risk for a variety of prenatal and delivery-related complications (*see Table 16.1*). Maternal risks associated with multigestational pregnancy include spontaneous abortion, stillbirth, preterm labor, preterm delivery, placental previa, anemia, urinary tract infection (UTI), pre-eclampsia, and postpartum hemorrhage. Up to two-thirds of twin pregnancies will result in loss of one twin in the first trimester. The rate of fetal demise is twice as high as for single pregnancies. The risk of both anemia and UTI is 2–3 times normal. There is a threefold increased risk of pre-eclampsia and a fivefold increased risk of postpartum hemorrhage in multigestational pregnancy.

Fetal risks include developmental abnormalities, growth abnormalities, and preterm delivery complications, including a death rate that is three times the normal.

**Table 16.1** Complications of multigestational pregnancy

Maternal complications
Spontaneous abortion
Stillbirth
Preterm labor
Preterm delivery
Placenta previa
Gestational diabetes
Anemia
Urinary tract infection
Pregnancy induced hypertension/pre-eclampsia
Postpartum hemorrhage
Fetal complications
Perinatal mortality
Developmental abnormalities
Growth abnormalities
Preterm delivery complications

Perinatal mortality is three times that of single pregnancy risk. The risk of both major and minor malformations is double that of single pregnancies. The average gestational age at delivery is 36 weeks for twins and 33 weeks for triplets, significantly increasing the likelihood of complications from prematurity.

## Diagnosis

The diagnosis of multiple gestations is generally made via ultrasound (US) during the course of prenatal care. Careful prenatal care combined with the prevalence of obstetrical US has greatly diminished the number of unanticipated multigestational deliveries. Although the diagnosis is generally made ultrasonographically, the provider's index of suspicion may be heightened by historical or physical examination findings during the course of prenatal care. Key findings are summarized in Table 16.2.

## History

Prior to conception or early in the prenatal course, a past history or family history of multigestational pregnancy should be explored. In addition, a history of assisted reproduction should be noted, when present. Patients with multigestational

**Table 16.2** Findings associated with multigestation

History
Family history of multigestational pregnancy (including patients who are themselves twins)
Past history of multigestational pregnancy
Assisted reproduction
Maternal symptoms
Nausea
Vomiting
Headache
Shortness of breath
Abdominal distention
Constipation
Physical findings
Uterus size larger than dates
Excess maternal weight gain
Multiple palpable fetuses
Multiple audible fetal heart tones
Laboratory/diagnostic studies
Significantly decreased hemoglobin
Elevated maternal serum $\alpha$ -fetoprotein
Ultrasound documentation of multiple fetuses

pregnancy may report an increase in pelvic pressure, nausea, vomiting, headache, shortness of breath, distention, and constipation. Although none of these symptoms is specific to multigestational pregnancy, the number of symptoms and/or the severity of the complaint may be increased in such pregnancies.

## ***Physical Examination***

Because the symptoms noted here are neither sensitive nor specific for multigestational pregnancy, suggestive physical findings may be important in identifying patients with multiple fetuses. Increased maternal weight gain is a nonspecific but suggestive finding. Uterine size greater than expected for gestational age may also be a critical finding. Two palpable fetuses or multiple fetal heart tones, although less common, should prompt immediate US evaluation.

## ***Laboratory and Diagnostic Studies***

As previously noted, obstetrical US is the definitive study. In skilled hands, US may demonstrate multiple gestations as early as 4 weeks' gestation. Other suggestive laboratory findings include decreased hemoglobin and elevated maternal serum  $\alpha$ -fetoprotein (levels approximately 2–3 times higher than for singleton pregnancies even in the absence of fetal abnormalities).

## **Management**

### ***Prenatal Care***

In general, the course of prenatal care is similar to that of singleton pregnancies with additional care directed toward specific increased risks associated with multigestational pregnancy. The diagnosis should be confirmed as early as possible. For patients at high risk (e.g., assisted reproduction), this may include US documentation as early as 4 weeks' gestation. Because of the increased potential for genetic abnormalities, providers may consider offering genetic diagnosis for patients over the age of 33. The frequency of prenatal visits may be increased to monitor for signs/symptoms of preterm contractions or preterm labor.

Patients should be counseled concerning the increased need for careful dietary intake. Folic acid supplementation should be started at the first prenatal visit and iron supplementation may also be appropriate. Maternal weight gain should be closely monitored with a target weight gain of 35–45 lb over the course of pregnancy.

Fetal growth should be closely monitored starting early in the third trimester (or earlier if patient is determined to be at risk for abnormal fetal growth). US studies every 4 weeks will allow for documentation of adequate and symmetric fetal growth.

## ***Labor and Delivery***

Management of multigestational deliveries is associated with several significant challenges. Providers with limited experience or without access to necessary obstetrical and neonatal support should arrange for appropriate backup or transfer prior to the onset of labor. Patients with three or more fetuses are generally not candidates for vaginal delivery and arrangements should be made early in the prenatal course for appropriate cesarean section.

Because the method of delivery may vary with the presentation of the infants at the time of labor, patients with twin pregnancies should be admitted at the first signs of labor, bleeding per vagina, or rupture of membranes. On admission, US should be performed to confirm the position and presentation of each infant. All twin deliveries should be attended by one pediatric team (with all necessary neonatal resuscitation equipment) for each infant.

By convention, the first twin is designated twin A and the second, twin B. Possible variations of presentation include (a) twin A vertex, twin B vertex (~40 %), (b) twin A vertex, twin B breech (~40 %), or (c) twin A breech, twin B any presentation (~20 %). Vaginal delivery can only be attempted if twin A is in a vertex presentation; therefore, pregnancies with twin A in a breech position at the time of labor will require cesarean section delivery. If twin A is in a vertex presentation at the time of labor and there are no other contraindications, vaginal delivery may be attempted. In addition to the usual indications for cesarean section associated with any pregnancy (*see Chap. 22*), cesarean section is indicated in twin deliveries that demonstrate twin–twin transfusion.

Because the position of twin B cannot be absolutely known prior to delivery of twin A, all attempted vaginal deliveries should be performed in a setting equipped for a cesarean section if necessary. All patients should have intravenous access, available typed and crossmatched blood, and a complete blood count prior to delivery.

Vaginal delivery of the first twin is managed in a similar manner to singleton deliveries. Following the delivery of twin A, US should be performed to confirm the presentation of twin B. If twin B is in a vertex presentation, delivery is again managed in a manner similar to singleton pregnancies. If twin B is found to be in a breech position following delivery of twin A, external version may be attempted to position the infant in a vertex presentation. If twin B remains in a breech position, cesarean section should be performed.

# Chapter 17

## Postdates Pregnancy

### Contents

Background .....	133
Diagnosis.....	134
History.....	134
Physical Examination.....	135
Diagnostic Studies .....	135
Management.....	135
Assessment of Fetal Well-Being .....	135

### Key Points

1. Term pregnancy is defined as 37–42 weeks’ gestation.
2. Accurate pregnancy dating is critical to assessment and management of post-dates pregnancy.
3. Timing of delivery should be prior to 42 weeks’ gestation and earlier if antenatal testing is nonreassuring.

### Background

When a firm estimated date of delivery (EDD) is established early in pregnancy, providers can anticipate that most pregnancies will result in spontaneous delivery at term. Term in this setting is defined as 38–42 weeks’ gestation. Under some circumstances, however, pregnancy may continue beyond 42 weeks, requiring assessment and management as a postdates pregnancy. Although the exact number of pregnancies that continue beyond term is not well established (3–12 %), approximately 10 % of all pregnancies will result in induction of labor (although not all for

postdates pregnancy). The cause of prolonged gestation is not well understood but patients with one postterm delivery have a 50 % likelihood of subsequent postterm delivery.

A significant first step in identifying postdates pregnancies is confirmation of gestational dating. As noted in Chap. 3, a variety of measurements may be used to establish the EDD, including the last menstrual period (LMP) and obstetrical ultrasound (US). Confirmation of the EDD is critical to appropriate management of postdates pregnancy. For this reason, all such data should be reviewed carefully and confirmed.

Although pregnancy is not considered postdates until 42 weeks of gestation, planning for management should begin at or near the EDD. Careful fetal monitoring and management for delivery is critical as postdates pregnancy is associated with an increased risk for operative delivery, macrosomia, shoulder dystocia, meconium aspiration, and fetal mortality (twice baseline at 42 weeks, six times baseline by 44 weeks).

## Diagnosis

Approximately one-half of all postdates pregnancies are caused by inaccurate gestational dating. For this reason, confirmation of the appropriate gestational age is critical.

## History

The patient's menstrual history should be reviewed, including the timing and normality of the LMP. Under a variety of conditions, the episode of bleeding considered to be the LMP may be inaccurate. Oligomenorrhea, prior use of contraception such as oral contraception or medroxyprogesterone, and pregnancy-related first-trimester bleeding may all alter the accuracy of menstrual history. First-trimester bleeding per vagina is very common and such bleeding may be interpreted as menstrual bleeding when, in fact, it was not. Early pregnancy bleeding is reviewed in Chap. 9.

The date of the first positive pregnancy test may be helpful in narrowing the possible dates of pregnancy. A review of the prenatal record should include obstetrical US results, if available, fundal height measurements, fetal quickening, and first noted fetal heart tones by US (4–6 weeks), handheld Doppler (10–12 weeks), or fetoscope (18–20 weeks). A pelvic examination with bimanual assessment of uterine size early in pregnancy may also provide confirmatory support for EDD.

Prior obstetrical history should be reviewed as a past history of postdates delivery is associated with an increased risk of subsequent postdates delivery.

## ***Physical Examination***

Primary confirmation of postdates pregnancy is generally provided by a careful history. Physical examination is generally supplementary at term and should not alter an otherwise well-established EDD.

## ***Diagnostic Studies***

In the absence of adequate prenatal data to establish EDD, late-pregnancy US may provide a broad estimate of gestational age. US accuracy diminishes with increasing gestational age, however, and late-pregnancy results should be interpreted with caution. Although the exact accuracy of dating by US cannot be established, the “1 week per trimester” rule of thumb is a reasonable estimate of accuracy. US studies performed in the first trimester are accurate to within 1 week; those performed in the second trimester are accurate to within 2 weeks; those performed in the third trimester are accurate to within 3 weeks.

## **Management**

Postdates pregnancy presents two related challenges to providers: (a) assessment of continued fetal well-being and (b) assessment of need for induction.


Induction of labor is discussed in Chap. 20. As a general rule, the risk associated with postdates pregnancy after 42 weeks’ gestation provides support for a policy of induction at or before that time. An overview of management is provided in Fig. 17.1.

## ***Assessment of Fetal Well-Being***

A variety of tests to assess fetal well-being are available. These tests range from patient-performed outpatient monitoring to formal monitoring with US examination. Prenatal care providers should be familiar with each of these options, their role in the management of postdates pregnancy, and the strengths and limitations of each study. Assessment of fetal well-being should begin between 40 and 41 weeks of gestation and should continue until delivery.

### **Fetal Kick Count**

Thirty to sixty minutes postprandial, lying on her left side, the patient monitors and counts fetal movements (“kicks”). Normal frequency is approximately five kicks per hour. Fewer than ten kicks in 2 h is considered abnormal. Although a reasonable



**Fig. 17.1** Management of postdates pregnancy

adjunct to other methods of monitoring, fetal kick counts alone are probably insufficient to ensure fetal well-being. All reports of decreased fetal movement should be followed up by a nonstress test (NST) or a biophysical profile.

### Amniotic Fluid Volume

Amniotic fluid volume can be calculated weekly. Although not a direct measure of fetal well-being, oligohydramnios (low fluid volume) is an indication for delivery. Because the result of this test may alter clinical management, amniotic fluid volume has become standard practice.

### Ultrasound Estimate of Fetal Weight

Like the amniotic fluid volume, assessment of fetal weight is not a direct measure of fetal well-being. However, just as with amniotic fluid volume, the outcome of the study may impact management and is therefore commonly included in the

assessment of postdates pregnancies. Although ultrasound estimates of fetal weight are subject to an error of up to 500 g, estimated fetal weight may alter delivery planning. Macrosomia is an indication for possible elective caesarian section (5000 g in nondiabetic patients; 4500 g in patients with diabetes).

### **Nonstress Test**

During the NST, patients are monitored with external tocodynamometer and a Doppler fetal heart rate monitor. A reassuring NST consists of at least two accelerations in fetal heart rate in 15–20 min. Each acceleration should last at least 15 s and increase at least 15 beats per minute from baseline. (Fetal heart rate monitoring is reviewed in Chap. 26.) Nonstress testing should be repeated twice weekly beginning at 40–41 weeks' gestation.

Nonreassuring NSTs may be associated with threats to fetal well-being, may represent a period of fetal sleep, or may be related to external factors such as medication. Nonreassuring NSTs should prompt additional follow-up.

### **Contraction Stress Test**

Fetal stimulation associated with uterine contraction has been observed to induce decelerations of fetal heart rate in circumstances where fetal well-being is threatened. This observation led to the development of the contraction stress test. Oxytocin is used to induce uterine contractions. The fetal heart rate is monitored, in turn, for decelerations indicative of fetal stress. Contraindications to contraction stress test include preterm labor risk, classical cesarean section scar, and placenta previa.

Oxytocin is started at 0.5–1.0 mU per minute and increased every 15 min until a pattern of three contractions every 10 min is established. Late decelerations with 50 % or more of contractions are considered a positive test and require further evaluation. Infrequent late decelerations should prompt close monitoring and possible further evaluation. Variable decelerations should be followed by US evaluation to assess amniotic fluid status. A normal or negative test (no late or variable decelerations) is reassuring and should prompt routine fetal monitoring.

### **Biophysical Profile**

A biophysical profile is a multicomponent assessment of fetal well-being. Each of the five components is given a score of 0–2 with a maximum possible score of 10. A score of 8 is reassuring; 6 is suspicious; 4 indicates a need for acute intervention.

**Table 17.1** Biophysical profile scoring

Measure	Normal (2 points)	Abnormal (0 points)
Amniotic fluid	At least one pocket $\geq 2$ cm	No pocket $\geq 2$ cm
Fetal heart rate	Reactive two or more episodes; acceleration ( $\geq 15$ beats per minute over baseline) lasting at least 15 s within 20 min	Nonreactive
Fetal tone	Active limb extension and flexion	Slow extension, no return to flexion, no movement or partially open fetal hand
Gross movement	At least two separate limb or body movements within 30 min	Fewer than two separate limb or body movements within 30 min
Breathing movements	At least one episode lasting $>20$ s within 30 min	No episode of sufficient duration within 30 min

The scoring matrix is summarized in Table 17.1. Components of the biophysical profile include the following:

1. An NST is performed. A reassuring NST is scored 2. A nonreassuring NST is scored 0.
2. Amniotic fluid index (AFI) score is obtained. An AFI score of 5 or higher with at least one  $2\text{ cm} \times 2\text{ cm}$  pocket of amniotic fluid present is scored 2 points. AFI less than 5 or no pocket of fluid is scored 0.
3. Sustained fetal breathing is monitored. Sustained fetal breathing of at least 30 s is scored 2. Absence of fetal breathing activity or duration less than 30 s is scored 0.
4. Fetal movement is monitored. At least three limb or trunk movements are scored 2. Less than three movements is scored 0.
5. Fetal tone is measured. Fetus flexion at rest with at least one movement of extension followed by return to flexion is scored 2. Extension at rest or lack of at least one extension/flexion movement is scored 0.

## **Part III**

# **Labor and Delivery**

# Chapter 18

## Normal Labor

### Contents

Background .....	141
Prelabor .....	141
Labor .....	142
Assessment.....	143
History.....	143
Physical Examination.....	143
Laboratory Studies .....	144
Management.....	145

### Key Points

1. Labor is defined as uterine contractions resulting in progressive cervical change.
2. Assessment of labor begins with confirmation of gestational age.

### Background

The experience of labor and delivery is the culmination of the prenatal period. It is the period with the most potential for both the most joy and the most anxiety.

### Prelabor

Prior to labor there is generally a sequence of predictable events that mark the physiological preparation for delivery of the infant. Beginning 4–8 weeks prior to delivery, the patient may begin to experience slight, irregular, and non-sustained

**Table 18.1** Bishop scale cervical scoring

Indicator	Score			
	0	1	2	3
Dilation (cm)	0	1–2	3–4	5–6
Effacement (%)	0–30	40–50	60–80	80–100
Fetal station	−3	−2	−1/0	≥+1
Consistency	Firm	Medium	Soft	
Cervical position	Posterior	Mid	Anterior	

contractions. These contractions, referred to as Braxton–Hicks, are marked by only mild discomfort in most circumstances and do not lead to cervical change.

Approximately 2 weeks prior to delivery, the fetal head will often settle into the pelvic brim. This settling is referred to as lightening and the patient may report that the baby has “dropped.” There is potentially a measurable decrease in fundal height and the patient may report a decrease in pregnancy symptoms related to intra-abdominal pressure. The woman may also, however, report an increase in symptoms related to fetal pressure within the pelvis.

Beginning several days to several weeks prior to delivery, the cervix will begin to undergo preparatory changes that will include softening and may also include some degree of effacement and dilation. Dilation up to 3 cm may occur during this phase and is generally more pronounced in multiparous patients. A standardized measure of cervical condition exists and is often used in the evaluation of patients for possible induction of labor, when necessary. The Bishop scale is summarized in Table 18.1.

As the cervix begins to dilate and efface, the cervical mucus plug that has occupied the os during the course of pregnancy comes out. This is occasionally associated with a small amount of blood referred to as “bloody show.” The loss of the mucus plug and bloody show are generally signs that the onset of labor is imminent.

## Labor

Labor is divided into three separate stages, which are summarized in Table 18.2. The first stage of labor is from the onset of contractions through full cervical dilation and effacement. Because the early cervical changes may be protracted and unpredictable in their course, the first stage of labor is divided into early- or latent-phase labor and active-phase labor. Although no absolute distinction can be made between these two phases of the first stage of labor, patients are generally considered in latent-phase labor until cervical dilation reaches approximately 4 cm. The second stage of labor begins with full cervical dilation and continues until delivery of the infant. The third stage of labor begins with the delivery of the infant and is complete with the delivery of the placenta. Although the duration of each stage is highly variable, the duration tends to shorten with each subsequent pregnancy.

**Table 18.2** Stages of labor

First stage	Onset to full dilation	Primigravid 6–18 h (mean, 12)	1.2 cm/h	Effacement, dilation, station
		Multiparous, 2–10 h (mean, 6.5)	1.5 cm/h	
Second stage	Full dilation to delivery of infant	Primigravid, 1–3 h		Descent, rotation
		Multiparous, 5–30 min		
Third stage	Delivery of infant to delivery of placenta	Up to 30 min		Delivery of placenta

## Assessment

### *History*

Assessment of possible labor begins with an abbreviated history. Accurate pregnancy dating is critical to the appropriate management of labor. If uncertain dating makes preterm labor a possibility, the patient must be managed as if preterm. A complete discussion of the management can be found in Chap. 7. In addition to gestational dating, the history should include a review of the prenatal course and any complications that arose during pregnancy. Pre-existing medical conditions, including any allergies, should be reviewed. A history of the contractions should include onset, frequency, duration, and intensity. Patients should be asked about bleeding or rupture of membranes. Fetal movement should be confirmed.

### *Physical Examination*

The physical examination should include vital signs, abdominal examination of the abdomen including Leopold's maneuvers, and a clinical estimate of fetal size. A manual examination of the cervix should be performed to determine dilation, effacement, station, and presentation.

### **Dilation**

Cervical dilation is measured in centimeters and ranges from 0 (closed) to 10 (complete). Standardized instruments exist that allow providers to “feel” various degrees of dilation and providers should occasionally test their own assessment against these instruments. Hand size varies significantly, but 1 cm dilation is approximately equivalent to a fingertip. A measurement of 3 cm is approximately equivalent to two

fingers side by side. A 5 cm dilation is approximately equivalent to spreading index and middle fingers in a “victory” sign, and 10 cm is roughly equivalent to fully spread index and middle fingers.

### **Effacement**

Effacement represents thinning of the cervix over the fetal head (or presenting body part in non-vertex presentations). This can be visualized as equivalent to pulling a tight turtleneck sweater over one’s head. Effacement is described in percentages, with 0 % effacement marking no change and 100 % effacement representing no appreciable thickness to the cervix. Although dilation and effacement often occur in tandem, either may occur without evidence of the other.

### **Station**

Station refers to the position of the fetal head in the birth canal. Zero station is defined as the level of the ischial spines. Positions above the ischial spines are measured as negative values, whereas positions below the ischial spines are measured as positive values. Traditionally, the distances were divided in thirds. A fetal head one-third of the distance between the ischial spines and the outlet is +1, two-thirds +2, and at the outlet +3. The same applies in reverse for negative station measures. Some authorities now recommend measuring in centimeters from the spines, which translates to a -6 to +6 scale. Considerable head molding may occur during descent and providers should be careful to measure distance to the fetal head and not to the fetal caput.

### **Presentation**

During the examination, the presenting body part should be examined and confirmed. If a cephalic (head-first) presentation cannot be confirmed on manual examination, an ultrasound should be performed.

### **Laboratory Studies**

Prenatal labs should be reviewed and any missing laboratory values should be ordered. In particular, hemoglobin, platelets, and evidence of infection (including group B strep) should be noted. Additional labs may be required if the patient is presenting with a complication of pregnancy or labor.

## Management

In the first stage of labor, management is generally expectant. Patients should be admitted (preferably as late in the first stage as possible under most circumstances). Blood pressure should be checked every 2–4 h, and fetal heart rate should be monitored every 30 min, unless abnormalities arise. Patients may be allowed to ambulate, but intake by mouth should be limited. Adequate anesthesia should be provided at the request of the patient. Recent studies support the use of early anesthesia if requested by the patient.

The second stage of labor is marked by descent of the fetal head through the birth canal. The usual sequence of head movements is engagement, flexion, descent, internal rotation, extension, external rotation, and expulsion.

Delivery will usually occur without any necessary assistance on the part of the provider. At the time of delivery of the head, gentle counterpressure may be applied to control delivery and minimize perineal trauma. Delivery of the anterior shoulder is effected with gentle downward traction followed by delivery of the posterior shoulder in an upward motion.

With delivery of the head, a quick check is made to ensure that no nuchal cord is present and bulb suction of the infant's nose and mouth should be performed. The mother should breathe rather than push during this activity. With delivery of the body, the umbilical cord is clamped and then cut. The infant should be held above the level of the uterus while at this point to minimize transfusion of blood from the cord to the infant.

Delivery of the placenta also requires little assistance from the provider under normal circumstances. Gentle traction on the umbilical cord may assist with release, but excess traction may lead to uterine inversion and excess postpartum hemorrhage.

# Chapter 19

## Induction and Augmentation

### Contents

Background .....	147
Preparation .....	148
Pharmacological Options .....	148
Mechanical Options .....	148
Induction .....	149

### Key Points

1. Induction is defined as artificial initiation of labor.
2. Augmentation is defined as artificial stimulation of labor.
3. Induction should always be performed for a specific indication.

### Background

Under most circumstances, the initiation of labor occurs without assistance at the appropriate time. Under some circumstances, however, induction of labor is indicated prior to the onset of labor via natural processes. Table 19.1 presents a list of indications for induction. Induction should always be performed for a specific indication and, when possible, under circumstances where cervical status is favorable for delivery.

**Table 19.1** Indications for induction

Pre-eclampsia
Chronic hypertension
Diabetes mellitus
Heart disease
Postdates
Rh incompatibility
Fetal abnormalities or demise
Chorioamnionitis
Premature rupture of membranes
Intrauterine growth restriction

## Preparation

Prior to induction patients should be carefully assessed for contraindications to induction (see Table 19.2). In addition, cervical status should be assessed. A Bishop cervical score (see Chap. 18) of 8 predicts a success rate for induction approximately equivalent to spontaneous labor.

If the cervical status is not favorable for induction and the need for delivery is not immediate, cervical preparation may be a helpful adjunct in preparing for delivery. Several modalities are available to enhance favorable cervical status, including mechanical and pharmacological options.

### ***Pharmacological Options***

Prostaglandin formulations are available including PGE1 (misoprostol) and PGE2 (dinoprostone). Both mimic the prostaglandin activity physiologically present with the onset of spontaneous labor. Misoprostol is available as a pill that can be placed intravaginally (25 µg every 3–6 h). Dinoprostone is available in a gel formulation (0.5 mg/2.5 mg gel). A single dose is placed intracervically and can be repeated in 6–12 h. Contraindications to prostaglandin use include unexplained bleeding per vagina, rupture of membranes, and prior cesarean section. In addition to those general contraindications, dinoprostone is contraindicated in patients with a history of asthma, glaucoma, or myocardial infarction. Side effects include fetal stress and decelerations, hypertonicity, nausea, vomiting, and fever.

### ***Mechanical Options***

Two mechanical options are also available for cervical ripening. These are placement of a Foley bulb and hygroscopic dilators (laminaria). A Foley catheter with a 25–50 cc balloon may be inserted above the internal os, inflated, and withdrawn to the internal os.

**Table 19.2** Contraindications to induction of labor

Pelvic abnormalities incompatible with vaginal delivery
Placenta previa
Classical cesarean section scar
Full-thickness myomectomy
Hysterotomy
Breech presentation not compatible with vaginal delivery

Within 12 h, the cervix can be expected to dilate 2–3 cm. This dilation will be apparent clinically when the Foley bulb falls out of the cervix. The laminaria is placed in the cervix and as the fluid is absorbed the laminaria swells to three to four times its original size within 6–12 h.

Preparation for induction should also include preparation for failure of induction and therefore should only be performed in a setting where access to operative delivery is available.

## Induction

Induction is generally performed with oxytocin. Specific protocols vary between institutions and providers should be aware of the specific protocol at their own institution. One potential regimen is 10 units of oxytocin in 1 L fluid. The oxytocin is attached to an infusion pump and is slowly titrated up until adequate contractions are achieved or contraindications to continue induction develop. Complications include hyperstimulation with possible tetanic contractions, abruptio placentae, uterine rupture, precipitous delivery, cord prolapse, and fetal stress. Because of the potential serious nature of these side effects, patients must have vital signs monitored regularly and must have both fetal heart rate and tocodynamometer monitoring during the course of induction. Details of fetal heart rate monitoring are reviewed in Chap. 26.

# Chapter 20

## Pain Management in Labor

### Contents

Background .....	151
Nonpharmacological Management .....	152
Pharmacological Management.....	152
Narcotic Analgesia.....	152
Local Anesthesia .....	153
Epidural–Spinal Anesthesia .....	153

### Key Points

1. Pain in labor is a multifaceted experience with physiological, psychological, and social components.
2. Pain management in labor requires a multifaceted approach, including pharmacological and nonpharmacological options.
3. Appropriate pain management will greatly enhance the experience of labor.

### Background

Virtually all labor is accompanied by pain. Uterine contractions, cervical dilation, fetal descent, and perineal stretching (and, when it occurs, laceration) are all associated with pain. Although pain accompanies all labor, the patient's perception of and response to pain is highly variable. Providers must be aware of all factors that contribute to the patient's pain and adequately address each of these factors in order to achieve appropriate pain control.

Although the patient's perception of pain depends on a variety of factors, the physiological basis of pain is reasonably well described. In the first stage of labor,

most pain is secondary to uterine contractions, intermittent ischemia, and cervical dilation. The primary neurological innervation associated with these components is located at the level of T10–L1. In the second stage of labor, vaginal and perineal distentions are the primary source of pain. Innervation is at the level of S2–S4, the pudendal nerve.

Options for pain management include pharmacological and nonpharmacological modalities with which providers should be familiar. Among the nonpharmacological options are hydrotherapy, hypnotherapy, positioning, and support. Pharmacological options include epidural–spinal anesthesia, narcotic pain management, and local anesthesia.

## **Nonpharmacological Management**

Ideally, nonpharmacological modalities are a part of the management of all labor. Such management begins in the prenatal period with appropriate education concerning the process of labor and delivery and a discussion of reasonable expectations concerning pain in labor. In addition, patients will often benefit from a prenatal course in one of the many nonpharmacological approaches to pain management.

Although the specifics of each approach vary somewhat, several key principles are shared among all approaches. These principles include relaxation, suggestion, concentration, and preparation. With appropriate preparation and intrapartum support, the efficacy of nonpharmacological approaches is quite high, approaching 80 % in some populations.

Patients should be educated concerning the duration of each stage of labor, the expected events of each stage, and the active role that laboring mothers have in contributing to the success of each stage. Patients should receive education concerning preadmission management of pain, appropriate timing of presentation to the labor and delivery suite, and warning symptoms that require prompt assessment.

At the time of labor, patients are counseled to anticipate contractions, to focus on each contraction in turn, to develop a point of focus for each contraction, and to use relaxation techniques including visualization and breathing techniques. As with all skills, practice is an important component of the success of this technique and introduction should begin early in the course of pregnancy to allow for adequate preparation.

## **Pharmacological Management**

### ***Narcotic Analgesia***

Narcotic analgesics are nonspecific agents for pain management and can be successfully utilized for pain management in labor. Narcotics, in general, cross the placenta and will affect the fetus as well as the mother. Fetal side effects may include

sedation and respiratory depression. Narcotic analgesia should be used with caution late in the course of the first stage of labor to minimize these side effects in the newborn. All delivery suites should have ready access to naloxone for use in infants with evidence of narcotic-related side effects at the time of delivery.

A variety of analgesic options are available for use in labor and the specific agents used will vary from one institution to another. Providers should be familiar with the commonly employed agents at their institution. Among the acceptable agents commonly used in labor are 60 mg of codeine via intramuscular injection and 50–100 mg via intramuscular injection or 25–50 mg intravenously of meperidine.

### ***Local Anesthesia***

A variety of local anesthetic options are available and may be utilized during the course of labor. Local anesthesia is also often used postpartum prior to repair of lacerations or episiotomies. Commonly used options include tetracaine, lidocaine, and bupivacaine.

### ***Epidural–Spinal Anesthesia***

The most commonly employed pharmacological pain management modality in the United States is epidural or combined epidural–spinal anesthesia. Approximately 60 % of all laboring patients receive such anesthesia. Epidural anesthesia involves introduction of pharmacological agents directly into the lumbar epidural space via a specialized needle and catheter system. Combination epidural–spinal anesthesia involves epidural anesthesia along with the use of medication delivered to the subarachnoid space usually as a single dose. This bolus allows for relatively rapid effect. The addition of an epidural catheter allows for continued pain management over the course of labor, which as previously noted, may be protracted. A significant benefit of the combined approach is the ability of the patient to ambulate for a longer time period than with epidural anesthesia alone.

Epidural anesthesia is delivered via catheter placed at the L3–L4 epidural space. The appropriate landmarks are identified and a long epidural needle is introduced to the epidural space. A hollow needle is introduced first, followed by placement of the epidural catheter through the needle. The needle is then withdrawn and the catheter remains in place. Medication can be delivered continuously and/or as individual boluses through the catheter. Medication often includes a combination of analgesic and anesthetic agents.

Combination therapy involves introduction of the epidural needle followed by placement of a spinal needle into the subarachnoid space. A bolus of medication is delivered to the subarachnoid space. The spinal needle is withdrawn, the epidural

catheter is placed, the epidural needle is withdrawn, and epidural anesthesia may follow as needed via the epidural catheter.

Patients should be closely monitored for changes in blood pressure and all patients should be on fetal heart rate monitors and tocodynamometers. Side effects associated with epidural anesthesia include maternal fever (which is probably not infectious in nature) and a 3 % risk of placement in the subarachnoid space. Subarachnoid placement may be associated with postanesthesia headache. Recent studies have demonstrated that early placement of epidural anesthesia does not appear to be related to delay in delivery or to an increased rate of cesarean section. Although local backache may be a transient side effect of epidural anesthesia, studies have shown no increase in long-term backache measured at 12 months' postpartum.

# Chapter 21

## Operative Delivery

### Contents

Background .....	155
Forceps Delivery .....	156
Use of Forceps .....	157
Vacuum-Assist Delivery .....	157
Cesarean Section.....	158

### Key Points

1. Operative delivery is defined as any procedure undertaken to facilitate delivery of an infant.
2. Operative delivery methods include vacuum-assist delivery, forceps delivery, and cesarean section.
3. Operative delivery should be undertaken for specific indications and those indications should be specifically noted in labor record.

### Background

Although most deliveries will result in spontaneous vaginal delivery, under some circumstances, additional assistance is required to deliver the infant. Operative delivery is defined as any procedure undertaken to facilitate the delivery of the infant. These procedures may include vacuum-assist delivery, use of forceps, and cesarean delivery.

## Forceps Delivery

The use of forceps has become increasingly uncommon in obstetrics and is now relatively uncommon. The use of forceps, however, remains a critical skill in modern obstetrics and a familiarity with the indications and general use of forceps is important for all providers of obstetrical care. The use of forceps should always be preceded by an assessment of the risks and benefits measured against the possibility of a cesarean section. Because of the skill and experience required to effectively utilize forceps and the potential complications associated with inappropriate use, forceps delivery should only be attempted for specific indications, when superior alternatives are not available or have been attempted, and by a provider with experience in both the identified indication and the appropriate use of forceps.

A variety of forceps models exist and providers should be familiar with the available types and the indications for their use. In general, all forceps consist of two pieces with curved blades and locking handles. The curve of the blades is designed to accommodate the fetal head and the maternal pelvis. The blades are not interchangeable and must be positioned correctly to assist with the two principle activities of traction and fetal rotation.

Complications associated with the use of forceps include extension of the episiotomy, laceration, uterine or bladder rupture, transient facial paralysis, and intracranial damage.

The use of forceps is classified in part on the position of the infant in the birth canal. Historically, forceps have been used with fetuses in a variety of positions and in varying degrees of descent in the birth canal. Modern use of forceps is primarily limited to two areas, outlet forceps use and low forceps use.

Outlet forceps is defined as the use of forceps for an infant that is crowning with the skull at the pelvic floor. In addition, the position of the infant head must be identified with the sagittal suture in the anterior-posterior, right-left occiput anterior or right-left occiput posterior positions. Use of outlet forceps should be limited to no more than 45° of fetal rotation.

Low forceps is defined as the use of forceps for an infant whose skull has reached at least +2 station but that is not yet at the pelvic floor. Because low forceps use is, by definition, associated with less advanced infant progression through the birth canal, delivery may be associated with either less than or more than 45° of rotation.

Indications for forceps delivery include failure to progress with a prolonged second stage of labor, maternal cardiac or pulmonary disease, or nonreassuring fetal heart tracings. As previously noted, however, each of these indications should be considered in relation to the possibility of cesarean section as an alternative to forceps use.

## ***Use of Forceps***

Prior to the use of forceps, providers must first assess the adequacy of labor, maternal pelvic adequacy, and fetal position and station and must identify the specific indication for forceps use. There must be adequate uterine contractions and no evidence of cephalopelvic disproportion, and the fetal head must be at or below +2 station with an appropriate presentation.

The steps involved in the use of forceps to effect forceps delivery are as follows:

1. Identify specific indication.
2. Rule out contraindications to forceps delivery, including assessment of maternal pelvic adequacy (ischial spine prominence, sacral contour, and suprapubic arch size).
3. Assess risks and benefits of cesarean section as an alternative.
4. Determine fetal head presentation.
5. Determine fetal head station.
6. Prepare for cesarean section in case of failed forceps delivery.
7. Place forceps appropriately.
8. Gentle traction and or rotation for delivery.

## **Vacuum-Assist Delivery**

The use of vacuum assist is similar to that of forceps. Providers should be aware that the use of vacuum assist rather than forceps does not alter the necessary steps prior to delivery. Although the mechanics of placement and delivery may appear to be less complex than for forceps delivery, vacuum-assist delivery remains an operative delivery with associated risks and benefits and specific indications and contraindications.

A variety of vacuum-assist devices exist and providers should be familiar with the specific device utilized at their institution. In general, vacuum-assist devices consist of a cup applied to the fetal head, a handle for providing traction, a mechanical or electric device for producing vacuum pressure, and a meter for measuring pressure.

Indications for vacuum-assist delivery are similar to those for outlet forceps delivery. The nature of the vacuum-assist device does not allow for fetal head rotation and attempts to rotate the head may result in characteristic lacerations of the scalp. Contraindications to the use of vacuum-assist devices include cephalopelvic disproportion and abnormal presentation. As with forceps delivery, all vacuum-assist deliveries should be preceded by an assessment of the risks and benefits of cesarean section as an alternative operative option.

The steps involved in the use of vacuum-assist to effect delivery are as follows:

1. Identify specific indication.
2. Rule out contraindications to vacuum-assist delivery, including assessment of maternal pelvic adequacy (ischial spine prominence, sacral contour, and suprapubic arch size).
3. Assess risks and benefits of cesarean section as an alternative.
4. Determine fetal head presentation.
5. Determine fetal head station.
6. Prepare for cesarean section in case of failed forceps delivery.
7. Place vacuum-assist device appropriately.

Delivery with a vacuum-assist device is somewhat different than with forceps. The cup is applied over the sagittal suture approximately 3 cm in front of the posterior fontanelle. Negative pressure (vacuum pressure) is developed and gentle traction is applied with contractions. Traction should not be applied in the absence of contractions and no attempt should be made to rotate the position of the fetal head. In general, delivery should be expected within a few contractions and in no more than 30 min. If the infant has not been delivered, the attempt should be considered failed and cesarean section should be performed.

## **Cesarean Section**

Cesarean section is the operative delivery of the infant through an abdominal and uterine incision. The placenta and membranes are also delivered transabdominally. The indications for cesarean section include all instances when vaginal delivery is either contraindicated or not feasible. For a complete description of cesarean section, providers should consult a text on operative obstetrics.

**Part IV**  
**Complications of Labor and Delivery**

# Chapter 22

## Prolonged Labor

### Contents

Background .....	161
Complications of Labor .....	162
Prolonged Latent-Phase Labor .....	162
Failure to Dilate/Efface .....	163
Failure to Descend .....	165

### Key Points

1. Complications of labor include prolonged transition from latent- to active-phase labor, failure of cervical dilation, and failure to descend.
2. Diagnosis of an abnormality of labor requires a firm understanding of the normal progress of labor.
3. Each complication of labor requires individual assessment and management.

### Background

Most pregnancies will proceed with a minimum of abnormality and delivery will occur without significant complications. All deliveries have the potential for complications; providers should be aware of and prepared for the potential complications associated with delivery of the infant.

## Complications of Labor

Labor is defined as uterine contractions resulting in progressive cervical dilation, effacement, and eventual delivery of the infant. The normal course of labor is reviewed in Chap. 19. In general the progression of labor depends on three identifiable factors: adequate uterine contraction (both frequency and force), fetal size and position, and adequate pelvic anatomy to allow descent. Routine labor management includes sequential assessment of labor progress via manual examination of the cervix and presenting fetal body part. Although labor is predictable and progressive in most patients, under some circumstances the normal progression is disturbed. These may include a delay in the transition from latent- to active-phase labor, failure of cervical dilation to occur, and occurrence of dilation without fetal descent.

### ***Prolonged Latent-Phase Labor***

Latent-phase or early labor is the period marked by contractions and initial cervical dilation. The contractions are generally frequent and less strong than those of active labor and the progress of cervical dilation may be variable. Although average latent-phase labor lasts between 5 and 8 h, there is considerable variability. Often, the management of latent-phase labor occurs outside the medical facility. Ideally, patients without obstetrical complications or medical risk factors would arrive at the hospital in active labor, having self-managed the latent phase of labor.

Under some circumstances, however, patients will present for management while in latent-phase labor. When the latent phase of labor has continued significantly beyond the expected duration (>20 h in nulliparous patients and >14 h in multiparous patients), management decisions must be made. Most patients with prolonged latent-phase labor will progress to active labor and subsequent vaginal delivery.

### **History**

Management begins with a review of the patient's history. Review of the gestational age, prenatal course, and prior obstetrical history, if any, should be performed. Although most instances of prolonged latent-phase labor are idiopathic, use of sedation and alcohol and prior episodes of prolonged labor may all be associated with a prolonged latent phase.

### **Physical Examination**

Assessment of cervical status is the key physical finding. Cervical dilation, effacement, and fetal descent should all be noted. Rupture of membranes should also be noted, as management will vary if membranes are ruptured. Documentation at regular

intervals will assist in determining the rate of labor progression and the degree to which the current pregnancy deviates from the norm. As feto–pelvic disproportion may contribute to a prolonged latent-phase assessment of fetal size, presentation and pelvic adequacy should be noted.

### **Laboratory/Diagnostic Studies**

Generally, diagnostic studies are of limited value in the management of prolonged latent-phase labor. Obstetrical ultrasound may assist in assessment of fetal size and examination of pooled vaginal fluids, if any, may contribute to assessment of possible rupture of membranes. Fetal heart tones should be monitored intermittently to assess fetal well-being.

### **Management**

Most patients with prolonged latent-phase labor require no specific intervention. Of patients with latent-phase labor, 10–15 % will show little if any cervical change. These patients have not yet started true labor and may be sent home to rest or walk, with precautions concerning when to return. Rest and hydration will result in active labor in the majority of patients (80–85 %) who are kept in the hospital. A small percentage (5–10 %) will demonstrate active uterine contractions but insufficient cervical dilation. These patients may benefit from the use of oxytocin to augment labor. Patients with ruptured membranes should be admitted and monitored for signs or symptoms of infection. Details concerning the management of such patients can be found in Chap. 8.

### ***Failure to Dilate/Efface***

With the onset of active labor, most patients can be expected to follow a predictable pattern of cervical dilation and effacement. As noted in Chap. 19, expected dilation is approximately 1 cm per hour for primigravid patients and 1.2–1.5 cm per hour for multiparous patients. Total duration of active first-stage labor is approximately 10 h (6–18 h) for primigravid and 5 h (2–10) for multiparous patients. Documented failure to dilate at the expected rate despite the presence of organized uterine contractions is a second complication of labor.

The underlying etiology for failed cervical dilation is not well understood. Broadly understood, the problem may be with the fetus (size, presentation), with the birth canal (feto–pelvic disproportion), or with the uterine forces necessary to complete expulsion of the fetus. Evaluation of failure to dilate requires assessment of each of these components.

## History

The history may contribute to assessment of risk factors associated with either the fetus or the birth canal. The patient's prenatal course should be reviewed, with a particular emphasis on malpresentation and risk factors for macrosomia such as gestational diabetes. Past obstetrical history should also be reviewed for prior failure to dilate, past history of gestational diabetes, or prior macrosomic infants. Feto-pelvic disproportion is largely a diagnosis of exclusion; however, those patients with bony abnormalities of the birth canal can be expected to have recurrent difficulties.

## Physical Examination

Physical examination contributes significantly to the diagnosis and management of delayed cervical dilation. Serial cervical examination to assess dilation, effacement, and station should be performed and the results plotted on a normal labor curve. Identification of abnormal presentation may be apparent on physical examination. Abnormal presentations such as occiput posterior, brow, or face presentation occur in approximately 5 % of all deliveries and should generally be apparent on examination. Breech presentation with abnormal presenting fetal body parts may also be determined on pelvic examination. An assessment of fetal size should be performed, as ultrasound assessment of fetal size at term may be inaccurate. Although the reliability of manual assessment of pelvic adequacy has been questioned, a brief evaluation of the birth canal should also be performed as a part of the pelvic examination.

Critical to the assessment and management of prolonged dilation is an assessment of the adequacy of uterine contractions. Although external monitors may be useful for determining the frequency of uterine contractions, determination of the strength of those contractions requires the placement of an intrauterine pressure catheter (IUPC).

## Laboratory/Diagnostic Studies

In general, laboratory and diagnostic studies are limited in the management of delayed cervical dilation. An obstetrical ultrasound may assist in the assessment of fetal size or presentation.

## Management

Management of delayed cervical dilation requires assessment of which, if any, identifiable factors are contributing to the delay. The management of malpresentation is covered later. The indications for operative delivery are reviewed in Chap. 21.

In the absence of clearly contributory factors such as malpresentation or macrosomia, adequacy of uterine contractions should be assessed. Placement of an IUPC allows for calculation of the adequacy of uterine contractile activity. The most common and simplest measure of uterine activity is the Montevideo unit, measured as the increase in intrauterine pressure with contractions (maximum pressure–baseline pressure) over a 10-min period. The Montevideo units for each contraction are calculated and all contractions in a 10-min period are added together. A total of 200 Montevideo units are considered evidence of adequate uterine contractile activity.

For patients without adequate uterine contractile activity, oxytocin augmentation should be administered until adequate contractions are established. Cervical dilation should be periodically documented thereafter. Failure to dilate may be diagnosed with 2 h of adequate uterine contractions and no cervical change. If cervical dilation is occurring, management depends on the status of the infant. Slow but steady cervical dilation (with or without oxytocin augmentation) should be allowed to progress unless evidence of fetal stress is noted.

Patients for whom inadequate uterine contractile activity is the only apparent source of incomplete cervical dilation will generally have an excellent outcome, with two-thirds eventually delivering vaginally.

### ***Failure to Descend***

Despite full cervical dilation, the fetus may fail to descend through the birth canal. Although inadequate uterine contractile activity contributes to many of these cases, feto–pelvic disproportion makes up roughly half them. Evaluation is similar to that for failure to dilate. Particular attention should be paid to clinical evidence of pelvic adequacy, fetal size, and malpresentation. If clinical evidence suggests feto–pelvic disproportion, consideration should be given to cesarean section delivery. In the absence of clinical evidence of feto–pelvic disproportion, assessment of uterine activity adequacy and oxytocin augmentation, if necessary, would be indicated.

# Chapter 23

## Shoulder Dystocia

### Contents

Background.....	167
Diagnosis.....	168
Management.....	169

### Key Points

1. Clinically, shoulder dystocia may be diagnosed when delivery of the head is followed by an inability to deliver the shoulders.
2. Shoulder dystocia is a serious complication of delivery and must be managed rapidly to minimize maternal and fetal morbidity.

### Background

Shoulder dystocia is an uncommon but serious complication of delivery. Clinically, shoulder dystocia may be diagnosed when delivery of the head is followed by an inability to deliver the shoulders. Shoulder dystocia generally requires additional maneuvers to free the shoulders and effect delivery of the infant. Although the exact mechanism is not well studied, the postulated mechanism is impaction of the anterior shoulder against the maternal symphysis pubis or impaction of the posterior shoulder on the sacrum. Rarely, dystocia may be the result of or may be made worse by impaction against the soft tissue of the birth canal.

The risk for shoulder dystocia is approximately 1 in 100 for normal-size infants. A variety of risk factors (see Table 23.1) have been associated with an increased risk for dystocia, including prior history of shoulder dystocia, known anatomic abnormalities of the birth canal, gestational diabetes, postdates pregnancy, macrosomia, and protracted labor. Although most cases cannot be identified on the basis of identifiable

**Table 23.1** Risk factors for shoulder dystocia

Assisted delivery
Protracted labor
Postdates pregnancy
Macrosomia
Diabetes
Constitutional short stature
Abnormal pelvic anatomy
Prior shoulder dystocia
Prior macrosomic infant

risk factors, infant size is clearly related to an increased risk for dystocia. Macrosomic infants ( $>4000$  g) have a five- to tenfold increase in risk (absolute risk 5–9 %).

The complications of shoulder dystocia include direct trauma to the mother and/or infant, hemorrhage and, less commonly, possible complications of the delivery itself. Direct trauma to the mother may result in laceration, extension of episiotomy, and postpartum hemorrhage. Approximately 10 % of deliveries with shoulder dystocia result in postpartum hemorrhage (for management, *see* Chap. 28). Approximately 3–4 % of deliveries will result in fourth-degree lacerations or extensions of an existing episiotomy. Although the connection between birth trauma and subsequent neonatal outcomes is not clear, approximately 10 % of all deliveries complicated by shoulder dystocia will result in brachial plexus palsy. Of these, approximately 10 % will be persistent. An increased risk for clavicular fracture is also associated with shoulder dystocia.

The management of shoulder dystocia is critical to minimize the medical complications associated with its presentation. Despite appropriate management, shoulder dystocia is associated with an increased risk for clavicular fracture, humeral fracture, fetal hypoxia, and fetal death.

## Diagnosis

As noted, shoulder dystocia is a clinical diagnosis made at the time of delivery. No such diagnosis can be made prior to the delivery itself. Prenatal patients should, however, be screened for historic risk factors associated with dystocia, including gestational diabetes, prior macrosomic infant, prior shoulder dystocia, known pelvic anatomic abnormalities, or prior deliveries complicated by feto–pelvic disproportion. Risk factors from the current pregnancy should also be reviewed, including macrosomic infant, gestational diabetes, or risk factors for previously undiagnosed macrosomia, including abnormally large weight gain or abnormally large fundal height.

The diagnosis of shoulder dystocia is made at the time of delivery. Following delivery of the fetal head, the fetal shoulders are delivered via gentle downward traction. With delivery of the anterior shoulder, the posterior shoulder and the

remainder of the infant body is delivered via upward movement. When shoulder dystocia occurs, the head is delivered but delivery of the shoulder is impaired and cannot be achieved with reasonable levels of traction. On occasion, the provider may notice that the head is delivered with a contraction but subsequently retracts with the cessation of the contraction (like a turtle retracting its head back into the shell). When normal traction fails to deliver the shoulders, the diagnosis of dystocia should be made and management should be immediately instituted.

## Management

Patients at risk for shoulder dystocia should be managed from the onset of labor with the anticipation that shoulder dystocia will occur. All delivery room personnel should be aware of the risk, all necessary equipment should be available in the room, and delivery should be performed with sufficient support staff available to immediately begin management if necessary. In addition, some experts recommend immediate delivery of the anterior shoulder with the head for patients who are at high risk for shoulder dystocia. This maneuver, although widely recommended, has not been well studied in clinical trials.

With the diagnosis of shoulder dystocia, management should begin immediately, as outlined in Table 23.2. Each intervention is completed in a stepwise manner until the infant is delivered:

1. If sufficient assistance is not available in the room, help should be summoned immediately and staff should be made aware of the situation.
2. Following failure of gentle traction to deliver the shoulders, the patient should be positioned with hips flexed and abducted (McRoberts maneuver) and suprapubic pressure should be applied while the patient pushes. It should be noted that fundal pressure is contraindicated; pressure should be downward and administered just above the symphysis pubis.
3. If delivery is not achieved with these maneuvers, an episiotomy may be performed to decrease soft-tissue dystocia and facilitate delivery.

**Table 23.2** Management of shoulder dystocia

Get assistance
Flex and abduct hips
Suprapubic pressure (NOT fundal pressure)
Shoulder rotation
• Anterior shoulder forward
• Posterior shoulder backward
• Posterior shoulder forward
Reposition patient on all fours
Emergency maneuvers

4. If delivery is still not achieved, internal rotation of the infant's shoulders should be attempted. The anterior shoulder is rotated forward (pressure is applied from behind the anterior shoulder directed forward from the infant's perspective). This is followed by rotating the posterior shoulder backward (pressure on the anterior surface the shoulder directed backward from the infant's perspective). This is performed while maintaining the forward pressure on the anterior shoulder.
5. If delivery has not been achieved, then forward rotation of the posterior shoulder is attempted followed by delivery of the posterior arm/shoulder. The posterior arm is swept forward across the infant's chest, elbow flexed.
6. Repositioning the patient on all fours may sufficiently alter the position and forces to allow for delivery of the infant.
7. If all of the above have failed to deliver the infant, immediate emergency maneuvers will be required to effect delivery. These may include clavicular fracture and/or replacing the head in the birth canal followed by cesarean section (Zavanelli maneuver).

# Chapter 24

## Malpresentation

### Contents

Background .....	171
Occiput Positions .....	172
Diagnosis .....	172
Management.....	172
Nonoccipital Presentations.....	172
Breech Presentation .....	173
Compound Presentation .....	174

### Key Points

1. Normal delivery is marked by a characteristic fetal presentation and a stereotyped series of fetal repositions.
2. Failure to present in the usual occiput anterior position may lead to prolongation and complications of labor and may be incompatible with vaginal delivery.
3. Careful assessment of fetal presentation is critical to the diagnosis and management of abnormal presentations.

### Background

Although fetal position during the prenatal period is variable and subject to change (especially prior to 36 weeks' gestation), most infants will arrive head first, neck flexed with the occiput (either right or left occiput) in an anterior position. Variations from this position and presentation do occur, however, and providers should be aware of possible variants. Assessment of fetal presentation should be a routine component of late prenatal care and with all patients at the time of labor. Complications with the progress of labor, as noted earlier, should prompt reevaluation of fetal presentation and position.

## Occiput Positions

Occipital position is described in relation to the anterior surface of the mother. In the usual dorsal lithotomy position, this will place the anterior surface upward. In the normal presentation, the fetus will present vertex first with the occipital portion of the skull in an anterior location (occiput anterior; occiput toward the symphysis pubis, upward in the usual dorsal lithotomy position). The occiput may, however, be either posterior (occiput away from the symphysis or downward in the dorsal lithotomy position) or transverse (occiput horizontal or to the side in the dorsal lithotomy position). With the onset of labor, most infants will already be in the occiput anterior position. Approximately 20 %, however, will be positioned in an alternative position at the beginning of labor. For most of these infants, occiput posterior or occiput transverse is only a temporary position that will revert to occiput anterior during the course of delivery. Fewer than 5 % of infants will present with occiput posterior or occiput transverse at the time of delivery.

### ***Diagnosis***

Diagnosis of fetal position is made via manual examination at the time of delivery. Examination should reveal anterior and posterior fontanelles as well as the normal fetal suture lines. These three landmarks should be sufficient to determine the orientation of the fetal head.

### ***Management***

Occiput posterior and occiput transverse positions are both compatible with vaginal delivery although their presence is associated with a higher rate of assisted or surgical deliveries. If fetal size is within normal limits and there is no evidence of pelvic abnormalities, rotation of the fetal head may be considered. The use of forceps should be limited to patients without the above abnormalities and should only be performed by a provider with considerable experience with the use of forceps. In the absence of such an experienced provider, in the presence of macrosomia, fetopelvic disproportion, or with prolonged failure of dilation or descent, cesarean delivery may be indicated.

## Nonoccipital Presentations

Although occipital (or vertex) presentation is the most common presentation, other presentations are possible. Brow presentations (partially deflexed neck) are uncommon and are generally self-limited. Approximately 50 % of such presentations will

revert to vertex presentation with the continuation of labor. Brow presentation at the time of delivery occurs less than 1 in 1000 births. Diagnosis is made by manual examination and management consists of continued management with an expectation of vaginal delivery. If the patient has been on oxytocin, this should be discontinued to minimize the possibility of dystocia. Failure of brow presentation to revert to vertex presentation is associated with a high likelihood of dystocia and is an indication for cesarean delivery. Face presentations represent fully deflexed neck position with the face descending the birth canal in the lead position. Face presentation is more common than a brow presentation but is still uncommon, only occurring approximately 2 per 1000 deliveries. The retroflexion of the fetal neck combined with the presentation of the chin as the presenting body part significantly increases the likelihood of cephalopelvic disproportion and is not generally compatible with vaginal delivery. If the chin is in a posterior position (mentum posterior), cesarean delivery is indicated. If the chin is in an anterior position, vaginal delivery may be attempted, but such an attempt carries a higher-than-expected risk of failure and subsequent cesarean delivery.

## Breech Presentation

Although most infants will descend the birth canal in a head-first position, a small number will present with an alternative presenting part. Such abnormal positions can be determined during prenatal care and as such should be noted at all visits in the last 2 months of the prenatal course. Breech presentation at term but prior to labor may be amenable to manipulation/rotation (external cephalic version) to position the infant's head in a downward position. External version should be done at approximately 37 weeks' gestation to minimize the likelihood of reversion and should be done in a setting that allows for management of labor, including tocolysis. External version is an indication for the use of rhogam in Rh-negative patients. Between one- and two-thirds of such procedures are successful (defined as vertex presentation at the onset of labor).

Breech presentations may be classified by the presenting body part and the position of the fetal body. Frank breech implies buttock presentation with flexion at the hips and extension of the knees (in essence, folded in half at the hips with the buttock in the birth canal and the head at the uterine fundus). Complete breech is buttock presentation with both hips and knees flexed (the typical "fetal position" with buttock presentation and the head near the uterine fundus). An incomplete breech—sometimes referred to as a footling breech—implies presentation of one or more of the lower extremities. In the United States, delivery of breech presentation is generally via cesarean section, although optimal management remains controversial. If vaginal delivery is to be attempted, all of the following should be true: at or near term in labor with no evidence of fetal stress, frank breech presentation with full neck flexion, normal fetal size (2500–3800 g), no known congenital abnormalities, and no known pelvic abnormalities. Only providers with experience in the use of piper forceps and with cesarean backup ready at the time of delivery should perform such an attempt.

## Compound Presentation

Compound presentation is the presentation of an arm or leg along with the presenting part. It is most commonly associated with either small infants or large birth canals. The most common variant is presentation of a hand along with the head. Diagnosis is made on manual examination at the time of labor and management generally consists of expectant vaginal delivery. Care should be exercised to determine that the umbilical cord has not prolapsed along with the compound body part. In addition, providers should be prepared to proceed to cesarean section in the case of fetal compromise, failure of dilation or descent, or dystocia. Repositioning of the presenting body part is generally not recommended.

# Chapter 25

## Fetal Heart Rate Monitoring

### Contents

Background .....	175
Normal Fetal Heart Tracings .....	176
Evaluation of Fetal Heart Rate Baseline .....	176
Tachycardia .....	177
Bradycardia .....	177
Evaluation of Fetal Heart Rate Variability .....	177
Acceleration .....	177
Early Deceleration .....	178
Variable Deceleration .....	178
Late Decelerations .....	178

### Key Points

1. The common use of continuous fetal heart rate monitoring requires that providers be aware of the interpretation of variations in fetal heart tracings.
2. Normal fetal heart rate is 120–160 beats per minute (bpm) with evidence of short-and long-term variability.
3. Fetal heart rate acceleration must be distinguished from fetal tachycardia and is generally considered a favorable finding.
4. Abnormalities of fetal heart tracings may be related to either rate or deceleration.

### Background

The advent of electronic fetal heart rate monitoring has dramatically changed intrapartum management within the United States. The almost universal presence of such monitoring during the course of most deliveries presents providers with a variety of challenges. Controversy exists concerning the clinical benefit of

continuous electronic fetal monitoring. Such controversy, however, does not eliminate the need for obstetrical providers to be familiar with the basics of electronic fetal monitoring, normal and abnormal findings, and appropriate management for abnormal tracings.

## Normal Fetal Heart Tracings

Routine fetal heart tracing should be evaluated for baseline heart rate as well as variation from that baseline rate. Normal baseline fetal heart rate during pregnancy is between 120 and 160 bpm. The baseline heart rate may be determined by examining a fetal heart tracing of sufficient length to determine the heart rate to which the tracing consistently returns. It may be helpful to use a ruler or other straight edge along the course of a fetal heart rate tracing to help determine the baseline value. Fetal heart rate tracing should demonstrate a degree of variability over the course of time. A fetal heart rate tracing with little evidence of variability requires careful monitoring and evaluation if persistent.

Once the baseline fetal heart rate has been determined, variation from this baseline should also be noted. It should be apparent that this variation from baseline may be in either direction. Variation upward (toward higher fetal heart rate) is referred to as *acceleration*, whereas variation downward (toward lower fetal heart rates) is referred to as *deceleration*. In addition to the absolute direction of movement, a note should be made of patterns of fetal heart rate activity that may be indicative of careful follow-up or intervention. Nonreassuring fetal heart rate patterns are outlined in Table 25.1.

## Evaluation of Fetal Heart Rate Baseline

The baseline fetal heart rate may, under some circumstances, vary from the normal range of 120–160 bpm. When the baseline is determined to vary from this normal range, a careful review of potential causes should be performed.

**Table 25.1** Nonreassuring fetal heart rate patterns

Fetal tachycardia (persistently >160 bpm)
Fetal bradycardia (persistently <120 bpm)
Variable deceleration (decelerations with onset mid-contraction)
Late deceleration with or without short-term variability (decelerations with onset after peak of contraction)
Prolonged severe bradycardia (persistently or recurrently <100 bpm)
Sinusoidal pattern (smooth, rounded, wavelike pattern)

*bpm* beats per minute

### ***Tachycardia***

Tachycardia is defined as a baseline at or above 160 bpm. Mild tachycardia is defined as 160–180 bpm. Severe tachycardia is defined as more than 180 bpm. Fetal tachycardia may be associated with fetal hypoxia; maternal fever; drug or medication use; infection; fetal cardiac abnormalities; anemia; and hyperthyroidism. Persistent tachycardia should prompt review of potential causes and intervention if indicated.

### ***Bradycardia***

Fetal bradycardia is defined as a baseline at or below 120 bpm. A variety of conditions may produce bradycardia in the range of 100–120 bpm. If variability is good and no other abnormalities are noted, careful monitoring may be sufficient. Prolonged or severe fetal bradycardia may be associated with cord compression or prolapse, anesthesia, uterine tetany, or rapid descent of the fetus through the birth canal.

## **Evaluation of Fetal Heart Rate Variability**

Once the baseline fetal heart rate has been determined, variation from this baseline should also be noted. This variation from baseline may be in either direction. Variation upward (toward higher fetal heart rate) is referred to as *acceleration*, whereas variation downward (toward lower fetal heart rates) is referred to as *deceleration*. In addition to the absolute direction of movement, note should be made of patterns of fetal heart rate activity that may be indicative of careful follow-up or intervention.

### ***Acceleration***

In contrast to a persistent rise in baseline fetal heart rate (tachycardia), fetal heart rate acceleration is generally a favorable finding and may be associated with fetal stimulation (e.g., with contractions or cervical examinations). Fetal heart rate acceleration following variable deceleration (*see* below) is a common finding and is generally considered a good prognostic indicator.

### ***Early Deceleration***

Early decelerations are defined by a slow onset that coincides with the onset of contractions. The decelerations are thought to correspond to fetal head compression and are considered reassuring. The slow onset is matched by a similarly slow recovery producing a symmetric shape that corresponds with the duration of the contraction.

### ***Variable Deceleration***

As implied by its name, the onset, shape, and recovery of variable decelerations is less uniform than for either early or late decelerations. Interpretation of variable deceleration is likewise dependent on the associated clinical factors and the specific findings noted on the tracing. In general, variable decelerations have a relatively rapid onset and recovery with a shape that resembles a “V.” As noted earlier, variable decelerations are often associated with accelerations immediately preceding onset and immediately following recovery, yielding a pattern that resembles shoulders. Variable decelerations are thought to be associated with umbilical cord compression and their interpretation is therefore dependent of the potential causes of such compression. Mild decelerations are of less than 30 s in duration and are no lower than 80 bpm at their nadir. Moderate decelerations last between 30 and 60 s and reach 70–80 bpm at the nadir. Severe variable contractions last longer than 1 min and/or reach less than 70 bpm at the nadir. Several findings on the tracing are considered nonreassuring in the assessment of variable decelerations. These include increasing frequency or severity, delayed recovery, decreased variability, and loss of associated accelerations (“shoulders”).

### ***Late Decelerations***

Late decelerations are characterized by an onset at or after the peak of the associated uterine contraction. Distinguishing late decelerations from persistent variable decelerations may be difficult under some circumstances. Late decelerations are thought to be related to uteroplacental insufficiency and are often indicative of fetal hypoxia. Conditions associated with an increased risk for late decelerations include diabetes, hypertension/pre-eclampsia, and postdates pregnancy.

# Chapter 26

## Maternal Fever in Labor

### Contents

Background .....	179
Diagnosis.....	180
History .....	180
Physical Examination.....	180
Diagnostic Studies .....	180
Management.....	181

### Key Points

1. Labor may be complicated by maternal infection with associated maternal and neonatal risk.
2. Management of maternal fever requires a knowledge of the common sources of infection and appropriate antibiotic coverage for those organisms.
3. Management of maternal infection requires an awareness of the risks and benefits of antibiotic use in pregnancy.

### Background

Maternal fever ( $T \geq 38^{\circ}\text{C}$ ,  $100.4^{\circ}\text{F}$ ) during the course of labor is surprisingly uncommon given the number of potential pathogens in the genitourinary (GU) and gastrointestinal (GI) tracts and the nonsterile conditions of labor and delivery. Despite the relative rarity of maternal fever, providers should monitor maternal temperature regularly and be prepared to intervene appropriately if maternal fever develops. The use of epidural anesthesia is the most common source of fever in labor.

## **Diagnosis**

### ***History***

The prenatal history should be reviewed for prior infection as well as risk factors for intrapartum infection. In particular, the results from any recent testing for gonorrhea, chlamydia, group B streptococcus, urinary tract infection, and bacterial vaginosis should be noted. Although herpes and syphilis infections are not usually associated with fever, both should be noted if present. Prior antibiotic use should be documented.

The nature and timing of rupture of membranes should also be noted. In addition, manipulation should be reviewed including artificial rupture of membranes, frequent manual cervical checks, or placement of fetal scalp electrodes, intrauterine pressure catheters, or bladder catheter placement.

Although less common, respiratory sources should be considered in laboring patients with fever. History should also include a review of the patients' PHx and exposures prior to labor to identify additional possible etiologies.

### ***Physical Examination***

Physical examination should include documentation of maternal temperature, blood pressure, and pulse. Maternal fever may be associated with fetal tachycardia; therefore, fetal heart rate should also be documented. Examination of the lungs for abnormal lung sounds should be performed. Abdominal examination may reveal abdominal tenderness. Although vaginal discharge may be difficult to determine during labor, sterile pelvic examination may also be indicated.

### ***Diagnostic Studies***

Although not all maternal fevers are infectious, all such fevers should be assumed to be of infectious etiology until proven otherwise. Blood and urine cultures should be sent. If not previously performed, gonorrhea, chlamydia, and group B streptococcus testing should be performed. A complete blood count should be reviewed. Although the maternal white blood cell count may be elevated in pregnancy, an elevated white blood cell count in the setting of maternal fever may be indicative of infection.

## Management

As noted, fever should be assumed to be infectious until proven otherwise. Administration of appropriate antibiotic therapy may be of benefit to both the mother and the fetus. If a specific etiology is known, antibiotic choice should be dictated by the sensitivity of that infectious agent. If no specific etiology is noted, broad-spectrum antibiotic coverage sufficient to cover routine GU and GI flora should be initiated. Maternal fever in labor should prompt early and thorough evaluation of the infant following delivery.

# Chapter 27

## Postpartum Hemorrhage

### Contents

Background .....	184
Complications Causing Hemorrhage .....	184
Uterine Atony .....	184
Lacerations .....	184
Retained Placenta .....	185
Coagulopathy .....	185
Uterine Inversion .....	185
Diagnosis .....	185
Management .....	185
Laceration .....	186
Persistent Bleeding .....	186

### Key Points

1. Postpartum hemorrhage may result from lacerations, retained placenta, uterine inversion, or coagulopathy.
2. Management of postpartum hemorrhage begins prior to delivery with assessment of precedent risk factors including macrosomia, polyhydramnios, precipitous labor, grand multiparity, anesthesia, augmentation, and caesarian delivery.
3. Postpartum hemorrhage is a critical postpartum complication that requires rapid identification and management.
4. Management of postpartum hemorrhage should proceed in a stepwise manner until hemorrhage is controlled.

## Background

Postpartum hemorrhage complicates approximately 5 % of all deliveries. It is the second leading cause of maternal mortality, causing approximately one-sixth of all such deaths. All deliveries are associated with blood loss. Postpartum hemorrhage is defined as blood loss in excess of 500 cc. Actual blood loss during the course of routine delivery may exceed 500 cc if carefully measured. Although the definition of postpartum hemorrhage remains unchanged, from a practical standpoint, postpartum hemorrhage is often understood as hemorrhage that persists beyond expectation. Early postpartum hemorrhage is defined as blood loss occurring in the first 24 h postpartum. Late postpartum hemorrhage is defined as blood loss occurring between 24 h and 12 weeks' postpartum.

Following delivery of the infant, normal bleeding is controlled by separation of the placenta, uterine contraction with constriction of placental bed vessels and normal hemostatic pathways. Postpartum hemorrhage occurs when one or more of these events are disrupted. Postpartum hemorrhage may be caused by a variety of obstetrical complications including uterine atony, lacerations, retained placenta, and obstetrically related coagulopathy. Most cases of postpartum hemorrhage are caused by obstetrical complications; however, providers should also be aware that pre-existing coagulopathies may also manifest as postpartum hemorrhage.

## Complications Causing Hemorrhage

### *Uterine Atony*

Following routine delivery, myometrial contraction results in vascular constriction and control of bleeding. A variety of conditions may result in diminished myometrial contraction and subsequent uterine atony. Factors associated with an increased risk of uterine atony include (a) anatomic conditions such as leiomyosis; (b) uterine distention from such conditions as multigestation, polyhydramnios, or macrosomia; (c) labor-related factors such as prolonged or precipitous delivery; (d) management factors such as anesthesia, augmentation/induction, or caesarian delivery; (e) maternal factors such as multiparity; and (f) postpartum complications such as infection. Uterine atony is responsible for 50 % of postpartum hemorrhage cases.

### *Lacerations*

Delivery often results in trauma to the birth canal and may result in lacerations to the uterus, cervix, vagina, or perineum. Significant lacerations are associated with both precipitous and operative delivery. Although bleeding from such lacerations is generally self-limited or controlled with routine repair, lacerations are responsible for up to 20 % of postpartum hemorrhage cases.

## ***Retained Placenta***

Retained placenta represents the third significant cause of postpartum hemorrhage. Approximately 10 % of cases are related to this cause.

## ***Coagulopathy***

Although relatively uncommon, a number of obstetrical complications may lead to coagulopathy, which may in turn lead to persistent postpartum bleeding. Factors associated with coagulopathy include fetal demise, amniotic fluid embolus, pre-eclampsia/eclampsia, sepsis, and abruptio placenta.

## ***Uterine Inversion***

Under some circumstances, the uterine fundus may invert, preventing myometrial contraction and vascular constriction.

## **Diagnosis**

Diagnosis is generally straightforward and consists of persistent bleeding that exceeds expected levels following delivery. Although the exact blood loss may be difficult to quantify, any suspicion of excess blood loss should lead to an immediate investigation of potential causes. In addition, providers should have a low index of suspicion for initiating general management steps, as postpartum hemorrhage may be both rapid and severe.

## **Management**

Management of postpartum hemorrhage begins prior to delivery. Patients with pre-disposing risk factors should be identified and complications should be anticipated. For patients with significant predisposing risk factors, intravenous access and cross-matched blood products should be arranged prior to delivery. The risk of postpartum hemorrhage may also be reduced with appropriate management of delivery. The delivery should be controlled, operative deliveries should be minimized, and delivery of the placenta should be performed with gentle traction applied to the umbilical cord. Recent studies have suggested that the early administration of oxytocin (with the delivery of the anterior shoulder) may also reduce the risk of persistent bleeding.

Despite appropriate predelivery and postpartum management, postpartum hemorrhage may occur. Because postpartum hemorrhage may represent a life-threatening complication, initial steps should be taken to ensure hemodynamic stability. Blood should be typed and crossmatched. Intravenous access should be established preferably with two large-bore access sites. Patient blood pressure should be closely monitored. Significant drops in blood pressure should lead to initiation of fluid support with either intravenous fluid or blood products. In addition, appropriate labs should be sent including a complete blood count and a coagulation panel.

While performing the measures mentioned above, a review of risk factors should be performed and common causes explored. Uterine tone should be assessed. A comprehensive inspection of the perineum, vagina, and cervix should be performed. Under some circumstances, exploration of the uterine cavity (either manually or via ultrasound) may also be indicated.

## ***Laceration***

Significant lacerations will require repair. The presence of laceration does not preclude the possibility of either uterine atony or coagulopathy. Repair of lacerations does not necessarily ensure the cessation of bleeding and such bleeding must be managed as noted here if it persists.

## ***Persistent Bleeding***

Management of persistent bleeding will generally follow a stepwise approach:

1. Uterine massage: fundal massage of the uterus will often result in myometrial contraction and control of the bleeding. It should be noted that massage may need to be repeated. Alternately bimanual uterine massage (one hand in the vagina, one at the fundus) may also be effective. Simultaneous with massage, IV access should be obtained if not already done, oxygenation should be assessed, and consideration should be given to sending labs to assess coagulation and blood count status as well as type and crossmatch for possible transfusion
2. Oxytocin may be started with the delivery of the anterior shoulder. Oxytocin may be administered as either 10 U intramuscularly or 40 U in 1 L of normal saline delivered intravenously.
3. Misoprostol may be used either in conjunction with oxytocin or as a secondary agent. A single sublingual or rectal dose of 400–800 µg is recommended. Dinoprostone (20 mg vaginal or rectal suppository) may be used as an alternative to misoprostol.
4. Methylergonovine efficacy is similar to oxytocin but is associated with more significant side effects, including a significant rise in blood pressure. The usual dose is 0.2 mg intramuscular. This may be repeated after 2–4 h if necessary.

5. Carboprost (15-methyl-PGF2 alpha): 250 µg IM every 15–90 min with a maximum 2 mg dose.
6. Long-acting oxytocin analog: available outside the United States but not available in this country at this time

If bleeding persists despite the measures just given, immediate evaluation for possible surgical or embolization intervention is indicated.

# Chapter 28

## Perineal Laceration and Episiotomy

### Contents

Episiotomy .....	189
Background .....	189
Perineal Laceration .....	190
Background .....	190

### Key Points

1. Laceration and episiotomy are common complications of the delivery process.
2. Lacerations and extension of episiotomies may be minimized with careful management of delivery.
3. Laceration and episiotomy repair is an essential skill for all providers who deliver babies.

### Episiotomy

#### *Background*

Episiotomy is a planned incision of the perineum designed to facilitate delivery of the infant. Although routine episiotomy is not generally considered indicated, a variety of conditions may require episiotomy. Such conditions include shoulder dystocia, assisted delivery, or an anticipated macrosomic infant. Studies concerning the use of episiotomies to reduce the likelihood of laceration extension to third or fourth degree have shown conflicting results. The roles of episiotomies under the conditions just described, however, have generally been recognized to assist with delivery of the infant.

## Procedure

Following appropriate anesthesia (epidural anesthesia if present or local anesthesia if not), preparation is made for surgical incision of the perineum. With early crowning, a sharp incision is made through the perineal tissue. Median episiotomies are directed posterior toward the rectum with caution to avoid the anal sphincter and rectum. Mediolateral episiotomies are directed posteriorly approximately 45° left or right of midline.

## Perineal Laceration

### Background

Either with or without a planned episiotomy, delivery of an infant may result in laceration of the vagina, perineum, or rectum. Lacerations may involve the vagina, perineum, cervix, or uterus, as well as the vestibular tissue. Careful inspection of each of these areas should occur when postpartum hemorrhage persists beyond the expected interval. Perineal lacerations are graded (first to fourth degree) based on the degree of tissue involvement and the repair varies by laceration type. Generally, an episiotomy is equivalent to a second-degree laceration, but clinical conditions may require a more extensive episiotomy or secondary extension of the episiotomy may increase the degree of involvement. Repair of lacerations and episiotomies are generally similar and are summarized in Table 28.1.

**Table 28.1** Grading of vaginal/perineal lacerations

Degree	Description	Repair
First degree	Superficial laceration involving the skin (vaginal or perineal). These may also be superficial periurethral laceration	Generally no repair is necessary unless persistent bleeding from the site is noted
Second degree	Deeper laceration involving perineal tissue up to but not including the capsule of the anal sphincter	Approximation of laceration tissue with suture repair of laceration
Third degree	Laceration involving the anal sphincter but sparing the rectal mucosa	Approximation and suturing of lacerated ends of the anal sphincter followed by repair of the more superficial tissue as with second-degree laceration
Fourth degree	Laceration involving the rectal mucosa	Repair of the rectal mucosa followed by repair of the sphincter and more superficial tissue as noted above

## Diagnosis

### History

Any delivery may result in laceration; however, some deliveries may increase the risk of laceration. Rapid deliveries, especially those for which control of the exiting head or shoulders could not be maintained, increase the risk. Assisted deliveries (with either forceps or vacuum-assist devices) are often associated with laceration, are often accompanied by a planned episiotomy, and may also result in a lacerated extension of the episiotomy. Larger infants may increase the risk of laceration. Deliveries complicated by shoulder dystocia are at increased risk for episiotomy and/or laceration. Prior cesarean section increases the risk of uterine rupture/laceration.

### Physical Examination

All deliveries should be followed by thorough inspection of the outlet tract to identify any possible lacerations. Such lacerations may be present in the vagina, the perineum surrounding vestibular tissue, the cervix, or the uterus itself. Careful inspection with appropriate visualization (including retraction when necessary and appropriate lighting) will allow for determination of the presence and degree of lacerations, if any. All identified lacerations should be fully inspected to determine the full extent of tissue damage. This includes both the depth of involvement and the length of the laceration.

### Management

Management of a laceration depends on the location and degree of tissue involvement. General principles of management are included in Table 28.1.

*First-Degree Lacerations:* first-degree lacerations will rarely require repair. Careful inspection should be performed to determine that persistent bleeding does not occur at the site, however.

*Second-Degree Lacerations:* second-degree lacerations will often require repair. Once the extent of the laceration is determined, the area is infiltrated with local anesthesia such as 1 % plain lidocaine. Anatomic approximation of the lacerated tissue is critical, although exact approximation may be difficult owing to uneven, irregular, or damaged tissue margins. Repair is usually performed with medium-weight absorbable suture. Repair begins above the apex of the laceration and proceeds toward the vaginal opening to the hymeneal ring. Deep tissue of the perineum between the hymeneal ring and the rectum is then approximated, followed by repair of the superficial tissue and skin.

*Third-Degree Lacerations:* these lacerations will require repair in almost all cases. The first step involves identification of the lacerated ends of the anal sphincter. Once the ends are secured, repair of the anal sphincter and capsule is performed. The remainder of the repair is similar to that of a second-degree laceration

*Fourth-Degree Lacerations:* fourth-degree lacerations are the most significant of the perineal lacerations, with the highest likelihood of both short- and long-term complications. For this reason, repair of fourth-degree lacerations should only be performed by providers with considerable experience and expertise. Consultation with an experienced provider is recommended if personal experience is limited. Repair begins with repair of the rectal laceration, proceeds to repair of the anal sphincter, and is completed with the repair described for second-degree lacerations.

## **Part V**

# **Postpartum Management**

# Chapter 29

## Newborn Evaluation

### Contents

Background .....	195
The Examination .....	196
History .....	196
Physical Examination .....	196
Laboratory and Diagnostic Studies .....	198
Vaccination .....	198

### Key Points

1. The newborn examination forms the basis for all subsequent management. It is therefore comprehensive in nature.
2. The newborn evaluation includes a review of prenatal and peripartum history, as well as newborn nursery course and physical examination.

### Background

The initial newborn examination occurs immediately postpartum and will be repeated each day of the newborn's hospital stay. This examination forms the basis for all subsequent management by providing an assessment of development and congenital abnormalities, if any. This examination is therefore comprehensive in nature.

## The Examination

### *History*

The newborn history consists primarily of a review of the prenatal and delivery course, including complications, if any. Particular attention should be made of the family history of congenital abnormalities, maternal medical conditions, and prenatal exposures including infection, medications, tobacco, alcohol, and illicit drugs.

### *Physical Examination*

As noted, the newborn physical examination will serve as the baseline comparator for all subsequent examinations. It should, therefore, be comprehensive, detailed, and guided by an understanding of the most common areas of abnormality.

### **Vital Signs**

Vital signs include temperature, pulse, respiratory rate, length, weight, and head circumference. Temperature can be checked in a variety of locations and the specific location should be noted along with the reading. Pulse and respiratory rate are both measured most accurately with the infant resting quietly, preferably in a parent's arms or lap. Length is often most easily measured by marking the disposable paper on the exam table. A mark can be made at the crown of the head. The infant's legs can be fully extended and a mark is made at his or her heel. The infant is then removed and the distance between the two marks is recorded. The additional weight of clothing and diapers can be significant for infants, so weight should be measured with the infant fully disrobed. Head circumference is measured as the circumference from the brow (above the eyebrows) to the temple (above the ears) and around the occiput (roughly equivalent to the position of a hat band).

### **General Observation**

General observations should include whether the child appears healthy, comfortable, and normal.

### **Head and Neck**

The face should be observed for rashes. The ear canals should be checked for patency and the ears for position. Also, the preauricular pits should be noted when present. One should check the mouth and soft palate for defects and make note of

the mucosal lining for both moisture and oral thrush, if present. Both anterior and posterior fontanelles should be open. The neck should be palpated for adenopathy. When the child is gently raised from the table, the head lag should be noted.

## **Eyes**

All infants should be checked for red reflex and for normal eye movement in all directions. Reaction of pupils to light should be noted.

## **Cardiovascular**

Although it is often difficult for students to distinguish heart sounds in a rapid infant cardiac cycle, note should be made of S1 and S2 in all infants and murmurs, if present. Congenital heart defects may not be apparent at birth and may be picked up for the first time in the physician's office. Palpate peripheral pulses with particular note made of femoral pulses (both quality and symmetry).

## **Pulmonary/Thoracic**

Normal breath sounds and, if present, adventitial (rales, rhonchi, wheezes) sounds should be noted. Note should be made of the chest wall contour, especially at the sternum; the clavicle should be palpated for uneven contour, which may indicate a fracture. The provider should examine the breasts and palpate for breast tissue.

## **Abdomen**

Particular note should be made of the umbilical stump if present. This generally detaches by 2–4 weeks of age. The umbilical region should also be examined for umbilical hernia, noted as either a palpable defect below the umbilicus or as a visible bulging of the area below the umbilical stump.

## **Genital Examination**

Males should be examined for the presence of both testicles. When applicable, the site of circumcision should be inspected. In uncircumcised males, the foreskin should be retracted to examine the glans. In females, patency of the vagina should be noted. The inguinal region should be examined for the presence of congenital hernias.

## **Anus**

The anus should be checked for patency and note should be made of rashes that might represent either diaper contact dermatitis or candidiasis.

## Spine

The entire course of the spine should be examined for evidence of spina bifida. Particular attention should be paid to the upper- and lowermost portions of the spine.

## Skin

Note should be made of the tone of the skin, as well as the presence of any congenital birthmarks. Particular note should be made of the face, scalp, posterior neck, and sacral spine.

## Extremities

All extremities should be examined for symmetry and shape. Note should be made of muscle tone and symmetry of movement and normal posture. Hips should be examined for evidence of hip dysplasia via the Barlow and Ortolani tests. The Barlow test is performed with the hips flexed to 90° and adducted. Downward pressure is applied to the knees. In infants with unstable hips, an audible and/or palpable click is noted. The Ortolani test is performed with the hips flexed to 90°. The hips are then gently adducted and then abducted. Again, note is made of an audible and/or palpable click.

## *Laboratory and Diagnostic Studies*

All states mandate routine neonatal screening for a variety of abnormalities, including phenylalanine and thyroid-related disease and hemoglobinopathies. When appropriate, newborns will also be screened for syphilis and hyperbilirubinemia. Additional laboratory studies may be indicated based on the prenatal and maternal history. Most newborns will also undergo a newborn hearing screen.

## *Vaccination*

Most children will have received the first hepatitis B vaccine prior to discharge from the newborn nursery. Providers should confirm that this occurred.

# Chapter 30

## Routine Hospital Postpartum Management

### Contents

Background.....	199
Postpartum Day 1.....	200
History.....	200
Physical Examination.....	200
Laboratory Studies.....	201
Management.....	201
Postpartum Day 2.....	202
History.....	202
Physical Examination.....	202
Laboratory Studies.....	203
Management.....	203

### Key Points

1. Postpartum management serves to identify early complications, if any, of the postpartum period.
2. Postpartum management includes educational as well as medical components.
3. Preconception management may begin in the immediate postpartum period.

### Background

Immediate postpartum management falls largely within the domain of labor management and was discussed in detail in Chap. 19. The postpartum management of patients serves to identify early complications of the postpartum period as well as providing the basis for ongoing management of both the mother and new infant.

## Postpartum Day 1

### ***History***

Evaluation on postpartum day 1 should begin with a brief review of the prenatal and labor and delivery course. Particular attention should be paid to those issues that may impact immediate postpartum care. Maternal laboratory values from the prenatal period should be reviewed. Particular note should be made of Rh status, maternal infection (urinary tract infections, sexually transmitted disease, and rubella and varicella immune status). Medical complications of pregnancy, such as hypertension, diabetes, and infection, should be noted.

Complications of delivery including prolonged labor and delivery method and immediate postpartum complications such as hemorrhage, uterine atony, or maternal fever should be reviewed. The management of these conditions and current status of each should be noted.

A variety of medical conditions such as an abnormal Pap smear may be uncovered in the prenatal course of management. These conditions will usually be deferred until after pregnancy. When present, these medical conditions should be noted. In addition, the management of a variety of medical conditions may have been modified during the course of pregnancy. Examples of this might include a change in hypertension medication or the discontinuation of antiseizure medications. When such changes have been made, the postpartum history should include prepregnancy treatment management as well as the management regimen implemented during pregnancy.

The history of postpartum day 1 focuses on a few key elements. Bowel and bladder function should be reviewed. When present, pain location and severity should be noted. Patient activity, including ambulation, should be noted. Patients should be asked about bleeding, discharge per vagina, and subjective fever.

### ***Physical Examination***

The postpartum examination is comprehensive but focuses primarily on a few key elements. Vital signs should be reviewed for maternal tachycardia and blood pressure. In patients for whom fluid status is being monitored (e.g., those with pre-eclampsia or those who are postoperative), weight should be recorded daily. Cardiovascular examination should include a note of cardiac murmurs. Pulmonary examination should note the presence of rales, rhonchi, or wheezing.

The most important elements of the physical examination are the abdomen and the perineum. On abdominal examination, note should be made of the size and consistency of the uterus as well as any tenderness if present. In general, the uterus should feel firm and the fundal height should be at or below the umbilicus. For patients who underwent operative delivery, note should be made of the surgical

wound status. In the perineum, the external genitalia should be examined for swelling and tenderness. For patients who had an episiotomy or laceration, the site of the repair should be noted.

On postpartum day 1, all patients will have both bleeding and discharge per vagina. Note should be made of the quantity and quality of the bleeding. Although the distinction between normal and abnormal bleeding postpartum is sometimes difficult to make, normal postpartum bleeding should, in general, be no more than heavy menstrual bleeding. Clots may be noted in the early postpartum course but should resolve relatively quickly. In addition to bleeding, all postpartum patients will have a normal discharge per vagina referred to as lochia. Lochia represents a mixture of decidua tissue and blood in varying contents that changes in a predictable manner over time. In the first 3–4 days postpartum, this lochia includes considerable red blood cells and therefore appears red (lochia rubra). On postpartum day 3 or 4, as bleeding diminishes, the lochia becomes more pale or straw colored. This is referred to as lochia serosa. As the quantity of discharge diminishes and the presence of leukocytes increases, the lochia becomes clear (lochia alba). Lochia alba is variable in quantity and may last up to 8 weeks postpartum.

## ***Laboratory Studies***

As noted, all prenatal laboratory values should be reviewed. In the postpartum period, key laboratory values include hemoglobin or hematocrit (to assess for anemia) and rapid plasma reagent test (if not performed near term). In patients with fever or other signs of possible infection, a complete blood count with differential may assist in evaluation. White blood cell count must be interpreted with caution as it is often elevated in the postpartum period even in the absence of infection. In addition, urine, blood, and wound cultures should be obtained when appropriate.

## ***Management***

In addition to those items just noted, management will consist of assessment of and education concerning newborn care. Newborns can be expected to perform five basic functions: sleeping, eating, crying, urinating, and defecating. In addition to noting the presence or absence of each, providers should educate new parents about normal expectations for each, signs or symptoms of concern in each area, and appropriate follow-up for such warning signs. Although it is beyond the scope of this chapter to discuss each of these in detail, a few basic facts should be noted. All infants can be expected to urinate within the first few hours of life and note should be made of the number of wet diapers. If questions arise concerning the adequacy of urine output, these diapers can be weighed to determine the quantity of urine produced. All infants can be expected to stool prior to discharge, although they

may not have done so by the time of the first postpartum rounds. The initial stools consist of meconium, a grainy, green material with little or any odor. These will gradually transition to more typical stools as the infant increases oral intake of either breast milk or formula.

All infants cry and parents should be made aware of the fact that this is neither abnormal nor of significant concern as long as the infant can be consoled and assessment is made that possible infant needs (hunger, stool, comfort) are met. Infants can be expected to sleep up to 18 h each day. Parents should be made aware, however, that this likely represents short periods of sleeping interspersed with periods of being awake. That is to say, it will not feel to most parents as if their newborn is sleeping most of the time. Parents should be educated to sleep when their infants sleep in anticipation of being awake at times when they might not usually anticipate being awake.

In general, all mothers without a specific contraindication to breastfeeding should be encouraged to breastfeed their newborn. This may begin in the immediate postpartum period while the patient is still in the delivery room setting. Breastfeeding itself stimulates the production of breast milk and earlier initiation enhances the likelihood of success. In the first several days, the principle available breast product is colostrum rather than breast milk. Parents should be assured that this is sufficient for most infants' needs in the period prior to the onset of breast milk production. New mothers may benefit from the assistance of skilled teaching in the breastfeeding technique by nursing staff, specialized lactation consultants, physicians, or other family members with breastfeeding experience.

## **Postpartum Day 2**

### ***History***

For most patients with uncomplicated vaginal deliveries, postpartum day 2 represents the day of discharge to home. In addition to the elements just reviewed, providers should inquire about arrangements for transportation home (many institutions will not allow discharge of an infant unless the parents have a car seat) and arrangements at home.

### ***Physical Examination***

The examination on postpartum day 2 is similar to that of the first day. Abnormal findings noted on day 1 should be reviewed and particular attention should be paid to those areas on day 2.

### ***Laboratory Studies***

In general, there are no additional laboratory studies necessary on postpartum day 2, unless prior abnormal values require follow-up.

### ***Management***

As noted, postpartum day 2 is often the day of discharge from the hospital. Management should focus on transition of care from the hospital setting to the home setting and any needs the patient may have in arranging for this transition.

# Chapter 31

## Complications of the Hospital Postpartum Period

### Contents

Background .....	205
Fever .....	206
Infection .....	206
Endometritis .....	206
Urinary Tract Infections .....	207

### Key Points

1. Fever in the postpartum period may be a sign of significant infection.
2. Although maternal temperature elevation may be normal in the first 24 h, persistent, markedly elevated, or unexplained fevers must be evaluated.

### Background

For most pregnancies, the postpartum period is uncomplicated and the routine management is discussed in Chap. 30. Occasionally, however, the hospital postpartum period is complicated by a variety of developments. Among these complications, postpartum hemorrhage, fever, and infection are the most common. Postpartum hemorrhage is discussed in Chap. 28. This chapter focuses on a discussion of an approach to the postpartum patient with fever and infection.

## Fever

A variety of conditions may contribute to an elevated maternal temperature, especially in the first 24 h following delivery. Many of these conditions do not require specific intervention and are not considered a true fever. Maternal fever in the postpartum period is defined as a temperature of 38 °C (100.4 °F) on two occasions at least 24 h apart.

True maternal fever is relatively common, occurring in up to 10 % of all pregnancies. Risk factors for postpartum fever include preceding maternal infection, prolonged rupture of membranes, and operative delivery. Although most postpartum fever will have an identifiable and treatable etiology, postpartum infection contributes to approximately 5 % of maternal deaths.

When infection is identified, most consist of mixed anaerobic and aerobic flora of the genitourinary and gastrointestinal tracts. Common organisms include staphylococcus, streptococcal species, *E. coli*, gonorrhea, gardnerella, and mycoplasma. Because of the local trauma inherent in delivery, most women will demonstrate asymptomatic colonization in the immediate postpartum period. As noted, however, true infection occurs in only 10 % or less of women. Sterility of the intrauterine cavity can generally be demonstrated within 1 month of delivery.

## Infection

Postpartum infection is generally one of three conditions: endometritis, urinary tract infection (UTI), or wound infection of the laceration or episiotomy. Of these, endometritis is, by far, the most common. UTI and wound infection are considerably less common, although each must be considered in the postpartum patient with true fever.

### *Endometritis*

Endometritis is defined as an infection of the endometrium. As noted earlier, colonization of the endometrium following delivery is almost universal, but infection following routine delivery is considerably less frequent (~10 %). Risk factors that increase the likelihood of endometritis include operative delivery, prolonged rupture of membranes, prolonged labor, pre-existing maternal infection including preceding chorioamnionitis, manual manipulation, and internal monitoring during pregnancy (intrauterine pressure catheter or fetal scalp electrode).

The diagnosis is suspected in the presence of maternal fever with foul-smelling lochia. The uterus is generally soft and tender and the patient may demonstrate cervical motion tenderness. Laboratory studies include a complete blood count with

differential, blood cultures and a urinalysis (to exclude the possibility of an UTI). The white blood cell count may be elevated with endometritis, but patients without infection may also demonstrate elevations in white blood cell count up to 20,000. Bacteremia is present in up to 10 % of patients with endometritis, making blood cultures useful if positive but less helpful if negative.

In general, endometritis may be caused by a variety of aerobic and anaerobic streptococcal species, Gram-negative coliforms, chlamydia, or mycoplasma. Treatment should include broad-spectrum intravenous antibiotics with activity against suspected organisms. One such regimen would include clindamycin (Gram-positive and anaerobic coverage) plus an aminoglycoside (Gram-negative coverage) plus ampicillin (if *Enterococcus* is suspected).

### ***Urinary Tract Infections***

The combination of considerable manipulation and bacterial seeding with the urinary stasis associated with late-stage pregnancy makes UTI in the postpartum period a relatively common occurrence. Approximately 3 % of pregnancies will be complicated by postpartum UTI. Diagnosis is made via urinalysis and urine culture. Treatment is directed by culture results or presumptive treatment for common pathogens (*E. coli*, *Proteus mirabilis*, *S. saprophyticus*). UTI diagnosis and management are discussed in detail in Chap. 13.

# Index

## A

Abdomen  
examination  
    premature rupture of membranes, 67  
newborn, 197  
pain with maternal infection, 103  
postpartum, 200

Abortion. *See also* Recurrent spontaneous abortion; Spontaneous abortion  
complete, 78  
    management, 80  
threatened, 78  
    management, 80

Abruptio placenta, 87–89  
    history, 88  
    laboratory studies, 88  
    management, 88–89  
    ultrasound, 88

ACE. *See* Angiotensin-converting enzymes

Acetaminophen  
    pregnancy, 30, 34

Adventitial sounds  
    newborn, 197

AFI. *See* Amniotic fluid index

AFP. *See*  $\alpha$ -fetoprotein

Alcohol  
    IUGR, 48  
    preconception counseling, 14  
    pregnancy, 32

Aldosterone  
    pregnancy, 8–9

Allergy  
    over-the-counter medications  
    pregnancy, 34

$\alpha$ -fetoprotein (AFP), 51  
    follow-up prenatal visits, 26  
    IUGR, 49

Aluminum hydroxide  
    pregnancy, 30

Aminoglycosides  
    pregnancy, 30, 106

Amniocentesis, 41–43

Amniotic fluid  
    postdates pregnancy, 138t

Amniotic fluid index (AFI)  
    IUGR, 51

Ampicillin  
    endometritis, 207  
    PROM, 70

Analgesics  
    narcotic  
        labor pain, 152–153  
        pregnancy, 30

Anesthesia  
    epidural/spinal  
        labor pain, 153–154  
    local  
        labor pain, 153  
    spinal  
        labor pain, 153–154

Angiotensin  
    pregnancy, 8–9

Angiotensin-converting enzymes (ACE)  
    pregnancy, 30

Angiotensin receptor blockade agents  
    pregnancy, 30

Antibiotics  
    pregnancy, 30, 103t, 105–106  
    PROM, 70

Antibody screen  
  follow-up prenatal visits, 26

Antihistamines  
  pregnancy, 34–35

Anus  
  newborn, 197

Aspirin  
  pregnancy, 30, 34

## B

Bacterial vaginosis, 108

Barlow test  
  newborn, 198

β-blockers  
  pregnancy, 30

β-human chorionic gonadotropin  
  early pregnancy bleeding, 76  
  spontaneous abortion, 79

β-methasone  
  preterm labor, 62  
  PROM, 70

Biophysical profile  
  postdates pregnancy, 137–138, 138t

Bishop scoring scale  
  cervix, 142t

Bismuth subsalicylate  
  pregnancy, 30

Blood clots  
  postpartum, 200

Blood cultures  
  postpartum, 201

Blood flow  
  pregnancy, 8

Blood pressure  
  mothers  
    early pregnancy bleeding, 75–76  
  postpartum, 200

Blood sugar monitoring  
  pregestational diabetes, 125

Bloody show, 142

Body temperature  
  menstrual changes, 4

Bony pelvic configuration  
  assessment, 23

Bradycardia  
  fetus, 177

Breast  
  menstrual changes, 4  
  newborn, 197

Breastfeeding, 202

Breathing movement  
  postdates pregnancy, 138t

Breath sounds  
  newborn, 197

Breech presentation  
  Cesarean section, 173  
  complete, 173

Brow presentation, 172–173

Bupivacaine  
  labor pain, 153

## C

Calcium carbonate  
  pregnancy, 30

Calcium channel blockers  
  preterm labor, 62

Cardiac output  
  pregnancy, 7–8

Cardiac volume  
  pregnancy, 8

Cardiovascular system  
  newborn, 197

Cefotetan  
  pregnancy, 30

Cephalexin  
  urinary tract infection, 111

Cephalosporins  
  pregnancy, 30

Cervix  
  Bishop scoring scale, 142t  
  dilatation, 143  
  early pregnancy bleeding, 76  
  manual examination, 143  
  maternal infection, 102  
  menstrual changes, 4  
  mucus plug, 142  
  preparatory changes for labor, 142  
  ripening, 148

Cesarean section, 158  
  breech presentation, 173

Chlamydia, 109, 207

Chlorpheniramine  
  pregnancy, 34–35

Chlorpromazine  
  pregnancy, 30

Chorioamnionitis  
  PROM, 70

Chorionic villus sampling (CVS), 41, 43

Chorions  
  multiple  
    multigestational pregnancies, 128

Ciprofloxacin  
  pregnancy, 105

Circumcision  
  newborn, 197

Clavicle  
    newborn, 197

Clindamycin  
    bacterial vaginosis, 108  
    endometritis, 207  
    group B strep, 112  
    pregnancy, 30

Clinical pelvimetry, 23

Coagulopathy  
    postpartum hemorrhage, 185

Cocaine  
    IUGR, 48

Codeine  
    labor pain, 153

Complete abortion, 79  
    management, 80

Complete breech presentation, 173

Congenital infections  
    IUGR, 47

Constipation  
    pregnancy, 9

Contraction stress test  
    postdates pregnancy, 137

Corpus albicans, 5

Corticosteroids  
    pre-eclampsia, 119

Cough medication  
    over-the-counter  
        pregnancy, 35

Counseling. *See* Preconception counseling

Crying  
    newborn, 201

CVS. *See* Chorionic villus sampling

Cytomegalovirus infections  
    IUGR, 47

**D**

Decongestants  
    pregnancy, 35

Delivery. *See also* Operative delivery  
    estimated date of, 49  
        establishment, 133  
        prediction, 21  
    forceps, 156–157  
    multigestational pregnancy, 131  
    placenta, 145  
    vacuum-assist, 157–158  
    vaginal  
        breech presentation, 173

Dextromethorphan  
    pregnancy, 35

Diabetes, 117–122. *See also* Pregestational diabetes  
    congenital abnormalities, 122t  
    diagnosis, 122–123  
    pregnancy, 30  
    White's classification, 124t

Diabetes mellitus. *See* Gestational diabetes mellitus (GDM)

Diclectin  
    pregnancy, 30

Diet  
    pregestational diabetes, 124

Dilate/efface  
    failure to  
        prolonged labor, 163, 165

Dinoprostone  
    labor induction, 148

Diphenhydramine  
    pregnancy, 34–35

Dipstick leukocyte esterase testing urinary tract infection, 110

Dizygotic (DZ) twins, 128

Down syndrome (Trisomy 21), 42, 47

Doxylamine  
    pregnancy, 30

Dropped, 142

Drugs  
    illicit  
        IUGR, 48  
        pregnancy, 33

Ductus arteriosus  
    premature closure, 61

Dysmorphic growth/genetic abnormalities, 39–43  
    diagnosis, 41  
    history, 40–41  
    physical examination, 41  
    screening, 40t

DZ. *See* Dizygotic twins

**E**

Ear canals  
    newborn, 196

Early pregnancy bleeding, 71–80  
    ectopic pregnancy, 75–77  
    evaluation and management, 73f  
    spontaneous abortion, 78–80  
    vagina, 72–75

Early satiety  
    pregnancy, 9

Ectopic pregnancy, 75–77  
 diagnosis, 76  
 history, 76  
 laboratory studies, 77  
 management, 77–78  
 physical examination, 76  
 ultrasound, 76–77

EDD. *See* Estimated date of delivery

Efface, 144  
 failure to  
 prolonged labor, 163–165

EIA. *See* Enzyme immunoassay test

Elevated liver enzymes. *See* HELLP  
 (hemolysis, elevated liver enzymes, low platelets)

ELISA. *See* Enzyme-linked immunosorbent assay

Endometritis  
 postpartum, 206–207

Endometrium  
 menstrual cycle, 4

Epidural/spinal anesthesia  
 labor pain, 153–154

Episiotomy, 189–190  
 procedure, 190

*Escherichia coli*, 109  
 postpartum, 206, 207

Estimated date of delivery (EDD), 49  
 establishment, 134  
 prediction, 21

Estrogen  
 follicular production, 5

Exercise  
 preconception counseling, 16  
 pregnancy, 8

Extremities  
 newborn, 198

Eyes  
 newborn, 197

**F**

Face presentation, 173

Failure to descend  
 prolonged labor, 165

Failure to dilate/efface  
 prolonged labor, 163–165

Family  
 preconception counseling, 13

FAS. *See* Fetal alcohol syndrome (FAS)

FDA. *See* Food and Drug Administration

Femoral pulse  
 newborn, 197

Ferning  
 PROM, 68

Fertility  
 physiology, 5–6

Fetal alcohol syndrome (FAS), 32  
 IUGR, 48

Fetus  
 bradycardia, 177  
 developmental assessment, 25  
 growth  
 multigestational pregnancies, 130  
 heart rate  
 acceleration, 176, 177  
 baseline evaluation, 176–177  
 deceleration, 177  
 early deceleration, 178  
 late deceleration, 178  
 monitoring, 176–178  
 nonreassuring patterns, 176t  
 postdates pregnancy, 138t  
 premature rupture of membranes, 68  
 variability evaluation, 177–178  
 variable deceleration, 178

heart tracings, 176

infection, 104

kick count  
 postdates pregnancy, 135–136

lung maturity testing  
 preterm labor, 62

Rh isoimmunization, 96

risks  
 multigestational pregnancy, 128  
 size estimate during labor, 143  
 tachycardia, 177  
 tone  
 postdates pregnancy, 138t  
 tracking growth, 50f  
 well-being assessment  
 postdates pregnancy, 135–137

Fever  
 in labor  
 mothers, 180–181  
 postpartum, 206

Fibronectin testing  
 preterm labor, 59

First prenatal visits, 20–25, 20t  
 estimating gestational age, 24  
 family history, 22  
 history, 21  
 history since last menstrual period, 22  
 laboratory and diagnostic testing, 23  
 medical history, 21–22  
 menstrual history, 21

obstetrical history, 21  
physical examination, 22–23  
treatment and follow-up, 24–25

**Fitness**  
preconception counseling, 16

**Fluoroquinolones**  
pregnancy, 105

**Foley bulb**, 148

**Folic acid**  
IUGR, 48

**Folic acid supplements**  
preconception counseling, 16

**Follicles**  
menstrual cycle, 4

**Follicle stimulating hormone (FSH)**, 5, 6  
ovulation, 6

**Follow-up prenatal visits**, 25–26  
interval history and physical exam, 25–26  
interval laboratory and diagnostic studies, 26

**Fontanelles**  
newborn, 197

**Food and Drug Administration (FDA)**  
OTC pregnancy safety ratings, 33

**Forceps**  
delivery, 156–157  
low, 156  
outlet, 156

**Fosfomycin**  
urinary tract infection, 111

**Frank breech presentation**, 173

**FSH. *See* Follicle stimulating hormone**

**Fundus**  
height  
IUGR, 51

**G**

**Gardnerella**  
postpartum, 206

**Gastric emptying**  
pregnancy, 9

**Gastritis**  
pregnancy, 30

**Gastroesophageal reflux**  
pregnancy, 9

**GBS. *See* Group B strep**

**GDM. *See* Gestational diabetes mellitus**

**Genetics**  
abnormalities, 39–43  
preconception counseling, 12, 14  
screening, 40

**Genitalia**  
newborn, 197

**Gentamicin**  
pregnancy, 106  
PROM, 70

**Gestational age**  
confirmation, 62  
estimating, 24, 51  
developmental milestones, 24  
last menstrual period, 24  
physical examination, 24  
ultrasound, 24

**Gestational diabetes mellitus (GDM)**, 122, 126  
diagnosis, 123t  
follow-up prenatal visits, 26  
IUGR, 51  
management in labor, 126  
postpartum management, 126  
preconception management, 126  
during pregnancy, 126  
risk factors, 123t

**Gingiva**  
pregnancy, 9

**Glans**  
newborn, 197

**Glomerular filtration**  
pregnancy, 8

**Glucose challenge test**  
follow-up prenatal visits, 26

**Glucose tolerance testing**  
diabetes, 122, 123t

**Glucosuria**  
pregnancy, 9

**Gonadotropin-releasing hormone (GnRH)**, 6

**Gonadotropins**, 6

**Gonorrhea**, 109  
postpartum, 206

**Gram-negative coliforms**, 207

**Gravida/Para figures**, 21

**Group B strep (GBS)**, 111–112  
diagnosis, 112  
PROM, 70  
treatment, 112  
universal screening, 112

**Growth. *See also* Intrauterine growth restriction (IUGR)**  
movement  
postdates pregnancy, 138t

**Guaiifenesin**  
pregnancy, 35

**H**

**HAART. *See* Highly active antiretroviral therapy**

**hCG. *See* Human chorionic gonadotropin**

**Head**  
 delivery, 145  
 examination  
   recurrent spontaneous abortion, 94  
 newborn, 196–197  
   circumference, 196  
   lag, 197

**Heart.** *See also* **Fetus**  
 examination  
   recurrent spontaneous abortion, 94  
 pregnancy, 7–8  
 rate  
   pregnancy, 8  
 sounds  
   newborn, 197  
 tracings  
   fetus, 176

**HELLP** (hemolysis, elevated liver enzymes, low platelets), 114

**Hemoglobin**  
 follow-up prenatal visits, 26  
 pregnancy, 9

**Hemolysis.** *See* **HELLP** (hemolysis, elevated liver enzymes, low platelets)

**Hemorrhage.** *See* **Postpartum hemorrhage**

**Heroin**  
 IUGR, 48

**Hips**  
 newborn, 198

**HIV.** *See* **Human immunodeficiency virus**

**Hospital postpartum management**, 199–203  
 day 1, 200–201  
   history, 200  
   laboratory studies, 201  
   management, 201–202  
   physical examination, 200–201  
 day 2, 202–203

**Hospital postpartum period**  
 complications, 205–207

**24-hour urine collection**  
 proteinuria, 116

**Human chorionic gonadotropin (hCG)**, 41  
 early pregnancy bleeding, 77  
 first prenatal visit, 24  
 urine, 23  
 spontaneous abortion, 80

**Hypertension**, 113–119  
 diagnosis, 114–115  
 history, 115–116  
 laboratory studies, 116  
 management, 116–117  
 physical examination, 116  
 pregnancy, 30

**Hypoglycemics**  
 oral  
   pregestational diabetes, 124  
   pregnancy, 30

**I**

**Illicit drugs**  
 IUGR, 48  
 pregnancy, 34

**Incomplete abortion**, 78  
 management, 80

**Incomplete breech presentation**, 173

**Indomethacin**  
 preterm labor, 61

**Induction**, 149  
 contraindications, 149t  
 indications, 148t  
 labor, 147–149  
   mechanical options, 148–149  
   pharmacological options, 148  
   preparation, 148–149

**Insulin**  
 pregestational diabetes, 123–125  
 pregnancy, 30

**Intrauterine devices (IUD)**  
 early pregnancy bleeding, 75

**Intrauterine growth restriction (IUGR)**, 45–53  
 antenatal tracking and diagnosis, 49  
 complications, 48, 52  
 constitutional factors, 48–49  
 diagnosis, 51–52  
 fetal-genetic factors, 47  
 management, 52–53  
 maternal factors, 47–48  
 physical examination, 49–50  
 risk factors, 46t, 47–48  
 risk reduction, 52–53  
 screening, 51  
 toxic exposures, 48  
 uterine-environmental factors, 47

**Intrauterine pressure catheter (IUPC)**, 164

**Iron**  
 pregnancy, 9

**IUD.** *See* **Intrauterine devices**

**IUGR.** *See* **Intrauterine growth restriction**

**IUPC.** *See* **Intrauterine pressure catheter**

**K**

**Kick count**  
 fetus  
   postdates pregnancy, 135–136

Kidneys  
    pregnancy, 8

*Klebsiella*, 109

**L**

Labetalol  
    pregnancy, 30

Labor, 141–145. *See also* Preterm labor  
    assessment, 143–148  
    augmentation, 147–149  
    complications, 162  
    history, 143  
    laboratory studies, 144  
    management, 145  
    maternal fever, 179–181  
    multigestational pregnancy, 130  
    pain management, 151–154  
        nonpharmacological, 152  
        pharmacological, 152–153  
    physical examination, 143–144  
    prelabor, 141–142  
    prolonged, 161–165  
        failure to descend, 165  
        latent-phase, 162–163  
    stages, 143t

Lacerations  
    postpartum hemorrhage, 184, 185

Laminaria, 148–149

Last menstrual period (LMP), 19, 21  
    postdates pregnancy, 134

Late pregnancy bleeding, 81–89  
    abruptio placenta, 87–89  
    management, 83f  
    placenta previa, 85–87  
    vagina, 82–85

Lecithin-sphingomyelin ratio  
    prenatal labor, 62–63

Length  
    newborn, 196

Leopold's maneuvers, 67, 68t, 143

Leukocytes  
    pregnancy, 9

Levofloxacin  
    pregnancy, 105

LH. *See* Luteinizing hormone

Lidocaine  
    labor pain, 153

Liver enzymes. *See* HELLP (hemolysis, elevated liver enzymes, low platelets)

LMP. *See* Last menstrual period

Local anesthesia labor pain, 153

Lochia  
    postpartum, 201

Lomefloxacin  
    pregnancy, 105

Low forceps, 156

Low platelets. *See* HELLP (hemolysis, elevated liver enzymes, low platelets)

Lung maturity testing  
    fetus  
        preterm labor, 62

Luteinizing hormone (LH), 5, 6  
    ovulation, 6  
    surge, 6

**M**

MAC. *See* *Mycobacterium avium* complex

Macrolides  
    pregnancy, 30

Magnesium hydroxide  
    pregnancy, 30

Magnesium sulfate  
    eclampsia, 119  
    preterm labor, 61  
    seizure, 119

Malpresentation, 171–174  
    breech presentation, 173  
    compound presentation, 174  
    nonoccipital presentations, 172–173  
    occiput positions, 172

Maternal. *See* Mothers

Maternal triple screen. *See* Obstetrical triple screen

Medications  
    IUGR, 48  
    over-the-counter, 33  
    pregnancy, 27–35

Menarche  
    average age, 4

Menstrual cycle, 4  
    follicular phase, 5  
    luteal phase, 5  
    secretory phase, 5

Menstrual period  
    last, 19, 20  
        postdates pregnancy, 134

Menstruation  
    physiology, 4–5

Meperidine  
    labor pain, 153

Methotrexate  
    early pregnancy bleeding, 78

Methylergonovine  
  postpartum hemorrhage, 186

15-methylprostaglandin  
  postpartum hemorrhage, 186

Metronidazole  
  bacterial vaginosis, 108  
  trichomoniasis, 108

Misoprostol  
  labor induction, 148

Monozygotic (MZ) twins, 128

Mothers  
  age  
    fetal genetic abnormalities, 41  
  early pregnancy bleeding  
    blood pressure, 74  
    pulse, 74  
  fetal transfusion, 98f  
  fever in labor, 179–187  
    diagnosis, 185  
    history, 180  
    management, 185–186  
    physical examination, 180  
  infection, 102–103  
  Rh-sensitized  
    pregnancy management, 99

Mouth  
  newborn, 196

Multigestational pregnancy, 127–131  
  complications, 128t  
  diagnosis, 129–130  
  fetal risks, 128–129  
  findings, 129t  
  history, 129–130  
  labor and delivery, 131  
  laboratory and diagnostic studies, 130  
  management, 130–131  
  physical examination, 130  
  prenatal care, 130–131

Multiple chorions  
  multigestational pregnancies, 128

Multiple gestation  
  IUGR, 47

Multiple placentae  
  multigestational pregnancies, 128

Mycoplasma postpartum, 207

MZ. *See* Monozygotic twins

**N**

Narcotic analgesia  
  labor pain, 152–15  
  pregnancy, 30

Nausea  
  pregnancy, 9, 30

Neck  
  examination  
    newborn, 196–197  
    recurrent spontaneous abortion, 94

Newborn, 195–198  
  crying, 201  
  examination, 195–198  
  history, 196  
  laboratory and diagnostic studies, 198  
  oral thrush, 197  
  physical examination, 196  
  Rh isoimmunization, 96–97  
  sleep, 201  
  urine, 201  
  vaccination, 198

Nifedipine  
  preterm labor, 62

Nitrazine testing  
  PROM, 68

Nitrofurantoin  
  urinary tract infection, 111

Nonsteroidal anti-inflammatory drugs  
  (NSAID)  
  pregnancy, 30, 34

Non-stress test (NST)  
  postdates pregnancy, 136

Norfloxacin pregnancy, 105

NRTI. *See* Nucleoside reverse transcriptase inhibitors

NSAID. *See* Nonsteroidal anti-inflammatory drugs

NST. *See* Non-stress test

Nutrition  
  preconception counseling, 16

**O**

Obstetrical triple screen, 41–42  
  confirmation/follow-up, 42  
  follow-up prenatal visits, 25

IUGR, 49  
  neural tube defects, 42

Ofloxacin  
  pregnancy, 105

Oligohydramnios, 61  
  IUGR, 47

Operative delivery, 155–158  
  Cesarean section, 158  
  forceps delivery, 156–157  
  vacuum-assist delivery, 157–158

Oral hypoglycemics  
  pregestational diabetes, 123–125  
  pregnancy, 30

Oral thrush  
  newborn, 197

Ortolani test  
  newborn, 198

Outlet forceps, 156

Over-the-counter medications  
  pregnancy, 33–35

Ovulation, 6

Oxytocin  
  failure to dilate/efface, 165  
  incomplete abortion, 79  
  induction, 149  
  postdates pregnancy, 137  
  postpartum hemorrhage, 186  
  prolonged latent-phase labor, 163

**P**

Pain  
  abdominal  
    maternal infection, 103  
  management in labor, 151–154  
    nonpharmacological, 152  
    pharmacological, 152–154

Pain medications  
  over-the-counterpregnancy, 33–34

Palate  
  softnewborn, 196

PCP. *See Pneumocystis carinii* pneumonia

Pelvimetry  
  clinical, 23

Pelvis  
  assessment, 23  
  examination  
    early pregnancy bleeding, 75  
    recurrent spontaneous abortion, 94  
    urinary tract infection, 110  
  inflammatory disease  
    early pregnancy bleeding, 76

Penicillin  
  group B strep, 112  
  pregnancy, 30  
  PROM, 70

Peptic ulcer  
  pregnancy, 30

Perineum  
  lacerations  
    diagnosis, 191–192  
    grading, 190t  
    history, 190  
    management, 191  
    physical examination, 191  
  postpartum, 200

Periodontal disease  
  pregnancy, 9

Peripheral vascular resistance  
  pregnancy, 8

PGE1  
  labor induction, 148

PGE2  
  labor induction, 148

Phosphatidylglycerol  
  prenatal labor, 63

Physiology, 3–9  
  fertility, 5–6  
  menstruation, 4–5  
  pregnancy, 7–8, 7t

PI. *See Protease inhibitors*

Pituitary  
  function, 6

Placenta  
  abnormalities  
    IUGR, 47  
    delivery, 147  
    retained  
    postpartum hemorrhage, 185

Placenta previa, 85–87  
  history, 85–86  
  laboratory studies, 86  
  management, 86  
  physical examination, 86  
  ultrasound, 86

Postdates pregnancy, 133–138  
  diagnosis, 135  
  fetal well-being assessment, 135  
  history, 134  
  management, 135–138, 138t  
  physical examination, 135

Postpartum hemorrhage, 183–187  
  complications causing, 184–185  
  diagnosis, 185  
  management, 185–187  
  persistent, 186–187

Postpartum infection, 206–207  
  urinary tract, 207

Postpartum management, 195–207. *See also*  
  Hospital postpartum management  
PPROM. *See Preterm premature rupture of membranes*

Preconception counseling, 11–16  
  anatomic factors, 12  
  behavioral issues, 13  
  contents, 15t  
  education, 14  
  environment, 13  
  intervention, 16

Preconception counseling (*cont.*)  
 patient, 12–13  
 physiological factors, 12  
 screening, 14, 15  
 underlying disease, 12

Preconception management  
 pregestational diabetes, 123–124

Pre-eclampsia, 117–119  
 complications, 116t  
 management, 117–119, 118f  
 prevention, 117  
 risk factors, 115t  
 seizure prevention, 119

Pregestational diabetes, 123–125  
 blood sugar monitoring, 125  
 diet, 124–125  
 insulin, 125  
 management, 124–125  
 preconception management, 123–124

Pregnancy. *See also* Early pregnancy bleeding; Ectopic pregnancy; Late pregnancy bleeding; Multigestational pregnancy; Postdates pregnancy  
 cardiovascular changes, 7–8, 7t  
 confirmation, 18–19  
 gastrointestinal changes, 7t, 9  
 hematological changes, 7t, 9  
 human teratogens, 30–31, 31t  
 infection, 101–112  
   group B strep, 111–112  
   symptoms, 102–103  
   urinary tract, 109–111  
   vaginitis/vaginosis, 106–109  
 medications, 27–35  
   acute medical conditions, 29  
   acute obstetrical conditions, 29  
   chronic medical conditions, 28–29  
   over-the-counter, 33–35  
   safety ratings, 31t  
   therapeutic categories and  
     considerations, 30  
 outline, 12f  
 physiology, 7–9, 7t  
 renal/urinary changes, 7t, 8–9  
 vascular changes, 7

Prelabor, 141–142  
 Premature rupture of membranes (PROM), 65–70  
 abdominal examination, 67  
 ascending infection, 67  
 diagnosis, 66–68  
 external genitalia, 67  
 fetal heart rate, 68  
 history, 66–67

laboratory, 68  
 management, 69–70, 69f  
   preterm, 70  
   at term, 70  
 physical examination, 67–68  
 preterm, 57  
 risk factors, 66t  
 vital signs, 67

Prenatal visits, 17–26. *See also* First prenatal visits  
 follow-up, 25–26  
 interval history and physical exam, 25–26  
 interval laboratory and diagnostic studies, 26  
 frequency, 19–26

Presentation. *See also* Malpresentation  
 breech, 172  
 brow, 172  
 face, 173  
 labor, 144  
 vertex, 173

Preterm labor, 55–63  
 diagnosis, 58  
 environmental factors, 57  
 fetal lung maturity assessment, 62–63  
 history, 58  
 intake assessment, 58–59  
 laboratory studies, 59  
 management, 59–62  
   prior to 34 weeks gestation, 60–62  
   at 34 to 37 weeks, 55  
 patient-related factors, 57  
 periodontal disease, 9  
 physical examination, 58  
 postconception factors, 57  
 preconception factors, 57  
 risk factors, 56t

Preterm premature rupture of membranes (PPROM), 57

Progesterone  
 early pregnancy bleeding, 77  
 luteal production, 4

Prolonged labor, 161–165  
 failure to descend, 165

Prolonged latent-phase labor, 162–163  
 history, 162  
 laboratory/diagnostic studies, 163  
 management, 163  
 physical examination, 162–163

PROM. *See* Premature rupture of membranes

Proteinuria  
 test, 116

*Proteus*, 109

*Proteus mirabilis*

postpartum, 207

Pseudoephedrine  
    pregnancy, 35

Pudendal nerve, 152

Pulmonary system  
    newborn, 197

Pulse  
    femoral  
        newborn, 197

    mothers  
        early pregnancy bleeding, 74

    newborn, 197

Pupils  
    newborn, 197

Pyridoxine  
    pregnancy, 30

**Q**

Quinolones  
    pregnancy, 30

**R**

Recurrent spontaneous abortion, 91–94  
    diagnosis, 93, 94

    family history, 93

    genetic abnormalities, 93

    history, 93–94

    management, 93

    physical examination, 94

    risk factors, 92t

Red cell volume  
    pregnancy, 9

Red reflex  
    newborn, 197

Renal pelvis  
    pregnancy, 8

Renin  
    pregnancy, 8

Respiratory rate  
    newborn, 196

Retained placenta  
    postpartum hemorrhage, 185

Rh isoimmunization, 95–100  
    diagnosis, 97

    fetal consequences, 96

    management prior to, 97–99, 98f

    newborn consequences, 96–97

    physical examination, 97

Rhogam  
    incomplete abortion, 80

Rh-sensitized mothers  
    pregnancy management, 99–100

Ritodrine  
    preterm labor, 61

Rubella  
    IUGR, 47, 52

**S**

Satiety  
    pregnancy, 9

Screening  
    dysmorphic growth/genetic abnormalities, 40t

    first prenatal visit, 23

    genetics, 40

    group B strep, 122

    IUGR, 49

    preconception counseling, 14–15

Serum test  
    first prenatal visit, 23

Sexually transmitted disease (STD)  
    recurrent spontaneous abortion, 93

Sexual partners  
    early pregnancy bleeding, 76

Shoulder dystocia, 167–170  
    diagnosis, 168–169

    management, 169–170, 169t

    risk factors, 168t

Skin  
    newborn, 198

Sleep  
    newborn, 201

Smoking  
    early pregnancy bleeding, 76

    IUGR, 48

    preconception counseling, 16

Socioeconomic status  
    preconception counseling, 13

Soft palate  
    newborn, 196

Spinal anesthesia  
    labor pain, 153–154

Spine  
    newborn, 198

Spontaneous abortion, 78–80. *See also*  
    Recurrent spontaneous abortion

    diagnosis, 79

    laboratory studies, 79

    management, 80

    physical examination, 79

    risk factors, 78–79

    ultrasound, 79

Staphylococcus  
    postpartum, 206

*Staphylococcus saprophyticus*, 109  
    postpartum, 206

Station, 144  
 STD. *See* Sexually transmitted disease  
 Sternum  
     newborn, 197  
 Stools  
     newborn, 201–202  
*Streptococcus agalactiae*, 111–112  
 Sulfa drugs  
     pregnancy, 30  
     urinary tract infection, 111  
 Sulfisoxazole  
     urinary tract infection, 111  
 Sulfonamides  
     pregnancy, 105  
 Systolic ejection murmur  
     pregnancy, 8

**T**  
 Tachycardia  
     fetus, 177  
     postpartum, 200  
 Temperature  
     menstrual changes, 4  
     newborn, 196  
 Teratogens  
     pregnancy, 30–31, 31t  
 Terbutaline  
     preterm labor, 61  
 Testicles  
     newborn, 197  
 Tetracaine  
     labor pain, 153  
 Tetracyclines  
     pregnancy, 30  
 Thoracic system  
     newborn, 197  
 Threatened abortion, 78  
     management, 80  
 Thrush  
     oral  
         newborn, 197  
 TMP-SMZ. *See*  
     Trimethoprim-sulfamethoxazole  
 Tobacco  
     IUGR, 48  
     pregnancy, 32  
 Tocolysis  
     contraindications, 62  
 Toxoplasmosis  
     IUGR, 47, 52  
 Trichomoniasis, 108  
 Trimethoprim-sulfamethoxazole  
     (TMPSMZ)  
     pregnancy, 105  
 Triple screen. *See* Obstetrical triple screen  
 Trisomy, 42, 47  
 Turner's syndrome, 47  
 24-hour urine collection  
     proteinuria, 116

**U**  
 Umbilical stump  
     newborn, 197  
 Ureters  
     pregnancy, 8  
 Urinary stasis  
     pregnancy, 8  
 Urinary tract infection, 109–111  
     diagnosis, 1110  
     history, 110  
     laboratory examination, 110  
     physical examination, 110  
     treatment, 111  
 Urine  
     culture  
         postpartum, 201  
         urinary tract infection, 110  
     24-hour collection  
         proteinuria, 116  
         human chorionic gonadotropin  
             first prenatal visit, 23  
         newborn, 202  
 Uterus  
     blood flow  
         pregnancy, 8  
     measurement, 23  
     postpartum hemorrhage  
         atony, 184  
         inversion, 185  
         massage, 186  
     size during first prenatal visit, 22–23

**V**  
 Vaccination  
     newborn, 198  
     preconception counseling, 16  
 Vacuum-assist delivery, 157–158  
 Vagina  
     delivery  
         breech presentation, 173  
     discharge, 66  
     early pregnancy bleeding, 72–75

- history, 74
- laboratory studies, 75
- physical examination, 74–75
- ultrasound, 75
- lacerations
  - grading, 190t
- late pregnancy bleeding, 82–85
  - history, 83–84
  - laboratory studies, 85
  - physical examination, 84
  - ultrasound, 84
- menstrual changes, 4
- newborn, 197
- postpartum, 200
- Vaginitis/vaginosis, 106–109
  - history, 106
  - KOH whiff, 108
  - laboratory studies, 107
  - microscopic examination, 107
  - pH testing, 108
  - physical examination, 107
  - treatment, 108
- Varicella
  - IUGR, 52
- Vertex presentation, 173
- Vital signs
  - newborn, 196
- premature rupture of membranes, 67
- Vitamins
  - preconception counseling, 16
- Vomiting
  - pregnancy, 9

**W**

- Warfarin
  - IUGR, 48

**Weight**

- newborn, 196
- pregnancy, 25

**White blood cells**

- postpartum, 201
- pregnancy, 9

**Wound cultures**

- postpartum, 201

**Y**

- Yeast vaginosis, 108

**Z**

- Zavanelli maneuver, 170