




Mortality Country Fact Sheet 2006


World Health Organization

Under-5 mortality rates per 1 000 live births by sub-national region Thailand, 1987



Summary	Year	Males	Females	Both sexes
Population (millions)	2005	32	33	64
Life expectancy (years)	2004	67	73	70
Under-5 mortality (per 1 000 live births)	2004	23	20	21
Adult mortality (per 1 000)	2004	265	154	
Maternal mortality (per 100 000 live births)	2000		44	

Source: World Health Statistics 2006




About the map
 Note:


- The interval of each of the categories have been derived by taking the difference between the minimum and maximum among the regional rates and dividing it equally into 5. The formula is: (maximum of regional rate - minimum of regional rate)/5
- Rate for 5 years preceding the survey

Source: DHS Thailand 1987

Under-5 mortality



Source: i) WHO mortality database ii) World Health Statistics 2006




Note: rate for 5 years preceding the survey
 Source: DHS 1987 ,

Under-5 mortality: for highest and lowest quintiles Thailand, DHS

Wealth/assets quintiles	Lowest	Highest	Ratio

Sex	Males	Females	Ratio
	56	42	1.3
Urban/Rural	Rural	Urban	Ratio
	52	34	1.5
Mother's education quintiles	None	Higher	Ratio
	76	21	3.7

Note: rate per 1 000 live births for 10-year period preceding the survey
 Source: DHS



Note: rate for 10-year period preceding the survey
 Source: DHS



World Health Organization

Mortality Country Fact Sheet 2006

Causes of death in children under-5

Distribution of causes of death among children under 5 years of age Thailand, 2000-2003			Annual estimated proportions of death by cause for neonates Thailand, 2000		
Causes	Deaths ^b		Causes	Deaths ^c	
	(%)	Regional average (%)		(%)	Regional average ^c (%)
Total neonatal deaths	100	100	Total neonatal deaths	100	100
Neonatal causes ^a	45	44	Neonatal tetanus	1	4
HIV/AIDS	6	1	Severe infection ^a	20	27
Diarrhoeal diseases	16	20	Birth asphyxia	27	23
Measles	0	3	Diarrhoeal diseases	1	3
Malaria	0	1	Congenital anomalies	11	6
Pneumonia	11	18	Preterm birth ^b	33	30
Injuries	5	2	Others	8	7
Others	16	10			

a. Includes diarrhoea during neonatal period
b. Sum of individual proportions may not add up to 100% due to rounding.

a. Includes deaths from pneumonia, meningitis, sepsis/septicaemia and other infections during the neonatal period.
b. Includes only deaths directly attributed to prematurity and to specific complications of preterm birth such as surfactant deficiency, but not all deaths in preterm infants.
c. Sum of individual proportions may not equal 100% due to rounding.

Causes of Death

Top ten causes of death, all ages Thailand, 2002				Life expectancy at birth among males (years) Thailand, 2004	
Causes	Deaths		Years of Life Lost (%)	30 40 50 60 70 80 90 100	
	(000)	(%)		WHO South-East Asia Region	Thailand
All causes	419	100	100	65	70
HIV/AIDS	56	14	23		
Ischaemic heart disease	28	7	4		
Cerebrovascular disease	24	6	3		
Diabetes mellitus	19	5	3		
Road traffic accidents	18	5	7		
Chronic obstructive pulmonary disease	17	4	1		
Liver cancer	13	3	2		
Lower respiratory infections	12	3	4		
Nephritis and nephrosis	11	3	2		
Trachea, bronchus, lung cancers	9	2	1		

Source: [Death and DALY estimates by cause, 2002](http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls)
<http://www.who.int/entity/healthinfo/statistics/bodgbddeathdalyestimates.xls>

Life expectancy at birth among females (years) Thailand, 2004	
30 40 50 60 70 80 90 100	
WHO South-East Asia Region	65
Thailand	70

Maternal mortality ratio (per 100 000 live births) Thailand, 2000	
0 500 1000 1500 2000	
WHO South-East Asia Region	500
Thailand	100

Legend:
 WHO South-East Asia Region
 Thailand

Source: World Health Statistics 2006