

SHORT COMMUNICATION

SEA-LEVEL PRESSURE OBSERVATIONS FROM DUTCH SHIPS 1854–1938 INCORPORATED IN COADS RELEASE 1C CLIMATOLOGY

H. WALLBRINK,* F. KOEK, G. P. KÖNNEN and T. BRANDSMA

Royal Netherlands Meteorological Institute (KNMI), De Bilt, The Netherlands

Received 9 August 2002

Revised 20 January 2003

Accepted 20 January 2003

ABSTRACT

The quality of the 5.7 million punched sea-level pressure observations from Dutch ships for 1854–1938 has been under debate due to ambiguities in the gravity correction. The observations were omitted from the 1985 COADS (Comprehensive Ocean–Atmosphere Data Set) Release 1 monthly summaries. We re-examined the Dutch data with the help of original meteorological ship logs. It is concluded that the Dutch procedure for pressure reduction was consistent throughout time. The Dutch data are incorporated in the recent (2001) COADS Release 1c. This increases the number of COADS pressure observations for 1854–1938 by about 28%. For the 1854–80 subperiod the increase is more than a factor two. Copyright © 2003 Royal Meteorological Society.

KEY WORDS: historical ship data; sea-level pressure; data bases; COADS; marine data; Dutch ship observations

The pre-1939 meteorological data in the Comprehensive Ocean–Atmosphere Data Set (COADS; Woodruff *et al.*, 1987, Elms *et al.*, 1993; recently renamed I-COADS (Diaz *et al.*, 2002)) consists, for a substantial part, of observations from Dutch vessels. They are part of the Dutch contribution to historical data of COADS for that period, which is a big set of 1854–1938 data that were chiefly taken from Dutch ships. This Dutch data set is called COADS Deck 193. For the period 1854–1938, Deck 193 represents 22% of the COADS observations; for the subperiod 1854–80 the Dutch contribution is almost 60% (Figure 1). However, in the 1985 version of COADS (known as Release 1), the 5.7 million Dutch sea-level pressure (SLP) observations of Deck 193 were omitted from the $2^\circ \times 2^\circ$ monthly summary trimmed groups (MSTG). The reason was that only standardized pressure data were considered at the time in COADS (see Woodruff *et al.* (1998)), whereas the Dutch data of Deck 193 were not corrected for gravity. As the inclusion of the Deck 193 data would significantly improve the temporal and spatial density in the COADS pressure grid boxes, it was considered in the late 1990s to incorporate them in the next (2001) COADS update of historical data, now known as COADS Release 1c.

By the end of 1999 it was reported (T. Basnett, personal communication) that the Deck 193 pressure data in the $20^\circ\text{S}–20^\circ\text{N}$ belt exhibited a strange behaviour with respect to the 1961–90 climatology of the Hadley Centre global mean SLP data set (version GMSLP2.1f; Basnett and Parker, 1997). The bias of the Deck 193 data with respect to the mean 1961–90 pressure over the entire $20^\circ\text{S}–20^\circ\text{N}$ belt (1012.0 mbar) was reported to be time dependent. This raised suspicion about the consistency over time of the correction procedures applied to the raw logbook data, particularly with regard to the gravity correction. An irreparable offset would seriously degrade the value of the Deck 193 data. A check by us with the raw Deck 193 data confirmed the

*Correspondence to: H. Wallbrink, Royal Netherlands Meteorological Institute, PO Box 201, 3730 AE De Bilt, The Netherlands; e-mail: Hendrik.Wallbrink@knmi.nl




Figure 1. Annual total number of observations and annual number of Dutch observations (Deck 193) in the COADS database 1850–1940

time dependency of the bias. In Figure 2, the dashed line shows that the Deck 193 offset with respect to the 1012.0 mbar level fell by a factor two between 1871 and 1890, after which it stabilized around a value that roughly corresponds to the magnitude of the gravity correction (2.5 mbar if averaged over 20°S–20°N).

It is well documented that the Dutch, from the very start of the meteorological data acquisition in the 19th century onwards, routinely never applied the gravity correction to the SLP readings, either for land station data or for ship data (Onnen, 1844; KNMI, 1941). Only after the approval of the recommendations of the WMO 1929 conference in Copenhagen about standardization of SLP data (Secrétariat du Comité Météorologique International, 1930), did the Netherlands switch to the present system. For marine data, this occurred in the second half of 1938.

Deck 193 is the result of a large KNMI data-punching project, which ran in the period 1935–41. The total number of observations in Deck 193 is 6.4 million, of which 5.7 million include pressure. Although the data are of multinational origin, observations from Dutch vessels make up the vast majority: 95.3% of the Deck 193 observations are from Dutch logbooks, 3.4% are from Swedish logs and the remaining 1.3% is from other countries (KNMI, 1941). In the 20°S–20°N belt, as well as in the 19th century, the fraction of Dutch observations in Deck 193 is even larger. The anomalous behaviour of the Deck 193 SLP data (Figure 2, dashed) seemed to suggest that some of the observations, particularly in the early period, had been erroneously de-corrected for gravity prior to the punching. As the fraction of non-Dutch observations in Deck 193 is so small, it was at the same time clear that the effect could never be attributed to errors in the gravitation adjustment of the foreign observations alone.

A detailed comparison of Deck 193 with the original sources is no longer possible. At the end of World War II almost all of the 20 000 Dutch meteorological ship logs from the period 1854–1938 were moved to Germany. At this present time they cannot be located and must be presumed lost (Wallbrink and Koek, 2000). Fortunately, there are 128 meteorological ship logs (205 worldwide ship voyages) covering the period 1851–60 preserved at KNMI.

We first looked into the instrument descriptions of the surviving original meteorological ship logs. All logs were found to contain a detailed section at the start of each journey describing the instrumental




Figure 2. Annual mean differences in the 20°S – 20°N belt between the 1961–90 pressure climatology and the Dutch pressure data 1854–1938, calculated by two different methods. Dashed line: average pressures over all Dutch observations in the 20°S – 20°N belt (not gravity corrected) minus 1012.0 mbar, the latter being the mean 1961–90 pressure averaged over the entire (360°) 20°S – 20°N belt. Solid line: gravity-corrected Dutch observations 1854–1938 minus mean climatological pressures 1961–90, first averaged over $10^{\circ} \times 10^{\circ}$ grid boxes, and then averaged over the sea area between 20°S and 20°N . The calculation has been done on a monthly basis, and then averaged over the calendar years. The value of the gravity correction averaged over 20°S – 20°N is indicated by a bar. The 1961–90 climatology used is the Hadley Centre GMSLP2.1f climatology (Basnett and Parker, 1997)

corrections that had to be applied to the pressure readings, such as the capillarity correction, the bias with respect to a standard barometer and the height of the instrument on board. With no exception, the instrumental sections indicate the use of a mercury barometer. This is in accordance with the formal regulations of KNMI, which forbade including readings from aneroid barometers in weather reports (KNMI, 1937).

We compared the pressure data of the logbooks with the Deck 193 data. It appeared that in all tables of daily weather an extra pressure column was added with blue pencil, next to the original pressure readings. Apparently, this column was added during the processing of the logbooks in the 1935–41 KNMI punching project. The values in the extra column turned out to match exactly with the values punched in Deck 193. By applying the temperature correction and the correction factors mentioned in the instrumental sections of the logs on the original pressure readings, we were able to reproduce exactly the values written in blue, and hence those punched in Deck 193. No evidence was found from any of the 128 logs that gravity adjustments were ever applied to the Dutch data.

After these checks, we compared the Deck 193 data directly with the 1961–90 standard normals of the GMSLP2.1f monthly climatology. For this, we first corrected all Deck 193 data for gravity, then calculated anomalies per $10^{\circ} \times 10^{\circ}$ grid box and per month and finally averaged for each calendar year the monthly anomalies over the boxes between 20°S and 20°N . In the averaging procedure, boxes with

less than three observations were excluded. As shown in Figure 2 by the solid line, we did not find evidence for a bias in the Dutch data with respect to the 1961–90 climatology. From this, together with our checks of the metadata described above, we concluded that the Deck 193 data are suitable for incorporation in COADS. The only reduction that was needed for this is to apply the gravity correction throughout.

What remains to be explained is the anomalous trend in the Dutch pressures in the initial calculation (Figure 2, dashed line). Its cause is the climatological SLP difference between the Indian Ocean and the Atlantic Ocean (Figure 3). Putting the border between the Indian and Pacific Oceans at 110°E, the mean 1961–90 pressures averaged over the 20°S–20°N belts of the Indian and Atlantic Oceans are 1011.1 mbar and 1013.7 mbar respectively (Basnett and Parker, 1997). Hence, the pressure over the Indian Ocean is 2.6 mbar lower. The completion in 1870 of the Suez Canal resulted in a gradually increasing presence in the 20°S–20°N belt of Indonesia-bound Dutch ships in the Indian Ocean relative to the Atlantic, and hence in more observations from the oceanic regions of lower pressure. This is enhanced by the accompanying shift in the Indian Ocean shipping lanes toward more northern regions, where the pressure is lower (see Figure 3). The result is a decrease of the annual Deck 193 pressure if averaged over all observations in the Atlantic and Indian Oceans. As the vast majority of the Deck 193 observations in the 20°S–20°N belt originate from voyages between Europe and Indonesia, the decrease remains apparent in the averaged Deck 193 pressure over the entire (360°) 20°S–20°N belt (see Figure 2, dashed).

The magnitude of this ‘Suez Canal effect’ on the Deck 193 pressures happens to be comparable to the magnitudes of the standard corrections that should be applied to raw mercury barometer readings, such as the temperature correction, the cistern correction, the capillarity correction or the gravitation correction (Letestu, 1966). Again, putting the border between the Indian and Pacific Oceans at 110°E, we find that, for the 1854–75 period, 57% of the annual Deck 193 SLP observations in the 20°S–20°N belt came from the Atlantic, 25% came from the Indian Ocean, and 18% came from the Pacific. For 1876–1938 these percentages are 24%, 59% and 17% respectively, indicating a reversal of the ratio Indian/Atlantic observations with respect to the earlier period. As the dashed line of Figure 2 indicates, this reversal leads to a lowering of the Deck 193 averaged pressure by 1.4 mbar.

The conclusion of this analysis is that comparison of Deck 193 with single-value climatology is not an appropriate procedure to assess the quality of the Deck 193 data. The solid line in Figure 2 demonstrates that the reported time-dependent offset in the Deck 193 data disappears if, in the comparison with the 1961–90 climatology, the spatial distribution in the climatological pressure fields is properly taken into account.




Figure 3. Mean annual pressure distribution in the 20°S–20°N belt, according to the 1961–90 Hadley Centre GMSLP2 pressure climatology (Basnett and Parker, 1997). The pressure over the Indian Ocean is, on average, 2.6 mbar lower than the pressure over the Atlantic Ocean

As the issue of the Dutch barometer corrections and the trend in the SLP bias is now settled, the previously untapped Dutch SLP data were incorporated in 2001 in COADS Release 1c.

ACKNOWLEDGEMENTS

We thank Scott Woodruff of COADS for discussions and for providing the data for Figure 1. We thank Tracy Basnett of the Hadley Centre for supplying the GMSLP2.1f data set, and for initiating this research.

REFERENCES

- Basnett TA, Parker DE. 1997. Development of the global mean sea level pressure data set GMSLP2. Hadley Centre of the UK Met Office. Climate Research Technical Note 79. (Available from the Hadley Centre for Climate Prediction and Research, Meteorological Office, London Road, Bracknell, RS12 2SY, UK.)
- Diaz H, Folland C, Manabe T, Parker D, Reynolds R, Woodruff S. 2002. Workshop on the advances in the use of historical marine climate data. *WMO Bulletin* **51**: 377–380 (also in *CLIVAR Exchanges* **25**: 71–73 (2002)).
- Elms JD, Woodruff SD, Worley SJ, Hanson C. 1993. Digitising historical records for the Comprehensive Ocean–Atmosphere Data Set (COADS). *Earth System Monitor* **4**(2): 4–10.
- KNMI. 1937. Handleiding voor het verrichten van meteorologische waarnemingen op zee. KNMI publicatie nr 118. 's Gravenhage. Algemene Landsdrukkerij (available from KNMI library).
- KNMI. 1941. Gegevens betreffende het archief maritime meteorologische waarnemingen op ponskaarten (45 rijen) volgens het Hollerith-en Powersysteem, (voorloopige uitgave). KNMI publicatie nr 123. 's Gravenhage. Algemene Landsdrukkerij (available from KNMI library).
- Letestu S. 1966. *International Meteorological Tables*. WMO-No.188.TP.94. World Meteorological Organization: Geneva, Switzerland.
- Onnen PL. 1844. Meteorologische waarnemingen te Buitenzorg op het eiland Java, Nieuwe verhandelingen der Eerste Klasse van het Koninklijk Nederlandsch Instituut van Wetenschappen, Letterkunde en Schoone Kunsten te Amsterdam, **10**: 1–36 (available from KNAW library; address: PO Box 41950, 1009 DD Amsterdam; reprint available from KNMI library, publication VII.r.7).
- Secrétariat du Comité Météorologique International. 1930. Procès = Verbaux des Seances de la Conférence Internationale des Directeurs, du Comité Météorologique International et de Diverses Commissions a Copenhague Septembre 1929, No. 3. Kemink & Zoon N.V.: Utrecht.
- Wallbrink H, Koek F. 2000. Gang van zaken 1940–1948 rond de 20000 zoekgeraakte scheepsjournalen. KNMI publicatie nr 192, Hisklim 2 (available from KNMI library).
- Woodruff SD, Slutz RJ, Jenne RL, Steurer PM. 1987. A comprehensive ocean atmosphere data set. *Bulletin of the American Meteorological Society* **68**: 1239–1250.
- Woodruff SD, Diaz HF, Elms JD, Worley SJ. 1998. COADS Release 2 data and metadata enhancements for improvements of marine surface flux fields. *Physics and Chemistry of the Earth* **23**: 517–527.