Purina® Accuration® Starter Complete WC



Specie: Beef Type of Feed: Complete Form of Feed: Textured

General Description:

A complete feed formulated to provide intake modifying properties for weaning and receiving cattle.

PURINA® ACCURATION® STARTER COMPLETE WC

COMPLETE FEED FOR STARTER FEEDLOT CATTLE
CAUTION: USE ONLY AS DIRECTED

GUARANTEED ANALYSIS

| Crude Protein, (Min) | 14.00 % |
|---------------------------------------|-----------------|
| (This includes not more than 2.30% ed | quivalent crude |
| protein from non-protein nitrogen.) | • |
| Crude Fat, (Min) | 3.00 % |
| Crude Fiber, (Max) | 20.00 % |
| Calcium (Ca), (Min) | 0.50 % |
| Phosphorus (P), (Min) | 0.50 % |
| Salt (NaCl), (Min) | 0.50 % |
| Potassium (K), (Min) | 1.00 % |
| Vitamin A, I.U./Ib, (Min) | 1700 |

NON-MEDICATED FEEDING DIRECTIONS:

Feed 2% of bodyweight per head daily along with good quality forage to weaning and receiving cattle.

CAUTION:

Store in a dry, well-ventilated area protected from rodents and insects. Do not feed moldy or insect-infested feed to animals as it may cause illness, performance loss or death.

IMPORTANT:

This product is to be fed only to the animal species as directed on this label. Follow these management practices: 1. Do not allow animals to run out of feed. 2. Feeders/bunks should be well protected and managed to prevent accumulation of fines and wet, moldy feed. 3. When making a ration change, allow 7 to 10 days for animals to adjust to the new ration. 4. Provide adequate feeder space for each animal. 5. Fermented feeds (high moisture grains and silages) should be limited to 10% of the diet for young or stressed cattle. 6. Stressed cattle should be placed in clean, uncrowded environments. 7. Provide salt free choice. 8. Provide fresh, clean water near the feeding area. In groups of cattle, there are certain animals that experience chronic bloat or other digestive disturbances and consequently, are poor performers. In addition, excess feed consumption, severe weather changes resulting in erratic feed consumption, and poorly managed feeding practices can increase the incidence of bloat in all cattle. If bloat occurs, please review feed management practices.

Available Options:

| Product | Product No. | Active Drug Ingredient(s) |
|--|--------------------------------|-----------------------------------|
| Accuration® Starter Complete WC | 00049103 – Bulk; 0049104 - Bag | None |
| *Accuration® Starter Complete WC AS77 | 00049105 – Bulk; 0049106 - Bag | Chlortetracycline, Sulfamethazine |
| *Accuration® Starter Complete WC AS140 | 3002426-501 – Bulk | Chlortetracycline, Sulfamethazine |

^{*}Must be fed according to feed label. Please see Purina® representative for information on individual product options and availability.

