7.1 DRUGS USED IN OBSTETRICS

7.1.1 PROSTAGLANDINS AND OXYTOCINS

Carboprost
250 micrograms/1mL injection

• **Dinoprostone** 3mg tablet

(Prostin E2) 1mg/2.5mL vaginal gel

2mg/2.5mL vaginal gel

• Ergometrine 500 micrograms/mL injection

• Ergometrine/oxytocin

(Syntometrine)

500 micrograms/5 units/mL injection

Gemeprost 1mg pessaries

• Oxytocin 10 units (10,000 milliunits/mL injection

See Induction of Labour guidelines on SharePoint for details of doses and administration

7.1.2 MIFEPRISTONE

Mifepristone 200mg tablets

See unit guidelines for indications and doses.

Often given in conjunction with misoprostol (see unit

guidelines)

• Misoprostol 200microgram tablets

7.1.3 MYOMETRIAL RELAXANTS

Atosiban
6.75mg in 0.6mL injection

7.5mg/mL (5mL) concentrate for intravenous infusion

• Indomethacin 100mg suppository (unlicensed use)

75mg m/r tablets (unlicensed use)

See PRETERM LABOUR MANAGEMENT GUIDELINES on SharePoint for dose and administration details

Terbutaline 500 micrograms/1mL injection

Indication: Uterine hypercontractility due to induction agents – unlicensed useSee Management of Hyperstimulation on Trove for guidelines on administration and dose

METHOTREXATE FOR ECTOPIC PREGNANCY (SINGLE DOSE REGIMEN)

Please note this is an unlicensed use of methotrexate.

Methotrexate injections are prepared by the Pharmacy Department and doses will be supplied within office hours Monday to Friday only.

See unit policy for dose and administration guidelines

CHOLESTASIS IN PREGNANCY

- Oral vitamin K from 32 weeks until delivery Phytomenadione 10mg od
- If pruritus present try (in order)
 - emollients (Cetraben)
 - antihistamine chlorphenamine
 - ursodeoxycholic acid (400mg bd)
 - dexamethasone 12mg od po (on advice of Consultant Obstetrician only)

7.2 TREATMENT OF VAGINAL AND VULVAL CONDITIONS

7.2.1 PREPARATIONS FOR VAGINAL AND VULVAL CHANGES

Refer also to HRT section 6.4.1

Topical HRT for vaginal atrophy

Vaginal oestrogen preparations are effective in the short-term treatment of urogenital symptoms. Systemic absorption can be significant and if these are to be used for more than a few weeks by a woman with an intact uterus, a progesterone should be added to treatment.

Estriol 0.01% cream
Ortho-Gynest[®] with applicator

 Estriol 500microgram pessaries Ortho-Gynest®

• Estrol 0.1% vaginal cream Ovestin®

 Estradiol 25microgram vaginal tablets Vagifem[®]

7.2.2 VAGINAL AND VULVAL INFECTIONS

Fungal infections

• Clotrimazole Topical cream 1%

Vaginal cream 10% Vaginal tablet 500mg

Other infections

Metronidazole
0.75% vaginal gel with applicators

NB. Must be prescribed as **vaginal gel** on presriptions to ensure that the correct preparation is supplied

Also see local antibiotic guidelines

7.4 DRUGS FOR GENITO-URINARY DISORDERS

7.4.1 URINARY RETENTION

Alfuzosin
2.5mg tablets, 10mg m/r

tablets

Tamsulosin
400microgram m/r capsule

7.4.2 DRUGS FOR URINARY FREQUENCY, ENURESIS AND INCONTINENCE

First line

Oxybutynin
2.5mg, 5mg tablets

5mg m/r tablets

36mg (3.9mg/24hours) patch

Duloxetine
20mg, 40mg capsules (Yentreve®)

NB not to be confused with Cymbalta® - 30mg and 60mg strength capsules for treatment of major

depression and diabetic neuropathy

Second line

Tolterodine
1mg, 2mg tablets

4mg m/r capsules

• Trospium 20mg m/r tablets

Solifenacin
5mg, 10mg tablets

For prescribing according to NICE TA290 (here)

• Mirabegron 25mg, 50mg tablets

Nocturnal enuresis

• **Desmopressin** 200microgram tablets

7.4.3 DRUGS USED IN UROLOGICAL PAIN

Lidocaine/lignocaine

(Instillagel)

2% syringes, 11mL

Potassium citrate
Various preparations – check local availability.

Effercitrate 1.5g tabs is the preparation used in

COCH

7.4.4 BLADDER INSTALLATIONS AND UROLOGICAL SURGERY

Catheter patency solutions

• Sodium chloride 0.9% Uro-Trainer 100mL

Solution G Uro-Trainer 100mL

7.4.5 DRUGS FOR ERECTILE DYSFUNCTION

Alprostadil
Various preparations available

Prescribing restrictions apply

Sildenafil Prescribing restrictions apply