

FOIA MARKER

This is not a textual record. This is used as an administrative marker by the William J. Clinton Presidential Library Staff.

Collection/Record Group: Clinton Presidential Records
Subgroup/Office of Origin: Council of Economic Advisers
Series/Staff Member: Subject Files
Subseries:

OA/ID Number: 21606
FolderID:

Folder Title:
[Global Climate Change and Emissions Trading Policy] [loose] [1]

Stack:	Row:	Section:	Shelf:	Position:
S	21	5	1	3

GCC and Emissions Trading Policy

9

Enclosures filed in
Oversize Attachments # 21606

NAM 18790



Acid Rain Program

[Overview](#) | [Effects](#) | [SO2 Program](#) | [NOx Program](#) | [Emissions](#) | [Conservation](#)

The table below displays the supporting data for the SO2 Allowance Price Chart. Prices reported by the brokerage firms and the Fieldston Publications' market survey are rounded to the nearest whole dollar.

Month/Year	Allowance Price in Dollars		
	Emissions Exchange	Cantor Fitzgerald EBS	Fieldston Publications
8/94	150	145	150
9/94	150	147	150
10/94	145	145	150
11/94	145	144	145
12/94	142	135	140
1/95	141	138	137
2/95	136	135	135
3/95	133	133	133
4/95	132	132	132
5/95	132	132	132
6/95	132	131	130
7/95	130	130	130
8/95	130	130	130
9/95	127	126	128
10/95	125	122	128
11/95	119	117	120
12/95	105	109	111
1/96	92	95	98
2/96	74	79	81
3/96	70	69	83
4/96	81	76	85
5/96	79	79	84
6/96	81	80	83
7/96	82	81	82
8/96	83	82	82
9/96	88	87	86
10/96	92	90	86
11/96	91	92	92
12/96	91	90	94

1/97	97	97	96
2/97	106	102	99
3/97	115	110	111
4/97	113	115	115
5/97	94	98	100
6/97	89	90	93
7/97	87	88	90
8/97	89	91	91
9/97	101	104	92
10/97	105	104	102
11/97	104	107	110
12/97	98	100	102
1/98	96	98	98
2/98	101	101	101
3/98	104	113	105
4/98	133	139	134
5/98	140	148	138
6/98	--	189	193
7/98	--	197	188
8/98	--	189	208
9/98	--	169	177
10/98	--	176	183
11/98	--	189	194
12/98	--	196	195
1/99	--	208	206
2/99	--	200	215
3/99	--	210	217
4/99	--	208	210
5/99	--	212	211

[Acid Rain Program Home](#) | [Price Chart](#) | [Contact Us](#)

<http://www.epa.gov/acidrain/ats/pricetbl.html>

Reasons why advance estimates of SO₂ trading prices were higher than the market price turned out to be:

- (0) Some estimates came from industry, which was self-interested
- (1) Nobody knew that railroad rates would fall sharply, thereby reducing the cost of low-sulphur coal from Wyoming.
- (2) Local restrictions on emissions (Title I of PM?) somewhat reduced the pressure on the federal cap (effect on $p = 25\%$?).
- (3) Early estimates were based on an imaginary wrong bill. (I call this the "real bill doctrine.") E.g. EPA ended up giving out an extra 3.5 million bonus allowances (allocated to plants installing scrubbers) beyond the level planned. The latest estimate (from an EPA/ICF study) was closest to the "real bill." It estimated a price of \$160 ($< 1/2$ earlier estimates).
- (4) Engineering models tended to use fixed coefficients, and made little allowance for substitution of fuels (low-sulphur coal).
- (5) [A rather speculative hypothesis] Perhaps firms are reluctant to buy the right to pollute, due to fear of stigma. This would also explain the low volume of trade in the permit market.

9/24/97 Barry Galet - ICF 703-934-3064, 301-585-1392, 301-585-5804

- not clear how much coal from Wyoming could be used in east
- scrubber costs have declined significantly
 - ① developers of scrubbers have become more experienced
 - ② operation costs ↓ bc utilities are more experienced
- utilities tend to be conservative → concerned about availability of permits, so bought scrubbers
- some stigma to buying permits possible
- Note: EPA early estimates were really low - still haven't hit those values yet

EPA contacts: Kennon Smith 202-233-9164
Sam Nepakitano 202-233-9751
Brian McLean

$\text{CO}_2 + \text{SO}_2$ aren't perfectly analogous

→ gains in SO_2 are easy bc w/o regulation, there is no incentive to cut SO_2 while gains in CO_2 are harder bc w/o regulation, there is an incentive to minimize energy costs

400
 50
 000
 2000
 20000 million

20 b



Acid Rain Program
U.S. Environmental Protection Agency

Fax Cover Sheet

To: *Keith Monk*

From: *Joe Kruger*

Phone:

Phone: *202-564-9125*

Fax:

Fax:

Date:

Pages (Incl. cover sheet): *2*

Keith,

*If you have any questions, call me or
Melanie Dean (202-564-9181).*

Joe


For more information about the Acid Rain Program,
visit our web site at www.epa.gov/acidrain, or
call the Hotline at (202) 564-9620.

The list below displays the supporting data for the SO2 Allowance Price Chart. Prices reported by the brokerage firms are rounded to the nearest whole dollar.


The first column is the Month/Year and the second is the Allowance Price in Dollars. EE is Emissions Exchange and CF is Cantor Fitzgerald EBS.

	<u>EE</u>	<u>CF</u>
	150	145
9/94	150	147
	145	145
	145	144
12/94	142	135
	141	138
	136	135
3/95	133	133
	132	132
	132	132
6/95	132	131
	130	130
	130	130
9/95	127	126
	125	122
	119	117
12/95	105	109
	92	95
	74	79
3/96	70	69
	81	76
	79	79
6/96	81	80
	82	81
	83	82
9/96	88	87
	92	90
	91	92
12/96	91	90
	97	97
	106	102
3/97	115	110
	113	115
	94	98
6/97	89	90
	87	88
	89	91
9/97	101	104
	105	104
	104	107
12/97	98	100


02/94



SO2 Permit Price 8/84-12/97



	EE	CF
8/94	150	145
9/94	150	147
10/94	145	145
11/94	145	144
12/94	142	135
1/95	141	138
2/95	136	135
3/95	133	133
4/95	132	132
5/95	132	132
6/95	132	131
7/95	130	130
8/95	130	130
9/95	127	126
10/95	125	122
11/95	119	117
12/95	105	109
1/96	92	95
2/96	74	79
3/96	70	69
4/96	81	76
5/96	79	79
6/96	81	80
7/96	82	81
8/96	83	82
9/96	88	87
10/96	92	90
11/96	91	92
12/96	91	90
1/97	97	97
2/97	106	102
3/97	115	110
4/97	113	115
5/97	94	98
6/97	89	90
7/97	87	88
8/97	89	91
9/97	101	104
10/97	105	104
11/97	104	107
12/97	98	100



 Randall W. Lutter
11/21/97 05:03:46 PM


Record Type: Record

To: Joseph E. Aldy/CEA/EOP

cc:

Subject: ERP section on acid rain

----- Forwarded by Randall W. Lutter/CEA/EOP on 11/21/97 05:01 PM -----


Richard B. Belzer
11/21/97 01:07:25 PM

Record Type: Record

To: Randall W. Lutter/CEA/EOP

cc:

Subject: ERP section on acid rain

Art asked me to take a look at short section (pp. 9-10) on the acid rain program because I've been reviewing NAPAP's 1996 Integrated Assessment Report to Congress. Despite the fact that CAA90 requires that this report include costs and benefits, neither OMB nor CEA were included in the project and CEA was excluded from the list of "key agencies" from which concurrence is said to be necessary. Fran Sharples is your point of contact. Good luck, Mr. Phelps.

For the most part, the science in the report is quite a bit better than the usual government product -- and that's because it appears to have been written almost entirely by consultants.

Unfortunately, the section on human health benefits (i.e., PM) will drive you certifiably insane. I'll fax it to you if you want an excuse for going postal. A jury of your peers (AEA members all) will acquit.

There is some science in the report that could be memorialized in the ERP as y'all see fit. I list these items below (with citations from the draft report):

1. 1995 SO₂ reductions estimated at 3,887,000 tons. 45% from scrubbers, 55% from fuel switching. Less than 1% of fuel switching is natural gas. Power River coal is only 13% of all tons (24% of fuel switching). 48% of fuel switching is within Appalachia -- i.e., substitution of lower-sulfur coal from within the region. (NAPAP at 10.) These figures are inexplicably different from other figures placing 1995 reductions at 5,600,000 tons (see NAPAP at 18), and this latter figure is the one used elsewhere in the report.
2. Actual costs = \$726 million/yr annualized (discount rate and term not specified; we're working to clarify). Ex ante estimates ranged from \$573m to \$1,338m per year. (NAPAP at 11.)
3. Scrubber cost savings have arisen both in capital and O&M costs. (NAPAP at 12.)

4. Various estimates of cost savings so far from market-based program, most recent is Ellerman (1997) at \$225m - \$375m per year. (NAPAP at 14.)
5. How well the banking and trading program is working are still subject to debate. Issues mentioned include State PUC regulation and the design of the allowance auction. Lack of enforceable property right is not discussed.
6. Actual SO₂ reductions in 1995 exceeded required reductions by 19%. (NAPAP at 18.) Utility emissions were 5.3m tons, well below 8.7m tons allowed by law (NAPAP at 19.)
7. No consensus exists on what SO₂ emissions would have been absent Title IV. SO₂ emissions were declining anyway. Table 4 reports an array of studies with SO₂ reductions attributable to Title IV ranging from 3.12m tons to 5.69m tons. The most recent estimate is 3.68m tons, which would be 2/3 of 5.6m tons total. This estimate appears to attribute some portion of rail deregulation to Title IV, but not enough detail is provided to check this out.
8. NAPAP at 23 reports substantial reductions in ambient SO₂ in 1995, on top of a strong secular decline over the 1986-94 period, and large declines in wet SO₄ deposition. However, NAPAP shows no reduction in ambient SO₄. (Needless to say, this result makes the claimed human health benefits of the program even more disturbing.)
9. FYI, the following BENEFITS are claimed in the NAPAP report:
 1. It's too early to detect changes in aquatic systems (NAPAP at 31).
 2. Forest ecosystems appear to be equally or more sensitive to N than S, and NO_x controls haven't kicked in yet (NAPAP at 42) so no improvements are expected. Embarrassingly, N deposition turns out to be beneficial where soils are N-deficient (NAPAP at 47).
 3. S/N deposition is not a factor in agriculture.
 4. S deposition is really bad for granite and limestone buildings, not to mention tombstones. No idea whether reduced deposition is helping (NAPAP at 53ff).
 5. Ambient SO₄ is responsible for 40-65% of regional haze. SO₄ is down 15% from "previous years," but no change was observed in 1995. 15% reduction was predicted to improve visibility 8%, but no improvements actually observed (NAPAP at 56).
 6. Human health effects will be Big. Real Big. See NAPAP at 62ff. Enough so that the executive summary calls the epidemiologicval studies evidence of a causal relationship (NAPAP at xii), and a table summarizing benefits says that human health benefits alone exceed costs (NAPAP at 30).

1. As an aside, took at figure 21. It would appear to be a concession that your lag is genuine.

TECHNICAL PAPER

ISSN 1017-3289 J. Air & Waste Manage. Assoc. 47:395-402
Copyright 1997 Air & Waste Management Association

Economic Benefits of Improvements in Visibility: Acid Rain Provisions of the 1990 Clean Air Act Amendments

Lauraine G. Chestnut

Hagler Bailly Services, Inc., Boulder, Colorado

Robin L. Dennis

U.S. Environmental Protection Agency, Research Triangle Park, North Carolina

ABSTRACT

The Acid Rain Provisions (Title IV) of the Clean Air Act Amendments of 1990 call for about a 10 million ton reduction in annual SO_2 emissions in the United States. Although the provisions apply nationwide, most of the reduction will occur in the eastern half of the country, where use of high-sulfur coal for electricity generation is most common. One potentially large benefit of Title IV is the expected improvement in visibility conditions in the eastern United States due to the reductions in secondary sulfate aerosols. This paper combines available economic estimates of willingness to pay for improvements in visibility with current estimates of the difference between expected visibility conditions in the eastern United States with and without Title IV, to estimate the expected visibility benefits of Title IV. The results suggest an annual value of \$2.3 billion (in 1994 dollars) in the year 2010, as a result of visibility improvements due to Title IV in residential areas of the eastern United States. The results also suggest a possible additional annual value for eastern U.S. residents of as much as \$1.2 billion for visibility improvements at national parks in the Southeast.

IMPLICATIONS

The 1990 Clean Air Act Amendments require an assessment of the costs and benefits of Title IV, the Acid Rain Provisions. Improvement in visibility conditions in the eastern United States, relative to what would have been expected to occur without Title IV, is one of the potential benefits. This paper applies available economic estimates of the value society places on changes in visibility aesthetics in residential and recreational areas, to determine the expected monetary value of predicted changes in visibility conditions in the eastern United States as a result of the Title IV-required reductions in SO_2 emissions. The results are useful to policy makers as part of the assessment of the cost-effectiveness of Title IV.

INTRODUCTION

Society has long recognized that there is a value to preserving visibility. Section 169A of the Clean Air Act, added in 1977, establishes a national goal of both remedying and preventing anthropogenic impairment of visibility in major national parks and wilderness areas. Visibility is also considered a welfare effect in setting secondary national ambient air quality standards for urban and rural areas. Several states (including Colorado, Vermont, and Oregon) have set or are considering establishing visibility standards entirely unrelated to other air pollution effects.

Visibility benefits assessments to date indicate that visibility aesthetic benefits of air pollution control can be substantial. This paper draws on and updates the literature review for NAPAP in Chestnut and Rowe¹ on the economic valuation of changes in visibility. This paper presents general visibility value functions based on the best available economic estimates of use values for visibility in residential settings, where people live, work, and recreate, and for preservation values for visibility at major national parks in the eastern United States. These value functions are applied to estimates of the differences between visibility levels expected in the eastern United States, with and without the Title IV-required reductions in SO_2 emissions, which have been estimated using EPA's Regional Acid Deposition Model (RADM).

Although the method and estimates are based on a substantial amount of previous research, there are still important uncertainties in estimates of visibility changes and their value to the public. The visibility change estimates are based on predicted levels of SO_2 emissions, which depend on many factors, including economic growth, future energy demand, and many other factors difficult to predict. Estimating the relationship between emissions, ambient aerosol levels, and visibility conditions also involves inherent uncertainties. The visibility values are primarily based on results from contingent valuation studies. There are important issues and uncertainties about the accuracy of those results and about

Chestnut and Dennis

interpretation of available preservation value results for visibility at national parks.

Economic Valuation of Visibility Changes

Visibility has a value to individual economic agents primarily through its effect on the viewing activities of consumers. Consumer values for changes in regional haze can be divided into use and nonuse values. Use values are related to the direct effect on the individual's well-being from experiencing various visibility conditions. Nonuse values are the values an individual holds for protecting visibility for use by others now and in the future (bequest value) and for knowing that it is being protected regardless of current or future use (existence value).

For this analysis, we further separate visibility effects in terms of residential and recreational settings. Residential settings include urban, suburban, and rural areas where people live, work, and participate in everyday recreation such as ball games, walking, picnics, etc. We define recreational benefits as related to major state and federal recreational sites such as state and national parks and wilderness areas. Therefore, we define the following categories of benefits:

Residential use values relate to effects on individuals at work, home, and recreation near their home.

Residential nonuse values relate to effects on other individuals, or purely for the sake of improved visibility.

Recreational use values relate to expected effects when one visits a major recreational site such as a national park or wilderness area.

Recreational nonuse values relate to bequest and existence values for visibility conditions at major recreational sites.

Based on available empirical literature, Chestnut and Rowe¹ conclude that residential use values probably account for more than half of all values for changes in visibility due to regional haze in the eastern United States. This is because most people spend most of their work and recreation time near their homes and because of the substantial numbers of individuals affected by visibility changes in residential settings (as defined above). Chestnut and Rowe also indicate that recreational nonuse values tied to bequest and existence value motives are likely to exceed recreational use values (as defined above). If a large number of individuals hold even small nonuse values for visibility at these sites, such values can exceed on-site use values when summed across the total affected population. Finally, there is insufficient empirical evidence to develop estimates of residential nonuse values, and there is little evidence to suggest that such values are significant. Therefore, residential nonuse values are omitted from this analysis.

Contingent Valuation Method Issues

Most quantitative information about visibility values available at this time is based on results obtained using the contingent valuation method (CVM). The CVM involves using surveys to ask respondents how much they would be willing to pay for specified changes in visibility conditions. It is a technique designed to obtain estimates of willingness to pay (WTP) for goods and services, such as visibility, for which direct prices are not revealed in the market. Without this technique many such nonmarket goods and services remain unvalued in quantitative analyses. This technique has been widely reviewed and debated,^{2,3} and many application and interpretation issues remain unresolved. We do not intend to present a thorough discussion of CVM issues, but we note some key issues.

In general, concerns about using CVM results for quantitative analysis fall into two categories. One category of concerns is that the estimates may be too high because, for example, the exercise is hypothetical rather than based on actual expenditures, and because respondents may have trouble separating the good or service in question from other related goods and services.² The second category of concerns is that the responses may not be quantitatively meaningful because, for example, respondents may not have formulated values for goods like visibility and because responses may reflect a general willingness to contribute to a "good" cause, but may not be directly related to the specific quantity presented in the questions.⁴ CVM research continues to progress in addressing these concerns, especially in the area of reducing suspected upward bias in the results, but available evidence has not yet definitively supported or refuted many of the concerns.² Mitchell and Carson² conclude that WTP estimates useful for quantitative analysis can be obtained for many types of public goods with carefully designed and executed CVM studies.

Empirical Estimates of Residential Use Values

Three CVM studies have estimated use values for changes in visibility in residential urban areas in the eastern United States. McClelland et al.⁵ estimated values for Atlanta and Chicago; Tolley et al.⁶ estimated values for Chicago, Boston, Washington, DC, Atlanta, Cincinnati, Mobile, and Miami. Rae⁷ estimated values for Cincinnati. We also discuss the results of two CVM studies and one hedonic property value study conducted in California cities. The California results provide useful comparisons of results obtained using different valuation approaches and are for coastal locations where visibility levels are within the range of those that occur in the eastern United States.

The Two Cities Study by McClelland et al.⁵ has been discussed in a report prepared for the U.S. EPA. This study

makes some important contributions to the literature in this area by addressing some of the criticisms of the previous research. We, therefore, place considerable emphasis on the results of this study, particularly given that the results are generally consistent with or lower than results from previous studies.

A mail survey was conducted in 1990 for the Two Cities Study in Chicago and Atlanta. About 500 completed responses were obtained. Respondents were shown photographs illustrating three different air quality levels in their area and were told how many days per year each level currently occurs on average. Respondents were asked what their household would be willing to pay per year to have air quality on 25 of the poor days improve to the best air quality level shown. This amounted to about a 14 percent improvement in average annual visual range. Respondents were asked to say what percentage of their response was attributable to concern about health effects, soiling, visibility, or other air quality impact. On average, respondents attributed about 18% of their total WTP for changes in air quality to visibility.

The authors conducted two analyses and adjustments of the responses. One was to estimate and eliminate the potential response bias in WTP estimates caused by nonresponse to the WTP questions by some respondents (including what have been called protest responses). The other was to account for the potential skewed distribution of errors because of the skewed distribution of responses (the long tail at the high end). Both of these adjustments caused the mean WTP estimate to decrease. The annual average household value (in 1994 dollars) for visibility was \$44 before the adjustments and \$20 after the adjustments. The adjustment for the potentially skewed distribution of errors brought the mean WTP to about \$28. This adjustment serves a purpose similar to the "trimming" that is proposed by Mitchell and Carson.² The adjustment for nonresponse to the WTP question reduced the mean WTP further from \$28 to \$20. This adjustment is a new idea implemented for the first time by these authors, who interpreted the adjustments as providing a lower bound on the "true" WTP value. The analysis of the WTP responses also found that income, education, and age were significant in predicting WTP responses. No statistically significant differences were found between the two cities, although different scenes (specific to each city) were used in the photographs.

The Two Cities Study makes an important contribution to addressing the potential for upward bias that may result if respondents mix health and other benefits of pollution reduction with values for visibility improvements when they answer the WTP questions. This potential mixing of visibility and health benefits has been a significant criticism of the earlier Six Cities Study,⁶ which

has led some analysts to the conclusion that the Six Cities results are probably too high.¹ Previous work by some of the same researchers who conducted the Two Cities Study⁶ has demonstrated that some respondents are unable to abstract from the other effects of air pollution when answering questions about visibility. The approach taken in the Two Cities Study was, therefore, to ask for a total value for changes in air quality and then ask the respondent to allocate this value among the different air pollution effects that might concern him, including visibility. This is similar to an approach recommended by Carson et al.⁹ in which respondents are asked to value changes in air quality that represent various combinations of health and visibility impacts. While the mechanics of the Carson et al. approach are different, the thought process required by the respondent is similar. In both cases, the degree to which respondents are embedding values for health and other nonvisibility concerns in their responses for visibility is greatly reduced.

It is difficult to compare the results for the different valuation studies because they are for different changes in visibility in different locations. In an effort to examine for consistent values and patterns across studies, Chestnut and Rowe¹ used the following function to put the mean WTP per household results from the different studies into a common framework:

$$HHWTP/year = b \cdot \ln(VR2/VR1) \quad (1)$$

where:

HHWTP/year	= annual WTP per household in area i for visibility changes in that year
VR ₁	= the starting annual average visual range
VR ₂	= the annual average visual range after the change in emissions
ln	= natural log
b	= estimated coefficient

This function implies that WTP is constant for a given percentage change in visual range and that WTP is zero when there is no change in visual range. Chestnut and Rowe selected this function because it is simple and consistent with results of perceptions studies which suggest that percentage changes in visibility measures are a good way to characterize an individual's perceptions of visual air quality; but other functional forms are also plausible. This function takes into account differences in starting and ending levels of visual range and is also consistent with the economic assumption of diminishing marginal utility for visibility enhancement.

Fitting all available estimates of WTP for changes in visibility conditions in residential areas into eq 1 requires

Chestnut and Dennis

putting the change in visibility conditions valued in each study into the same numeric terms. Chestnut and Rowe selected the change in annual average visual range because most of the studies presented respondents with hypothetical changes in annual conditions, defined either as a change in the annual average or a change in the distribution of good, fair, and poor visibility days. Chestnut and Rowe report that there is no definitive evidence that the presentation of a distribution, rather than an average level, consistently results in higher or lower WTP responses. Evidence on this is limited, however, due to the relatively small number of studies completed to date.

Verification of the validity of this functional form is difficult, given the limited number of observations available from studies conducted to date. Most studies have estimated WTP for one or two different scenarios of change in visibility conditions. Comparisons of results across studies are limited in that there are usually several significant differences in study design, each difference having the potential to cause differences in the results. The area of greatest uncertainty is for small changes in visibility as the function approaches the intercept. Most studies have estimated values for fairly substantial changes in visibility and few have considered both very small and very large changes within the same study.

Table 1 summarizes the results of visibility valuation studies that have some relevance to estimating residential use values for changes in visibility in the eastern United States. This is an update of a similar table provided by Chestnut and Rowe,¹ who also provide a detailed discussion of all of these studies, which is not repeated here. What is most critical for our current purposes is to assess how the McClelland et al.⁵ results compare to results of previous studies. A comparison of the b coefficients (as defined in eq 1) reported in Table 1 shows how the Two Cities results compare. The fully adjusted b coefficient based on the Two Cities Study results is at the low end compared to the Six Cities results, which actually varied considerably across the different cities. The Two Cities Study result is somewhat higher than the Brookshire et al.¹⁰ result for Los Angeles.

The last study in Table 1 is a property value study conducted in Los Angeles and San Francisco that provides estimates of WTP for better air quality at the residence location, based on differences in property values statistically attributed to differences in visual air quality for the residence locations in the sample. The implied results for a 20% improvement in visual range substantially exceed the CVM results in most cases. This is consistent with the expectation that the property value results will reflect values for all aspects of air quality, including concerns about health as well as visibility. These results, based on a very different estimation technique using real market data,

are reasonably consistent with the Two Cities results, which suggest a household annual WTP for a 20% change in visual range of about \$160 (fully adjusted) when all air quality concerns are included.

We selected the b coefficient of 160, as defined in eq 1, based on the fully adjusted McClelland et al. results, in 1994 dollars, for calculating the central estimate of residential use values for Title IV. This study was well designed and addressed many of the criticisms raised about previous contingent valuation studies for residential use values related to visibility. We selected a low estimate of b equal to 120, based on the Brookshire et al. Los Angeles study results, which is the lowest average WTP estimate obtained in any of the studies to date in residential areas with baseline visibility conditions at all comparable with those in the eastern United States. We selected a high estimate of b equal to 220, based on the partially adjusted McClelland et al. results.

Empirical Estimates of Preservation Values for Visibility at National Parks

The National Parks Visibility Values Study¹¹ has been completed since the Chestnut and Rowe¹ literature review. This is the only study available at this time that provides estimates of preservation values (use and nonuse values combined) for protection of visibility in national parks in the eastern United States. The only similar study that has been conducted focused on the Grand Canyon and other parks in the Southwest.¹² The National Parks Visibility Values Study addressed some important methodological issues raised concerning the previous preservation values study and obtained estimates of WTP that were roughly one-third the size for a comparable change in visual range at national parks in the Southwest.

Both studies found that total preservation values far exceeded on-site use values estimated previously, suggesting that the nonuse component of the WTP responses is large. The WTP estimates were obtained in both studies by asking respondents what they would be willing to pay for specific improvements in visibility at national parks. Samples were drawn from the general population who lived near and far from the parks in question and included those who visited the parks and those who did not. It was expected that the responses include the values people held for the opportunity to enjoy the protected visibility during possible future visits to the site, for others who may visit now and in the future, and just because they want visibility to be improved in these areas regardless of human use. Responses to follow-up questions in the National Parks Visibility Values Study suggest that nonuse motives may be significant because respondents indicated that, in giving their WTP responses, motives other than their own opportunity to enjoy the site were important. Also,

Table 1. Comparison of residential visibility valuation study results.

Study	City	Mean WTP (\$1994)	Starting VR (miles)	Ending VR (miles)	b coefficient	WTP for 20% change VR (\$)
Eastern CVM Studies						
McClelland et al. ⁵	Atlanta and Chicago	Unadj. \$44	17.6	20	346	\$63
		Partial \$28			222	\$41
		Full \$20			159	\$29
Tolley et al. ⁶	Chicago	-\$361	9	4	416	\$76
		\$346	9	18		
		\$430	9	30		
Tolley et al. ⁶	Atlanta	-\$301	12	7	469	\$86
		\$289	12	22		
		\$432	12	32		
Tolley et al. ⁶	Boston	-\$222	18	13	422	\$77
		\$212	10	28		
		\$262	18	38		
Tolley et al. ⁶	Mobile	-\$240	10	5	312	\$57
		\$257	10	20		
		\$302	10	30		
Tolley et al. ⁶	Washington, DC	-\$358	15	10	635	\$116
		\$366	15	25		
		\$465	15	35		
Tolley et al. ⁶	Cincinnati	-\$88	9	4	120	\$22
		\$87	9	19		
		\$98	9	29		
Tolley et al. ⁶	Miami	-\$152	13	8	256	\$47
		\$136	13	19		
		\$160	13	29		
Rao ⁷	Cincinnati	\$198	11.4	16.4	602	\$110
California CVM Studies						
Brookshire et al. ¹⁰	Los Angeles	\$130	2	12	119	\$22
		\$333	2	28		
		\$183	12	28	1329	\$242
Loehman et al. ¹¹	San Francisco	-\$211	18.6	16.3		
		\$124	18.3	18.6		
California Property Value Study						
Trijonis et al. ¹⁸	Los Angeles					\$245 - \$657
Trijonis et al. ¹⁸	San Francisco					\$496 - \$552

Note: VR = Visual Range

both of the preservation value studies found that those who had never visited, and never planned to visit, these sites still had positive values, on average, for visibility improvements at these sites. WTP responses were, however, substantially higher for those who had visited, or expected to visit the sites.

For the National Parks Visibility Value Study¹¹, a survey was conducted in 1988 with a sample of residents in Arizona, California, Missouri, New York, and Virginia. A total of 1,647 completed responses were obtained. National parks in three regions were considered in different survey

versions: California, southwestern United States, and southeastern United States. Respondents giving WTP estimates for each region were selected from a state within the region and four states outside the region. Respondents were shown photographs illustrating four levels (current 10th, 50th, 75th, and 90th percentiles) of visibility conditions at a prominent national park in each region (Yosemite, Grand Canyon, and Shenandoah). Respondents were asked what they would be willing to pay each year per household to have average visibility conditions at all national parks in one of the regions improve from the 50th to the 75th or to

Chestnut and Dennis

Table 2. Average annual household WTP estimates for visibility protection at national parks in the Southeast.

Region	Change in annual average visual range	Mean annual WTP In-state (\$1994)	Mean annual WTP out-of-state (\$1994)
Southeast (n = 346)	25 km to 50 km	\$71	\$37
	25 km to 75 km	\$90	\$58
	25 km to 10 km	\$81	\$51

Source: Chestnut and Rowe, 1990.¹¹

the 90th percentiles, or to prevent a degradation to the 25th percentile. Respondents were asked in a follow-up question whether their WTP was entirely for visibility rather than for other park protection concerns, and, if not, what percentage was just for visibility. The average response for all regions was that about 60% was just for visibility.

Table 2 shows the average annual household WTP responses for in-state and out-of-state residents for visibility changes at national parks in the southeastern region, after adjusting for the percentage reported as for visibility only and for identified protest responses. Analysis of the WTP responses showed that significantly higher responses were obtained from respondents who lived in the region or had higher household income. Responses were lower for older respondents and male respondents. We calculated b coefficients for visibility changes based on these results and obtained approximately $b = 85$ for in-state residents and $b = 50$ for out-of-state residents. These b coefficients suggest an annual value per household of about \$15 and \$9, respectively, for a 20% improvement in

visual range.

Estimates of Changes in Visual Range Due to Title IV

There are three steps involved in estimating the changes in visual range in the eastern United States that can be expected to result from the Title IV-required reductions in SO_2 emissions. This paper focuses on a comparison of visual range levels predicted for the year 2010 with and without Title IV. The steps taken to estimate the changes in visual range for this analysis are summarized in Table 3, along with the primary sources and models that these estimates rely on.

ICF Resources¹³ used the Coal and Electric Utilities Model (CEUM) to estimate future SO_2 emissions with and without the Title IV emissions limits. This analysis focused on the utility sector, where 85% of the Title IV emissions reduction is expected. The estimates were based on modeled least-cost compliance options, including consideration of the interaction between the demand for different types of fuels and the costs of supplying and delivering the fuels, as well as the interaction between utilities' marginal costs of compliance and the projected amount of allowance "banking". CEUM uses a series of selected economic, energy market, and utility sector assumptions, that play an important role in estimating emissions with and without Title IV. Factors such as substitute fuel prices, energy demand, and economic growth can all have significant effects on decisions by utilities about building new capacity or retrofitting plants for alternative fuel use. Emissions levels are directly related to levels of electricity production, fuels used, and compliance options employed.

SO_2 emissions estimates by county were partitioned into 80-km² grids and used as input to the Regional Acid Deposition Model (RADM),¹⁴ which combines meteorological and atmospheric chemical modeling to predict the transport, transformation, and ultimate deposition of SO_2 and the secondary sulfate that forms as a result of SO_2 emissions. Hourly visual range estimates are based on the intermediate estimates of atmospheric sulfate concentrations and relative humidity that RADM provides for each regulatory scenario.

Visual range estimates for each scenario were calculated as follows. An extinction coefficient was calculated as the sum of extinctions due to gas (Rayleigh) scattering and particulate (sulfate) scattering. The changes in visual range estimated for the Title IV scenarios were based on changes in sulfate levels; light extinction from other aerosols was assumed to remain constant. The sulfate

Table 3. Quantification steps for this assessment of visual range changes due to Title IV.

Quantification steps	Selected quantification method
1. Changes in SO_2 emissions	Use ICF Resources ¹³ estimates of 1985 emissions, 1997 emission with Title IV, and 2010 emissions with and without Title IV (prepared for EPA)
2. Changes in atmospheric sulfate concentrations in the eastern United States	Use EPA's Regional Acid Deposition Model (RADM) ¹⁴ runs for each of the SO_2 emissions scenarios
3. Changes in visual range associated with estimated changes in sulfate concentrations	Based on the CAPITA ¹⁵ method for calculating the aerosol part of the extinction coefficient, the Koschmeider equation for converting extinction to visual range, and the IMPROVE network results ¹⁶ for calibrating the RADM extinction estimates to visibility monitoring data

Table 4. Residential visibility benefits of Title IV (1994 dollars).

Region	1997 with Title IV versus current			2010 with Title IV versus 2010 without Title IV		
	% change median annual visual range	Total annual residential value (millions)	Average % value per change household	Total annual median annual visual range	Average value per residential value (millions)	household
New England	6.3%	\$59.3	\$10.85	6.6%	\$69.9	\$11.42
Mid-Atlantic	7.7%	\$145.7	\$8.16	17.1%	\$441.5	\$22.16
South Atlantic	5.3%	\$07.9	\$5.58	30.3%	\$689.3	\$39.07
South Central	8.3%	\$108.3	\$12.03	27.2%	\$391.6	\$38.93
Midwest	4.5%	\$193.3	\$8.61	15.1%	\$683.8	\$27.22
Total 31-States		\$594.5	\$9.06		\$2,276.2	\$28.00

aerosol part of the extinction coefficient was calculated according to the CAPITA method:¹⁵

$$B_{sp} = f(RH) * SE(\text{sulfate}) * [SO_2] * 10^{-4} \text{g}/\mu\text{g} \quad (2)$$

where:

B_{sp} = extinction in m^{-1}

SE = m^2/g

SO_2 = sulfate aerosol concentration in $\mu\text{g}/\text{m}^3$

The RADM estimates of light extinction were calibrated to be consistent (for current conditions) with visibility data available from the IMPROVE network.¹⁶ Visual range was then calculated by applying Koschmeider's equation to the light extinction estimates.

RESULTS

Table 4 shows the average estimated percentage changes in visual range by region and the associated monetary value estimates for residential use values attributable to the Title

IV-required SO_2 emissions limits. Annual estimates for 1997 and 2010 reflect expected average population growth, based on U.S. Census middle forecasts. In 1997, the annual benefits are spread fairly evenly across the regions and amount to about \$595 million (in 1994 dollars), which equals about \$9 per household. In 2010, the estimated benefits are much higher in every region except New England, with the

highest benefits per household in the south Atlantic and south central regions. The total annual benefit in 2010 is about \$2.3 billion, which equals an average of about \$28 per household.

Table 5 shows the visibility benefits estimates for national parks in the Southeast due to Title IV. These estimates are based on RADM results for Shenandoah and Great Smoky Mountains National Parks, which are located in the region where the largest improvements in visibility are expected. In 2010, the estimated benefits per household are \$14 to \$17 for out-of-state residents, and \$23 to \$28 for in-state residents. We applied the in-state values to residents of Virginia, North Carolina, and Tennessee, and applied the out-of-state values to the remainder of the 31-state eastern United States to obtain total annual values. This does not include any value for households in the western United States, which means that these results understate total national values for visibility protection at these parks, given that in the National Parks Visibility Values Study western residents did give positive WTP values for visibility at eastern national parks. The primary uncertainty in interpreting the results for the national parks is whether they should be

added to the residential use values. To be conservative, we assumed that residential and national park values should not be added for in-state residents, because residential use values may include some value for relatively nearby national parks. However, values for out-of-state residents probably should be added to residential use values. This suggests a total annual 2010 benefit of about \$3.4 billion for residents of the eastern United States for changes in visibility in residential areas and at national parks attributable to Title IV.

Table 5. Visibility benefits for national parks in the Southeast due to Title IV.

Estimate	1997	2010
% Change in Annual Median Visual Range at Parks ¹	9% to 11%	30% to 40%
Values per In-State Household	\$7.40 to \$8.40	\$22.70 to \$28.40
Values per Out-of-State Household	\$4.50 to \$5.10	\$13.60 to \$17.00
In-State Total Annual Value (Millions)	\$54.0 to \$62.2	\$185.8 to \$232.2
Out-of-State Total Annual Value (Millions)	\$286.6 to \$322.5	\$961.7 to \$1202.1

¹ Estimates based on the RADM results for cells that fall within boundaries of Shenandoah and Great Smoky Mountains National Parks.

Chestnut and Dennis

ACKNOWLEDGMENTS

This research was supported by the U.S. Environmental Protection Agency, in part under Contract No. 68-D3-0005. The views and conclusions expressed in this paper are not necessarily those of the U.S. Environmental Protection Agency.

REFERENCES

1. Chestnut, L.G.; Rowe, R.D. *Economic Valuation of Changes in Visibility: A State of the Science Assessment for NAPAP, Acidic Deposition: State of Science and Technology Report 27*; Brown, G.M. Jr., Callaway, J.M., Eds.; National Acid Precipitation Assessment Program: Washington, DC, October 1990.
2. Mitchell, R.C.; Carson, R.T. *Using Surveys to Value Public Goods: The Contingent Valuation Method*, John Hopkins Press for Resources for the Future: Washington, DC, 1989.
3. *Valuing Environmental Goods: An Assessment of the Contingent Valuation Method*; Cummings, R.G., Brookshire, D.S., Schulze, W.D., Eds.; Rowman and Allanheld: Totowa, NJ, 1986.
4. Fischhoff, B.; Furbey, L. "Measuring values: A conceptual framework for interpreting transactions with special reference to contingent valuation of visibility," *J. Risk and Uncertainty* 1988, 1, 147-184.
5. McClelland, G.; Schulze, W.; Waldman, D.; et al. *Valuing Eastern Visibility: A Field Test of the Contingent Valuation (Draft)*, Cooperative Agreement #CR-815183-01-3, U.S. Environmental Protection Agency: Washington, DC, 1991.
6. Tolley, G.A.; Randall, A.; Blomquist, G.; et al. *Establishing and Valuing the Effects of Improved Visibility in Eastern United States*, U.S. Environmental Protection Agency: Washington, DC, 1986.
7. Rae, D. *Benefits of Improving Visual Air Quality in Cincinnati: Results of a Contingent Valuation Survey*, Project 1742, Electric Power Research Institute: Palo Alto, CA, May 1983.
8. Irwin, J.; Schenk, D.; McClelland, G. "Urban visibility values: Some experiments in the use of the contingent valuation method," in *Visibility and Fine Particles*; C.V. Mathai, Ed.; *Transactions of the Air and Waste Management Association Specialty Conference on Visibility and Fine Particles*, Air and Waste Management Association: Pittsburgh, 1990.
9. Carson, R.T.; Mitchell, R.C.; Ruud, P.A. "Valuing air quality improvements: simulating a hedonic pricing equation in the context of a contingent valuation scenario," in *Visibility and Fine Particles*; C.V. Mathai, Ed.; *Transactions of the Air and Waste Management Association Specialty Conference on Visibility and Fine Particles*, Air & Waste Management Association: Pittsburgh, PA 1990.
10. Brookshire, D.; d'Arge, R.; Schulze, W.; Thayer, M. *Methods Development for Assessing Air Pollution Control Benefits. Volume 3: Experiments in Valuing Nonmarket Goods. A Case Study of Alternative Benefit Measures of Air Pollution Control in the South Coast Air Basin of Southern California*, EPA-600/6-79-0016, U.S. Environmental Protection Agency: Washington, DC, February, 1979.
11. Chestnut, L.G.; Rowe, R.D. *Preservation Values for Visibility Protection at the National Parks*, Cooperative Agreement #CR-813-686, U.S. Environmental Protection Agency: Research Triangle, NC, 1990.
12. Schulze, W.D.; Brookshire, D.S.; Waltham, E.G.; et al. "The benefits of preserving visibility in the national parklands of the Southwest," *Natural Resources Journal* 1983, 23, 149-173.
13. ICF Resources, Inc., *Economic Analysis of the Title IV Requirements of the 1990 Clean Air Act Amendments*, U.S. Environmental Protection Agency, Acid Rain Division, 1994.
14. Dennis, R.L.; McHenry, J.N.; Barchet, W.R.; et al. "Correcting RADMS sulfate underprediction: Discovery and correction of model errors and testing the corrections through comparisons against field data," *Atmos. Environ.* 1993, 27A:975-997.
15. Husar, R. *User's Guide: Visibility Module which Converts Aerosol Concentration to Visibility Parameters: An Empirical Relationship for Calculating Visibility Distance from Particle Concentrations Calculated by Regional Models*, Internal Report.
16. Sisler, J.F.; Huffman, D.; Lattimer, D.A.; et al. *Spatial and Temporal Patterns and the Chemical Composition of the Haze in the United States: An Analysis of Data from the IMPROVE Network, 1988-1991*, ISSN No. 0737-5352-26, 1993.
17. Loehman, E.; Hoidt, D.; Chaikin, K. *Measuring the Benefits of Air Quality Improvements in the San Francisco Bay Area. Part I: Study Design and Property Value Study*, Draft Final Report, SRI International: Menlo Park, CA, September 1980.
18. Titijs, J.C.; Thayer, M.; Murdoch, J.; Hagemann, R. *Air Quality Benefits Analysis for Los Angeles and San Francisco Based on Housing Values and Visibility*, California Air Resources Board: Sacramento, CA, 1984.


About the Authors

L.G. Chestnut is an economist with Hagler Bailly Services, Inc., P.O. Drawer O, Boulder, CO 80306-1906. R.L. Dennis is a physical scientist at the Atmospheric Sciences Modeling Division, Air Resources Laboratory, National Oceanic and Atmospheric Administration on assignment to the Atmospheric Research and Exposure Assessment Laboratory, U.S. Environmental Protection Agency, Research Triangle Park, NC 27711 (MD-80). L.G. Chestnut can be reached by phone at (303) 449-5515 or by fax at (303) 443-5684.

Volume of SO2 Allowances in Economically Significant Transfers

(same as
MIT def.)

Volume of Allowances



Market Chart

ACTIVITY IN THE ACID RAIN PROGRAM SO₂ ALLOWANCE MARKET**Total Private Transfer Activity**


	Q1	Q2	Q3	Q4	Annual Total
1994		48 transactions 1.8 M allowances	15 transactions 1.7 M allowances	152 transactions 5.8 M allowances	215 transactions 9.2 M allowances
1995	181 transactions 10.6 M allowances	118 transactions 2.2 M allowances	137 transactions 0.8 M allowances	177 transactions 3.1 M allowances	613 transactions 16.7 M allowances
1996	447 transactions 3.4 M allowances	112 transactions 0.5 M allowances	196 transactions 1.1 M allowances	319 transactions 3.2 M allowances	1,074 transactions 8.2 M allowances
1997	514 transactions 4.1 M allowances	263 transactions 4.2 M allowances	253 transactions 2.0 M allowances	399 transactions 4.9 M allowances	1,429 transactions 15.2 M allowances

Private Transfers Between Economically Distinct Organizations

	Q1	Q2	Q3	Q4	Annual Total
1994		41 transactions 0.8 M allowances	5 transactions 7,000 allowances	20 transactions 90,000 allowances	66 transactions 0.9 M allowances
1995	55 transactions 0.5 M allowances	88 transactions 0.2 M allowances	90 transactions 0.4 M allowances	96 transactions 0.8 M allowances	329 transactions 1.9 M allowances
1996	148 transactions 0.9 M allowances	98 transactions 0.5 M allowances	137 transactions 0.9 M allowances	195 transactions 2.1 M allowances	578 transactions 4.4 M allowances
1997	166 transactions 1.7 M allowances	208 transactions 1.9 M allowances	174 transactions 1.6 M allowances	262 transactions 2.7 M allowances	810 transactions 7.9 M allowances


↑
see
attached
graph.

Allowances Transferred Between Organizations By Year



Allowances Transferred Between Companies

Allowances (in millions)



NAPAP

National Acid Precipitation Assessment Program

1996 INTEGRATED ASSESSMENT

– REPORT TO CONGRESS –

National Science and Technology Council
Review Draft

October 16, 1997

About the National Acid Precipitation Assessment Program

The National Acid Precipitation Assessment Program (NAPAP) is a federal interagency program established by Congress in 1980 to coordinate research and monitoring activities related to acid deposition among six federal agencies: the Environmental Protection Agency, the Department of Energy, the Department of the Interior, the Department of Agriculture, the National Oceanic and Atmospheric Administration, and the National Aeronautic and Space Administration. Much of the research sponsored through NAPAP in the 1980s improved our knowledge and understanding of acid deposition and its effects on ecosystems, human health, visibility, and materials. The 1990 Clean Air Act Amendments established the first federal Acid Deposition Control Program (under Title IV) with market-based incentives for emission reductions. Its aim is to achieve significant environmental benefits primarily through reductions in utility emissions of sulfur dioxide (SO₂) and nitrogen oxides (NO_x), the precursors to acid deposition. These pollutants are released mainly from the burning of coal, oil, and other fossil fuels. In addition, Congress mandated that every two years NAPAP provide a technical update of acid deposition and its effects. Beginning in 1996 and every four years thereafter, NAPAP is to report on the costs, benefits, and effectiveness of Title IV, as well as the reduction in deposition rates that must be achieved to prevent adverse ecological effects.

NAPAP also coordinates acid deposition information as a member of the Air Quality Subcommittee under the President's National Science and Technology Council, which is described below. To obtain additional information regarding NAPAP, see details on the back cover.

About the National Science and Technology Council

President Clinton established the National Science and Technology Council (NSTC) by Executive Order on November 23, 1993. This cabinet-level council is the principal means for the President to coordinate science, space, and technology policies across the federal government. NSTC acts as a "virtual" agency for science and technology to coordinate the diverse parts of the federal research and development enterprise. The NSTC is chaired by the President. Membership consists of the Vice President, Assistant to the President for Science and Technology, cabinet secretaries and agency heads with significant science and technology responsibilities, and other White House officials.

An important objective of the NSTC is to establish clear national goals for federal science and technology investments in areas ranging from information technologies, environment and natural resources, and health research, to improving transportation systems and strengthening fundamental research. The Council prepares research and development strategies that are coordinated across federal agencies to form an investment package that is aimed at accomplishing multiple national goals. To obtain additional information regarding the NSTC, contact the NSTC Executive Secretariat at 202-456-6100.

About the Office of Science and Technology Policy

The Office of Science and Technology Policy (OSTP) was established by the National Science and Technology Policy, Organization and Priorities Act of 1976. OSTP's responsibilities include advising the President in policy formulation and budget development on all questions in which science and technology are important elements; articulating the President's science and technology policies and programs, and fostering strong partnerships among federal, state, and local governments, and the scientific communities in industry and academia. To obtain additional information regarding OSTP, contact the OSTP Administrative Office at 202-395-7347.

PREFACE

The National Acid Precipitation Assessment Program (NAPAP) is a cooperative federal program first authorized in the Acid Precipitation Act of 1980 to coordinate federal acid rain research and report the findings to Congress. The culmination of efforts by NAPAP and others in the 1980s provided a scientific basis for Title IV of the 1990 Clean Air Act Amendments, also known as the Acid Deposition Control Program. Title IV provided market-based incentives for controlling electric utility emissions of SO₂. Title IX of the Amendments reauthorized NAPAP, but giving it a different mission. The NAPAP agencies (EPA, DOE, DOI, USDA, NOAA, NASA) would continue to coordinate acid rain research and monitoring, but would also periodically assess the costs, benefits, and effectiveness of Title IV.

NAPAP is responsible for providing credible technical findings to inform the public decision process. The NAPAP 1996 Report to Congress documents the findings of NAPAP's 1996 integrated assessment of Title IV. The assessment was based on a compilation of the best available scientific and economic information relevant to the issue. The scope of the Report includes (1) emissions, atmospheric processes, and deposition; (2) effects on surface waters, forests, exposed materials and cultural resources, human health, and visibility; (3) compliance costs and emission allowance trading; and (4) valuation of the benefits of reducing acid deposition.

The report is directed to Congress but provides valuable economic and scientific information to all public officials who are responsible for setting air quality policy. It attempts to present highly technical information that addresses current public policy issues in a format that can be understood by the non-scientific reader. The report is organized in a series of sections that generally follow the causal chain of events: utility emissions and compliance costs, air concentrations and deposition, effects, and economic valuation of benefits. Where more scientific or economic detail is desired, references are denoted in the text and provided at the end of the report.

ACKNOWLEDGEMENTS

Production of this document was managed by the staff of the NAPAP Office of the Director. Current and former members of the staff who were instrumental in the planning, development, and publication of this report are listed below.

Hundreds of scientists and technical specialists contributed over the years to the extensive knowledge base upon which the 1996 integrated assessment depended. NAPAP's 1996 integrated assessment report to Congress was planned, developed, written, edited, and reviewed in a collaborative effort by many scientists, economists, and technical specialists. Their efforts are greatly appreciated. The authors and contributors to the various sections of the report are listed on the following pages.

Special thanks are extended to the extramural peer reviewers, who are identified following the authors. In addition to providing a technical review, they were asked to consider the content of the report as it relates to public policy. Their comments were timely, relevant, and informative.

NAPAP Office of the Director

Dr. Michael S. Uhart
Ms. Noreen B. Clancy
Ms. Karen E. King
Ms. Ann Georgilas, NOAA

Former Office of the Director

Dr. Derek Winstanley
Ms. Judy L. Hickey
Dr. John M. Barnes
Ms. Elizabeth B. Bennett

Editors

Dr. Michael S. Uhart
Ms. Noreen B. Clancy
Ms. Katie Smythe, Science & Policy Associates

Editorial Consultant

Dr. Christopher Bernabo, Science & Policy Associates

Primary Authors and Contributors

Implementation Status

Rosemary Wolfe, Environmental Protection Agency

Emissions and Compliance Costs

Denny Ellerman, Massachusetts Institute of Technology

Joe Kruger, Environmental Protection Agency

Juan Pablo Montero, Massachusetts Institute of Technology

Concentration and Deposition

Bruce Hicks, National Oceanic and Atmospheric Administration

Mark Nilles, U.S. Geological Survey

Phil Bowman, U.S. Geological Survey

Aquatic Ecosystems

Tim Sullivan, E&S Environmental Chemistry

Kathy Tonneson, National Park Service

Webb Van Winkle, Oak Ridge National Laboratory

Rona Birnbaum, Environmental Protection Agency

Forest Ecosystems

Bob Cook, Oak Ridge National Laboratory

Dev Joslin, Tennessee Valley Authority

Chris Eagar, U.S. Forest Service

Jennifer Knoepp, U.S. Forest Service

Sandy McLaughlin, Oak Ridge National Laboratory

Carl Trettin, U.S. Forest Service

Materials and Cultural Resources

Victor Mossotti, U.S. Geological Survey

Mary Striegel, National Park Service

Visibility

Marc Pitchford, National Oceanic and Atmospheric Administration

Human Health

Larry Folinsbee, Environmental Protection Agency

Tom Grahame, Department of Energy

Ann Watkins, Environmental Protection Agency

Adverse Effects

Harry Quarles, Oak Ridge National Laboratory

Response Indicators

Carolyn Hunsaker, Oak Ridge National Laboratory

Benefits Valuation

Dallas Burtraw, Resources for the Future

Erin Mansur, Resources for the Future

Tracking and Analysis Framework

Cary Bloyd, Argonne National Laboratory

Max Henrion, Lumina

Rich Sonnenblick, Lumina

Peer Reviewers

Integration and Assessment

Edward Bennett, New York Department of Environmental Conservation
James Hammitt, Harvard School of Public Health
Robert Mendelsohn, Yale School of Forestry and Environmental Studies

Emissions and Compliance Costs

Stephen Brick, Consultant
Andrew Van Horn, Van Horn Consulting

Concentration and Deposition

Thomas Butler, Cornell University
Alan Hansen, Electric Power Research Institute

Aquatic Ecosystems

Nicholas Nikolaidis, University of Connecticut
Kent Thornton, FTN Associates

Forest Ecosystems

Alan Lucier, National Council of the Paper Industry for Air and Stream Improvement
William Smith, Yale School of Forestry and Environmental Studies

Materials and Cultural Resources

A. Elena Charola, Consultant
John Spence, North Carolina State University

Visibility

Paulette Middleton, Science and Policy Associates
Pradeep Saxena, Electric Power Research Institute

Human Health

Petros Koutrakis, Harvard School of Public Health
Morton Lippmann, New York Medical Center

Benefits Valuation

A. Myrick Freeman, Bowdoin College
Douglas Lipton, University of Maryland

CONTENTS

EXECUTIVE SUMMARY	x
INTRODUCTION	1
NAPAP Mission and Mandate	2
Assessment Process	2
Organization of the Report	6
IMPLEMENTATION AND COSTS OF TITLE IV	7
What is the status of implementation?	7
How well have utilities complied with the provisions under Title IV?	8
What are the compliance costs of Title IV and how do they compare to projections? ..	9
CHANGES IN EMISSIONS, CONCENTRATIONS AND DEPOSITION ...	17
What emission reductions have been achieved?	18
How have air concentrations been affected by these emission reductions?	22
How have deposition rates been affected by the emission reductions?	25
What are current deposition rates and what is their variability?	27
BENEFITS OF EMISSION REDUCTIONS	29
Have changes in effects been observed (1980-1995)?	29
Aquatic Ecosystems	31
Forest Ecosystems	42
Materials and Cultural Resources	53
Visibility	56
Human Health	63
What are the economic benefits related to the effects areas?	67
REDUCTIONS TO PREVENT ADVERSE ECOLOGICAL EFFECTS	80
What are "adverse ecological effects"?	80
What are the dose-response relationships for sulfur and nitrogen deposition?	80
How might ecological effects be reduced if deposition were reduced further?	80
EFFECTIVENESS OF TITLE IV: A MARKET-BASED APPROACH	83
Did the market-based approach reduce costs?	83
How did emission and deposition changes compare to projections?	84
Has reduced acid deposition had an effect on sensitive receptors?	85
OUTLOOK TO 2000	86
Research, Monitoring and Development Needs	86
Fundamental Needs	88
Near- and Long-term Needs	90
Emissions and Compliance	90
Concentration and Deposition	91
Aquatic Ecosystems	91

TABLE OF CONTENTS

Forest Ecosystems	93
Materials and Cultural Resources	94
Visibility	94
Human Health	95
Benefits Valuation	96
Linkages to Other Issues	96
Related Science Issues	97
Related International Issues	99
REFERENCES	101
APPENDICES	
A. Advances in Relevant Modeling	121
Integrated Models	121
Issue-Specific Models	139
B. Design and Performance of Pollution Trading Programs	145
C. Interpretation of "Adverse Effects" for NAPAP Report to Congress	151

LIST OF FIGURES

- Figure 1: Conceptual framework for NAPAP's integrated assessment.
- Figure 2: Allowance price trends.
- Figure 3: National annual emissions of sulfur dioxide by all sources (1980-1995).
- Figure 4: National annual Phase I utility emissions of sulfur dioxide (1980-1995).
- Figure 5: Geographic distribution of annual sulfur dioxide emissions by utilities (1980-1995).
- Figure 6: National annual anthropogenic emissions of nitrogen oxides (NO_x) by sources (1980-1995).
- Figure 7: Processes leading to acid deposition.
- Figure 8: Ambient air concentrations of sulfur dioxide (SO_2) from 1986-1995.
- Figure 9: Sulfur dioxide (SO_2), sulfate (SO_4), and nitric acid vapor (HNO_3) concentrations near State College, Pennsylvania.
- Figure 10: Sulfate concentrations in Shenandoah National Park and Great Smoky Mountains National Park and a composite of ten western sites.
- Figure 11: Decrease in ambient air concentrations of total sulfur: 1989-1995.
- Figure 12: National Atmospheric Deposition Program/National Trends Network (NADP/NTN).
- Figure 13: Percent difference between 1995 observed and estimated sulfate concentrations.
- Figure 14: Percent difference between 1995 observed and estimated nitrate concentrations.
- Figure 15: Percent difference between 1995 observed and estimated hydrogen ion concentrations (acidity).
- Figure 16: Annual wet and dry deposition of sulfur and nitrogen at State College, PA.
- Figure 17: Annual mean hydrogen ion concentrations (in precipitation) as pH for 1995.
- Figure 18: Total wet deposition of sulfate ion in 1995.
- Figure 19: Total wet deposition of nitrate ion in 1995.
- Figure 20: Total wet deposition of nitrogen from nitrate and ammonia in 1995.
- Figure 21: Relative response times to changes in emissions.
- Figure 22: Sulfate, Acid Neutralizing Capacity (ANC), base cations, and pH (acidity) variations in Biscuit Brook, Pennsylvania.
- Figure 23: Sulfate, Acid Neutralizing Capacity (ANC), base cations, and pH (acidity) variations in Lake Rondaxe, New York.
- Figure 24: The four broad forest regions used in this assessment.

- Figure 25. Generalized dose-response relationship for nutrient limitations associated with increasing soil and deposition acidity.
- Figure 26. Model of stone alteration processes.
- Figure 27. Aerosol and visibility monitoring sites.
- Figure 28. Sulfate contributions to visibility reduction by region.
- Figure 29. Nitrate contributions to visibility reduction by region.
- Figure 30. Annual particulate sulfate reduction in 1995 compared to the 1993-1994 mean, by region.
- Figure 31. Seasonal particulate sulfate reduction in 1995 compared to the 1993-1994 mean, by region.

LIST OF TABLES

- Table 1. Methods used to reduce sulfur dioxide emissions in 1995.
- Table 2. Estimates of Phase I compliance costs.
- Table 3. Scrubber costs.
- Table 4. Title IV reductions of sulfur dioxide.
- Table 5. Selected policy-relevant developments since 1990.
- Table 6. Characteristics used to select forest ecosystems for regional assessment of the effects of acid deposition.
- Table 7. Qualitative, relative rankings for benefit areas with respect to the primary assessment questions of interest to NAPAP.
- Table 8. Value of a statistical life and other health end points.
- Table 9. Research and monitoring needs for future NAPAP assessments.

EXECUTIVE SUMMARY

In Title IV of the 1990 Clean Air Act Amendments, Congress set out to decrease the adverse effects of acid deposition through reductions in annual emissions of sulfur dioxide (SO₂) and nitrogen oxides (NO_x) from utilities burning fossil fuel. The legislation called for placing a cap on utility emissions to achieve a total reduction of 10 million tons of SO₂ emissions below 1980 levels by 2010. In combination with reductions under Title I (compliance with National Ambient Air Quality Standards) and Title II (mobile sources), Title IV will contribute to the overall two million ton reduction of NO_x emissions from 1980 levels. *W C C*

In contrast to the typical command-and-control approach to regulation, Congress adopted a market-based control strategy for SO₂. Emission reduction goals were established, with time allowed to achieve these goals through flexible compliance methods, including an innovative SO₂ emission allowance trading and banking program. The SO₂ reductions were to be implemented in two phases: Phase I, began January 1, 1995; and Phase II will begin January 1, 2000. Phase I affects the highest emitting electric generating units and Phase II will include the remaining electric generating units.

The Act mandated the interagency National Acid Precipitation Assessment Program (NAPAP) to continue the coordination of federal acid rain research and monitoring. In addition, Congress asked NAPAP to evaluate the costs, benefits, and effectiveness of Title IV and to assess what further reductions in deposition rates are needed to prevent adverse ecological effects. This report to Congress is the first in a series of quadrennial integrated assessments of Title IV. The major goal of NAPAP's 1996 integrated assessment is to provide structured, technical information in a format that enables decision makers to evaluate the effectiveness of current public policy and provides a sound science base for future policy decisions.

Summary of Significant Findings

The first years that affected utilities were required to comply with the SO₂ and NO_x emission reduction programs were 1995 and 1996, respectively. Although utilities have significantly reduced their emissions, observable responses were not yet expected mostly due to inherent time lags between changes in emissions and responses by sensitive receptors, especially within ecosystems. Weather variability and its impact on dispersion, transport, and deposition of pollution contribute to the uncertainty in identifying a response of ambient air concentrations and visibility to emission reductions. The most significant, policy-relevant findings of the assessment follow.

- The market-based approach has reduced compliance costs below those of a command-and-control approach. Costs are below what was expected due to a multitude of factors, including lower costs for low-sulfur coal, railroad deregulation, technological innovation, and lower capital and operating costs for scrubbers.
- All affected utilities have fulfilled the compliance requirements of Title IV. In the first two annual reconciliations of allowances and emissions, SO₂ allowances matched or exceeded SO₂ emissions.
- Allowance trading, a unique aspect of Title IV, has been successful both in terms of the volume of trades and in its effectiveness in keeping compliance costs down. Economic analysis shows the market of SO₂ allowances is functioning, liquid, and effective and can serve as a model for other air pollution control programs.

- 1 • All of the Acid Deposition Control Program goals for SO₂ and NO_x were fully or
2 substantially achieved in 1995.
 - 3 - In the first year of compliance (1995), sulfur dioxide emissions for Phase I electric
4 utility units were 39% (3.4 million tons) below the 1995 allowable level specified
5 under Title IV (see footnote). Most of this reduction occurred in the Midwest. By
6 1996, more than half of the overall 10 million ton SO₂ emissions reduction target
7 had been achieved by Phase I utilities.
 - 8 - Implementation of Title IV NO_x reductions only began January 1, 1996.
9 Nevertheless, NO_x emissions from all sources in 1995 were 6% (1.5 million tons)
10 below 1980 levels. Utilities contributed to 53% of that reduction and to 40%
11 achievement of the two million ton goal under the Clean Air Act Amendments.
12
- 13 • In response to SO₂ emission reductions in 1995, statistically significant reductions in the
14 acidity of (represented by hydrogen ion content) and sulfate concentrations in precipitation
15 were reported at deposition monitoring sites in the Midwest, Mid-Atlantic, and Northeast
16 United States. However, there was no evidence of statistically significant decreases in
17 nitrate concentrations in precipitation in 1995; NO_x reductions under Title IV were not
18 required until 1996.
- 19 • It is too early to determine whether changes in aquatic ecosystems have resulted from Title
20 IV. But over the last 15 years, lakes and streams throughout many areas of the United
21 States have experienced decreases in sulfate concentrations in response to decreased
22 emissions and deposition of sulfur. For example, there is evidence of recovery from
23 acidification in New England lakes. In contrast, the majority of Adirondack lakes have
24 remained fairly constant while the sensitive Adirondack lakes have continued to acidify. In
25 1995, EPA reported to Congress that to fully recover Adirondack lakes additional
26 reductions in sulfur and nitrogen deposition would be required (U.S. EPA, 1995, Standard
27 Feasibility Study).
- 28 • Sulfur and nitrogen deposition have caused adverse impacts on restricted highly sensitive
29 forest ecosystems in the United States. High-elevation spruce-fir forests in the eastern
30 United States are the most sensitive. Most forest ecosystems in the East, South, and West
31 are not currently known to be adversely impacted by sulfur and nitrogen deposition.
32 However, if deposition levels are not reduced in areas where it is presently high, adverse
33 effects may develop in more forests due to chronic, multiple decade exposure.
- 34 • The gradual leaching of soil nutrients from sustained inputs of acid deposition could
35 eventually impede forest nutrition and growth in several areas. Potential risk depends on
36 numerous factors, including rate of cation deposition, soil cation reserves, age of forest,
37 weathering rates, species composition, and disturbance history. Recent reductions in
38 sulfur deposition should result in some near term improvements in sensitive forests. Large
39 improvements will occur at varying rates, depending on the factors listed above.
- 40 • Nitrogen deposition can significantly degrade forest ecosystems, especially in areas where
41 nitrogen levels are already high and the soil has reached or is approaching saturation.

Emissions data through 1995 were used in this assessment; quality controlled emissions data for 1996 were not available at the time of writing. Since that time, the Environmental Protection Agency has released year-end 1996 emissions data. In 1996, SO₂ emissions by utilities were 5.4 million tons, or 2.9 million tons (35%) below the 8.3 million ton allowable level determined by 1996 allowance allocations. In the first year of NO_x compliance (1996) of Title IV, coal-fired boilers reduced emission rates by an average of 40% below 1990 levels, emitting 33% less NO_x in the process (340,000 tons) and demonstrating an average of 18% overcompliance with required emission rate levels.

However, some nitrogen-deficient forests may experience increased growth rates in response to continued elevated nitrogen deposition.

- For materials and cultural resources, dry deposition (the deposition of particles between rain events) is now considered to be more damaging to stone than wet deposition (the deposition of particles during rain events).
- Due to uncertainties generated by meteorological variability and other factors, the linkage between reduced sulfate concentrations and improved visibility is difficult to ascertain without long term data. Nevertheless, based on the assumption that sulfate significantly contributes to visibility-reducing particles in the eastern United States, reduced SO₂ emissions are expected to reduce sulfate concentrations and, in turn, their contribution to haze.
- Reduction in sulfur and nitrogen emissions is expected to reduce fine particulate sulfate and nitrate. Indications from epidemiology studies of a causal relationship between ambient sulfate and nitrate particles with human health end points suggest decreased emissions should lead to reductions in premature mortality from cardiovascular and respiratory causes and in the number of days that asthma symptoms are experienced.
- There is significant evidence that quantifiable economic benefits could be relatively large in the areas of human health and visibility. The magnitude of potential benefits in these two areas alone exceed the costs of complying with Title IV. Ecosystem (nonuse) benefits are expected to be large but reasonable estimates are not attainable at this time.

Title IV is a good example of the ^{effectiveness} ~~efficiency~~ of a market-based approach to environmental regulation and illustrates cooperative implementation of an emission allowance and trading program by the Environmental Protection Agency and the utility industry. The accumulated evidence through the end of 1996 strongly suggests the effectiveness of Title IV is unprecedented in keeping emission control costs low while attempting to reduce potential damage from acid deposition to ecosystems, materials, visibility, and human health. A well-designed, market-based system may have clear environmental benefits when compared to a more traditional method of regulation, as has been seen with Title IV, where trading lowered costs and led to early emission reductions. ?

Extrapolation of what has been achieved so far in the efforts to reduce acid rain at low cost should be made with caution. The most valued feature of Title IV -- a utility's flexibility of compliance methods -- is also the least predictable feature. Utility units have pursued a variety of compliance options, including scrubber installation, fuel switching, energy efficiency, and allowance trading. The allowance market has given some sources the incentive to overcontrol their SO₂ emissions and bank their allowances for use in future years. Other sources have deferred control by acquiring allowances from sources that have overcontrolled their emissions. In the aggregate, utilities may choose to comply differently in the future than they did in 1995. Their choices in 1995 led to an early reduction of SO₂ emissions, nationally. Continuation of this early reduction throughout Phase I is expected but is not yet known. In theory, use of a national emissions trading program to deal with a regional air quality problem could cause some areas of the country to experience higher emission (or deposition) levels than would have occurred with no trading. However, preliminary studies based on actual emissions data along with emission and atmospheric modeling projections strongly indicate that this is not the case. Furthermore, all of the human health, surface water, and welfare benefits of compliance choices will not be fully realized for many years because of the inherently long response times in most of these systems.

Although the results from the first two years of implementation of Title IV have exceeded expectations, a number of important environmental and economic questions about the program

1 will demand attention in the coming years. For example, will the year-to-year variability in and
2 geographic distribution of emissions allowed by the flexible market-based approach have any
3 noticeable environmental effect? What will surface water monitoring data reveal about the long-
4 term impacts on aquatic ecosystems? How will the emissions trading market change as utilities
5 approach the tighter emission limits in the year 2000? These questions and many others must be
6 addressed in future years in order to assess whether Title IV is an effective approach to pollution
7 control.

INTRODUCTION

Decision makers inside and outside of government continue to face important issues relating to the emissions of sulfur and nitrogen, their effects upon ecosystems, visibility, materials, and human health, and, ultimately, the control of these emissions. They need easy access to the latest monitoring and research results and scientific discoveries on topics relevant to acid deposition to evaluate the effectiveness of current public policy and to establish a sound science base for future policy decisions.

The history of air quality management includes many examples of society's responses to pollution. During the Middle Ages in London the ever-present cloud of dust and soot and its effects on human health led to prohibitions on coal-burning. During the Industrial Revolution, air pollution was generally considered a municipal problem rather than a public health issue and was managed on a local level, which continued well into the 20th century. The emergence of air pollution in the United States as a public health issue in the 1950s led to the development of federally funded research programs, culminating in the Clean Air Act and establishment of the U.S. Environmental Protection Agency (EPA) in 1970.

Around the world, countries were developing institutional responses to combat the air pollution threat to human health. The effects of air pollution on ecology, however, were not yet considered a serious issue. Having been first documented in England at the end of the 19th century, acid rain and its ecological effects became regional issues in northwestern Europe and in the northeastern United States in the late 1960s. The mounting anecdotal evidence of the harmful effects on aquatic and terrestrial ecosystems launched acid rain as perhaps the first air pollution threat to the environment to receive international attention.

In the late 1970s, the President's Council on Environmental Quality asked a few scientists to initiate a long-term, interagency research and assessment program to study acid rain. With Administration support and Congressional action, the Acid Precipitation Act of 1980 became law. During its first ten years, the research and periodic assessments conducted by the National Acid Precipitation Assessment Program (NAPAP) furthered understanding of the scientific processes and effects of acid deposition. Peer reviews, workshops, and annual reports throughout the 1980s culminated in the NAPAP *State of Science and Technology Reports* published in 1991 and the NAPAP *1990 Integrated Assessment Report*. The monitoring and research conducted in the 1980s and the subsequent integrated assessment provided a significant part of the scientific knowledge base for Title IV of the 1990 Clean Air Act Amendments, known as the Acid Deposition Control Program.

The purpose of Title IV is to reduce the adverse effects of acid deposition through reductions in annual emissions of sulfur dioxide (SO₂) and nitrogen oxides (NO_x), the precursors of acid rain. Recognizing that the principal sources of the acidic compounds and their precursors in the atmosphere are emissions from the combustion of fossil fuels, policymakers initiated control measures to reduce emissions from electric utilities. However, rather than the traditional command-and-control approach to regulation, Title IV provided utilities alternative methods of complying with specific emission limits and deadlines. Alternatives included technological adaptation (e.g., scrubbers, and higher-efficiency boilers), fuel switching, and an innovative SO₂ emission allowance trading and banking program.

Title IV is the first national effort to use market-based incentives to achieve environmental goals. At the time the 1990 Amendments were enacted, it was widely recognized that the emissions trading program was experimental. Due to the innovative nature of using market-based incentives for environmental regulation, and to ease the burden on industry, Congress allowed implementation of the program to be phased in over the following 20 years. Congress built into

the same legislation a mechanism for evaluating how well the experiment is working, both during and after its implementation.

NAPAP Mission and Mandate

Under Title IX of the 1990 Clean Air Act Amendments, Congress reauthorized NAPAP to continue coordinating acid rain research and monitoring, as it had done during the previous decade, and to report the results to Congress biennially, beginning in 1992. In addition, Congress asked NAPAP to assess all available data and information and answer two questions:

1. *What are the costs, benefits, and effectiveness of Title IV?* This question addresses the costs and economic impacts of complying with the Acid Deposition Control Program as well as benefit analyses associated with the various human health and welfare effects, including reduced visibility, damages to materials and cultural resources, and effects on ecosystems.
2. *What reductions in deposition rates are needed to prevent adverse ecological effects?* This complex question addresses ecological systems and the deposition levels at which they begin to experience harmful effects.

The results of the assessment of the effects of Title IV and of the relationship between acid deposition rates and ecological effects were to be reported to Congress quadrennially, beginning with this 1996 Report to Congress. The objective of this report is to address the two main questions posed by Congress and fully communicate the results of the assessment to decision makers. Given the primary audience, it is not written as a technical document, although information supporting the conclusions is provided along with references.

Assessment Process

The preparation of NAPAP assessments every four years requires the interaction of many disciplines, institutions, and individuals. It is important, therefore, that a framework for the assessments be clear. A framework was presented as a strawman proposal for use in preparing NAPAP assessments at the international workshop on "Developing the Framework for the NAPAP Assessment" held November 1993. The general consensus of the workshop participants was that the Tracking and Analysis Framework, or TAF (**Figure 1, also see Appendix A**), or some similar methodology, was a useful and necessary tool to plan, organize, and implement NAPAP assessments. Openness and transparency are cornerstones of the assessment framework and process.

The Tracking and Analysis Framework's detailed structure for NAPAP assessments provides a path to evaluate how acid deposition control decisions, specifically Title IV, can affect reductions of emissions and deposition, health and environmental benefits, monetary and nonmonetary costs and benefits of Title IV, and the reduction in deposition rates needed to prevent adverse ecological effects. This framework serves as an organizing tool to help identify the inputs and outputs between the operational components of the assessment, facilitates communication and information flow, and allows researchers to focus on the connections between, as well as the processes internal to, the individual components.

Policy-relevant assessments bridge the information gap between policy decisions and the social, natural, and physical sciences. An integrated assessment analyzes the linkages among many disciplines and along the full causal chain -- from causes to effects to costs and benefits -- to provide the information necessary to formulate and evaluate national and international environmental policies and strategies.

Since the passage of the 1990 Amendments and the regulatory decisions described in Title IV, NAPAP has coordinated federal acid rain research and monitoring in fulfillment of its mandate. Emission and deposition monitoring have been continued to characterize human and environmental exposure to acid rain and its precursors. Human health and environmental monitoring has been extended to perform routine assessment of the effects of acid deposition and its precursors. More resource-intensive levels of data gathering, model construction, and model application have been employed to characterize the cause-and-effect relationships more accurately, including the impacts of specific Title IV controls.

The major goal of NAPAP's 1996 integrated assessment is to provide structured, technical information in a format that enables decision makers to evaluate the effectiveness of current public policy and provides a sound science base for future policy decisions. It must be credible, open, and communicated fully and responsibly. Secondary goals are to further develop a process for future assessments and to identify the near-term monitoring, research, and modeling needs leading to the 2000 assessment.

The NAPAP assessment process builds upon the experience and lessons of the first 15 years of the program. It uses appropriate data and results from the NAPAP *State of Science and Technology Reports* (NAPAP, 1991a), and *1990 Integrated Assessment Report* (NAPAP, 1991b) and subsequent peer-reviewed literature, as well as lessons learned from other assessment activities in the United States and other countries. To enable the assessment to evolve and to be technically sound and policy-relevant, the assessment team pursued the following objectives:

- Define the policy-relevant questions.
- Adhere to the scientific method, including peer review.
- Treat assumptions and caveats.
- Synthesize multidisciplinary information at all relevant scales.
- Consider feedback mechanisms.
- Use technically sound extrapolation procedures.
- Set confidence intervals on all conclusions and consider uncertainties.
- Consider how to link outputs from one discipline as inputs to another discipline.
- Use only nonproprietary data and models.
- Encourage open and easy access to data.
- Consider the marginal, cumulative, episodic, and chronic consequences of acid deposition.
- Consider integration among affected natural resources and socioeconomic values.
- Include flexible approaches to address changing knowledge, emphases, and policy needs.
- Encourage the use of multiple approaches, models, and valuation schemes.

NAPAP followed four broad steps to conduct this assessment and communicate the results to stakeholders.

1. Define the scope of the assessment.
2. Collect and synthesize issue-specific scientific data and information pertaining to the policy-relevant questions defined in the 1990 Clean Air Act Amendments.
3. Integrate and assess the scientific data and information in a format that facilitates addressing the policy-relevant questions.
4. Communicate the results effectively.

Scope of the Assessment. In order to match assessment requirements with NAPAP agency resources, organizational roles and responsibilities were established. This was done through a series of workshops and meetings held between 1992 and 1995 to define the scope of the assessment and build the organizational structure needed to conduct the assessment. Beginning with the two main questions posed by Congress, a list of key policy-relevant questions were developed with input from experts within the NAPAP agencies, academia, the private sector, nonprofit organizations, environmental groups, industry groups and congressional staff. At two

workshops held in October 1995, these policy-relevant questions were given realistic bounds within the time frame of the report. The smaller, specific issues were identified along with the end points, sources of information, and tools needed to address them. All of the issues were expressed as questions that several research synthesis teams that were identified at the workshops were charged with answering and which serves as the structure for this report. See accompanying text box for the list of questions.

-----Text Box Begins-----

NAPAP 1996 ASSESSMENT QUESTIONS

Question 1. What is the status of implementation, the effectiveness, and the costs and benefits of Title IV of the 1990 Clean Air Act Amendments?

- A. What is the status of implementation (compared to the legislated requirements)?
- B. What emissions reductions have been achieved?
 1. How do the actual reductions (1990-1995) compare with the Title IV benchmark projections (i.e., compared to the expected emissions under Title IV and the projections made in 1990 if Title IV were not implemented)?
 2. What are the reasons for these emissions reductions?
- C. How have air concentrations and levels of deposition been affected by these emissions reductions and how do the new levels compare to the benchmark projections (i.e., with and without Title IV)?
- D. As measured by compliance costs, how effective is the market based approach to emissions control compared to a command and control approach (e.g. benchmark projections)?
 1. What are the factors affecting participation in the banking and trading program?
 2. What market innovations have been achieved?
- E. What are the benefits of Title IV within the United States (in the following areas)?
 1. aquatic ecosystems
 2. forest ecosystems
 3. materials and cultural resources
 4. visibility
 5. human health
- F. Address the following economic benefits questions in relation to the effects areas:
 1. For each of these is there a well identified endpoint that affects human welfare?
 2. If so, for which of these economics (methodology) contribute to an understanding of the impact? Also, recognize that other criteria exists.
 3. For which of these is there relative confidence with economics (existing data)?
 4. For each of these, how are the endpoints linked to Title IV?
- G. The following effectiveness questions should be addressed sequentially:
 1. What are the current physical, chemical, or biological characteristics/states of the sensitive receptors?
 2. What role do these receptors play in maintaining ecosystems?
 3. How have these states changed since 1980 and what are the trends in these changes?
 4. What is the role of acidic deposition controls in these trends?
 5. What is the difference between current conditions (i.e., with implementation of Title IV) and benchmark scenarios (i.e., with and without Title IV)?
- H. What have been the values of the benefits of the emissions reductions compared to benchmark projections (monetary and nonmonetary)?
 1. What are the consequences of emissions allowance trading and banking on the environment?

2. What have been the monetary and nonmonetary values of the benefits of implementation compared to benchmark projections?

I. Where are the significant research gaps in the valuation of benefits, costs, upstream effects (in the pathways) and relative importance of how it affects our ability to do benefits and costs?

Question 2. What are the reductions in deposition rates that are needed in order to prevent adverse ecological effects?

A. What are current deposition rates and what is the variability in time of those rates?

B. How do we address the issue of "adverse" over time?

C. Are there resources whose responses are unique enough to be identified as specific indicators of changes in acidic deposition?

D. What are the dose-response relationships (observed and modeled for sulfur and nitrogen deposition in the effects areas of interest (see Question 1E)?

1. What is the regional extent and magnitude of these responses?

2. How are these key ecological responses generally related to and influenced by other factors (e.g., climate change, ozone, land-use changes, and carbon cycle)?

E. How might effects be reduced if deposition were reduced further?

F. What levels of deposition and associated effects (if any) will remain after implementation of Title IV?

-----Text Box Ends-----

Subsequent to the workshops, the NAPAP Interagency Committee refined the scope of the assessment with guidance from the Air Quality Research Subcommittee of the Committee on Environment and Natural Resources under the President's National Science and Technology Council. The Committee recommended, for several reasons, an abbreviated report be produced for 1996 and a more comprehensive assessment be conducted in 2000. Since 1995 was the first year of implementation of Title IV, it would be too early to observe the changes in most effects areas resulting from a single year of emission reductions. Weather variability and its impact on dispersion, transport, and deposition of pollution would contribute to the uncertainty in identifying a response of ambient concentrations and visibility to emission reductions over the brief span of time after Title IV took effect.

As a result, this 1996 assessment of the costs, benefits, and effectiveness of Title IV focuses on the observed regional changes in emissions, deposition, and effects for the period 1980-1995. Additional analyses is provided on areas of special interest (e.g., Adirondack Mountains of New York) where information is available. Because of the inherent danger in extrapolation from a one year period of record, this report contains no projections. Its approach to answering the question about any needed deposition reductions is to describe the ecosystem responses to reductions along a continuum, thereby allowing decision makers to determine the level of acceptable risk.

An abbreviated report was also necessary because of the limited time and resources available. No new research was to be conducted for the 1996 assessment; it was to be based on existing peer-reviewed data, information, and published federal agency reports. This revised scope of the assessment was presented at a workshop with the research synthesis teams in September 1996.

Synthesis and Evaluation. Seven research synthesis teams composed of NAPAP agency scientists and contractors with extensive experience in the acid rain issue were responsible for

collecting, synthesizing, and updating the scientific data and information necessary to answer their particular questions. The NAPAP *State of Science and Technology Reports* and *1990 Integrated Assessment Report* were used as a baseline. The data were synthesized and made available to the integration and assessment team in question and answer format.

Integration and Review Process. The integration and assessment team was composed of senior agency scientists, NAPAP staff, and a former Director of NAPAP. With the support of the leaders of each of the research synthesis teams, this team integrated the synthesized data into the report format by addressing the linkages and relationships among the many disciplines.

The independent experts who reviewed this report were made aware of the policy-oriented nature of the work and the need for high-quality science. Most of them were asked to review those portions of the report that pertained to their areas of expertise and the integrative aspect of the assessment. Three experts were tasked with evaluating the entire report, from a multidisciplinary, integrated viewpoint. The integration and assessment team incorporated their comments and suggestions into the report for agency review.

Communication of Results. The primary method of communicating the results of NAPAP assessments is through publishing the reports to Congress. It is the responsibility of NAPAP, through the Office of the Director, to inform stakeholders of the results of assessments and receive feedback. To expand the availability of the report beyond those that will receive the printed version and to improve dialogue with interested parties, NAPAP will place the report on the World Wide Web. In addition, the research synthesis teams have been encouraged to publish their individual synthesis and analyses in the open literature.

Organization of the Report

The report is organized to present the assessment results in a clear and logical manner. The status of implementation of Title IV by the EPA and the affected utilities and the costs of compliance is presented first, followed by an analysis of the observed changes, both past and present, in emissions of acid rain precursors, air concentrations and deposition of acidic species. The impact of acid deposition on aquatic and terrestrial ecosystems, visibility, materials and cultural resources, and human health is presented along with the observed and modeled responses by sensitive receptors. The responses are addressed in terms of the benefits of emissions reductions to physical injury and economic valuation. The next section addresses the effectiveness of Title IV, a market-based approach to environmental regulation, relative to the more traditional command-and-control approach to regulation. Finally, an outlook to what can be expected in NAPAP's 2000 assessment is presented, including the monitoring, research, and modeling activities that must be completed for the assessment.

IMPLEMENTATION AND COSTS OF TITLE IV

Title IV of the 1990 Clean Air Act Amendments requires the reduction of acid rain precursors – namely, emissions of sulfur dioxide (SO₂) and nitrogen oxides (NO_x) from electric utilities. This section discusses the status of the Acid Deposition Control Program as required under Title IV of the 1990 Clean Air Act Amendments as well as the compliance costs associated with Title IV. The results of the emission reduction program are discussed in the section on *Changes in Emissions, Concentrations and Deposition*.

The Acid Deposition Control Program has two major goals, which are to be completed by the year 2010.

1. Reduction of total SO₂ emissions by 10 million tons below 1980 levels.
2. Reduction of NO_x emissions from coal-fired boilers that will contribute to the overall target of a 2 million ton reduction below 1980 levels.

Under the SO₂ emission reduction program, utility emissions will be capped at 8.95 million tons per year in 2010, and non-utility industrial emissions will be capped at 5.6 million tons per year. There is no national cap on NO_x emissions.

What is the status of implementation?

SO₂ Implementation Program

- The SO₂ emissions reduction program started on schedule, which reflects the cooperation by participating electric utilities and the Environmental Protection Agency. Phase I was initiated in January 1995; Phase II begins in 2000.

In 1996, there were a total of 431 electric utility units under Phase I of the SO₂ implementation program: the original 263 highest emitting utility units in the country plus 168 substitution and compensating units that chose to comply with Phase I requirements early. In the first two years of implementation, Phase I utility units reduced SO₂ emissions significantly below the allocated allowable levels. SO₂ emissions in 1995 were 39% below the 8.7 million ton allowable level; in 1996, SO₂ emissions were 35% below the target of 8.3 million tons.

NO_x Implementation Program

- The first phase of the NO_x emission reduction program began in January 1996, one year later than originally scheduled due to a U.S. court action regarding language in the initial EPA rule. Phase II becomes effective in 2000.

The 1990 Clean Air Act Amendments require the 2 million ton reduction in NO_x emissions to be achieved by a combination of measures from stationary and mobile sources, as specified under Titles I, IV and V. Title IV, which covers coal-fired electric utility boilers, will contribute a large amount toward the overall emission reduction goal. In 1996, the first year of implementation, Group 1 units (dry-bottom, wall-fired, and tangentially fired boilers) reduced emissions by 33% below 1990 levels (U.S. EPA, 1997).

Beginning in the year 2000, NO_x emissions from Group 1 boilers are expected to be reduced by approximately 1,170,000 tons annually. Group 2 units (wet-bottom boilers, cyclones, cell-burner boilers, and vertically fired boilers) will provide additional NO_x reductions of 890,000 tons per year, thereby decreasing total NO_x emissions by more than 2 million tons annually.

However, because there is no national cap, NO_x emissions are expected to rise again after the full-scale reductions take effect in 2000 under Phase II. To meet the expected growth in electricity generation, existing units will be used more, even though their emission rates are limited, and new units, which are not required to offset their emissions, will be built and operated. NO_x emissions from mobile sources and some industrial sources are also expected to rise within the next 10 years (U.S. EPA, 1995a).

How well have utilities complied with the provisions under Title IV?

- All Phase I units have met their emission reduction obligations. There has been an unprecedented level of accuracy and compliance by utilities in reporting emissions due to installation of continuous emissions monitors on all affected units.
- Allowance trading, a unique aspect of Title IV, has continued to increase since the first trades were recorded in March 1994 and serves as a model for other emission reduction programs.

Significant milestones have already been achieved under Title IV. The first year of compliance for Phase I units was marked by a sharp decline in SO₂ emissions. In addition, all 431 Phase I units met their compliance obligations. In the first annual reconciliation of allowances and emissions, SO₂ allowances matched or exceeded SO₂ emissions generated in 1995 at each of the Phase I units. No excess emissions were reported. A total of 3.44 million tons of unused 1995 vintage SO₂ allowances were "banked" for future use. See the text box for further discussion on the allowance trading system.

-----Text Box Begins-----

Allowance Trading

Allowance trading is a unique feature of the Clean Air Act Amendments. Under this system, emission allowances are allocated to affected utility units based on their historic fuel consumption and a specific emission rate. Each allowance permits a unit to emit one ton of SO₂ during or after a specified year, known as the vintage year. However, emissions cannot violate other program or permit conditions under the Clean Air Act, even if the utility units have the allowances to cover the emissions. Once allocated, allowances are fully marketable. They can be bought, sold, or banked for future use. Sources can also secure allowances through EPA's annual public auction and through EPA awards of special program allowances. The allowance system gives companies the flexibility to transfer allowances and achieve cost savings by coordinating compliance strategies among their various utility units. For example, if a company believes that it will cost more to control a ton of emissions in five years than it will today, it will reduce its emissions more today and bank the allowances for future years.

-----Text Box Ends-----

The volume of allowance transfers has doubled annually since the first transactions were recorded in 1994. By the end of 1996, over 3,100 transactions involving more than 51 million allowances had occurred. Of these 51 million allowances, over 34 million were transferred by private parties; the remainder were transfers from EPA to private parties under various provisions of the Clean Air Act. Approximately 20% of the allowances transferred by private parties to date have occurred among organizations or companies that have no direct economic link; the number of transfers increased from 1.9 million in 1995 to 4.4 million in 1996. About half of these transfers involved utilities acquiring allowances from brokers, fuel companies or other electric utilities. More than 25% of all utilities and 50% of Phase I affected utilities have engaged in these allowance transfers. Utilities in 33 different states participated in these types of transactions (Kruger and Dean, 1997). The majority of activity in EPA's allowance tracking system,

however, continues to involve the shifting or redistribution of allowances within one utility's (or group of utilities') holdings, i.e., for the purpose of covering individual generating unit emissions or for allowance accounting purposes.

Through 1996, a total of 775,000 allowances were sold at four public auctions. In addition, EPA awarded more than 18,000 allowances from the Conservation and Renewable Energy Reserve to utilities that reduced their emissions through energy efficiency and renewable energy projects before the emission reductions deadlines. This growth in awards reflects the increased cumulative number of energy efficiency and renewable energy measures installed since 1992, the first year for which utilities can take credit for measures that earn bonus allowances based on energy savings or renewable generation (U.S. EPA, 1994). The 1990 Amendments also established a program to assist small diesel refiners in defraying the capital costs of installing desulfurization equipment at their refineries by providing SO₂ allowances based on the amount of diesel fuel produced.

In another program initiative, EPA has awarded allowances to industrial sources voluntarily entering the Acid Deposition Control Program. Promulgated in May 1995, the "opt-in" program allows additional combustion sources (e.g., fossil fuel-fired boilers, turbines, or internal combustion engines) not already affected by the program to create allowances through emission reductions. Opting in will be profitable if the revenue from selling allowances exceeds the combined cost of the emission reductions and of participating in the opt-in program. During the first year of implementation of the opt-in program, beginning in May 1995, two sources voluntarily entered the program.

The effectiveness of the allowance trading program has inspired consideration of its application for other emission reduction programs, such as for controlling greenhouse gases. (See **Appendix B** for more information on pollution trading programs.)

What are the compliance costs of Title IV and how do they compare to projections?

- Emission reductions have been achieved at a lower cost than estimated by many studies.

Estimates of total costs of Title IV continue to be revised downward. Much of the cost savings associated with more recent analyses can be linked to reductions in costs and improved performance of scrubbers. In addition, competition between different compliance options and the integration of the allowance and fuel markets may have a downward impact on compliance costs (EPRI, 1993 and 1995). Anecdotal evidence suggests that technology innovation is leading to cost savings (Burtraw, 1996; EPRI, 1996). Allowance prices are also considerably lower than many experts predicted (Hahn, 1994; GAO, 1994). These prices have been influenced by developments in the low-sulfur coal market, particularly greater compatibility than expected of western, sub-bituminous coals with eastern boilers designed for bituminous coals, reduced rail rates for delivering western low-sulfur coal to some utilities in the midwestern United States, and lower than expected future costs of scrubbers for Phase II of Title IV (EPRI, 1996 and 1997). Incentive or bonus allowances that encouraged retrofit flue gas desulfurization on Phase I units led to the building of flue gas desulfurization technologies, which has kept the marginal costs of compliance so far in Phase I, lower than estimated by prior projections.

The significance of these lower costs is twofold. First, they show that it is often difficult to estimate future technological improvements and the more efficient use of existing technologies. For example, significant cost reductions in renewable and energy-efficient end-use technologies are expected in the coming years, but the overall contribution of these technologies to reductions in future environmental compliance costs is difficult to estimate. Economic models that do not estimate these factors may overstate compliance costs for environmental programs. Second, these

lower costs again illustrate the benefits of a flexible approach to compliance that allows different technologies and fuels to compete against each other, and rewards firms for finding cost-effective measures that exceed emission-reduction targets.

Methods of Compliance

- SO₂ emissions reduction in 1995 was achieved by an almost equal split between scrubbing and fuel switching.

Table 1 pairs emission reductions at Phase I units with the method of emission control used; where the method was switching, the table lists the source of the lower sulfur coal. In the aggregate, slightly more than half of the reduction came from switching to lower sulfur coals, but the contribution from scrubbers was significant. Twenty-six units installed scrubbers under Phase I; these units accounted for 45% of the reduction accomplished in 1995 (Ellerman et al., 1997), and for 62% of the net surplus of allowances banked in 1995 (EPRI, 1997).

Low-sulfur western coal from the Powder River Basin continued to play an important role in reducing SO₂ emissions, but the main contribution to reducing emissions by fuel switching in 1995 came from the bituminous coal-producing regions, mostly in Central Appalachia. Although switching is often seen as changing the source of the coal from a high-sulfur region to a geographically distinct low-sulfur region, much of the reduction occurring in 1995 resulted from fuel switches within a region. This phenomenon was observed in all regions, including the predominantly high-sulfur coal-producing regions of North Appalachia and the Midwest, where the coal was typically mid-sulfur rather than the conventional low-sulfur coal.

Table 1. Methods used to reduce sulfur dioxide emissions in 1995. *Source: Ellerman et al., 1997*

1995 SO ₂ Reduction at Phase I Units		
Method/Region	Tons SO ₂ Removed (in thousands of tons)	% of Total
Scrubbing Total	1,755	45.1 %
New Title IV Scrubbers	1,734	44.6%
Other Scrubbers	21	0.5%
Switching Total	2,133	54.9 %
North Appalachia	205	5.3%
Central Appalachia	756	19.5%
South Appalachia	60	1.5%
Midwestern	406	10.4%
Powder River Basin	518	13.3%
Other Western Coal	146	3.8%
Imported Coal	22	0.6%
Natural Gas	20	0.5%
TOTAL	3,887	100.0 %

Associated Costs of Compliance

Preliminary results from a retrospective analysis show the total annualized cost of the 3.9 million ton reduction of SO₂ in 1995 was about \$726 million (Ellerman et al., 1997), which is at the lower bound of earlier predictions (Table 2). In particular, the observed cost of compliance by scrubbing has been markedly lower than expected. This experience is influencing today's allowance prices, which are more closely related to marginal compliance costs, rather than average costs that include the capital costs of flue gas desulfurization. Table 3 compares the expected SO₂ removal cost for a representative retrofitted unit (as projected by ICF in 1990 and EPRI in 1993), with actual SO₂ removal cost for the scrubbed units that were fully operational in 1995 (as reported by MIT in 1997). By 1995, Phase I scrubbers were removing sulfur at an average total cost of \$282 per ton, or about 40% less than what had been predicted by earlier estimates of average total overall cost. The cost savings arise from two factors: lower operating and maintenance costs, particularly fixed costs, and more intensive utilization of generating units with scrubbers (83% capacity factor in 1995 versus 65% as assumed in most earlier studies).

No attempt was made in Table 3 to quantify emissions reduced through demand-side energy-efficiency programs. However, some companies and state public utility commissions have analyzed the impacts of these programs on emissions and compliance strategies, and others have analyzed how the value of SO₂ emissions avoided by energy efficiency and renewable energy programs improve the cost-effectiveness of these technologies (U.S. EPA, 1994; Kruger et al., 1995).

Lower-than-expected fixed operating costs appear to be a result of reduced personnel costs due to greatly improved instrumentation and controls on new scrubbers. Increased competition in the industry has also added to pressures to reduce these costs. Lower variable costs appear to be because of reduced requirements for power to run scrubbers as a result of better controls. Utilities have also used some innovative sludge disposal techniques to reduce these costs (Ellerman et al., 1997).

Table 2. Estimates of Phase I compliance costs (in 1995 dollars). *Source: Ellerman et al., 1997*

Study	Total Annual Cost (in millions of 1995 \$)
ICF 1989 (low, constrained)	\$871
ICF 1989 (low, flexible)	\$599
ICF 1990 (low, flexible)	\$573
EPRI 1993	\$1,338
GAO 1994	\$1,163
EPRI 1995	\$894
Actual '95 (Ellerman et al., 1997)	\$726

Table 3. Scrubber costs (per ton SO₂ removed, in 1994 dollars). *Source: Ellerman et al., 1997*

Types of Costs (per ton SO ₂)	ICF 1990	EPRI 1993 ^a	Actual 1995 ^b
Capital Charge	\$285	\$262	\$203
Fixed Operation & Maintenance	\$66	\$83	\$14
Variable Operation & Maintenance	\$104	\$129	\$65
Total Cost	\$455	\$474	\$282

^a The representative retrofitted unit used in EPRI (1993) is a 300 megawatt unit with a retrofit difficulty factor of 1.27. The unit is assumed to remove 90% of the sulfur from a 3.97 lb coal and to be operating at a 65% capacity factor and a gross heat rate of 9,722 Btu/kwh.

^b Data is from a MIT questionnaire given to affected utilities as reported in Ellerman et al., 1997.

Since enactment, the projected cost of compliance with the SO₂ emission reductions has declined with each analysis. In 1990, EPA estimated the annualized cost of the SO₂ reductions to be \$4 billion by the year 2010, assuming full inter-utility trading. For its regulatory impact analysis in 1992, EPA estimated the annualized cost in 2010 at \$3.7 billion. In 1993, the Electric Power Research Institute estimated Phase II costs with full inter-utility trading to be \$2.2 billion. In 1994, the General Accounting Office estimated that with full inter-utility trading the annualized cost in 2010 should be less than \$2.0 billion.

Allowance prices have reflected these declining estimates of control costs. The price of an allowance has dropped from an estimated \$500-600 per ton when the Clean Air Act Amendments were passed to a low of \$68 per ton at the March 1996 allowance auction; the price at the end of 1996 was about \$90 per ton (**Figure 2**). A number of hypotheses have been offered as to why allowance prices are so much lower than expected.

- The low price and widespread availability of low sulfur fuel reduces allowance prices. Moreover, lower rail costs have made low-cost coal from the Powder River Basin available to midwestern utilities that are "on the margin" in choosing their compliance strategies.
- Scrubber costs, particularly variable costs, are much lower than expected. This has lead utilities to run units with scrubbers at high utilization rates, as long as the value of the extra allowances generated is greater than the variable costs of scrubbing.
- Projected future costs of scrubbing for Phase II could be lower than current costs.
- Costs incurred through fuel switching or scrubbing in Phase II, when the full effect of emission reductions will be realized, should be discounted by the rate of interest for comparison with current period allowance prices.
- The lead time for large capital expenditures on scrubbers required utilities to estimate future compliance costs and allowance prices. However, expectations in the early 1990s were that the marginal costs of compliance and allowance prices would be considerably higher than their actual levels, leading to overinvestment in scrubbers, overcontrol of emissions, excess allowances, and lower allowance prices.

- 1 • Competition between compliance options (e.g., low-sulfur fuels, more intensive
2 utilization of scrubbers, and bundling of higher sulfur fuels with allowances) has led to
3 lower marginal costs and allowance prices.
- 4 • Bonus allowances and regulatory incentives for scrubbing have led to excess allowances
5 and lower allowance prices.
- 6 • Substitution and compensating units depress allowance prices because they have lower
7 marginal costs for compliance.
- 8 • The Title IV auction design artificially depresses allowance prices.
- 9 • Uncertainty about possible future environmental regulations reduces the demand for
10 allowances.
- 11 • State public utility commission policies and regulations are a disincentive to allowance
12 trading and depress allowance prices.

13 These hypotheses are not mutually exclusive. They can be grouped into two categories: 1)
14 hypotheses that assume that low costs are due to underlying market fundamentals (the first five
15 listed), and 2) hypotheses that assume regulatory or other distortions of the market. While most
16 observers agree with at least some of the hypotheses in the first category, there is more debate
17 about hypotheses in the second category. Further discussion about several of these hypotheses
18 appears in the banking and trading discussion later in this chapter.

19 **Market-Based System**

- 20 • The market-based approach in Title IV has achieved significant savings when compared to
21 a command-and-control regulatory approach.

22 A particularly notable feature of Title IV is the cap on utility emissions and the choice of fully
23 tradable emission permits to achieve the environmental objective. The appeal and promise of such
24 market-based instruments is a lower cost of compliance than would be achieved under an
25 equivalent command-and-control instrument. Earlier studies of compliance costs with Title IV
26 have provided estimates of the potential savings, derived by both the differences in predicted costs
27 between scenarios that assumed more and less trading, and variations among these studies in the
28 amount of assumed emissions trading and in the corresponding costs:

- 29 • ICF's 1989 analysis predicted the cost savings for limited trading among utilities within
30 the same state to be approximately \$225 million (in 1988 dollars), about 33% of the
31 predicted cost of compliance in the "less trading" case. [ICF's 1990 analysis does not
32 provide a comparable estimate of cost savings from emissions trading because it assumed
33 flexible implementation ("trading to the fullest extent permitted") for both the higher- and
34 lower-growth scenarios modelled.] In addition, ICF89 estimated that banking provisions
35 might result in an additional annual reduction of emissions in Phase I of between 0.8 and
36 1.2 million tons, resulting in a present-value cost savings of \$150 million, about a third of
37 the predicted cost of avoided Phase II emissions reductions.
- 38 • EPA's 1992 Regulatory Impact Analysis predicted cost savings of 40%, or \$400 million
39 (in 1990 dollars), from a "national SO₂ trading program" as compared with a "traditional
40 SO₂ control program."

- The 1994 GAO report provided an estimate from intra-utility trading only, because very little inter-utility trading was expected: \$230 million (in 1992 dollars), or 18% of the predicted cost of compliance under a "traditional command-and-control" approach.

Most recently, MIT has made a preliminary estimate of the cost savings of 25-34% of the costs incurred in 1995, or \$225-\$375 million (Ellerman, 1997).

Several studies have estimated the cost savings from the allowance trading program as compared to a command-and-control approach for Phase II of Title IV. One study estimates the annualized cost of compliance with the Phase II SO₂ emissions cap in 2010 with full inter-utility trading to be less than \$2 billion, compared to an annualized cost of compliance without trading of \$4.9 billion (GAO, 1994).

Technology Innovations

- Increased technology efficiency and innovation have been achieved from the market-based approach.

Recent studies have highlighted the efficiency and innovation in pollution controls that have accompanied implementation of Title IV (Burtraw, 1996; EPRI, 1996). For example, scrubber costs have declined dramatically in the past six years and are now 40% or more below 1989 levels. At the same time, scrubber sulfur removal efficiencies have improved from 90-92% in 1988 to 95% or more in retrofits to utility units affected by Phase I.

There have also been innovations in blending of high- and low-sulfur coals. In the past, it was assumed that blending would cause problems for utility boilers, but technical difficulties have been largely worked out through experimentation prompted at least in part by Title IV compliance requirements and flexibility.

Banking and Trading Program

- The SO₂ allowance market has become more active and has begun to take on some of the attributes of a more established commodity market. However, there is still debate about the scope and magnitude of opportunities for further trading.

The sulfur dioxide allowance market is becoming an effective private market for trading allowances, and many of the early doubts about the emergence of active allowance trading have not been realized. A wide cross-section of electric utilities have engaged in allowance trading. Nearly half of the 1995 allowance issuance has been traded through time or space, which indicates that electric utilities are availing themselves of the opportunities for cost savings created by the emissions trading provisions of Title IV. On the other hand, there is still some dispute about whether there are barriers preventing even wider development of the market and additional cost savings.

Some analysts contend that state public utility regulations and policies have had an adverse effect on electric utilities' decisions to trade. Advocates of this position note that uncertainty about how commissions will treat the recovery of allowance costs may dissuade utilities from allowance trading. Moreover, where Public Utility Commission policies prevent utilities from keeping the profits from allowance trades, there is a reduced incentive to trade. However, one analysis argues that there is little evidence of such adverse effects (Bailey, 1996). As of January 1996, only 15 state Public Utility Commissions (out of the 35 states where trading has occurred) had explicitly addressed the issue of allowance trading through the issuance of a formal generic order and/or an informal guideline. Trading does not appear to have been impeded in the 20 states where state commissions have yet to address the rate-making treatment for allowances.

1 Another possible barrier to allowance trading is the design of the Title IV auction. Some observers
 2 have asserted that the price-discriminating auction design has depressed allowance prices and has
 3 thereby discouraged utilities from trading (GAO, 1994; Cason, 1994). However, a more recent
 4 analysis argued that regardless of whether the design of the auction was optimal, there is little
 5 evidence that it harmed the allowance market. In fact, the early auctions were probably a more
 6 accurate reflection of the underlying costs of SO₂ control than were the projections of most
 7 analysts (Joskow et al., 1996).

8 The broader question raised by these debates about market barriers is whether all cost-effective
 9 trades are taking place. Some analysts have estimated that the full cost-saving potential of
 10 allowance trading is not being reached (GAO, 1994; Burtraw, 1996). Others have argued that the
 11 relatively active allowance market provides evidence of cost-effective trades (Joskow et al.,
 12 1996).

13 The Title IV compliance data for 1995 provides an exact accounting of vintage allowances, and in
 14 particular, how many allowances were traded. The 8.69 million 1995 vintage allowances can be
 15 divided into three categories of use: (1) 4.77 million were used to cover emissions by generating
 16 units that were issued allowances equal to or greater than this number; (2) 0.53 million were used
 17 by almost 100 Phase I affected units (about 20%) to cover emissions in excess of the allowance
 18 issuance to each unit; and (3) 3.39 million allowances were not used in 1995, but were reserved
 19 for compliance in future years. The second use category corresponds to the minimum amount of
 20 spatial trading, and the third corresponds to inter-temporal trading or "banking." These two uses
 21 of vintage 1995 allowances, accounting for a total of 45%, provide the statistics concerning the
 22 effectiveness of Title IV's tradable permit program.

23 Several estimates have been made, based on transactions in EPA's allowance tracking system, of
 24 overall allowance trading levels (i.e., not just 1995 allowances used for compliance in 1995). For
 25 example, one estimate shows that at least 6.7 million allowances were traded as of March 1996
 26 (Joskow et al., 1996). Using a slightly different estimation technique, EPA has estimated that at
 27 least 7.2 million allowances were traded as of the end of 1996. EPA's estimate includes
 28 allowance transfers between utilities, between brokers or traders and utilities, between fuel
 29 companies and utilities, between different brokers and traders, and several other types of transfers
 30 (Dean and Kruger, 1997). The Electric Power Research Institute has estimated that through
 31 1996, 6.8 million allowances have been traded between different companies in about 960
 32 recorded transactions (EPRI, 1997). Of course, both of these estimates are lower bound
 33 estimates because only those trades entered into EPA's allowance tracking system are included.
 34 The real level of trading is higher, since many trades are reportedly not entered into the tracking
 35 system until they are needed for compliance (Ellerman et al., 1997; EPRI, 1997).

36 **Administrative Costs**

- 37 • Government administrative costs for Title IV implementation have been lower than under
 38 more traditional approaches.

39 The fundamentally different approach to air pollution control embodied by the allowance trading
 40 program can minimize many administrative costs associated with command-and-control and
 41 previous trading programs. For example, Title IV's performance-based approach eliminates the
 42 need to devise source-specific emission limits and to review control technologies and detailed
 43 compliance schedules. In addition, eliminating case-by-case review and approval of each trade
 44 (including determining the "useful life" of equipment, the intent of the sources regarding future
 45 emissions and activity levels, and "real" emission reductions achieved), greatly reduces the
 46 administrative and transaction costs associated with emissions trading programs (McLean, 1996).

1 The program's administrative costs of approximately \$12 million per year translate into a cost of
2 about \$1.50 per ton of pollution reduced. Most of these administrative costs are associated with
3 operating the emissions monitoring and reporting components of the program. To put these
4 expenditures into context, during the first five years of the program, government spending to set
5 up and operate the SO₂ allowance program totaled less than \$60 million out of a total \$3.5 billion
6 estimated government expenditure for air pollution control. Thus, the program is achieving 40%
7 of the emissions reductions under the Clean Air Act with only about 2% of the staff and other
8 resources (McLean, 1996).

CEMS costs?

CHANGES IN EMISSIONS, CONCENTRATIONS AND DEPOSITION

There is no universally accepted method for measuring long-term trends in key air pollutants. The intensive ambient air monitoring networks operated by EPA and the states are currently focused on urban areas where ambient air concentrations mainly reflect local emissions, which are relevant to the human health issue. Besides representing the inhalation exposure of humans, air concentrations play a significant role in visibility, the reflection of incoming solar radiation (radiative transfer), and the indirect measurement of dry deposition.

The intent of NAPAP is to take a more regional view by also using sites related to ecosystem effects and regionally representative deposition and visibility data. Analysis is based, for the most part, on direct observational evidence. A clear distinction is made between observed emissions data and model calculations. In addition, a distinction is made between the contribution to total atmospheric emissions of SO₂ and NO_x by utilities, both Phase I and II, and by other sources.

This section examines changes in acid deposition patterns and relates them to changes in emissions of precursor pollutants. Current and historical emissions of relevant pollutants are presented along with the resulting air concentrations and deposition at specific receptor locations. Quality-controlled emissions and deposition data through 1995 were available for this assessment. Year-end, quality-controlled emissions data for 1996 were not released in time to be incorporated into this assessment, but the data are inserted in footnotes where appropriate.

-----Text Box Begins-----

Emissions Monitoring

The advent of Title IV brought about a change in the method for tracking trends in current emissions. To ensure compliance, the statute required each affected combustion unit (boiler or turbine) to install a continuous emissions monitoring system – or for oil and gas units, an approved alternate measurement method – to record the concentration and mass of emissions on an hourly basis. Prior to 1994/95, emissions were estimated primarily from emissions factors and information filed with the Department of Energy's Energy Information Administration.

With the current system, emissions data are collected directly from the continuous emissions monitors, which are maintained according to rigorous quality assurance standards of accuracy. The data are then electronically transferred into EPA's Electronic Tracking System, ensuring the quality of the data and eliminating a source of errors potentially caused by repeated data entry. EPA then reconciles the data annually with the unit's SO₂ allowance holdings to ensure that the SO₂ emissions are equal to or lower than the allowances.

Any change in measurement methodologies will raise questions about whether observed differences in emission trends are due to the new method or to actual emission reductions. How continuous emissions monitoring compares to the methodology for calculating historical emissions is still under investigation. However, for the 263 original Phase I units, there is continuous emissions monitoring data before Phase I began (1994) and for the first year of Phase I (1995). Between 1994 and 1995, these 263 units reduced their SO₂ emissions by 40% (almost 3 million tons).

Continuous emissions monitors must undergo a relative accuracy test. The test ensures that the installed monitor measures the "true" value of the pollutant by comparing the monitor to a reference method, which is used to measure the pollutant in the stack gas concurrently with the monitor. Thus far, the continuous emissions monitoring data collected under the Acid Deposition Control Program are remarkably accurate. Although there was some concern that the Phase II utilities (generally smaller companies than those in Phase I) might not achieve the same rigorous quality assurance standards as their Phase I counterparts, 98%

of the almost 5,000 monitors tested for the Acid Deposition Control Program passed the required 10% relative accuracy standard (U.S. EPA, 1997).

The operational record for the continuous emissions monitors is excellent. The monitors only require maintenance a few days a year. Over half of all monitors used to report data for the Acid Deposition Control Program were up and running for more than 98% of the last year, and 92% of all monitors were running for more than 90% of the last year (U.S. EPA, 1997).

-----Text Box Ends-----

What emission reductions have been achieved?

Although SO₂ control provisions of Title IV only began in January 1995, dramatic emission reductions that year provide a unique opportunity to place some significance on the analysis of just one year of data. This cannot be said for NO_x provisions since they were not implemented until January 1996, which was too late for a meaningful analysis of Title IV effectiveness regarding NO_x. Discussion of SO₂ and NO_x emissions in this section is further divided into two categories of sources: electric utilities and all other sources.

SO₂ Emissions

- The United States has already achieved a large portion of the SO₂ emission reduction goals mandated under Title IV. Significant emission reductions have been achieved by mid-western states and in other areas that have electric utilities with historically high emission rates.

SO₂ emissions have been declining slowly since the early 1970s when the Clean Air Act was first passed. In 1980, the baseline year for Title IV, total emissions from all sources of man-made SO₂ were about 26 million tons. By 1995, total SO₂ emissions had dropped to 18.3 million tons, with utilities contributing about 12 million tons (**Figure 3**). There was a dramatic decrease in emissions in 1995, illustrating the first year of compliance under Title IV. The reduction of SO₂ emissions from 1994 to 1995 was approximately 19%, all of which can be attributed to reductions from Phase I utilities. In 1995, Phase I units achieved an overall reduction of 5.6 million tons from 1980 levels, which is a 51% drop in emissions.

There have been major emission reductions in some of the largest, highest-emitting areas of the country. This outcome is important for two reasons. First, underlying Title IV was the understanding that emissions in the Midwest are a significant cause of deposition in the northeastern United States and eastern Canada. The intent of Congress was to reduce these emissions through the 1990 Clean Air Act Amendments, which is now being realized. Second, reductions in these high-emitting areas support a fundamental premise of Title IV's market-based approach – i.e., that the highest-emitting plants have an incentive to make deep reductions in SO₂ emissions because they often face a lower cost per ton of SO₂ reduced due to compliance flexibility. So far, concerns expressed at the time Title IV was enacted that the highest emitters of SO₂ would simply buy allowances and continue to emit have proven to be unwarranted. Electric utilities have the flexibility to use banked allowances, which has spurred early reduction in emissions.

Market-based mechanisms, such as emissions trading, are often thought to be designed entirely for cost-saving purposes. However, the banking provision in Title IV illustrates that a properly designed market-based mechanism may have clear near-term environmental benefits when compared to a more traditional method of regulation.

Of course, the banking provision is also projected to extend the time for achieving the ultimate emission reduction target – the 8.95 million ton cap. Most banked allowances are expected to be used between the years 2000 and 2009, after the more stringent emissions limits of Phase II of the Acid Deposition Control Program take effect. Further research is needed to determine whether there is a net environmental or health benefit from the large early emissions reductions spurred by the banking provision, compared to a scenario with no early reductions but full achievement of the cap in the year 2000.

Electric Utilities

- As of 1995, electric utilities reduced their SO₂ emissions significantly beyond the level required; total utility SO₂ emissions dropped to 11.9 million tons, which was 31% below 1980 levels and within 3 million tons of the 8.95 million ton cap.

In 1995, the first year of compliance, the allowable level of SO₂ emissions from Phase I electric utilities was 8.7 million tons. However, only 5.3 million tons of SO₂ were emitted, which was 3.4 million tons (39%) lower than the level required (**Figure 4**) (see footnote). These unprecedented early emission reductions are largely believed to be a direct result of two key features of the banking provision in Title IV. First, Title IV provides a performance-based, rather than a technology-based standard for reducing pollution. It gives each unit a performance target in the form of an emission allowance allocation or budget, but does not require the unit to use a specific technology to reach this target. Second, the banking provision of the allowance trading system gives companies the flexibility to control emissions beyond their performance standards, in effect saving part of their emissions budget for future years. Many companies have adopted this strategy of overcontrolling their emissions during Phase I and are banking their extra allowances to meet the more stringent requirements of Phase II (Ellerman et al., 1997).

As shown in **Figure 5**, SO₂ emissions have decreased from 1980 levels in most of the 21 states containing Phase I units. In 1995, Ohio, Missouri, Indiana, Illinois, and Tennessee had the largest overall decline in emissions (3.4 million tons) from 1980 levels. SO₂ emissions increased in a few states, including Texas, North Dakota, Louisiana, and Oklahoma, which have no Phase I units; these states built new coal-fired powerplants that came on-line after 1980.

There is general consensus that the large reduction of SO₂ emissions in 1995 was a direct result of the Clean Air Act Amendments. A more difficult question is what led to reductions of SO₂ in the early part of the 1990s. Answering this question requires estimating SO₂ emission levels in the absence of Title IV. This estimation is fraught with uncertainty because it requires making difficult assumptions about how coal markets and electric utilities respond to a variety of interdependent economic and regulatory forces, and whether the efforts to pass legislation in the late 1980s had any effect on fuel supplies and purchases. Perhaps a more important aspect of this question is that it indirectly highlights the compliance flexibility inherent in Title IV, allowing electric utilities to take advantage of market forces and the changes in the relative prices of compliance options (Burtraw, 1996). Even if some of Title IV's compliance cost savings had accrued without the 1990 Amendments – because of changes in the railroad industry and low-sulfur coal markets or state acid rain regulations – these hypothetical cost savings might not have materialized under a more traditional regulatory structure (e.g., mandatory scrubbers at all utility units).

EPA reports that in 1996 these units emitted 5.4 million tons of SO₂, which was 2.9 million tons (35%) below the 8.3 million ton allowable level for 1996. All participating units were in compliance.

Certainly a major impact on Title IV costs and SO₂ emissions was the change in the economics of choosing low-sulfur coal. Railroad deregulation during the 1980s introduced competition into the transportation of low-sulfur coal from the Powder River Basin (Wyoming and parts of Montana) and led to large productivity improvements in rail transportation. The result has been continually lower rail rates that have made distant western coals more competitive in midwestern markets on a purely Btu basis (heat content), without regard to the significant difference in sulfur content. As a result, the frontier for western, low-sulfur coal has advanced steadily eastward to the disadvantage of local, higher sulfur midwestern coals.

Much of the emission reductions between 1985 and 1993 may have occurred even without Title IV, as declining costs for low-sulfur coal would have made fuel switching attractive to midwestern electric utilities for purely economic reasons (Ellerman and Montero, 1996). One study estimates that the economics of coal choice explains a significantly larger share of SO₂ reductions during this period than such factors as early compliance by electric utilities or state acid rain programs (Ellerman and Montero, 1996).

Perhaps anticipation of impending Clean Air Act legislation in the 1980s led western low-sulfur coal producers to invest in new low-sulfur coal properties and expanded existing low-sulfur coal mines (EPRI, 1996). This foresight helped them avoid upward pressure on low-sulfur coal prices in the early 1990s. On the other hand, some of this expansion of low-sulfur coal production in the West could have been spurred by the 1977 amendments to the Clean Air Act.

Anticipation of new legislation may also have spurred experimentation with boiler modification and coal-blending techniques that ultimately allowed electric utilities to burn different types of coal at lower costs than expected (Burtraw, 1996). For Phase I units that used flue gas desulfurization technologies, emissions were reduced by 2.0 million tons below their allowance allocation in 1995, while most of the remainder of the reductions were achieved by coal switching or blending (EPRI, 1997).

Ascertaining how much of an emission reduction would have happened if there had been no Title IV is a necessary step to estimate Title IV's costs and benefits. To the extent that some level of emission reductions may have occurred without the 1990 Clean Air Act Amendments, cost estimates should be revised downward to reflect "no cost" reductions. Benefits directly attributable to Title IV would also be less.

Table 4 provides estimates from a variety of studies of 1995 Phase I electric utility SO₂ emission reductions that are directly attributable to Title IV, as opposed to those that would have occurred even without Title IV. Most of these estimates were developed before the end of 1995, but this table provides a snapshot of the range of projections that were made regarding Title IV's effectiveness in reducing emissions. Actual data are now available for some of the modeling variables – such as actual electricity demand for 1995 – that were only estimated for most of these studies. Therefore, the emission rates assumed in the "without Title IV" case are now the key variables in estimating the 1995 emission reductions attributable to Title IV. By varying the emission rate, a range of estimates can be developed that differs slightly from those in Table 4 but that benefits from our actual knowledge of utilization in 1995. This new range has 3.12 million tons as the lower bound (assumes *none* of the reduction in emission rates for 1985-1995 is attributable to Title IV) and 5.69 million tons as the upper bound (assumes *all* of the reduction in emission rates is attributable to Title IV) (Ellerman et al., 1997).

1 Table 4. Title IV reductions of sulfur dioxide. *Sources: NAPAP, 1991; Ellerman et al., 1997*

Projected SO ₂ Emissions (millions of tons)			
Study	Emissions Without Title IV	Emissions With Title IV	Reduction Attributable to Title IV
NAPAP 1990	19.31	13.62	5.69
ICF 1989 (constrained)	16.64	13.08	3.56
ICF 1989 (flexible)	16.64	13.32	3.32
ICF 1990 (flexible)	16.64	13.52	3.12
EPRI 1993	16.29	11.93	4.36
EPRI 1995	14.65	11.47	3.18
MIT 1997	15.70	12.02	3.68

2 Other Sources

- 3 • Total SO₂ emissions from non-utility sources are 2.1 million tons below 1980 levels,
4 which exceeds the Title IV reduction goal of 1.5 million tons.
- 5 • For every predicted scenario, there were substantial reductions in SO₂ between 1990 and
6 1995, when compared with Title IV benchmark projections.

7 An estimated 8.4 million tons of SO₂ were emitted from all other sources – industrial, residential,
8 commercial, and transportation – in 1980. The largest contributors were industrial fuel combustion
9 and metals processing, which contributed about 2.9 and 1.8 million tons of SO₂, respectively.
10 NAPAP estimated 1985 emissions from non-utility sources to be 6.1 million tons, with 3.3 million
11 tons from fossil fuel combustion (Placet et al., 1991). Emissions from non-utility sources were
12 estimated to be 5.7 million tons in 1990, with industrial combustion making up 3.1 million tons
13 and metals processing 663,000 tons. In 1995, emissions from these non-utility source categories
14 were on the order of 6.3 million tons; industrial combustion accounted for 3.0 million tons, and
15 metals processing accounted for 692,000 tons.

16 **NO_x Emissions**

17 In 1980, the baseline year for Title IV, total emissions from all sources of man-made NO_x were
18 about 23.3 million tons. By 1995, total NO_x emissions amounted to about 21.8 million tons
19 (Figure 6). The 1990 Clean Air Act Amendments require NO_x emissions from coal-fired boilers
20 to be reduced by 2 million tons below 1980 levels by 2010; there is no national cap for NO_x
21 emissions.

22 Electric Utilities

- 23 • Annual NO_x emissions from electric utilities have decreased by 800,000 tons from 1980
24 levels.

Only a modest decrease in NO_x emissions from electric utilities has occurred between 1980 and 1995. The Phase I requirements for NO_x reductions became effective after this time period in January 1996. NO_x emissions from utilities were about 6.2 million tons in 1995, or approximately 11% below 1980 levels and 17% below 1990 levels (see footnote). This recorded decline in emissions is thought to be due to several factors, including early installation of Title IV NO_x controls at some facilities, installation of NO_x controls in the northeastern United States to meet Reasonably Available Control Technology standards under Title I, and a change in NO_x monitoring and estimation techniques between 1994 and 1995.

Other Sources

All other sources of NO_x emissions were estimated to be about 16.3 million tons in 1980. The largest non-utility contributors were on-road vehicles and industrial fuel combustion, contributing about 8.6 and 3.6 million tons of NO_x, respectively. In 1995, emissions from all non-utility source categories were estimated to be about 15.5 million tons, with 7.6 million tons coming from on-road vehicles and 3.1 million tons from industrial combustion.

-----Text Box Begins-----

Formation and Transport of Acid Deposition

The natural and anthropogenic sources of emissions consist primarily of gaseous sulfur dioxide (SO₂), nitrogen oxides (NO_x), a variety of volatile organic compounds (VOCs), and fine particulates (**Figure 7**). SO₂ and NO_x are the precursors of acid deposition. Once the primary, gaseous pollutants are emitted into the atmosphere they can be removed from the air and deposited continuously onto surfaces (dry deposition). Depending on meteorological conditions, they can be transported long distances by the wind before being deposited. Gaseous pollutants not removed by deposition are gradually transformed through a variety of reactions to secondary pollutants (e.g., fine particulate sulfate and nitrate, sulfuric acid aerosol, ozone), which can also be deposited as acids if dissolved in cloud droplets or rainwater (wet deposition).

The rate of transformation of primary to secondary pollutants varies significantly in time and space. The length of time airborne pollutants stay in the atmosphere is determined primarily by the meteorological environment into which they are emitted and by their interaction with cloud and precipitation systems. The area affected by the emission sources is determined to a large extent by the time that pollutants stay in the atmosphere before removal.

-----Text Box Ends-----

How have air concentrations been affected by these emissions reductions?

Sulfur

- Emission reductions imposed by Title IV have caused a reduction in air concentrations of SO₂. This reduction is most evident in the Northeast but apparently extends across the entire region east of the Mississippi River.

In 1996, the first year of implementation, Phase I utility units reduced NO_x emissions by 33% (340,000 tons) below 1990 levels. These units demonstrated an average of 18% overcompliance with required emission rate levels.

1 • Reductions in concentrations of particulate sulfate are also widespread, but are less
2 striking. At a few locations, particulate sulfate concentrations appear to have increased
3 when decreases would have been expected; the reasons remain obscure, and the data record
4 is too short to provide insight to this apparent inconsistency.

5 Ambient air concentrations of SO₂ decreased steadily in the years prior to implementation of Phase
6 I and decreased significantly in the year following implementation (see **Figure 8**). Between 1986
7 and 1995, national SO₂ concentrations decreased 37%; SO₂ emissions decreased 18%. Between
8 1994 and 1995, national SO₂ concentrations decreased 17%, while emissions decreased 13%
9 (U.S. EPA, 1996). These data were obtained from observations of air quality at 473 sites across
10 the United States. The sites are generally located in urban areas as part of a regulatory network
11 used to monitor six pollutants (including SO₂) covered by the National Ambient Air Quality
12 Standards under Title I of the Clean Air Act.

13 The benefits of the emission reductions are clearly evident in air concentration records derived at
14 sampling stations operated by the two major monitoring networks addressing dry deposition and
15 concentrations in surface air – the EPA Clean Air Status and Trends Network (CASTNet) and the
16 NOAA Atmospheric Integrated Research Monitoring Network (AIRMoN). Both of these
17 networks are composed of sites primarily in rural areas and are not part of the regulatory network.
18 The benefits are most evident in SO₂ concentration data obtained in the region most highly affected
19 by acid rain – the corridor downwind of the Ohio River Valley and its surrounding area, mainly the
20 northeastern United States. However, the length of record is currently too short to report with
21 confidence that the effects are as widespread or as large as has been predicted. Data are still being
22 collected and analyzed. The present conclusions represent only a preliminary interpretation of a
23 complicated data record in which year-to-year variability is a strong factor.

24 **Figure 9** shows long-term records of sulfur dioxide, sulfate and nitric acid vapor concentrations
25 in surface air derived at the AIRMoN-dry site near State College, Pennsylvania; other sites in the
26 Northeast show similar results. The Pennsylvania record is used here as an example because it
27 appears to be representative of the region as a whole (for sulfur species but not necessarily for
28 nitrogen, as will be discussed later) and also because there are nearby long-term wet deposition
29 records suitable for comparison against dry deposition data. The strong seasonality of both the
30 SO₂ and sulfate concentrations is clearly evident, though there are significant phase differences in
31 the timing of the annual peaks. SO₂ concentrations peak during the winter, while sulfate peaks
32 regularly during summer.

33 The time series of **Figure 9** shows that 1995 does indicate a downward trend in SO₂
34 concentrations. However, with the year-to-year variation (due to meteorology and other factors) in
35 the annual behavior, a clear statement about how much the emission reductions of 1995 have
36 influenced air concentrations is not easily derived. More data are required before the effects of the
37 year-to-year variability can be taken into account. For the present, the 1989–1995 reduction in
38 airborne concentrations of total sulfur (SO₂ plus particulate sulfate) derived from the data in
39 **Figure 9** for the State College site amounts to 23%. An independent examination of CASTNet
40 data (Sickles et al., 1997) results in much the same conclusion.

41 A third network, the Interagency Monitoring of Protected Visual Environments (IMPROVE)
42 network, also offers useful data. The three networks (CASTNet, AIRMoN-dry, and IMPROVE)
43 operate with different sampling systems and protocols, reflecting their different purposes and
44 design criteria. An analysis of 1984–1994 IMPROVE data (Malm et al., 1994; Cahill et al., 1996)
45 indicated an upward trend in sulfate concentrations in summer at the IMPROVE site in the
46 Shenandoah National Park (**Figure 10**). The data from the Great Smoky Mountains site do not
47 reveal an increase as great as that from Shenandoah, and there is no trend evident in the winter
48 data.

For several years in the 1980s, an AIRMoN site was also operated at Shenandoah National Park, yielding an opportunity to compare the performance of the two independent sampling systems and to test the statistical variability introduced by the fact that IMPROVE samples represent two days per week, whereas AIRMoN-dry (and also CASTNet) protocols call for continuous sampling. Although preliminary results reveal little bias between the two sampling systems, they indicate that the variability of the IMPROVE data is affected by the two-day-per-week protocol.

A spatial picture of the extent of the 1989–1995 reduction in total sulfur concentrations is derived from the CASTNet data set (**Figure 11**). It is clear that the reduction is a truly regional phenomenon, with annual average decreases in concentrations of more than 30% at many locations in the eastern United States. It is tempting to point to the emission reductions of the same period and to claim them as a reason for the lower concentrations. While it is certain that the emission reductions did improve regional air quality, it is not yet clear how closely the two correspond. To illustrate some of the underlying causes of uncertainty, it is of interest to note that cold winters impose a stronger demand for power, which is likely to be reflected in increases in both power plant emissions and regional air concentrations of sulfur. However, it is not yet clear how much considerations like this influence the interpretation of short air-concentration records, such as are currently available for sulfur. It is evident that meteorology -- if not regional climatology -- needs to be considered before final conclusions are reached about how well air concentrations track emission reductions.

Regardless of the year-to-year variability, the CASTNet data indicate that the 1995 reduction in airborne total sulfur (relative to 1989) is truly widespread. For the central Pennsylvania station, for example, the AIRMoN data in **Figure 9** show a 23% reduction in total sulfur from 1989 to 1995. This is almost exactly what **Figure 11** reveals from CASTNet data.

Nitrogen

- For nitric acid vapor in air, regional concentrations show evidence of a similarly widespread reduction as for sulfur species, although of markedly smaller magnitude and with some variation among sites. Control of NO_x emissions under Title IV was not required until 1996.

In general, nitric acid concentrations display a less uniform seasonality than sulfur concentrations, and tend to maximize in the summer when the related chemical reactions are most vigorous. **Figure 9** shows that nitric acid concentrations were slowly increasing at the Pennsylvania site until 1993. However, this upward trend is not representative of other stations in the Northeast, which show slight decreases. Any increase in airborne nitric acid concentrations is of concern because of the corresponding increase in deposition. It is not clear whether the cause is associated with changes in the use of nitrogenous fertilizers in the area, with difficulties in measuring nitric acid vapor, or with some other reason not associated with upwind point-source emissions. As in the case of sulfur, a longer record is needed before it is possible to quantify the changes associated with Title IV of the 1990 Clean Air Act Amendments and its emission reductions.

-----Text Box Begins-----

Monitoring Networks

Ambient air concentration data are collected in both urban and rural settings. The intensive ambient air monitoring networks operated by the EPA and states -- the National Air Monitoring System / State and Local Air Monitoring System (NAMS/SLAMS) networks -- are focused on urban areas where ambient air concentrations mainly reflect local emissions. Ambient air concentrations are also measured in rural areas where they are a necessary part of dry deposition networks.

Wet deposition data are obtained from a 191-site monitoring network initiated in the early 1980s to assess the magnitude of the acid rain problem and to determine spatial and temporal trends in atmospheric chemistry and deposition. The National Atmospheric Deposition Program / National Trends Network (NADP/NTN) is supported by more than 100 organizations including federal, state and local government agencies; universities; and industry. It is the only long-term deposition monitoring program in the United States with national coverage (**Figure 12**). NADP/NTN operates with two levels of sampling intensity: a large primary array of routine weekly sampling, and a smaller sub-network of more intensive research sites with daily sampling. The subnetwork is a part of the NOAA Atmospheric Integrated Research Monitoring Network (AIRMoN-wet).

Dry deposition cannot be monitored directly by any known method, except at great expense and under special circumstances. In existing monitoring operations, dry deposition is not directly measured. Instead, it is inferred using directly measured air concentrations and model estimates of deposition velocity from which the dry deposition rates are evaluated as the product of deposition velocity and air concentrations. There are two air concentration/dry deposition arrays set up by NAPAP and currently operating: AIRMoN-dry and CASTNet. CASTNet has broad coverage, providing information on spatial distributions; it is also used for model development and testing. AIRMoN-dry operates as a research program, developing dry deposition methodologies for use in CASTNet and other applications.

-----Text Box Ends-----

How have deposition rates been affected by the emission reductions?

Wet Deposition

- Wet deposition monitoring results clearly and objectively indicate that Phase I of Title IV has reduced acid deposition in the eastern United States. In 1995, the concentration of sulfate in precipitation in the East was 10–25% lower during 1983–1994.

A critical step in assessing the effectiveness of emission reductions is to evaluate spatial and temporal trends in sulfate, nitrate, and hydrogen ion (acidity) concentrations in precipitation (rain and snow). The effects of Phase I emission reductions on precipitation chemistry was assessed by determining the significance of the deviations of the 1995 bimonthly observations from estimates of 1995 precipitation chemistry based on more than a decade of pre-1995 observations. A seasonal trend model was developed to explain the historical trend of deposition monitoring data from 1983 through 1994 (Lynch et al., 1996). The model was used to extrapolate forward one year to estimate the expected precipitation chemistry in 1995. Allowing for early compliance by some utilities, the 1983–1994 period could be considered prior to Title IV and 1995 could be considered to be under Title IV. In effect, the difference between the 1995 observations and the 1995 projection is the difference in precipitation chemistry between Title IV and no Title IV, respectively. Results indicated that the observed 1995 chemistry data were different from the estimated chemistry at deposition monitoring sites located in the eastern United States.

Analysis of the 1995 wet deposition monitoring data demonstrates that the 1995 reduction in SO₂ emissions in the midwestern and northeastern regions of the United States resulted in a substantial reduction of the acidity (as represented by hydrogen ion concentration) and sulfate concentration of precipitation in those regions. **Figures 13, 14, and 15** display changes in precipitation chemistry over the eastern United States for 1995. The figures illustrate the departures of *observed* sulfate, nitrate, and hydrogen ion concentrations (in percent) from the *estimated* seasonal trends for 1995. These reductions occurred in the Ohio River Valley region where many of the Title IV Phase I units are located, as well as to the east of this region – across the Mid-Atlantic region and north through Maine. The spatial pattern of hydrogen ion decreases in the East correlated well with

the pattern of sulfate decreases, although the magnitude of the hydrogen ion concentration and percentage decreases was even larger than the sulfate decreases.

Sulfate and hydrogen ion concentrations in 1995 were 89% and 79% of the estimated concentrations for northeastern and southeastern sites, respectively. Overall, measured sulfate concentrations in the eastern states were about 10% less than estimates and as much as 25% less in areas in the Upper Northeast, Mid-Atlantic, and Ohio Valley. In contrast, at sites west of the Mississippi River, 1995 sulfate concentrations averaged about 7.5% higher than the estimates. Due to the fact that these sites are upwind of virtually all of the Phase I units and the average increase is within the range of natural variability, the increase is not indicative of any changes in emissions.

Unlike sulfate and hydrogen ion concentrations, nitrate concentrations in 1995 were greater than estimated concentrations at most of the sites in the eastern and western regions of the country. This is not unexpected, since implementation of Title IV NO_x reductions only began January 1996. Approximately 61% of the eastern sites and 57% of the western sites recorded higher nitrate concentrations in 1995 than estimated. Nitrate concentrations in 1995 were about 5% greater than the estimates in the eastern part of the country (**Figure 13**), and about 4% greater in the western part. Unlike sulfate concentrations, there was no evidence of sharp decreases in nitrate concentrations recorded in the Northeast. As expected, these results suggest that overall NO_x emissions in 1995 were not substantially affected by Phase I of Title IV.

Departures from the long-term average precipitation amount (1983–1994) cannot by themselves explain the observed decreases in sulfate and hydrogen ion concentrations in precipitation chemistry in 1995. Changes in precipitation volume do not selectively reduce only sulfate and hydrogen ion concentrations. Other ions would be similarly affected and data show that these ions were not affected. Also, smaller precipitation volumes are typically associated with higher concentrations, and most of the eastern states had below average precipitation in 1995. In fact, in 1995, the smaller precipitation volumes resulted in greater concentrations for virtually all ions except sulfate and hydrogen. The notable exceptions are the lower Mississippi River basin including east Texas, and an area from Illinois to lower Michigan. In these areas, the increases in both sulfate and nitrate ion concentrations are primarily the result of below average precipitation volumes. The conclusion is that sulfate and hydrogen ion concentrations decreased independently of precipitation volume in the Ohio River Valley and the states located immediately downwind of this region in the northern part of the Mid-Atlantic, indicating a reduction in acid precipitation in 1995.

Dry Deposition

- In 1995, dry deposition rates of sulfur at State College, Pennsylvania, decreased to their lowest level yet recorded. Deposition rates of nitrogen compounds remained at their historic level.

Current capabilities for understanding the processes controlling dry deposition are still exploratory. Clear answers cannot be given for the general area-wide deposition rates that affect sensitive ecosystems. With this limitation, scientists have focused on some specific areas, where the chances of success are greatest and where relevant information is most desired.

Work conducted in the United States (Meyers et al., 1997) on the deposition of SO₂ also indicates that the historic evaluations of dry deposition of gaseous sulfur may underestimate actual rates, perhaps by as much as 15–20%, depending on the site. The reason is complex, but involves the development of a better understanding of the processes controlling dry deposition and their

representation in analysis routines. Since any underestimates are additive in all years, they should not affect the trends in deposition.

Total Deposition: A Relevant Example

Data gathered at the research area near State College, Pennsylvania, has been identified as a good example of the sort of understanding that could be made available for the region as a whole, provided the scientific research is appropriately targeted (**Figure 16**). Total deposition of sulfur is the sum of measured wet deposition of sulfate and estimated (through calculations) dry deposition of sulfur dioxide and particulate sulfate. The data indicate annual dry deposition rates of sulfur are typically about the same as for wet deposition, with dry exceeding wet for drought years. Average total deposition rates of sulfur have been about 13–16 kg/ha/yr (an average of the last ten years), with 1995 being somewhat lower (about 11 kg/ha/yr). It is not clear how much of this drop in 1995 is properly attributable to emission reductions. All estimates of total deposition are site specific. They are likely to be underestimates because of underestimation of the deposition velocity for particulate sulfate (used to calculate dry deposition), but it should not seriously affect temporal trends.

For deposition of nitrogen compounds, the data indicate a trend in the ratio of dry to wet deposition, from about 0.2 in the mid-1980s to about 1.0 in 1995. The reason is not clear, but several potential explanations are currently being explored—all related to the fact that dry deposition of nitric acid vapor is a more local phenomenon than wet deposition of nitrogen as nitrate and ammonia. First, local farmers could have increased their use of nitrogenous fertilizer, which would affect regional air quality. Second, regional decreases in NO_x emissions could decrease the amount of wet deposition. As expected, total nitrogen deposition data do not indicate a sharp decline in 1995. The values appear to be maintaining a total deposition rate of about 8 kg/ha/yr of nitrogen at State College. Again, this is likely to be an underestimate due to current omission of organic nitrogen contributions and of dry nitrogen species other than nitric acid vapor. It is also possible, however, that some of these causes for underestimation may be offset by the likely presence of ammonium nitrate particles in the air. Ammonium nitrate is not measured well by the apparatus used in the AIRMoN-dry program. A possible explanation for the time trends evident at State College is that local use of nitrogenous fertilizers has increased, in which case ammonium nitrate is quite likely to have been present. If so, then the methodologies reported here would have overestimated nitrate deposition. Clarifying this situation is not within existing capabilities. More studies are needed if the related questions are to be answered.

What are current deposition rates and what is their variability?

- Precipitation data show that widespread declines in sulfate concentrations and acidity (hydrogen ion) accompanied by decreases in such cations as calcium, magnesium, potassium, and sodium have occurred since about 1980. The most significant declines in sulfates and hydrogen ion occurred in 1995. Nitrate and ammonium concentrations showed considerable variability.
- The acidity (as measured by hydrogen ion) of precipitation may not have declined as much during the period 1983–1994 due to the parallel decline in base cations, which act to buffer the acidity, in precipitation (hydrogen ion had declined; pH had increased) and also to an increase in nitrate. However, 1995 wet deposition data show a substantial drop in acidity (hydrogen ion) and in increase in pH commensurate with the sulfur concentration reductions.
- Ammonium, which along with nitrate is a significant contributor to nitrogen enrichment to ecosystems via atmospheric deposition, generally increased throughout the United States.

1 Based on NADP/NTN data, nitrate concentrations in precipitation did not exhibit a consistent
2 spatial or temporal pattern between 1983 and 1994, although the number of sites measuring
3 increases in nitrate exceeded the number where decreases occurred (Lynch et al., 1996). The
4 largest increases in nitrate concentration occurred in the western United States. Ammonium
5 concentrations increased at 80% of the deposition monitoring sites over the period 1983–1994, for
6 a total increase of 28%. Hydrogen ion concentrations (acidity) declined during 1983–1994,
7 although a lack of consistency between the magnitude of sulfate reductions and reduction in free
8 acidity suggests that the decline in acid-neutralizing cations may have offset some of the expected
9 decline in the acidity of precipitation.

10
11 Widespread declines in sulfate concentrations accompanied by decreases in base cations have been
12 measured since the late 1970s. The largest declines occurred from 1979 to 1982, with continued
13 declines throughout the 1980s (Sisterson et al., 1991). Consistent with these findings, Lynch et
14 al. (1995b) found concentrations decreased more rapidly during the early 1980s and less rapidly
15 thereafter until 1995, when significant reductions were measured. Nitrate and ammonium
16 concentrations showed considerable variability. The large decreases in the early 1980s were
17 similar to changes in SO₂ emissions, which also decreased rapidly between 1980 and 1983, then
18 remained nearly constant in many states during the remainder of the decade.

19
20 **Figure 17** shows the 1995 weighted mean pH of precipitation, while **Figures 18-20** present
21 the 1995 wet deposition rates for sulfate, nitrate, and total nitrogen (nitrate plus ammonia,
22 expressed as nitrogen).

23 A major factor influencing the statistical depiction of atmospheric deposition is the areal variability
24 of the signal. Wet deposition is largely a “spatially ergodic” process – that is, a sample obtained at
25 a selected location is likely to be representative of the area as a whole. Dry deposition is
26 substantially different, because it is driven by surface properties that are site specific. Deriving a
27 spatial average rate of dry deposition from single-location information presents a challenge that is
28 currently daunting. The capability to measure dry deposition rates across selected areas (using
29 highly advanced instrumentation on aircraft) has been developed only very recently. In the
30 meantime, the best contemporary numerical models use grid cells that contain many land-use types.
31 Comparing model outputs for a selected grid cell against dry deposition data obtained for a
32 particular land-use type within the same grid cell remains a difficult problem (Hicks et al., 1991).

BENEFITS OF EMISSION REDUCTION

As the amount of sulfur and nitrogen oxides emitted into the atmosphere is reduced, concentrations of pollutants in the air and deposition of acidic compounds to the earth's surface should be reduced. Reducing the amount of acid deposition is also expected to decrease the damage to ecosystems, materials, visibility, and human health. The emission reductions under Title IV only began in January 1995, and responses to those reductions vary (**Figure 21**). Concentrations of pollutants in the air respond to these reductions within hours, and these compounds are deposited to the earth's surface usually within days to weeks (e.g., during a rain event). Thus, receptors affected by air concentrations, such as human health and visibility, will respond quickly (from hours to days).

The effects of deposition on sensitive receptors can range from days to centuries. Ecosystems are complex systems that are simultaneously responding to a variety of inputs, such as climate, land-use patterns, and other pollutants besides sulfur and nitrogen oxides. These multiple stressors can result in chemical changes within the ecosystem, which can exhibit long lag times before manifesting a response. Therefore, in many effects areas, responses to the Title IV emission reductions will not be expected for many years. Monitoring the changes in the effects areas over time will be necessary to determine whether the expected benefits are realized.

Have changes in effects been observed (1980–1995)?

NAPAP has focused on five effects areas for this assessment: aquatic ecosystems, forest ecosystems, materials and cultural resources, visibility, and human health. The following issues are addressed in each effects area:

- effects of acid deposition on sensitive receptors over the period 1980-1995,
- significant developments since the NAPAP *1990 Integrated Assessment Report*, and
- dose-response relationships.

Highlights of the policy-relevant developments since 1990 are presented in **Table 5**.

TABLE 5. SELECTED POLICY-RELEVANT DEVELOPMENTS SINCE 1990

ISSUE	WHAT'S NEW (since 1990)	SIGNIFICANCE
Sulfur concentration in precipitation	Sulfur concentration levels in wet deposition decreased 10-25% over large areas of the eastern United States in 1995, with the largest decreases in and downwind of the Ohio River Valley. Similar decreases were found in sulfur concentrations in dry deposition.	The significant drop in acid deposition (and emissions) in 1995 provides a unique scientific (and economic) opportunity to validate atmospheric deposition models and dose-response relationships.
Aquatic effects from nitrogen	Nitrogen is now recognized as playing a greater role in watershed acidification.	There may be limited recovery in some sensitive systems because NO _x reductions under Title IV may not be sufficient enough to result in measurable improvement in chemical or biological changes.
Surface water trends	Concentrations of sulfate in lake and stream water have decreased in many areas, with evidence of recovery from acidification in New England. However, the majority of Adirondack lakes have remained fairly constant, while the sensitive Adirondack lakes continue to acidify.	Reductions in sulfur deposition to date may be insufficient in some areas to improve the acid-base status of acidified surface waters.
High elevation spruce-fir dieback	Strong linkage was confirmed between acid deposition and damage to forest ecosystems.	There is a higher likelihood of some short-term benefits from controls of acid deposition precursors.
Cultural resources	Dry deposition of SO ₂ and aerosols is now thought to be more damaging to stone than wet deposition.	The role of dry deposition and its relative impact on benefits to cultural resources must be reviewed.
Fine particle impacts on health	New studies associate PM _{2.5} exposure with premature mortality and other health impacts.	Title IV controls on emissions of precursors of fine sulfate and nitrate particles will reduce ambient PM _{2.5} concentrations.
Valuation of economic benefits	Expected benefits are greater than previously thought, especially in the areas of human health and visibility.	The magnitude of valued benefits in these areas exceed the costs of Title IV, independent of benefit estimates for other areas.

Aquatic Ecosystems

Observed Effects

Although it is too early to detect specific changes in aquatic systems from emission reductions under Title IV, significant progress has been made since the 1990 *Integrated Assessment Report* (NAPAP, 1991) in refining understanding of acidification processes and quantifying dose-response relationships. This information improves model forecasts of anticipated change in aquatic systems due to reduced emissions. Particular areas include naturally occurring organic acidity, the depletion of base cation reserves from soils, nitrogen dynamics in forested and alpine ecosystems, interactions between acid deposition and land use, and the role of aluminum in fisheries response. These advancements and observed trends in aquatic effects are discussed below.

Sulfate and Nitrate Concentrations

- Lakes and streams throughout many areas of the United States have experienced substantial decreases in sulfate and base cation concentrations in response to reduced levels of atmospheric sulfur deposition. However, acid neutralizing capacity and pH, which can affect aquatic biota, show little consistent change, particularly in the most sensitive ecosystems.

Monitoring data have been available since the early 1980s for many lakes and streams in acid-sensitive areas of the United States. The concentration of sulfate in surface waters has decreased in many areas over the past 10–20 years. This decrease has been caused by reductions in emissions and subsequent decreases in atmospheric deposition of sulfur on a regional basis throughout many parts of the United States during that time period. To some extent, these changes may be partly related to implementation of Title IV, but changes were also already occurring without Title IV, most likely as a result of pollution controls including those in response to the 1970 Clean Air Act and 1977 Amendments. In parts of the northeastern United States, approximately 15% reductions in sulfate concentrations of lakes and streams have been measured in recent years, and additional decreases undoubtedly occurred prior to implementation of long-term monitoring programs. Sulfate concentrations are expected to continue to decline in the Northeast and many other areas. Exactly how that will affect surface water acidity and biological recovery is uncertain and will require continued monitoring. On the other hand, lakes in New England do appear to show statistically significant recoveries in acid neutralizing capacity as a result of sulfate reductions (Stoddard et al., in press). This is based on analyses of long-term monitoring data and is discussed in more detail under Sensitive Receptors. Future data analyses should focus on the years following Title IV implementation to provide a period of record long enough to determine trends.

Concurrent changes in the concentrations of other chemical parameters have been generally less clear and less consistent than for sulfate and base cations. These other parameters are more strongly influenced by factors other than atmospheric deposition. The most consistent response of surface water chemistry to the recent observed decrease in sulfate concentration in lakes and streams has been a decrease of approximately the same magnitude in the concentration of calcium and other base cations (see text box). In most cases, pH, aluminum, and acid neutralizing capacity have not responded in a large or systematic fashion to changes in sulfate concentrations. Examples of monitoring data from two sites are shown in Figures 22 and 23. Seasonal and annual variability tend to be large in relation to the overall trend.

Seasonal variability in the concentration of key chemical parameters often varies by more than the amount of acidification that might occur in response to acid deposition. For example, a decrease in sulfate concentration in a lake or stream is not directly attributable to a decrease in acid deposition

because the decrease could be within the range of normal seasonal variability of sulfate concentration in that lake or stream. Such variability makes it difficult to quantify acidification and recovery responses, and also complicates attempts to evaluate sensitivity to acidification based solely on "index" chemistry (i.e., samples taken during broad-scale synoptic surveys at a time when water chemistry is expected to be relatively stable). Seasonal variability is particularly problematic in the assessment of deposition standards for nitrogen. Continued monthly monitoring of different types of lakes and streams located in sensitive regions will provide much needed data on seasonal variability. This will be especially important for evaluating the possible need to reduce nitrogen deposition in some regions during the winter/spring period prior to snowmelt, in view of the apparent seasonal nature of nitrogen impacts on water quality.

The observed changes in the concentration of nitrate in some surface waters have most likely been due to a variety of factors, including nitrogen deposition and climate. In many areas of the northeastern United States, nitrate concentrations in lakes and streams increased during the latter part of the 1980s. This trend was reversed in about 1990, however, despite relatively constant levels of nitrogen deposition during the past 10 years (Driscoll et al., 1995). This is because the amount of nitrate that leaches through soils to drainage waters is the result of a complex set of biological and hydrological processes that include: nitrogen uptake by plants and soil microbial communities, microbial transformations between different forms of inorganic and organic nitrogen, rates of organic matter decomposition, the amount of rain and snow received, and the amount (and form) of nitrogen that enters the ecosystem as atmospheric deposition. Most of these important processes are strongly influenced by climatic factors such as temperature, moisture, and snowpack development. The end result is that nitrate concentrations in surface waters, although clearly influenced by atmospheric nitrogen deposition, respond to many factors and can be difficult to predict (Mitchell et al., 1996).

-----Text Box Begins-----

Effects on Water Chemistry

If there is a change in the concentration of an anion contributed by acid deposition (e.g., sulfate or nitrate), then other anion and/or cation concentrations must also change such that the total concentration of anions always equals the total concentration of cations in the surface water. This is called the principle of electroneutrality, whereby positive and negative ions are found in equal amounts. Thus, if acid deposition causes sulfate and nitrate concentrations to increase, then some or all of the following changes will also occur:

- bicarbonate anion decreases, which causes a reduction in acid neutralizing capacity;
- base cations (calcium, magnesium, sodium, potassium) increase, which prevent or minimize acidification of drainage waters, but may deplete soil reserves and affect forest growth;
- hydrogen cation increases (decreases pH), which can adversely affect aquatic biota; and
- aluminum cation increases, which can negatively affect aquatic biota.

If a reduction in acid deposition causes sulfate and nitrate concentrations to decrease, then changes opposite to those enumerated above will also occur. An increase in the concentration of base cations in drainage water as a consequence of acid deposition has both positive and negative connotations. Removal of base cations from soils to balance sulfate or nitrate from acid deposition minimizes the extent of surface water acidification. Over time, however, base cation reserves in the soils can become depleted if they are lost from the soils faster than they are supplied from atmospheric inputs and weathering. This can delay acidification recovery.

-----Text Box Ends-----

Episodic Acidification

- Nitrogen is quantitatively as — or, in some cases, more — important than sulfur as a cause of both chronic and especially episodic acidification than was understood in 1990.
- Biological effects on fish during acidic episodes are largely attributable to increased concentrations of dissolved aluminum in stream water. This aluminum is transported from soils to streams by short-term increases in acidity, which are largely associated with nitrogen — specifically, nitrate leaching.

Episodic acidification is the process by which lakes and streams experience short-term (hours to weeks) decreases in acid neutralizing capacity and pH. Generally, these episodes occur during periods of high water flow, such as storms or snowmelt, and often occurs at the most biologically sensitive time of the year — spring spawning. Typically, several chemical changes accompany the decrease in acid neutralizing capacity that occurs during episodes. In regions with the most severe episodes, pH and calcium often decrease and dissolved aluminum increases, creating more adverse conditions for aquatic life (Wigington et al., 1991). A review of episodic acidification of surface waters in portions of the eastern United States (Wigington et al., 1993) clearly demonstrated that (1) episodic processes are a natural facet of watershed processes, (2) nitrate attributable to atmospheric deposition plays an important role in the episodic acidification of some surface waters, and (3) increased aluminum concentration has the greatest impact on biota. EPA's Episodic Response Project documented the occurrence of acidic episodes with a minimum pH of less than 5 and high aluminum concentration in the northern Appalachian region of Pennsylvania and the Catskill and Adirondack Mountains of New York (Wigington et al., 1996). A regional assessment of the acid-base status of native brook trout streams in the mountains of Virginia included examination of the relationship between the median acid neutralizing capacity observed during spring and the minimum acid neutralizing capacity observed during episodic acidification. The results indicated that the extreme episodic acid neutralizing capacity values were about 20% lower than the median values (Webb et al., 1993).

Episodic acidification has been found to be nearly ubiquitous in drainage waters. Nearly all lakes and streams that have been studied throughout the United States, Canada, and Europe experience loss of acid neutralizing capacity during hydrologic events. Chemical changes during episodes are controlled by a number of natural processes, including dilution of base cation concentrations, nitrification, flushing of organic acids from terrestrial to aquatic systems, and the neutral salt effect. Acid deposition contributes to episodic acidification, particularly via enhanced nitrate leaching. Under some conditions, episodes can be partly caused by increased sulfate concentrations. There is also the likelihood that chronic acidification by acid deposition can precondition a watershed, thereby increasing the severity of episodic acidification.

Nitrate is the principal acid anion in snowmelt in many areas of the United States. Nitrate in snowmelt runoff has been recognized for some time as an important component of biological damage resulting from atmospheric deposition. A number of factors can be involved in controlling the loss of nitrogen from forested and alpine watersheds, including atmospheric inputs, forest stand age and condition, land use, soil nitrogen and carbon pools, and flow paths of percolation and melt-water within the watershed. Except in cases of excessive nitrogen saturation, the effects of nitrogen deposition on surface waters are expected to be primarily episodic. However, data required to make regional assessments of episodic effects are generally not available.

Organic Acidity

- Naturally occurring organic acids are more important components of surface water acid-base chemistry (pH) and have more significant influences on the sensitivity of model

projections of acidification than was widely believed in 1990. Predictions of the pH of sensitive lakes in the Adirondacks have been revised accordingly.

Organic acids commonly exert a major influence on surface water acid-base chemistry, particularly in dilute waters having moderate-to-high concentrations of dissolved organic carbon. Some lakes and streams are naturally acidic as a consequence of organic acids in solution. The presence of organic acids also provides buffering to minimize pH changes in response to changes in the amount of mineral acid anions (e.g., sulfate and nitrate) in solution. Organic acids will most likely become more important components of acid-base chemistry as acid deposition continues to decline.

Recently revised model simulations that included organic acids (Driscoll et al., 1994; Sullivan et al., 1996) showed improvements in the agreement between modeled and historical acidity inferred from diatoms of up to one full pH unit for individual lakes. The revised MAGIC model predicts less pH recovery of Adirondack lakes in response to decreased sulfur deposition than the earlier version of the model used as the basis for the *1990 Integrated Assessment Report* (NAPAP, 1991). Other surface water chemistry indicators, such as acid neutralizing capacity, are not affected to the same extent by organic acids. See Appendix A for further discussion of MAGIC.

Base Cation Depletion

- Sensitive watersheds in the Adirondack Mountains have responded to atmospheric sulfur inputs primarily by enhanced base cation leaching, especially calcium and magnesium. This means that drainage waters have been *less* acidified than was earlier thought. Such a response has also raised concerns about long-term depletion of base cation reserves from forest soils, which could decrease future forest productivity and the rate of watershed recovery in response to lower levels of sulfur deposition.

Base cation depletion has been recognized as an important effect of acid deposition on soils for many years. However, scientific appreciation of the importance of this response has increased with the realization that many watersheds previously acidified by acid deposition are not exhibiting acid neutralizing capacity and pH recovery in response to recent decreases in sulfur deposition. As sulfate concentrations in lakes and streams have declined, so too have the concentrations of calcium and other base cations for several reasons. First, decreased movement of sulfate through watershed soils reduces leaching of base cations from soil surfaces. Second, the atmospheric deposition of base cations has decreased in recent decades (Hedin et al., 1994), probably due to a combination of air pollution controls that targeted large particles, changes in agricultural practices, and the paving of roads. The latter two of these factors generate dust that is rich in base cations (NAPAP, 1991 [SOS/T 6]). Most of these reductions of dust occurred in the 1960s and 1970s; dust has remained fairly stable for the past 20 years. Third, soils in some sensitive areas have experienced prolonged base cation leaching to such an extent that their base cation reserves have been depleted. Such depletion prolongs the acidification recovery time of watersheds and may adversely impact forest productivity (Kirchner and Lydersen, 1995; Likens et al., 1995) (See the Forest Ecosystems section for further discussion.)

Nitrogen Cycling

- Significant progress has been made in understanding nitrogen cycling in forested and alpine ecosystems, and the effects of nitrogen deposition on nitrate leaching and consequent acidification of surface waters. The ability to model nitrogen dynamics effectively on a regional basis is on the immediate horizon.

Increased atmospheric deposition of nitrogen does not necessarily cause adverse environmental impacts. In some cases, nitrogen acts as a fertilizer and is taken up by terrestrial biota increasing forest productivity (Kauppi et al., 1992). However, in some areas, especially at high elevations,

terrestrial ecosystems have become nitrogen saturated, and high levels of deposition have elevated the levels of nitrate in drainage waters (Aber et al., 1989; Stoddard, 1994). This enhanced leaching of nitrate depletes calcium and other base cations from forest soils and can cause acidification of drainage waters in base-poor soils. Nitrogen saturation means that the input of nitrogen (e.g., nitrate, ammonium) to the ecosystem exceeds the requirements of terrestrial biota (that is, more nitrogen comes in than can be taken up by plants and microorganisms) and a substantial fraction of the incoming nitrogen leaches out of the ecosystem in groundwater and surface water. Further discussion about nitrogen effects is presented in the section on Forest Ecosystems.

Prior to the mid-1980s, research on the effects of atmospheric deposition focused almost exclusively on sulfur. During the last 10 years, the research focus has shifted more to nitrogen. Although some high-quality research has been conducted in the United States on the environmental effects of atmospheric nitrogen deposition since 1990, such work has been conducted to a far greater extent in Europe.

In Europe, direct ecosystem-level data have been obtained on nitrogen saturation from atmospheric deposition and subsequent ecosystem recovery. Results from experimental manipulations and other studies have shown that nitrogen generally leaches into drainage waters in European forests only where atmospheric deposition of nitrogen is greater than about 10 kilograms per hectare per year (kg/ha/yr). Other important factors controlling nitrogen leaching include the ratio of carbon to nitrogen in the forest floor and soils, the age of the forest, and past forest management practices and disturbances.

Chronically high nitrate concentrations in lake or stream water, indicative of ecosystem nitrogen saturation, have been found in recent years at a variety of locations in the western and eastern United States, including the San Bernardino and San Gabriel Mountains within the Los Angeles air basin (Fenn et al., 1996), the Front Range of Colorado (Baron et al., 1994; Williams et al., 1996), the Allegheny Mountains of West Virginia (Gilliam et al., 1996), the Catskill Mountains of New York (Murdock and Stoddard, 1992; Stoddard, 1994), and the Great Smoky Mountains in Tennessee (Cook et al., 1994). Nitrogen saturation is occurring throughout high-elevation watersheds of the Colorado Front Range (Williams et al., 1996).

Total nitrogen deposition in high-elevation watersheds of the Colorado Front Range is about 4 kg/ha/yr, which is about double that in most other mountainous areas of the West and approaching the deposition levels found in some parts of the East. It is much lower, however, than the 10 kg/ha/yr threshold value for nitrogen leaching found in Europe. The observed high concentrations of nitrate in lakes and streams of the Front Range are likely due to leaching from tundra, exposed bedrock, and talus areas at high elevations. In the alpine areas of Colorado, about 50% of the nitrogen is deposited in the snowpack and released with the spring thaw. This large release of nitrogen moves through the watershed, removing base cations and reducing the acid neutralizing capacity of lakes and streams (Williams et al., 1996).

A comparison was recently made for 37 watersheds in the United States and Canada to examine the relationship between atmospheric nitrogen deposition inputs and nitrogen outputs in drainage water. Over one-third of the watersheds that receive nitrogen deposition higher than about 6 kg/ha/yr are leaching appreciable amounts of nitrogen (more than about 2 kg/ha/yr; Peterjohn et al., 1996). The highest inputs and outputs of nitrogen were found for the Fernow Experimental Forest in West Virginia.

Results from an experimental manipulation site in Maine illustrate the rapidity with which some forested watersheds in the northeastern United States may move toward nitrogen saturation in response to increased nitrogen loadings. The forest ecosystem continued to accumulate nitrogen, but the rate of accumulation decreased steadily with the experimental treatment (Kahl et al., 1993).

Land Use

- Human land use activities, especially disturbance of soils and forest management, have important effects on the responses of terrestrial and aquatic ecosystems to the atmospheric inputs of sulfur and, especially, nitrogen.

In the northeastern United States, land use practices and resulting vegetation patterns have changed from forests to agriculture and back to forests more or less continuously for about the past 250 years. These changes in human activity, and consequent changes in forest structure and dynamics, can influence the response of forested ecosystems to external stressors, such as atmospheric deposition of sulfur or nitrogen, exposure to ozone, natural disturbance factors (e.g., wind and fire), and climatic changes. Some landscape processes contribute to the acidification of soil and surface waters or reduce the base cation saturation of the soils, thereby increasing their sensitivity to acid deposition. Acid deposition is only one of many factors that affects acidity status although in acid-sensitive regions it remains one of the primary factors influencing the health of ecosystems.

Disturbances such as logging, blowdown, and fire affect surface water pH and acid neutralizing capacity. When they occur within a watershed these disturbances disrupt the normal flow of water. In some cases, they cause increased contact between runoff water and soil-surfaces, and often lead to increased base cation concentration and acid neutralizing capacity in surface waters. In most cases, recovery from disturbance will lead to a decrease in pH and acid neutralizing capacity, as the system returns to predisturbance conditions. Such acidification can be erroneously attributed to acid deposition.

The removal or cutting of forests has immediate effects on the chemistry of drainage water. Dry deposition of sulfur and nitrogen to the site is reduced. Leaching of nitrate increases and, in some cases, causes a pulse of surface water acidification, and base cations are lost from the soils.

The subsequent regrowth of the forest following deforestation also affects drainage water chemistry. Young, fast-growing forests are considerably more acidifying to the soil than older forests because young trees take up more of the base cations from the soil, leaving less to neutralize any acidic inputs. In contrast, old-growth forests need less nitrogen to sustain growth and, therefore, may be more susceptible to nitrate leaching into nearby surface waters.

Lake water acidification in the Adirondacks has been correlated with widespread changes in landscape cover that occurred in response to massive logging operations and fires around the turn of the 20th century and unprecedented forest blowdown during a large wind storm in 1950 (Sullivan et al., 1996). Although such changes — particularly forest regrowth — are unlikely on their own to cause lakes to become acidic, they can cause decreases in the base saturation of soils, thereby predisposing sensitive watersheds to subsequent acidification from acid deposition. The importance of these landscape changes has not been widely recognized.

Modeling studies and calculations performed for selected watersheds in Europe have suggested that acid deposition and landscape processes are of approximately equal importance as regulators of surface water acid-base chemistry within the watersheds investigated. In the United States, however, the importance of landscape processes in influencing surface water acid-base chemistry and the response of surface waters to acid deposition have not been well studied. Early model predictions (NAPAP, 1991) of the response of acid-sensitive watersheds in the United States implicitly assumed that changes in landscape processes either do not occur or are not important in determining surface water chemical response to changes in atmospheric deposition of sulfur and nitrogen. This assumption may have biased model projections of acidification and/or recovery of surface waters in response to changing levels of sulfur deposition. It may be responsible — at least in part — for the observed tendency for MAGIC model hindcasts of historical acidification to

1 exceed diatom-inferences of acidification of some Adirondack Mountain lakes. The need to include
2 land use in modeling efforts will most likely increase as efforts shift more to model-based
3 assessments of nitrogen effects. This is because nitrate leaching from forested watersheds is
4 largely controlled by age-dependent forest nitrogen uptake processes as well as atmospheric
5 deposition of nitrogen.

6 Future studies to address the quantitative response of surface water pH and acid neutralizing
7 capacity to changing levels of sulfur or nitrogen inputs should include the influence of landscape
8 changes on key watershed processes. This will require the collection of detailed, watershed-
9 specific information on past, current, and future forest management and the occurrence of major
10 landscape disturbances. In addition, research is needed on watersheds that contain multiple land
11 uses — e.g., forests, agriculture, and urban areas. Consideration of the cumulative impact of all
12 major processes that influence the acid-base chemistry of drainage waters will be required to
13 improve model projections of future dose-response relationships.

14 *Biological Response to Deacidification*

- 15 • Biological recovery of previously acidified lakes can be a slower process than chemical
16 recovery.

17 Studies in both the United States and Canada have provided new understanding of the feasibility
18 and complexity of biological recovery in response to chronic deacidification. The rate of biological
19 recovery can be slower because of several factors: (1) other environmental stresses, such as metal
20 contamination (e.g., as in Sudbury, Ontario) (Gunn, 1995; Havas et al., 1995; Jackson and
21 Harvey, 1995; McNicol et al., 1995a; Yan et al., 1996a, and 1996b); (2) barriers imposed by
22 water drainage patterns between lakes, hindering recolonization by some fish species (Jackson and
23 Harvey, 1995); (3) the influence of predation by fish species that are not identified as acid-sensitive
24 on the recovery of zooplankton and macroinvertebrate communities (McNicol et al., 1995a); and
25 (4) predation on tributary-spawned young trout when they move downstream into lakes inhabited
26 by predatory fish and birds (Schofield and Keleher, 1996).

27 Results from a liming study in the Adirondacks indicated that reestablishment of tributary-
28 spawning populations of brook trout may be possible with future reductions in acid deposition.
29 However, restoration of tributary-spawning habitat may not be sufficient to produce self-sustaining
30 populations because of high rates of predation on young trout after they move downstream into
31 lakes (Schofield and Keleher, 1996). Woods Lake continues to be examined as part of the
32 Adirondack Long Term Monitoring Project to evaluate the effectiveness of liming to provide a self-
33 sustaining brook trout population.

34 Sensitive Receptors

35 Several types of aquatic systems and/or regions are particularly sensitive to acid deposition. The
36 focus of this report is on four sensitive receptors: Adirondack and New England lakes, high-
37 elevation western lakes, aquatic biota, and estuaries and near-coastal waters.

- 38 • Overall, lakes and streams in areas of the United States that have experienced significant
39 recent decreases in sulfur deposition have shown decreases in sulfate and base cation
40 concentrations of approximately similar magnitudes. Other ions have been less affected by
41 the changes in deposition.

42 *Adirondack Mountains and New England*

43 In the Adirondack Mountains, where acid deposition has been considerable recently, the acid
44 neutralizing capacity of the majority of lakes has remained fairly constant, but has continued to
45

decline in the sensitive lakes (i.e., acidification has increased) despite relatively large decreases in sulfate concentrations in lake water. As a result, the anticipated recovery has not been realized.

In contrast, lakes in New England, especially those considered highly sensitive to acid deposition, have shown statistically significant recoveries in acid neutralizing capacity, based on analyses of long-term monitoring data (Stoddard et al., in press). Although the precise reasons for these differences are not known, it is believed that the extent to which base cation reserves in the soil have been depleted by acid deposition is an important factor. Monitoring of acid-sensitive waters should continue in order to identify the processes of recovery and to improve the models used to predict recovery times.

High-Elevation Western Lakes

- Episodic acidification associated with nitrogen deposition may be occurring to a significant degree in the watersheds of many high-elevation western lakes. These systems may be predisposed to episodic events.

Based on existing data, many high-elevation lakes in the West are currently experiencing nitrogen deposition sufficiently high to cause chronic nitrate leaching (i.e., nitrate concentrations in lakes and streams remain high, greater than 10µeq/L, throughout the year), and associated chronic and episodic acidification, albeit small in magnitude.

A number of factors predispose western systems to potential episodic effects, including:

1. The abundance of dilute to ultra-dilute lakes, exhibiting very low concentrations of base cations and, therefore, acid neutralizing capacity, throughout the year.
2. Large snowpack accumulations at high elevations, causing substantial episodic acidification via the natural process of base cation dilution.
3. Short retention times for many of the high-elevation drainage lakes, enabling snowmelt to rapidly flush lake basins with highly dilute melt water.

Lakes and streams in the Sierra Nevada, Cascade, and Rocky Mountains are highly sensitive to potential effects of acid deposition because of the predominance of granitic bedrock, thin acidic soils, large amounts of precipitation, coniferous vegetation, and extremely dilute waters. Although, there are no data to suggest that lakes in these areas have experienced chronic acidification to any significant degree, episodic effects could be occurring under current deposition regimes, and nitrate concentrations could be causing a small loss of acid neutralizing capacity on a chronic basis at some high-elevation sites. Unfortunately, the data that would be needed for such determinations have not been collected to a sufficient degree in acid-sensitive areas of the West to permit any regional assessment of either episodic or chronic nitrogen-driven acidification.

Aquatic Biota

- Acid deposition can cause long-term adverse effects on fish populations and other aquatic organisms in both lakes and streams. The migration of fish downstream to less acidic water may reduce some effects, but not long-term impacts on species composition.

New information is available on adverse effects on fish populations and communities in chronically acidified streams in Shenandoah National Park (Virginia). Fish species richness, population density, condition factor, age distribution, size, and bioassay survival were all reduced in streams with low acid neutralizing capacity, as compared to those with intermediate and high acid neutralizing capacity (Bulger et al., 1995; Dennis et al., 1995; MacAvoy and Bulger, 1995).

A study of 13 streams in the Adirondack and Catskill Mountains in New York and the Northern Appalachian Plateau in Pennsylvania showed long-term adverse episodic effects on fish populations. Streams with suitable chemistry during low flow, but low pH and high aluminum levels during high flow, had substantially lower numbers and biomass (weight) of brook trout than were found in nonacidic streams. Streams having acidic episodes showed significant fish mortality. Some brook trout avoided exposure to stressful chemical conditions during episodes by moving downstream or into areas with higher pH and lower aluminum. This migration of brook trout only partly mitigated the adverse effects of episodic acidification, however, and was not sufficient to sustain fish biomass or species composition at levels that would be expected in the absence of acidic episodes.

These findings indicate that stream assessments based solely on chemical measurements during low-flow conditions will not accurately predict the status of fish populations and communities in small mountain streams (Baker et al., 1990; Baker et al., 1996; Wigington et al., 1996). A study of coastal plain streams indicated that larval mortality of river herring due to episodic acidification may be substantial during wet years, which exhibit more frequent and more severe episodes (Gerritsen et al., 1996). Based on lake chemistry from the Adirondack Long-Term Monitoring Project, episodic acidification may also be biologically relevant to certain kinds of lakes, depending on the magnitude and duration of the spring snowmelt period. In 1993, 1994, and 1996, approximately 50% of the highly and moderately sensitive lake classes in this study became acidified during the snowmelt period. Additional research is needed to determine the impacts on the populations of sensitive fish species that inhabit these waters.

Rainbow trout are sensitive to acidification not because of acidity itself, but because of elevated aluminum concentrations due to low pH levels (less than 5.0). Aluminum accumulates on gills and disrupts gill ion transport and respiratory function (Marcus and Mount, 1991).

Some organisms can serve as specific indicators of changes in acid deposition. In aquatic ecosystems, loons are effective "bioindicators," linking recovery from lake acidification to higher trophic levels (fish). If loons nest on acidic lakes, their reproductive success is lower, in part due to the reduced survival of their young. This reduced survival is presumably linked to reduced fish biomass (McNicol et al., 1995b). Thus, loons are less likely to nest on acid lakes.

Estuaries and Near-Coastal Waters

- Changes in atmospheric deposition of nitrogen can have significant impacts on the biology and chemistry of estuaries and near-coastal waters.

Through a combination of monitoring, experimental research, and modeling, scientists better understand how atmospheric deposition of nitrogen -- in the form of nitrate and ammonium -- to both water surfaces and watersheds is affecting the biological and chemical status of estuaries and near-coastal waters. Nitrogen is the limiting nutrient for the growth of algae in many estuaries and near-coastal systems, rather than phosphorus, which typically limits algal growth in freshwater systems.

Defined as one of the United States' "Great Waters," the Chesapeake Bay is the nation's largest estuarine system with a watershed of almost 64,000 square miles encompassing one-sixth of the eastern seaboard. It has an important fish and shellfish industry, and serves as a nursery for marine commercial and sport fish. There has been considerable research and monitoring on the effects of nitrogen and phosphorus loading to the Chesapeake Bay. Changes in atmospheric nitrogen deposition can have significant impacts on aquatic biology. Excess nitrogen entering the Bay produces algal blooms that block sunlight needed for submerged aquatic grasses, and the decomposition of excess algae depletes life-sustaining oxygen needed by invertebrates inhabiting

bottom waters. The best estimates of atmospheric nitrogen loads to the Chesapeake Bay and other estuaries along the Atlantic and Gulf coasts range from 10% to 45% of the total nitrogen inputs from all sources. Additional research is needed to quantify the current effects and the expected benefits from reducing nitrogen deposition on estuary systems.

Dose-Response Relationships

- Chemical effects in surface waters due to changes in atmospheric deposition can exhibit lag times of one to many years. Lags in measurable effects on aquatic biota can be even longer. Continued monitoring of surface water chemistry into the future will be required to assess potential improvements that may occur as a consequence of emissions reductions already realized.

Researchers must be cautious in interpreting the observed surface water chemistry as a direct response to estimated changes in sulfur and/or nitrogen deposition. Some effects of changing deposition can take years before the ecosystem comes into balance with the changed or cumulative amount of sulfur and nitrogen inputs. For example, certain watershed soils -- e.g., those that are sulfur saturated -- may continue to release sulfur at a higher rate for an extended period of time subsequent to a decrease in atmospheric sulfur loading. Thus, sulfate concentrations in surface waters may continue to decrease in the future as a consequence of deposition changes that have already occurred. Also, if the soil's acid neutralizing base cation reserves continue to be depleted by long-term sulfur deposition, base cation concentrations in some surface waters could continue to decrease causing additional acidification, irrespective of any further changes in sulfate concentrations.

Quantitative dose-response relationships for sulfur have been determined, using a variety of approaches in a number of regions in North America and Europe. Such studies have included, for example, measured changes in water chemistry during periods when sulfur deposition changed appreciably, regional paleolimnological investigations (e.g., diatom-inferred change in pH and acid neutralizing capacity), watershed manipulation studies, and intensive process modeling.

Measured changes in surface water chemistry in areas that have experienced short-term (less than 20 years) changes in chemical constituents in response to changes in mineral acid inputs are available from a number of sources. Proportional changes in acid neutralizing capacity, base cations, and inorganic aluminum relative to changes in sulfate or sulfate plus nitrate concentrations have been documented for several lakes and streams (Sullivan and Eilers, 1994), including lakes in the Sudbury region of Ontario; the Galloway lakes area of Scotland; a stream site at Hubbard Brook, New Hampshire; and catchment manipulation experiments in Norway and Wisconsin. Most of the observed changes are coincident with decreased acidic deposition, but it is unclear whether acidification and recovery are symmetrical.

In most cases, the largest response to the change in dose (sulfate plus nitrate concentration) has been a concomitant change in base cation concentrations. The measured base cation response has typically been in the range of 50-90%, although values as low as 35% have been found for highly sensitive watersheds made up largely of exposed bedrock. The acid neutralizing capacity response is usually smaller (10-50%), as is the aluminum response (0-15%).

Considerable experimental research has been conducted over the past several years to quantify the environmental effects of atmospheric deposition, especially of nitrogen. This research has been initiated almost exclusively in Europe; little comparable work has been conducted in the United States. The experimental approach has shifted heavily into the area of broad, whole-ecosystem manipulations across gradients of atmospheric deposition, climate, and other environmental factors. This holistic approach is used to evaluate key processes in the broader context of

ecosystem structure and function, rather than as isolated process studies. The research needs to be sustained over an extended time to understand long-term impacts of atmospheric pollutants and other factors.

The whole-ecosystem manipulation experiments in Europe have been augmented by a number of detailed, process-level studies at the various manipulation sites. Key aspects include stable isotope tracer studies to quantify the partitioning of nitrogen into various ecosystem pools (i.e., soil, litter, trees, ground vegetation) and to measure changes in the quantities of stored nitrogen in these pools. Other studies focus on quantifying the rates of important ecosystem processes, including the nitrogen conversion processes, nitrification, and mineralization.

Results of both the broad-scale and detailed studies are being used to build and test mathematical models that simulate nitrogen processing, nutrient cycling, and water regulation in coniferous forest ecosystems under varying depositional and climatic regimes. Ultimately, these models will be used to predict nitrogen saturation, estimate the critical loads of nitrogen for forests, and specify emission controls needed to protect forests from the detrimental effects of excess nitrogen deposition.

The effects of other ecosystem stressors must also be considered in quantifying ecosystem effects attributable to acid deposition. Climatic fluctuations, especially precipitation input and its effects on water availability, act synergistically with a variety of indirect effects of acid deposition. The effects of climatic variability make it particularly difficult to predict acidification or deacidification responses. The obvious linkages between short-term climatic fluctuation and anthropogenic sources of nitrogen and sulfur are incorporated into the experimental approach followed by the European EXMAN program (Tietema and Beier, 1995). Both drought and nitrogen-plus-sulfur inputs are evaluated alone and in combination under a variety of conditions. The linkage with climatic change is taken further still in the European CLIMEX project, which entails simultaneous whole-ecosystem manipulation of temperature, atmospheric carbon dioxide, and acid deposition (Jenkins et al., 1992). Long-term climatic trends will also affect ecosystem responses to acid deposition, and atmospheric concentrations of sulfur oxides have been linked with regional or global impacts on temperature.

Forest Ecosystems

- At present, the mortality and decline of red spruce at high elevations in the Northeast and observed reductions in red spruce growth rates in the southern Appalachians are the only cases of significant forest damage in the United States for which there is strong scientific evidence that acid deposition is a primary cause.
- The interaction of acid deposition with natural stress factors has adverse effects on certain forest ecosystems. These effects include:
 - Increased mortality of red spruce in the mountains of the Northeast . This mortality is due in part to exposure to acid cloud water, which has reduced the cold tolerance of these red spruce, resulting in frequent winter injury and loss of foliage.
 - Reduced growth and/or vitality of red spruce across the high-elevation portion of its range.
 - Decreased supplies of certain nutrients in soils to levels at or below those required for healthy growth.
- Whereas forest ecosystems other than high-elevation spruce-fir are not currently manifesting damages directly attributable to acid deposition, these less sensitive forests throughout much of the United States are experiencing gradual losses of base cation nutrients due to acid deposition, which in many cases will reduce the quality of forest nutrition over the long term (decades).
- Nitrogen deposition is now recognized with sulfur as an important contributor to effects on forests in some ecosystems, which occurs through direct impacts via increased foliar susceptibility to winter damage, foliar leaching, leaching of soil nutrients, elevation of soil aluminum levels, and/or creation of nutrient imbalances. Excessive amounts of nitrogen cause negative impacts on soil chemistry similar to those caused by sulfur deposition in certain sensitive high-elevation ecosystems, and is a potential contributor to these impacts in some low-elevation forests.

Responses of forest ecosystems to emissions reductions will depend on the relationship between the location of the reductions in SO₂ and NO_x and the location of sensitive receptors (soils and forests types) and other factors, such as elevation. Some of the most sensitive soils and vegetation are at high-elevation sites in the eastern United States, where regional emissions are already high and have been so for decades. The frequent occurrence of acidic cloud water approximately doubles the exposures at these sites, compared to exposure levels typically encountered at lower elevations. In these sensitive high-elevation ecosystems, nitrogen deposition can be of equal or greater importance than sulfur deposition in causing adverse effects. In contrast, in many forests throughout the United States, where nitrogen is deficient, nitrogen deposition may actually be beneficial.

Because sensitive forest receptors exhibit both a short-term and long-term response to changes in deposition, the timing of those changes needs to be estimated. But during the 1990s, there has been no systematic attempt to monitor the changes in forest condition or in the chemistry of forest soils in response to changes in acid deposition. The scarce data available have been site specific, not regionally representative, and do not cover the 1990-1996 period.

Observed Effects

- Geographic regions of the United States – and the forest ecosystems within those regions – vary in their current status and their sensitivity to effects from the deposition of nitrogen and sulfur. Variations in potential future forest response and/or sensitivity result in part from differences in (1) deposition of sulfur and nitrogen, (2) sensitivities of ecosystems to sulfur and nitrogen, and (3) responses of soils to inputs of sulfur and nitrogen.

The response of ecosystems to sulfur and nitrogen deposition is frequently an indirect one in which soil chemistry plays a potential role. Research results on soil and plant responses to sulfur and nitrogen additions in greenhouse, laboratory, controlled-exposure, and field studies indicate that negative responses will occur in nutrient depleted soils. These negative responses include: (1) plant nutrient deficiencies, particularly of base cation nutrients like calcium, magnesium, and potassium; (2) elevated levels of aluminum and calcium / aluminum ratios in soil water that interfere with plant uptake of these nutrients; and (3) elevated levels of aluminum that are directly toxic to plant roots. The last is least common of the three because it requires particularly high levels of aluminum for most forest species.

-----Text Box Begins-----

Sensitive Forest Components

Predicting changes in health or function of forest ecosystems in response to changes in levels of acid deposition requires identifying ecosystem components that are affected by acid deposition and that also affect the function of forest ecosystems. One forest component that has become more prominently recognized as a valuable indicator of ecosystem change is foliar nutrient concentration (Cronan and Grigal, 1995).

Foliar Nutrient Concentration

Both the store of nutrients in the foliage and the immediately available store of nutrients in the soils can be quite small and potentially susceptible to large depletion caused by acid deposition. There is increasing evidence that reduced availability of calcium, both from short-term increased foliar leaching (Joslin et al., 1988) and long-term reduced soil supply, may alter photosynthesis to respiration ratios and cause reductions in carbon fixation rates and whole plant growth rates in red spruce (McLaughlin et al., 1993). Foliar calcium plays a key role in a wide variety of critical plant processes, including cell division, cell wall synthesis, disease resistance, cold acclimatization, and plant senescence. Changes in calcium content in the soil caused by acid deposition make both foliar calcium leaching losses and reduced calcium availability in soils of increasing interest in understanding the ability of forest ecosystems to respond to other environmental stresses.

Forest soils

These can also be critical and sensitive ecosystem components. In examining the response of soils to acid deposition, it is important to make a distinction between short-term response (from changes in intensity factors) and longer-term response (from changes in capacity factors) (Robarge and Johnson, 1992). *Intensity factors* refer to the concentration of a chemical in soil water, whereas *capacity factors* refer to the total content of bases, aluminum, and iron stored in the soil itself. The soil represents a large chemical pool compared to the chemicals in soil water. Changes in intensity factors result from significant and rapid shifts in certain small chemical pools in the soil and vegetation. Because the chemical pool size of capacity factors is large relative to changes caused by acid deposition or other natural processes, changes in capacity factors in the short term are typically small.

Changes in intensity factors can have a rapid impact on the chemistry of soil solutions. Increases in amounts of sulfur or nitrogen in acid deposition can cause immediate increases in acidity and mobilization of aluminum in soil solutions. Increased concentrations of aluminum and an increase in the ratio of

calcium to aluminum in soil solution have been linked to significantly reduced plant availability of essential base cations, increased plant respiration, and increased biochemical stress indicators.

Changes in capacity factors are the result of many processes acting over long time periods. For example, the base cation content of soils is the result of inputs of base cations (calcium, magnesium, sodium, potassium) from atmospheric deposition, vegetative decomposition, geologic weathering, and losses due to plant uptake and leaching. Increased leaching of base cations caused by acid deposition may result in nutrient deficiencies in soils if other inputs of base cations (e.g., atmospheric deposition, geologic weathering) cannot resupply these losses at the same rate. This appears to be the case in many sensitive forest ecosystems in the United States, especially since the atmospheric deposition of base cations has declined markedly since 1980 (see earlier section on Emissions, Concentrations and Deposition) and geologic weathering rates are very slow in many ecosystems. The situation is also complicated by the removal of base cations from forest soils by tree uptake and incorporation in plant material (Johnson et al., 1993). Subsequent harvest removal of base cations is generally of the same magnitude as leaching due to acid deposition.

Even though acid deposition causes cation nutrient depletion in soils, the nitrogen in acid deposition may stimulate tree growth in certain areas. In many forest ecosystems, nitrogen is the primary nutrient that limits growth (Richter and Markewitz, 1995). Thus, positive impacts (nitrogen fertilization) and negative impacts (cation leaching from soils) may be operating at the same time in the same system. The relative importance of these two impacts must be assessed to understand the level and timing of responses of forest ecosystems to emission reductions.

The intensity and capacity characteristics of forest ecosystems are significant because they highlight the importance of interactions between deposition of strong anions (sulfate and nitrate) and natural factors that influence soil acidity and, hence, the sizes and sensitivities of nutrient pools. These characteristics are important in evaluating forest effects in the context of the different soil types, stages of stand development, and forest management practices in relation to acid deposition levels.

-----Text Box Ends-----

Soil sensitivity to deposition varies based on differential ability of soils to retain the deposited sulfur and nitrogen. Soils with a low sensitivity have high capacity to adsorb sulfur and have plants and microbes that are low in nitrogen and readily take up and use deposited nitrogen. The sensitivity of soils is also determined by the size of their stores of base cation nutrients. Sensitive soils, which cannot retain much additional sulfur and/or nitrogen (i.e., are saturated) lose these base cation nutrients as sulfur and nitrogen in soil water pass through the forest and into streams. When soil base cation stores become low, naturally occurring but potentially toxic soil aluminum begins to replace the cations in soil water. Differences in ecosystem sensitivity are also tied to a number of plant characteristics, including the sensitivity of individual species to direct effects of deposition, and the nutrient requirements and growth rates of those species. The latter affect a plant's ability to use deposited nitrogen or to respond to nutrient losses.

In addition to the broad geographic patterns of deposition, increases in elevation can cause large increases in both sulfur and nitrogen deposition. Not only do rainfall and wind speed increase, and temperature and other natural stresses change with elevation, but both cloud water and dry deposition also increase dramatically. On average, concentrations of sulfur, nitrogen, and acidity in cloud water are two to five times higher than in rainfall. Acidic cloud water, plus high winds on mountaintops and the efficient collection of cloud water by tree needles make deposition to high-elevation sites several fold higher than nearby low-elevation sites.

Sensitive Receptors

For purposes of discussion of forest ecosystem effects in this assessment, the United States has been divided into four very broad regions based on their differential sensitivities in general and the differential amounts of deposition received (**Figure 24 and Table 6**). In addition, these regions have differing aesthetic, recreational, and economic characteristics that need to be considered when evaluating impacts from acid deposition. Three regions are east of the Great Plains: high-elevation spruce-fir forests; southern pine and pine-hardwood forests; and hardwood (deciduous) forests of the Midwest and the Northeast, which includes areas of low-elevation conifer forests, especially in the northernmost portions of the eastern United States. The fourth broad region, which includes all the ecosystems west of the Great Plains, encompasses a wide variety of plant communities, but significant impacts of acid deposition have only been detected on a limited area. Most of these western ecosystems are relatively unaffected either because they receive limited amounts of sulfur and nitrogen, or because they are relatively well buffered against their effects, or both. A summary of the observed effects on each of these four types of forest ecosystems follows.

Table 6. Characteristics used to select forest ecosystems for regional assessment of the effects of acid deposition.

("X" indicates a predominant characteristic of the forest type and "VAR" indicates that some forests have these characteristics and some do not.)

Characteristics	Type of Forest Ecosystem			
	High-Elevation Spruce-Fir	Southern Pine / Pine Hardwood	Eastern Hardwood	Western Conifers
Sensitive Ecosystem	X			VAR
Sensitive Soils	X	X	VAR	
High Regional Deposition	X	VAR		
High Local Deposition	X		VAR	X
Aesthetic or Recreational Value	X	VAR	X	X
Economic Value		X	X	VAR

High-Elevation Spruce-Fir Forests

High-elevation spruce-fir ecosystems in the eastern United States epitomize sensitive soil systems. Base cation stores are generally very low, and soils are near or past their capacity to retain more sulfur or nitrogen. Deposited sulfur and nitrogen, therefore, pass directly into soil water, which leaches soil aluminum and minimal amounts of calcium, magnesium, and other base cations out of the root zone. The low availability of these base cation nutrients, coupled with the high levels of aluminum that interfere with roots taking up these nutrients, can result in plants not having sufficient nutrients to maintain good growth and health.

1 Exposure to acidic clouds and acid deposition has reduced the cold tolerance of red spruce in the
2 Northeast (Vann et al., 1992; Sheppard et al., 1993a, 1993b), resulting in frequent winter injury of
3 current year foliage during the period 1960-1985. Repeated loss of foliage due to winter injury
4 caused crown deterioration and contributed to high levels of red spruce mortality in the Adirondack
5 Mountains of New York, the Green Mountains of Vermont, and the White Mountains of New
6 Hampshire (Craig and Friedland, 1991; Eagar and Adams, 1992). However, there is uncertainty
7 about the relative importance of sulfur, nitrogen, and acidity in causing this decline in cold
8 tolerance.

9 Acid deposition has contributed to a regional decline in the availability of soil calcium and other
10 base cations in high-elevation and mid-elevation spruce-fir forests of New York and New England
11 and the southern Appalachians through two mechanisms: leaching of base cations from foliage and
12 from the primary rooting zone, and the mobilization of labile aluminum (McLaughlin and Kohut,
13 1992; Eagar and Adams, 1992). The role of nitrogen deposition in this deterioration of soil base
14 cation availability is greater than previously realized.

15 Changes in soil chemistry and red spruce nutrition over the past few decades may have contributed
16 to both growth declines and deterioration of crown condition in the Northeast and at the higher
17 elevations of the southern Appalachians, based on physiology changes induced by acid deposition
18 in field and laboratory studies (McLaughlin and Kohut, 1992). While red spruce in the southern
19 Appalachians has not shown areawide increased mortality in recent decades (Peart et al., 1992),
20 approximately a two to threefold increase in significant crown deterioration at upper elevations was
21 observed between 1985 and 1989. The relative roles of acid deposition and severe winter weather
22 on red spruce crown deterioration in the southern Appalachians is unclear (Peart et al., 1992).
23 While ice damage has been associated with loss of canopy integrity in the Black Mountains
24 (Nicholas and Zedaker, 1992), it should be noted that significant loss of canopy integrity also
25 occurred in the Smoky Mountains where significant ice damage was not observed (Johnson et al.,
26 1992b). Foliar calcium levels and soil and root calcium / aluminum ratios are considered low to
27 deficient over most of the southern spruce-fir region and portions of the northern region (Joslin et
28 al., 1992; Cronan and Grigal, 1995). Changes in soil chemistry, trends in wood chemistry, and
29 changes in lake water chemistry suggest that a significant loss of calcium has occurred during
30 recent decades from the organic soil layers, which is the primary rooting zone for red spruce
31 (Lawrence et al., 1997; Shortle et al., 1997). Both tree uptake and storage of calcium and acid
32 leaching losses are responsible for these losses from the soil (Joslin et al., 1992). Fertilization
33 experiments in the southern Appalachians show positive growth responses and positive leaf
34 physiological responses to additions of calcium (Joslin and Wolfe, 1994; McLaughlin and Kohut,
35 1992). Stress in red spruce in the Northeast is indicated by an increase in the biochemical stress
36 marker putrescine in red spruce needles from areas of high acid deposition and low soil calcium /
37 aluminum ratios (Minocha et al. 1996; Shortle et al. 1997). Negative physiological responses have
38 also been observed at highly polluted sites in the South (McLaughlin et al., 1991) and in the North
39 (Amundson et al., 1992).

40 Recent evidence indicates that most southern Appalachian spruce-fir ecosystems are at or near
41 nitrogen saturation. Nitrogen saturation is often defined as a condition of the terrestrial ecosystem
42 where outputs from the system equal or exceed inputs, resulting in elevated levels of nitrogen in
43 soil solutions, groundwater, and streams draining the terrestrial system (see Aquatics section,
44 under the Nitrogen subheading). For these saturated systems, high nitrogen deposition is
45 transmitted through the ecosystem and results in base cation and aluminum leaching from soils,
46 elevated aluminum in soil water, which reduces base cation nutrient uptake, and elevated nitrate
47 and aluminum levels in streams (Joslin et al., 1992; Van Miegroet et al., 1992).

1 *Southern Pines and Pine-Hardwood Forests*

2 Significant impacts of acid deposition on forest health have not been detected in the southern pine
3 and pine-hardwood region. However, acid deposition is contributing significantly to the depletion
4 of base cations in many poorly buffered soils supporting southern pines and will most likely, over
5 the long term (decades), impede productivity. Short-term positive effects on growth are expected
6 for some nitrogen-deficient soils, while negative effects are expected to be limited to the most
7 acidic, base-depleted soils.

8 A synthesis of studies that originated as a part of NAPAP to evaluate the sensitivity of southern
9 pines to acid deposition and ozone has now been completed (Fox and Mickler, 1995). In chamber
10 studies, saplings of the three principal commercial pine species (loblolly, shortleaf, and slash pine)
11 were exposed to simulated acid rainfall and ozone. Maximum annual growth reductions in
12 saplings due to ambient ozone were quite small (2-5%), but yield could be significantly reduced
13 over longer time frames (Teskey, 1995). Growth rates of saplings typically responded positively
14 to ambient levels of acid rainfall. However, longer-term exposures are expected to have
15 cumulative negative effects on soil nutrition. Currently, an estimated 59% of the pine forests of the
16 Southeast have soils that are low enough in base cations to be susceptible to acidification by cation
17 leaching due to acid deposition, and 10% of the soils are considered extremely acidic (Richter and
18 Markewitz, 1995; after Turner et al., 1986). An estimated 10-15% of the commercial pine forests
19 in the South may be limited by availability of these nutrients (Binkley et al., 1989). Thus,
20 additional leaching may further limit pine growth on these soils. While these impacts will most
21 likely have negative long-term consequences, the inputs of atmospheric sources of nitrogen to
22 many soils with low nitrogen reserves, which currently support southern pine forests, should have
23 small cumulative positive effects on productivity for as long as decades (Allen, 1987; Binkley et
24 al., 1989; Dougherty, 1996).

25 *Eastern Hardwood Forests*

26 Studies have shown that acid deposition has caused increased cation leaching from some hardwood
27 ecosystems, and this leaching may affect the health of those systems (Likens et al., 1996; Mitchell
28 et al., 1996; Mitchell et al., 1994; Gilliam et al., 1996; Norton et al., 1994; Rustad et al., 1996;
29 MacDonald et al., 1992). Although sulfate is the primary anion causing base cation leaching,
30 nitrate is a significant contributor in watersheds that are nearly nitrogen-saturated as a result of acid
31 deposition (Peterjohn et al., 1996), and studies indicate that some hardwood ecosystems are close
32 to nitrogen saturation (Mitchell et al., 1996b). Nevertheless, in general, the health of eastern
33 hardwood forests has not been shown to be adversely affected by acid deposition. However,
34 broad-scale monitoring has not been conducted to confirm this finding.

35 The compositional properties of foliage may also be altered by acid deposition, resulting in changes
36 in organic matter turnover and nutrient cycling (Adams et al., in press). The sensitivity of
37 hardwood soils to acid deposition is largely controlled by inherent properties, climate, and land use
38 (Mitchell et al., 1996a). However, tools to assess present conditions or susceptibility to nutrient
39 depletion are not readily available or widely applicable (Desanker et al., 1994).

40 *Western Forests*

41 While most of the forest ecosystems in the western United States do not show significant impacts
42 from acid deposition, certain areas appear to be affected negatively, especially by high rates of
43 nitrogen deposition. Forested sites near urban areas in southern California and alpine meadows of
44 the Front Range in Colorado have been exposed to high levels of nitrogen and sulfur deposition for
45 the last 40 years (Kiefer and Fenn, 1997; Williams et al., 1996c). Many of these sites are showing
46 signs of nitrogen saturation, which include elevated levels of nitrogen in streams (see the previous

section on Aquatics) and changes in the cycling of nitrogen and carbon. Continued nitrogen deposition may also lead to shifts in species composition and biodiversity.

In southern California, ozone exposure and nitrogen and sulfur deposition occur coincidentally. Ozone exposure and acid deposition are greatest near Los Angeles, and decrease along a gradient running to the east/northeast toward the San Bernardino and San Gabriel Mountains (Fenn and Bytnerowicz, 1993). The greatest exposure to all pollutants occurs in the summer, when these forests are also subject to drought stress (Bytnerowicz and Fenn, 1996). These multiple stressors weaken trees, making them vulnerable to bark beetle attack, which ultimately causes mortality (Miller, 1992). In addition, ozone causes premature leaf abscission, and nitrogen deposition has a major impact on soil and forest floor processes. These factors combined can lead to changes in patterns of nitrogen and carbon cycling (Fenn, 1991).

In the alpine areas of Colorado, nitrogen deposition occurs during summer and winter (Sievering et al., 1996). Nitrogen additions to alpine meadows also change the cycling of nitrogen and carbon. Nitrous oxide emissions from soil increase and rates of methane oxidation decrease (Neff et al., 1994). Methane is a greenhouse gas and its removal from the atmosphere by oxidation is an important function of natural systems (Neff et al., 1994). In addition, sites with high nitrogen deposition have decreased surface water quality due to elevated nitrate levels.

Dose-Response Relationships

- Models predict that recent reductions in sulfur deposition will rapidly improve some aspects of soil chemistry related to intensity factors (see text box on Sensitive Forest Components) in sensitive forests (i.e., immediate but brief increases in the concentrations of base cations in solution and/or reductions in aluminum concentrations), but full recovery of these soils and forests from previous impacts will be quite slow (decades to centuries). Full recovery would require: (1) re-growth of dead stands, (2) return to historical growth rates for species like red spruce, and (3) replenishment of lost cation nutrient stores.
- Nitrogen deposition is of approximately equal importance to sulfur deposition in causing changes to occur in sensitive forest ecosystems. Such systems should rapidly show partial recovery in response to significant reductions in nitrogen deposition.

-----Text Box Begins-----

Soil-Related Effects

In sensitive soils – ones where base cation stores are very low and the ability of the ecosystems to retain sulfur or nitrogen is minimal – the response to changes in sulfur or nitrogen doses can be quite rapid. As levels of sulfur and nitrogen compounds in soil water decline, both aluminum and base cation levels decline as well, but aluminum declines slightly more rapidly. The opposite occurs as concentrations of sulfur and nitrogen compounds increase; aluminum increases more rapidly than base cations increase. Responses – both up and down – are particularly sensitive to changes in nitrogen inputs because in nitrogen-saturated ecosystems nitrate passes quickly into soil water and on to streams. Sulfur compounds are more likely to be retained by various reactions in the soil. While some rapid changes in soil water chemistry will occur, full replenishment of base cation stores will be a slow process.

The response of less sensitive ecosystems is considerably slower. These systems either still have adequate stores of base cation nutrients and/or still maintain considerable capacity to retain the sulfur and nitrogen deposited on them. Most forest ecosystems in the United States, such as southeastern pine plantations, are in this less sensitive category, and most are slowly losing base cations and slowly reaching their capacity to retain sulfur and nitrogen. However, in some systems, virtually all sulfur and/or nitrogen deposited is

either adsorbed on the soil or used by growing vegetation or microbes. Much of this sulfur and especially nitrogen may be later removed by forest harvesting. These systems are not negatively affected by increases in sulfur and nitrogen deposition, and even may respond with more rapid growth.

Forest-Related Effects

Understanding of how sensitive forest ecosystems respond to changes in acid deposition has significantly improved since the *1990 Integrated Assessment Report* (NAPAP, 1991). For discussion of general concepts, high-elevation spruce-fir forests are used as a model. In general, other forest ecosystems in the United States are not currently experiencing the same effects in mortality and growth as the spruce-fir ecosystem. However, the process of base cation leaching and a trend toward more acidic, aluminum-dominated soils is occurring slowly in many of these forested ecosystems. **Figure 25** illustrates the general dose-response relationship for nutrient limitations associated with increasing soil and deposition acidity.

The availability of calcium is increasingly recognized as important to forest health (McLaughlin and Kohut, 1992). In studies of spruce-fir forests, calcium deficiency symptoms increased in red spruce saplings at high-elevation sites with increased exposure to acid deposition. Low calcium availability was well correlated with high levels of soil aluminum. Acidic mists can reduce the efficiency of carbon assimilation in foliage, cause reduced growth, leach foliar calcium, and alter patterns of root growth. Fertilizer experiments in the field have now substantiated that foliar calcium levels at high-elevation sites are low enough to reduce growth of both red spruce saplings (Van Miegroet et al., 1993) and mature trees (Joslin and Wolfe, 1994).

Changes in foliar nutrient content, soil nutrient content, and the availability of calcium are involved in the observed responses of red spruce to acid deposition. While some rapid reduction in base cation leaching can be expected in response to reduction in emissions, at present there is not a solid basis for evaluating the relative importance of such intensity factors and longer term, soil-based capacity factors. Hence, accurate quantitative projections of response rates to emission reductions are not possible.

-----Text Box Ends-----

Regional Differences

When evaluating how forests respond to changes in the levels of acid deposition it is important to take into account specific soil and forest responses (see soil- and forest-related effects in corresponding text box). The different responses of various regions are described below.

High-Elevation Spruce-Fir Forests. Reductions in ambient concentrations of acid deposition and cloud water should rapidly reduce the susceptibility of red spruce to winter injury. However, data are not available to provide the specific reduction levels of various compounds (sulfate, nitrate, or acidity) that are required to fully protect foliage from past increases in susceptibility to freezing injury. A linear dose-response relationship exists between the acid dose from cloud water and the freezing injury (Fowler et al., 1989; Cape et al., 1991; Sheppard, 1994) both in the Northeast (Vann et al., 1992) and in the southern Appalachians (DeHayes et al., 1991, 1992; Thornton et al., 1994). While increased frequency of damage to red spruce foliage has been notable at high elevations in the North, only a very mild form of winter damage has been observed in the South (Anderson et al., 1991), probably because extremely low temperatures occur less frequently at southern high elevations.

While high-elevation spruce-fir forests should experience some immediate improvements in health with reductions in sulfur and/or nitrogen deposition, their growth and vitality will probably be limited by the availability of certain nutrients for decades. Results of modeling efforts using a nutrient cycling model (NuCM) for a highly acidic spruce-fir forest in the southern Appalachians

predicts that significant reductions in sulfur or nitrogen deposition would result in some rapid improvement in the amount of calcium available for uptake (Johnson et al., 1996) and small improvements in the calcium / aluminum ratio. While this model predicts significant improvements in the short-term supplies of calcium and other base cations, the recovery of long-term storage capacity will be limited because (1) existing high levels of aluminum saturation reduce the ability to store calcium introduced by atmospheric deposition, weathering, or decomposition of litter (Lawrence et al., 1995; Lawrence et al., 1997), and (2) decreased atmospheric deposition of base cations will limit inputs (Likens et al., 1996).

The future health of spruce-fir ecosystems may also be quite sensitive to increased temperatures due to global warming (McLaughlin et al., 1987, 1997). High-elevation spruce-fir soils are particularly vulnerable to increased acidification by their release of stored nitrogen as a result of warming. These soils typically have very large stores of nitrogen in their organic matter (Fernandez, 1992; Johnson et al., 1991; Joslin et al., 1992). Rates of nitrogen mineralization, nitrate release, nitrate leaching, and nitrogen in streams are already very high in these ecosystems (Sasser and Binkley, 1989; Joslin et al., 1992). Increases in nitrate resulting from soil warming have been observed in field studies (Joslin and Wolfe, 1993; Mitchell et al., 1996b). Warmer temperatures will increase foliar respiration, an effect that has also been stimulated by calcium deficiency and increasing exposure of red spruce to acid deposition in the southern Appalachians (McLaughlin et al., 1992). Analyses of tree-ring data from mature high-elevation red spruce trees in the southern Appalachians show that growth effects associated with warmer temperatures and higher rainfall have become increasingly negative during the past three decades (McLaughlin et al., 1997).

Southern Pines and Pine-Hardwood Forests. The productivity of southern pines is expected to be more limited by ambient ozone levels than by ambient levels of acid deposition. Short-term (one- to three-year) experiments to evaluate the interactive effects of acid deposition and ozone on seedlings and saplings have indicated that ambient levels of acid deposition have not exacerbated the negative effects of ozone on growth (Flagler and Chapelka, 1996). Studies on older trees in acid soils are needed to evaluate the long-term impacts.

Adverse effects of ozone on growth have typically been associated with reduced carbon gain in photosynthesis and premature loss of foliage. Estimated annual effects of ozone on seedlings under ambient exposures are typically small (2-5%) and difficult to detect statistically, but are a basis of concern for longer term, cumulative effects on growth (Teskey, 1995). In addition, studies with mature loblolly pine trees have demonstrated short-term reductions in stem growth at ambient ozone levels that were amplified by both moisture stress and high air temperatures (McLaughlin and Downing, 1995; 1996). The latter study suggests that future ozone effects may be amplified if global change produces a warmer, drier climate.

In some cation-deficient soils, potentially positive short-term growth responses to deposited nitrogen may be followed by longer-term negative influences of acid deposition due to continued leaching of base cations (Flagler and Chappelka, 1995; McLaughlin et al., 1994; Richter and Markewitz, 1995). Predictions of growth responses across the region are not available yet, however, because the effects have not been quantified.

Reducing the levels of acid deposition will reduce the rate of cation depletion of southern pine forests. However, recovery of soil cations by natural processes is expected to be slow because both mineral weathering and cation recycling are very low (Richter and Markewitz, 1995). Since neither growth losses from past acid deposition levels nor growth enhancement from nitrogen fertilization have been quantified to date, it is not possible to estimate either the direction or the rate of change in these forests in response to emission reductions.

Eastern Hardwood Forests. Hardwood soils are known to be sensitive to acid deposition, but the dose-response relationships have not been determined. The sensitivity is expressed in terms of low buffering capacity (i.e., low resistance), but low levels of acidic input can cause nitrogen saturation and base cation depletion. Changes in atmospheric deposition may cause rapid changes (months to years) in the chemical properties of soils in hardwood forests. Models may be useful in assessing dose-response relationships, but they must be tested and validated.

Lack of specific dose-response relationships or modeling tools to assess sensitivities or thresholds precludes large-scale assessments of hardwood forest responses to altered levels of acid deposition. Watersheds exposed to chronic, low-level acidic inputs have shown symptoms of nitrogen saturation and base cation depletion, and experimental additions of nitrogen and sulfur have shown that nitrogen saturation can be induced in a short period (i.e., three years) (Norton et al., 1994). Assessment of ecosystem responses to dosed additions of nitrogen and sulfur is further complicated by the exposure history of the site or watershed, land use history, natural disturbance regime, and climate. Models have been useful in evaluating acidification processes and cation leaching (Johnson et al., 1996); however, further model development and validation are needed.

Western Forests. Several sensitive areas in the western forests are becoming nitrogen saturated due to chronic nitrogen deposition. Pine forests in the San Bernardino Mountains of southern California receiving high levels of nitrogen deposition (~25 kg/ha/yr) are already nitrogen saturated, while those receiving moderate levels (6-16 kg/ha/yr) show evidence of approaching saturation (Kiefer and Fenn, 1997). Exposure to high levels of sulfur, nitrogen, and ozone, along with drought stress, weakens the trees and leads to premature mortality.

Alpine meadows along the Colorado Front Range are also showing signs of nitrogen saturation, but at lower levels of nitrogen input (3-5 kg/ha/yr) than those in southern California (Sievering et al., 1996). The increase in nitrogen deposition at sites in Colorado has coincided with climatic changes, including decreased mean annual temperature, increased annual precipitation, and decreased daily solar radiation. The increase in precipitation accounts for about 50% of the increase in nitrogen deposition. These changes in climate are driven by local conditions and may not be related to regional and global climate trends (Williams et al., 1996c). In the alpine areas of Colorado, about half of the nitrogen is deposited in snowpack and then is released with the spring thaw. This large release of nitrogen leaches through the soil, removes base cations from the soil, and degrades the quality of streams and lakes (Williams et al., 1996b).

-----Text Box Begins-----

The Calcium/Aluminum Ratio as a Forest Ecosystem Indicator

The health of forests can be assessed by monitoring such indicators of stress as crown condition, growth, mortality, and basal area. The calcium / aluminum ratio of the soil solution is a possible ecological indicator for identifying approximate thresholds at which the risk of forest damage (tree mortality or reduced growth) from aluminum stress and nutrient imbalances increases (Cronan and Grigal, 1995). This ratio can also be used to assess forest ecosystem changes over time in response to acid deposition as well as to forest harvesting and other processes contributing to acid soil infertility. This ratio may not be a reliable indicator of stress in (1) areas with both high atmospheric deposition of ammonium and magnesium deficiency via antagonism involving ammonium rather than aluminum, and (2) areas with soil solutions of calcium concentrations greater than 500 micromoles per liter.

If the calcium / aluminum ratio is 1.0, then forests have a 50% risk of adverse impacts.
If the ratio is 0.5, then the risk is 75%; if the ratio is 0.2, then the risk is nearly 100%.

The overall uncertainty of the calcium / aluminum ratio associated with a given probability estimate is $\pm 50\%$. Cronan and Grigal (1995) suggest four sequential indices to corroborate the calcium / aluminum ratio of the soil solution.

- Soil base saturation less than 15% of effective cation exchange capacity (Reuss, 1983; Cronan and Schofield, 1990)
- Soil solution calcium / aluminum molar ratio less than 1.0 for 50% risk
- Fine root tissue calcium / aluminum molar ratio less than 0.2 for 50% risk
- Foliar tissue calcium / aluminum molar ratio less than 12.5 for 50% risk

-----Text Box Ends-----

Agriculture

Based on crop-effects research conducted by NAPAP and other research programs in the 1980s, acid deposition at ambient levels in the United States was not found to be responsible for regional crop yield reduction. In fact, sulfur and nitrogen deposition in precipitation can provide a portion of the crop requirement for these elements essential for plant growth (NAPAP, 1991 SOS/T 18). Ozone, on the other hand, represents a significant stress factor in agricultural production in the United States. NO_x reductions under Title IV should decrease the creation of ozone and its subsequent damage to agriculture. However, data on changes in ozone resulting from Title IV are not currently available.

Materials and Cultural Resources

- Approximately 900,000 properties of aesthetic and historic value in the United States are potentially at risk for damage by air pollution, including sulfates, other particulate matter, and ozone. This figure does not include the 10-30 million tombstones and grave site memorials. The value at risk of such cultural assets (e.g., statues, monuments) is greater than that for purely operational resources (e.g., buildings, fences, bridges) because the value of cultural resources is mainly in their appearance, which is particularly vulnerable, and because of the emotional capital attached to such properties by an extended population.
- Structures made of limestone and marble are particularly sensitive to acid deposition. The salts produced on stone surfaces as the result of acid deposition are now considered to be potentially more damaging than are the dissolution effects of the actual acid delivered to the stone. While the incremental erosion of calcareous stone due to acid rain is generally a small fraction of the background erosion by clean rain (clean rain effect), the soiling effects and the loss of structural integrity of the stone associated with salt deposition generally represent the most deleterious effects.

Observed Effects

Acid deposition, particularly dry deposition of fine sulfate aerosols (Haynie and Lemmons, 1990) and fine particulate matter, damages materials used both for operational assets and for cultural resources potentially affecting their structural and aesthetic values. While the reduction of sulfur dioxide emissions associated with Title IV is expected to decrease the rate of deterioration of materials, the effects of such a reduction may not be currently visible because latent incremental effects often require decades to appear above the natural soiling and weathering effects.

An implicit assumption by NAPAP of materials effects research is that the benefits of Title IV's pollution reduction for cultural materials (such as statues, monuments, and historic buildings) greatly exceed those for purely operational assets (such as commercial buildings, fences, and roads). The value at risk is greater for cultural resources because such resources are mainly valued for their unique character and historic significance. These attributes are normally associated with the surface appearance and morphology of the construction materials, rather than with their bulk properties. In contrast, the preservation value of purely operational assets is not significantly affected by pollution because their utility is normally associated with the bulk properties of the materials used for their construction (shelter, strength, etc). It follows that pollution compromises the general utility of cultural resources in less time than it diminishes the utility of purely operational assets. However, there is no program for systematically monitoring the condition of cultural properties for potential damage from air pollution. Moreover, while the impact on cultural resources is not generally reversible, any negative impact of pollution on operational assets (e.g., paint on a bridge) can be reversed because replacement is a viable option.

Most benefit-assessment research has focused on the materials that are predominately used for cultural resources. Such materials include granite, calcitic sandstone (i.e., sandstone cemented with calcite rather than silicate minerals), bronze, mortar, limestone, and many types of marble. Calcareous stone, which is the stone type used for many cultural resources, has received precedence in materials effects research and, thus, is the focus of discussion in this report.

Studies of marble tombstone weathering in North America demonstrate that air pollution (probably sulfur dioxide) has been responsible for more deterioration of carbonate building stone and statuary than other weathering processes (Meierding, 1993). Indeed, most studies have focused more on the effects of sulfates than on the effects of nitrates because the products of nitrate deposition have not been found to be as pervasive as sulfates in stone deterioration processes and the relationship between NO_x emissions and ozone damaged materials is not well understood.

The interactions between pollutants and calcareous stone are complex and involve multiple processes that are highly interdependent. Laboratory and field studies of the degradation mechanisms have demonstrated that salts resulting from the wet and dry deposition of acids and the deposition of particulate matter are more damaging to calcareous stone than the dissolution effects of the actual acids delivered to the stone. Because the dissolution due to acid is generally insignificant relative to the dissolution effect of clean rain, the focus of research on stone deterioration has shifted to an understanding of alteration effects that degrade the aesthetic and structural qualities of the resource. Geographic location and the natural environment of the resource also affect the extent and magnitude of damage. For example, stone degradation rates are negligible in dry areas, such as southwestern United States. Degradation processes for wet and dry deposition are discussed in the accompanying text box.

-----Text Box Begins-----

Deposition Effects on Stone

The direct effects of wet deposition (acid delivered in rain water) on calcareous stone are relatively inconsequential. This is because once the stone is wet, the dissolution effect of pure water generates chemical species (OH^- and HCO_3^-) that neutralize the acid in the rain before it can attack the stone surface. The acid neutralizing capacity of the moisture layer on the stone depends on the pattern of rain delivery, the contact time of water on the stone, and the flow of water on the surface of the stone. The complete neutralization of acid delivered in the rain is feasible because the contact time of water on the stone can be many hours and because water, the species actually dissolving the stone, is never depleted from the solution in contact with the stone.

Sulfur dioxide deposited between rain events (dry deposition) forms gypsum deposits (McGee, 1996), which can deteriorate and weaken the stone. When the stone is damp (from dew, condensation, etc.), the dissolution effect of pure water increases the acid neutralizing capacity of the moisture layer facilitating the dry deposition of sulfur dioxide on the damp surface by enhancing the solubility of the gas. The gypsum deposits generally diminish the aesthetic value of the resource by trapping particulate material and by forming darkened soiling patterns in areas that are unwashed by rain. In some cases, catastrophic failure of the stone is the final result. An accurate quantitative assessment of the effects is not possible at this time because the deterioration mechanisms generally are driven by the action of flowing water over the stone. The intricacies of the water flow pattern over the stone are exceedingly sensitive to the water delivery pattern, stone morphology, and surface conditions of the stone. Currently, techniques are not available to model the processes that are driven by water flow, even for the simplest systems.

-----Text Box Ends-----

In a recent reinterpretation of the NAPAP-sponsored laboratory and field observations of materials effects, the most conspicuous dynamics of stone decay were codified (Mossotti et al., 1997). An elementary System-wide Unified Reduced Form (SURF) model was developed to show the dominant processes of stone erosion and soiling (see the corresponding text box). The SURF model is not intended to represent a conclusive understanding of stone decay, but rather to serve as an operational framework that provides perspective on the impact of environmental stresses on exposed materials. Under actual field conditions, stone weathering involves interactions between ever-changing meteorological variables -- such as rain, wind, temperature, humidity and other factors -- and continually changing complex stone surfaces.

Some interactions occur at the microscale, while others are affected by the macro properties of the stone's surface. These interactions are not independent. When they positively reinforce each other, the system can be driven to failure. For example, the way water flows over a carved stone surface (a macroscale variable) influences the mobilization of salts and particulate materials on the

surface and can change the stone's chemistry and porosity (microscale variables). A slight change in the overall shape of the surface, either due to weathering or by design, may change the way water flows over the surface and could radically influence the erosion and soiling patterns.

Sensitive Receptors

Both cultural materials and operational materials are sensitive receptors at risk to damage by acid deposition. Cultural properties are considered to be the most at risk to damage from acid deposition because of the greater value placed on the potential loss of these resources as was previously discussed. Some of the specific materials used for cultural properties include granite, calcitic sandstone (i.e., sandstone cemented with calcite rather than silicate minerals), bronze, mortar, limestone, and many types of marble. The focus of most of the research has been on calcareous stone (limestone and marble) and, thus, is the focus of discussion in this report.

Dose-Response Relationships

A dose-response relationship has been used to partition the effects of natural and anthropogenic acidity on galvanized steel (Spence et al., 1992; Spence and McHenry, 1994). The model has been used to predict regional corrosion rates of fencing, wire, and flat plates in eastern North America. Field studies on well-characterized calcareous stone have been used to develop preliminary dose-response relationships (Reimann, 1994; 1995a; 1995b). Laboratory studies in controlled environments have provided quantitative measurements of sulfur dioxide deposition to calcareous stone (Spiker et al., 1992; 1995). Although these basic field and laboratory data on materials effects have been systematically collected over the past 15 years, there have been many difficulties with reconciling observations from poorly controlled field experiments with results from highly sanitized laboratory experiments.

A preliminary damage function for marble has been developed based on studies of the effects of air pollution on gravestones in North America (Meierding, 1993). However, this linear damage function does not take geographic location into consideration. The slope of the damage function will differ for those areas with less pollution and more rain. Weathering rates for heavily polluted urban areas may be greater than measured because many badly weathered gravestones may have been broken or removed. Other improvements to the damage function may arise from better modeling of historic sulfur dioxide concentrations at more locations throughout the United States.

-----Text Box Begins-----

Degradation Process Model

The model in **Figure 26** describes three types of processes that lead to the deterioration of the stone:

I. Primary deposition--The way that dirt and pollutants are introduced to the surface of the stone. Once pollutants are present on the surface, they can react with the stone's surface or they can be mobilized, usually by water, and redeposited elsewhere on the surface. Processes that redeposit the pollutants are known as secondary mobilization processes, shown in Panel II. Implicit in the connections between Panel I and Panel II is the notion that the primary deposition processes feed the secondary mobilization processes. Processes that change the physical or chemical nature of the stone are called physiochemical alteration processes and are represented in Panel III. Chemical changes usually occur through chemical reactions. Physical changes arise through processes such as erosion or abrasion.

Boxes A, B, and C represent the collection of variables that drive the processes represented in Panels I, II, and III. There are three main types of variables: those that describe the conditions to which the stone is exposed in its ambient environment (Box A), those that describe the physical condition of the stone (Box B), and those that define the chemistry of the stone (Box C). The variables in Box A include the ambient sulfur dioxide concentration, the level of airborne particulate material, the meteorological conditions to

which the system is subjected, and the rain, daily moisture, and rising damp delivery patterns. Variables in Box B that represent the physical state of the stone include morphology, orientation of the surface, sub-surface fracturing, and degree of grain consolidation. Box C represents variables that represent the chemical state of the stone, such as elemental composition and crystalline structure.

An integral feature of the SURF model in Figure 26 is the multiplicity of feedback loops which couple the stone variables and the deterioration processes in the system. For example, a physiochemical change may affect the stone's morphology, but a change in morphology subsequently affects the physiochemical processes. The importance of feedback to the operation of the system cannot be overestimated. If the feedback loops are negative, the state of the system tends to adapt to changes in the input variables and to resist changing its rate of deterioration. If the feedback loops are positive, the state of the system tends to move toward a particular limit. Extreme examples of such limits include highly encrusted soiled stone, well-washed virgin clean stone, and complete mechanical failure of the stone.

Two additional properties of the system are not represented in Figure 26: the system dynamics are iterative, and the system has a long memory. In iterative systems with memory, the inputs at any given time are influenced by the cumulative outputs over an extended period in the past. Such systems, which run continuously, are exceedingly sensitive to initial conditions, and tend to be attracted to extreme limit-states.

-----Text Box Ends-----

Visibility

- Particulate sulfate formed in the atmosphere by the conversion of SO_2 is responsible for 40-65% of the haze in the eastern United States, based on a combination of measurements and calculations. Reduced SO_2 emissions under Title IV are expected to decrease sulfate concentrations and their contribution to haze.
- In 1995, sulfate concentrations were 15% lower than previous years at sites in the eastern United States. Although this decrease was predicted to improve visibility by 8% there was no measurable difference at the seven eastern sites that directly measure visibility. The relative influences of emission reductions, meteorological variations, and changes in nonsulfate particulate concentrations in the observed visibility are currently not well understood.

SO_2 emissions have an indirect effect on visibility. Some of the SO_2 emitted is converted in the atmosphere to sulfate aerosol, which is responsible for a large share of the haze in the eastern United States. Visibility levels can be directly measured at sites where optical instruments are used. At other locations, where only aerosol composition data are available, calculations can be made of visibility levels and the contribution of the five major aerosol species (sulfates, nitrates, organic carbon, elemental carbon, and crustal elements) to reduced visibility.

Observed Effects

The Interagency Monitoring of Protected Visual Environments (IMPROVE) network is the principal source of aerosol and visibility information (Malm et al., 1994). Thirteen of the IMPROVE sites, are located in the East, where emission changes under Title IV are expected to be the greatest (Figure 27). Twelve of those sites are in rural areas. At least three of the 13 eastern sites have been operational continuously since 1988: Acadia National Park, Maine; Shenandoah National Park, Virginia; and Great Smoky Mountains National Park, Tennessee. The Shenandoah and Great Smoky sites actually have data beginning in 1983, prior to the establishment of the IMPROVE network. Ten additional aerosol monitoring sites were initiated during 1992 and 1993. All 13 sites collect samples that allow for the analysis of the five major aerosol species for the fine

1 particle fraction (diameter of less than 2.5 μ m) and the mass concentration of the coarse particle
2 fraction (diameter of 2.5-10.0 μ m). Seven of the sites also directly measure visibility.

3 **Figures 28 and 29** show the average seasonal and annual fractional contributions of sulfates
4 and nitrates to the light extinction coefficient based on data and calculations from Sisler et al.
5 (1996). On an annual basis, sulfate aerosols contribute from just over 40% to nearly 65% of the
6 haze seen in the East. Sulfate contributions are highest in the summer (up to 70%) and lowest in
7 the winter (as low as 30%). Nitrates contribute 5-12% of the haze on an annual average basis,
8 with much lower contributions in summer (less than 5%) than in winter (greater than 20%).
9 Nitrate concentrations in urban areas are generally higher than in surrounding rural areas as a result
10 of the influence of urban transportation sources. It is important to note that a seasonal change in
11 visibility is likely to include days with no change and other days with significant change. The
12 same can be true for spatially averaged visibility. Within a region, the overall change in visibility
13 may appear small but may actually include certain areas that experience very noticeable changes in
14 haze levels (e.g., near sources that have reduced emissions).

15 The *1990 Integrated Assessment Report* (NAPAP, 1991) predicted that a 10-million-ton reduction
16 in SO₂ emissions would result in a 37% average reduction in sulfate concentrations and, therefore,
17 a 37% reduction in the sulfate contribution to light extinction. It also indicated that sulfate was
18 responsible for about 57% of the total light extinction on average, so that a 21% change in total
19 light extinction (57% of 37%) would result from the 10-million-ton SO₂ reduction. Annual
20 average sulfate reductions were predicted to vary spatially from more than 44% in the upper Ohio
21 River Valley to about 30% across much of the East moving away from the Ohio River. This is due
22 to having significantly greater SO₂ emissions in the Ohio River Valley than in other parts of the
23 eastern United States. Summer (April to September) was predicted to have somewhat greater
24 reductions in the sulfate concentrations (40%) than in the winter season (37%).

25 Spatial and seasonal patterns of sulfate change using the IMPROVE monitoring data were
26 examined to compare with predictions in the *1990 Integrated Assessment Report* (NAPAP, 1991).
27 The 1995 SO₂ emission levels were about five million tons lower than those of the previous few
28 years as a result of Title IV emission controls within the electric utility industry (see Figure 3).
29 Assuming that the nonutility SO₂ emissions were unchanged, about a 20% reduction would occur
30 in the annual average sulfate contribution to light extinction, which would correspond to about a
31 10% reduction in total light extinction. Because most of the emission reductions occurred near the
32 beginning of 1995, the three year data record at the 13 eastern IMPROVE sites is useful in short-
33 term trends analysis. The two sites with the longer monitoring records can be used to examine
34 long-term trends and to investigate variability during periods of relatively small annual SO₂
35 emission changes.

36 Interannual variations in meteorology confounds any attempt to make simple comparisons between
37 predicted and measured changes in sulfate concentration. Measured sulfate concentrations are
38 influenced by both emission changes and meteorology. Even years that may seem to be "typical"
39 can have meteorological differences that would have significant effects on the sulfate
40 concentrations, independent of SO₂ emission rates. A common approach to handle these
41 meteorological variations is to average 10-20 years of data. Obviously, this is not possible when
42 looking for the effects of emission changes over the previous five years or less. A more
43 sophisticated approach (not used here) would be to use deterministic or statistical models to try to
44 account for influence of meteorological variations.

-----Text Box Begins-----

Visibility Impairment

By interfering with the transmission of light, pollutants associated with acid deposition reduce the clarity of the atmosphere. Much of the concern about visibility is related to the aesthetic damage from air pollution – the inability to see the form, texture, and color of scenic features.

Unlike certain acid deposition effects that involve air/land/water interactions that build up over time or that are delayed, atmospheric optical effects are instantaneous properties of air quality. The fundamental physics relating light extinction and other optical parameters to atmospheric gases and particles is well established.

Under a variety of viewing conditions, “visibility reduction” or “haziness” is approximately proportional to a parameter called “atmospheric light extinction.” Light extinction refers to the fraction of light that is attenuated per unit distance by the atmosphere. Light extinction is a simple linear sum of four components: light scattering and light absorption by both gases and particles. In a homogenous atmosphere, visual range is inversely proportional to light extinction. However, neither visual range or light extinction necessarily relates directly to the aesthetic qualities of an individual scene. A new “haziness” index has been developed for this (Pitchford and Malm, 1994). Measured in deciview units, this index was designed to be more linear with respect to perceived visual change over its entire range, analogous to the decibel scale for sound. A one deciview change represents a small change in scenic quality that would be noticed by most people regardless of the initial visibility condition. Pollution free air corresponds to 0 deciviews, with higher values indicating greater visibility impairment.

Particles (aerosols composed of sulfate and nitrate as well as other chemicals) tend to dominate light extinction, except under extremely clean conditions when natural light scattering by air molecules predominates. Fine particles (diameter less than $2.5\mu\text{m}$) must be distinguished from coarse particles (diameter greater than $2.5\mu\text{m}$), because fine particles usually dominate visibility effects. Aerosols are basically composed of five types of chemical species: sulfates, organics, elemental carbon, ammonium nitrate, and soil dust. It is important to note that hygroscopic particles (those that absorb water, such as sulfates, ammonium nitrate, and some organics) include particle-bound water in amounts that vary with the relative humidity and can significantly add to the impairment of visibility under humid conditions (relative humidity greater than 80%).

-----Text Box Ends-----

Trend Analysis

Trends analysis has been conducted at the two IMPROVE monitoring locations with relatively long-term sulfate aerosol concentration data: the Shenandoah and Great Smoky Mountains National Parks. All 13 sites with three years or more of data were examined to assess short-term changes in concentrations that may be caused in part by emission changes. Spatial variations in short-term changes in sulfate concentrations were examined, but could not be interpreted without corresponding spatially resolved total SO_2 emissions data (including both utility and nonutility), which were not readily available. Seasonal sulfate and sulfur aerosol concentration values were used to identify seasonal patterns. Corresponding seasonal total SO_2 emissions values (current and historical) are needed to better interpret these patterns, but such data were not available in time for this report.

Long-term Trends. An examination was conducted on nearly 10 years of fine particulate sulfur concentration data (1983–1992) from 12 IMPROVE sites, two of which were in the East (Eldred and Cahill, 1994). One of the more surprising results of this analysis was an average annual increase in concentrations of 4% at the Shenandoah and Great Smoky Mountains sites. Both of

these sites had small decreasing trends during the winter and moderate increasing trends during the spring and fall. The overall annual fine particulate sulfate concentration increased by an average of 2–3% at the two eastern sites. What makes these trends unusual is the decreasing SO₂ emissions in the eastern United States during this same period. Trends at sites in other U.S. regions seemed better correlated with SO₂ emission change expectations. These included sites in the southwestern United States with decreasing fine particle sulfur changes in areas where copper smelter and coal-fired utility emissions were reduced. The researchers did not discuss why fine particulate sulfur was increasing at the two eastern sites.

In a more recent paper, a longer period of record was examined (1983 to summer 1995), and focused on possible explanations of the fine particulate sulfur trends in the eastern United States, especially at the Great Smoky Mountains monitoring site (Cahill et al., 1996). The possible effects of the changes in sampling protocols between two monitoring programs were discussed as well as the occasionally large unexplained discrepancies between paired sulfate ion and elemental sulfur measurements that occur in the summer under very humid conditions. The discrepancies were highest in the summers of 1993 and 1994, with mean sulfate concentrations that were 21% and 31% higher than the mean sulfur concentrations at the Great Smoky Mountains site. The discrepancy did not appear in the summer of 1995 and remains a subject of further investigation. The authors make the case that these issues do not significantly affect their sulfur trend conclusions.

The effect of including an additional 2.5 years of data was to reduce the summer and winter trends, especially at the Great Smoky Mountain site. At that site the three additional summers (1993, 1994 and 1995) lowered the summer trend from +3.9% ($\pm 2.4\%$) per year to +2.1% ($\pm 1.3\%$) per year, and the two additional winters (1993 and 1994) changed the winter trend from -2.6% ($\pm 3.1\%$) per year to -0.9% ($\pm 1.7\%$) per year. By calculating summer trends separately for the earlier half of the data record (1984–1990) and the later half (1990–1995), the authors showed that the 12-year trend of increasing fine particulate sulfate resulted from large increases in the first half of the period followed by either flat trends (Shenandoah -0.1% ($\pm 1.7\%$) per year) or improving trends (Great Smoky Mountains -3.8% ($\pm 2.8\%$) per year) during the second half.

Figure 10 shows the winter and summer averaged fine particulate sulfur concentrations for the Shenandoah and Great Smoky Mountains, and a composite of the 10 western IMPROVE sites. Much of the year-to-year variations in the seasonal averages are thought to result from interannual meteorological effects on production, dispersion, and deposition of particulate sulfate from gaseous SO₂ and sulfate concentrations, since the emissions are thought to be less variable. The changes in seasonal averages were caused principally by changes in the highest concentration days since the low- and mid-level sulfate concentrations were nearly the same year to year.

The researchers suggest a few reasons for the apparent inconsistency between increasing fine particulate sulfur and decreasing national annual average SO₂ emissions of about 1% from 1985 to 1994 (Cahill et al., 1996). One plausible explanation is the increasing use of electricity for air conditioning. SO₂ emissions may be increasing over the years during the hottest days, and yet may not affect the overall reductions in the annual average SO₂ emissions. Hot days are thought to be associated with more rapid production of fine particulate sulfate from SO₂ and with greater pollution buildup caused by stagnant air motion compared to colder days. Therefore, any increment of additional SO₂ emissions on hot days may produce much more fine particulate sulfate than the same increment on other days. This explanation is consistent with the seasonal trends information developed by the authors, but should be tested further with temporally resolved SO₂ emissions; such data were not readily available for this report.

A second possible explanation given by the researchers involves the spatial distribution and influence range of SO₂ emissions. The trends in the 11-year annual fine sulfur concentrations at

the Great Smoky Mountains monitoring site were comparable to the annual SO₂ emission rates for the Tennessee Valley Authority power plants that dominate SO₂ emissions in the near-upwind region of that site. During the important high sulfur concentration periods, it is reasonable to expect stagnant air conditions and a greater than normal influence by nearby sources compared to other periods. This explanation also requires further testing with spatially and temporally resolved SO₂ emissions data when they are available.

Short-term Trends. Additional analysis using the IMPROVE aerosol data was conducted as a case study specifically for this report. The analysis involved calculating the ratio of the average seasonal sulfate concentrations in 1995 to the average seasonal sulfate concentrations for 1993 and 1994 combined. Restricting the analysis to these three years enabled the inclusion of data from all 13 eastern IMPROVE monitoring sites. This improved spatial resolution will ultimately allow comparisons to local SO₂ emission levels so that analysis can be conducted similar to those stated above for the Great Smoky Mountains site (Cahill et al., 1996). (Seasonal emissions data were not available prior to preparation of this report.)

A serious shortcoming of this analysis is that interannual variations in meteorology can easily overwhelm the effects caused by minor changes in emissions. **Figure 10** illustrates that substantial year-to-year variations are common. Ultimately, this difficulty may be mitigated to some extent if SO₂ emissions near some of the monitoring sites are substantially changed and if the speculation by the researchers for the Great Smoky Mountain site is correct that seasonal trends are dominated by the activity of nearby SO₂ emission sources (Cahill et al., 1996). No attempt is made in this analysis to sort out the confounding effects of interannual meteorological variations.

Annual averaged SO₂ emissions for the eastern United States were considerably reduced between 1993 and 1995 (see Figure 3). The use of sulfate aerosol data from the 13 monitoring sites provide much better spatial coverage than the three sites with longer periods of record. Sulfate ion data were used for this analysis instead of elemental sulfur data, as was used in the long-term trend analyses. Though not yet completely resolved, current thinking by those responsible for the IMPROVE aerosol monitoring program is that the sulfate ion values are likely to be more correct for those very humid periods in summer when there is a discrepancy between measured sulfate and sulfur.

The more northerly sites showed a greater reduction in fine particulate sulfate concentrations in 1995 (compared to the two previous years) than the more southerly sites (**Figure 30**). This pattern is not as clear in the seasonal patterns (**Figure 31**). An average of the ratios across sites indicates a network reduction of fine particulate sulfate in 1995 compared to the previous two years of about 24% in winter, 13% in spring, 10% in summer, and 18% in fall.

The overall conclusion that fine particulate sulfate concentrations at the eastern IMPROVE sites were about 15% lower in 1995 compared to the previous two years is sensitive to having selected the sulfate ion instead of elemental sulfur. Had elemental sulfur been selected, the 1995 concentrations would have been only 5% lower than the previous years.

A 15% decrease in one year could have been caused solely by interannual variations in meteorology. For instance, between 1989 and 1990, annual sulfate at the Great Smoky Mountains site increased by about 25%, while emissions near that site increased by less than about 12%, and overall SO₂ emissions in the East were slowly decreasing (Cahill et al., 1996). It seems less likely that such a large change would occur at a 13 site average across the East. Seasonal averaged data predating 1993 indicate that 1995 is consistently one of the lowest sulfate concentration years on record. Given consistency across space (13 sites in the East) and time, it is reasonable to hypothesize that a reduction in sulfate concentrations is beginning to emerge as a result of Title IV SO₂ reductions. However, a longer data record is needed to quantify the magnitude of this signal.

1 The selection of 1995 as the year to compare with previous years was due to 1995 being the first
2 year of implementation of Title IV, with accompanying significant decreases in SO₂ emissions
3 from the previous years. However, this is a poor approach for examining the effects of emission
4 changes on visibility. The timing and location of emission rate changes were not available when
5 this analysis was conducted. When these data are fully aggregated, a more appropriate method
6 would be to identify reasonable source influence regions for each of the monitoring sites by season
7 and to examine relationships between emissions within those regions and sulfate concentrations.
8 Another important approach to understanding the relationship between emissions and local
9 visibility is to examine the extent to which emissions are attributed to local air pollution emissions
10 or from regional transport of air pollutants.

11 The visibility change that would be predicted to result from an annual eastern sulfate reduction of
12 15% is about 8% change in light extinction, which corresponds to a change of about 0.8 deciview.
13 This magnitude of improvement in visibility would probably not be perceptible to the casual
14 observer. However, this change on average, involves some time periods and locations with much
15 less change and others with much greater change. From this perspective, it is likely that the
16 contribution of sulfates to haziness was noticeably lower in 1995.

17 Visibility was directly measured at seven of the 13 eastern IMPROVE sites. Comparisons of the
18 1995 annual averages to the previous two years – in a manner similar to that described above for
19 sulfate concentrations – resulted in no overall change (less than a 1% decrease or improvement).
20 The results ranged from 15% improved visibility (1.4 deciview change) at Great Smoky Mountains
21 to 9% decreased visibility (0.9 deciview change) at Mammoth Cave, Kentucky. A lack of
22 correspondence between these visibility changes and the changes in sulfate contributions to light
23 extinction is not surprising given that more than 40% of the contributions to light extinction are
24 related to non-sulfate parameters, which were primarily unchanged.

25 The nitrate contribution to visibility levels is relatively small, except during the winter (see **Figure**
26 **29**). Using the same technique as described above for sulfate, analysis of short-term trends in
27 nitrate concentrations resulted in no significant change (less than 1%) in the 13 site annual or
28 winter seasonal averaged nitrates for 1995 compared to the previous two years. Annual nitrate
29 trends at individual sites varied from an increase of 18% to a decrease of 22%, with no apparent
30 geographic pattern. Site-to-site variations of the nitrate trends were even greater during the winter,
31 ranging from a 47% increase to a 33% decrease. These variations are likely to result principally
32 from interannual meteorological variations, including temperature, which is critical to determining
33 the nitrate partitioning between particle and gas phases in the atmosphere.

34 Dose-Response Relationships

35 There are two approaches to determining visibility dose-response relationships. One is a
36 technically rigorous but complex situation-specific method, while the other is a simplified approach
37 using broad assumptions that can be more generally applied.

38 In the more rigorous approach, aerosol water growth relationships and Mie theory are applied to
39 determine the optical characteristics of aerosol from information on the composition, shape, and
40 number of particles as a function of the particle size and the relative humidity of the atmosphere.
41 Sophisticated radiative transfer models using the aerosol optical characteristics, the lighting and
42 scene characteristics, and the spatial distribution of the pollutants are used to calculate the path and
43 wavelength of image forming and nonimage forming light that reaches a specific observer at a
44 specific time and date from all points in the scene being viewed. Computer image processing
45 methodology can show the visual effects of any modeled pollution situation on a clear condition
46 photograph of the scene of interest. This technique was used in the *1990 Integrated Assessment*
47 *Report* (NAPAP, 1991) and *State of the Science and Technology Report 24* (Trijonis et al., 1991).

1 While this sort of analysis is useful for assessing specific cases, it is impractical to address regional
2 haze issues, where visibility is experienced in a nearly infinite variety of situations and where the
3 detailed characteristics of the pollution, lighting, and scene conditions are rarely known.

4 The simplified approach uses aerosol species-specific extinction efficiencies with water growth
5 functions to determine the light extinction coefficient of the aerosol from its composition and the
6 relative humidity of the situation of interest. Extinction efficiencies and the relative humidity
7 dependence of the aerosol are based upon typical results using the more sophisticated methods
8 indicated above. Light extinction coefficients determined by this approach are easily converted to
9 the deciview scale of the haziness index. This index was specifically designed so that anywhere
10 along its scale, haziness changes that are about equally perceptible correspond to the same deciview
11 difference (see text box for discussion of deciview units). For example, a 3 deciview difference
12 caused by changed air quality should result in about the same perceived change in haziness,
13 whether under pristine or highly polluted conditions. This characteristic of the deciview scale
14 requires that the scene being viewed have a sufficient number of sensitive scenic features for the
15 baseline haze level. No one scene is likely to have such scenic features for all conceivable haze
16 levels. However, the nearly infinite variety of scenes available where regional hazes are concerned
17 ensures that many will have the desired characteristic for any haze level.

18 Based upon viewing scenic photographs with computer generated hazes, changes as small as about
19 0.5 deciview can be visually detected if a sensitive scenic feature is prominent in the scene. A 1
20 deciview change is more often detected because the requirement for sensitive scenic feature is less
21 restrictive; however, such a change is subtle. One must bear in mind when considering how much
22 of a change is perceptible that visibility is an effect that is instantaneously manifested, while
23 assessment results generally are in terms of temporal averages. It is important to note again that a
24 seasonal change of 1 deciview is likely to include days with no change and other days with
25 prominent changes. A similar argument can be made for spatial averaging. Predicted regional
26 averaged visibility changes may seem small, while areas within the region and near sources with
27 changed emissions can experience much more noticeable changes in haze levels.

Human Health

- At the end of the 1980s, NAPAP reported that the human health benefits resulting from reductions of atmospherically formed particles and ozone that accompany the control of SO₂ and NO_x emissions were considered to be potentially significant, but very uncertain. Current scientific knowledge about the health effects of particulate matter indicates that the extent of such benefits may be substantial.

Sulfate and nitrate particles constitute a portion of particulate matter that varies geographically and that consists primarily of fine particles. Although no consensus exists on the causal relationship between sulfates and nitrates and adverse human health effects, many epidemiological studies suggest a statistical association between various pollutants and demand for hospital services, and mortality and morbidity rates.

Epidemiological studies evaluate the association between air pollution and adverse human health effects, typically in an urban setting. Statistical techniques (usually a type of regression analysis) are used to estimate dose-response relationships between pollution levels and health effects. These types of studies are useful in that they allow the estimation of the incidence of health effects related to varying levels of air pollution. However, epidemiological studies alone have significant limitations when trying to prove the existence of a cause-and-effect relationship between pollutant dose and a health response.

Reductions in levels of fine particulate matter represent a potential source of major health benefits as a result of Title IV's reductions in SO₂ and NO_x emissions. SO₂ and NO_x emissions interact with other chemical species in the atmosphere to form secondary sulfates and nitrates, which comprise a significant portion of atmospheric fine particulate matter. In the eastern United States, which is largely urban and where the impact of Title IV emission reductions will be greatest, sulfate aerosols represent 30-40% of average ambient levels of fine particulates. In the West, sulfate constitutes only 10% of fine particulate matter, whereas nitrate reportedly constitutes 15-20% (U.S. EPA 1996a) and, because of inaccuracies in measurement techniques, nitrate levels may be significantly underestimated.

Determining the specific effect of sulfate aerosols on human health is difficult because epidemiological studies are limited in their ability to separate the effects of sulfates from the effects of other components of fine particulate matter (U.S. EPA, 1995). Many studies of human health benefits from emission reductions are based upon epidemiological findings that are focused on health effects from exposure to particulate matter or fine particulate matter. However, a growing number of studies are focusing on the health effects of exposure to specifically sulfates (U.S. EPA, 1995).

The epidemiological studies indicate that some or all of the constituents of fine particulate matter (including sulfates and nitrates) are harmful to human health. Laboratory studies suggest that some types of sulfates, especially sulfuric acid aerosols, are harmful to the respiratory system when subjects are exposed to sulfates alone. This evidence suggests that sulfates may be contributing, at least in part, to the health effects observed in association with fine particulate matter (U.S. EPA, 1995).

Some of the epidemiological studies of the health effects of sulfates, specifically and particulate matter, in general, have examined the extent to which other factors – such as temperature, humidity, ambient levels of other pollutants, smoking, and time spent indoors – may confound the suggestion that the observed health effects are associated with Title IV pollutants. One reanalysis of particle epidemiology concluded that many of these studies correctly controlled for confounding factors, which supports the reported statistical association between levels of fine particulate matter

and health effects (Samet et al., 1997). However, this issue remains unresolved. A recent critical review of relevant epidemiological studies proposes that a case can still be made that some uncontrolled meteorological factor may be acting as a confounding factor and may thereby provide a plausible explanation for the apparent particle-related health effects (Vedal, 1997).

The Clean Air Act establishes the mechanism for National Ambient Air Quality Standards for particulate matter with the goal of protecting public health. EPA is required to periodically review and revise this standard, as appropriate, to primarily protect against adverse health effects, and secondarily to protect against welfare effects (e.g., impacts on vegetation, crops, ecosystems, visibility, materials, etc.). EPA recently completed a comprehensive review of the significant new body of epidemiological evidence, which can be found in the Particulate Matter Criteria Document (U.S. EPA, 1996a). Based on its review of the scientific criteria and standards, EPA recently revised the primary (health-based) and secondary (welfare-based) National Ambient Air Quality Standards for particulate matter, retaining the previous standards for particles 10 microns or less in size (PM₁₀) with some revisions to the form of the 24-hour standard, and adding new daily and annual standards for fine particles (particles 2.5 microns or less in size - PM_{2.5}). EPA estimates that the impact of the new standards could range from minimal in areas that already meet them to a substantial (45-50%) reduction in particulate matter associated morbidity and mortality in areas with high levels of particulate matter (U.S. EPA 1996b).

Members of the Clean Air Scientific Advisory Committee, an independent scientific review panel, reached a consensus that a new National Ambient Air Quality Standard should be established for fine particles. However, there was no consensus on the level, averaging time, or form of a fine particulate standard. The panel members' diversity of opinion reflects the many unanswered questions and uncertainties associated with establishing a cause-and-effect relationship between fine particulate matter and premature mortality (CASAC, 1996). There is also a lack of consensus on the existence of a level below which there will be no adverse health effects. However, there is a general consensus that a reduction in fine particles as a result of the Clean Air Act Amendments of 1990 will have a positive impact on public health.

Observed Effects

- Based on epidemiological studies, reductions in fine particulate matter (including sulfates and nitrates) as a result of the 1990 Clean Air Act Amendments are likely to improve public health.

Epidemiological studies indicate that potential health benefits from decreased SO₂ and NO_x emissions – and thereby decreased levels of fine particulate matter (assuming other fine particulate matter constituents do not change in concentration) – include reductions in: premature mortality from cardiorespiratory disease, particularly in the elderly (U.S. EPA, 1996a and 1996b); hospital admissions for respiratory and cardiovascular causes (Burnett et al., 1995; U.S. EPA 1996a); asthma attacks and medication use; respiratory symptoms; and school-loss and work-loss days. Whether such reductions will have a significant impact on rate of occurrence of chronic obstructive pulmonary disease, which includes chronic bronchitis and emphysema, is less clear. Air pollution has a limited role, if any, as a cause of asthma, but it clearly aggravates existing asthma. The association of respiratory emergency visits with air pollutants – specifically sulfates (winter), ozone (summer), and particulate matter (year-round) – even at very low levels, suggests that further reductions in these pollutants will lead to fewer asthma exacerbations (Bates, 1995).

Indications from epidemiology studies of a causal relationship between ambient sulfate and nitrate particles with human health end points suggest that Title IV's decreased emissions should lead to a reduction in premature mortality from cardiovascular and respiratory causes and fewer asthma symptom days. Using available epidemiological evidence, an EPA study estimated the annual

what was
baseline ÷
the study?

number of cases in 1997 of sulfate related health effects that would be prevented under Title IV (U.S. EPA, 1995). The study assumes that sulfates play a causative role in producing adverse health effects and estimates the following cases could be prevented (20th to 80th percentile range): 400 to 5,700 cases of premature mortality; 1,600 to 6,600 new cases of chronic bronchitis; 500 to 870 cardiac hospital admissions and; 800,000 to 2,370,000 asthma symptom days. EPA is currently conducting a cost-benefit analysis of all the 1990 Clean Air Act Amendments (not just Title IV), which will also include benefit projections for additional pollutants controlled by the Amendments.

Human health benefits are also associated with decreased concentrations of gaseous SO₂, gaseous NO₂ (including the associated changes in ozone levels), and air toxics (reductions of these are associated with reduced SO₂ and NO_x emissions). While the potential health benefits of reductions in air toxics are not well understood and the effects of NO₂ may be minimal, the effects of SO₂ and ozone (ozone levels are related to NO_x emissions) have been studied extensively. Increased ozone levels are associated with increased hospital admissions for asthma, increased asthma attacks, and increased medication use (U.S. EPA, 1996d). Asthma attacks are more likely to occur near point sources of gaseous ambient SO₂. Every year, 1-3% of asthmatics living near a pollution source will experience respiratory symptoms and changes in lung function as a result of SO₂ exposure (Freudenthal et al., 1989).

Sensitive Populations

- For this assessment, the sensitive populations are children, especially those with asthma, and the elderly, especially those with cardiorespiratory disease who may be more likely to die prematurely during episodic pollution events.

Individuals most at risk to the acute effects of sulfates and other fine particles are typically those with preexisting conditions (as a result of current or previous illness), genetics (inherited characteristics), age, or exposure. To be at risk, an individual must be exposed to particulates including sulfates and nitrates. Exposure to ambient airborne pollutants is generally increased by being active and being outdoors where concentrations are typically higher (increased breathing rates and increased exposure). Fine particles outdoors efficiently penetrate indoors where they contribute to human health effects. This makes indoor exposures important because of the large percentage of time spent indoors, despite the somewhat lower activity levels.

Among the diseases that are exacerbated by particulate air pollution are chronic obstructive pulmonary disease (mainly chronic bronchitis and emphysema), asthma, heart disease, and respiratory infections (acute bronchitis, pneumonia). Children and elderly adults may also be at increased risk for these diseases—children, because of their greater outdoor activity levels, and the elderly, because of their relatively poorer average health status. Effects may not be limited to these groups; new and ongoing studies indicate a potential statistical association between exposure of healthy people to fine particulates and potential shortening of their lives (Pope et al., 1995).

Concentration-Response Relationships

- Concentration-response information is available for numerous health end points, including premature mortality, hospitalization for respiratory causes, and respiratory symptoms.

Concentration-response functions have been adapted from the epidemiological literature (U.S. EPA, 1995), although it is still not known whether a threshold value for particulate matter exists. Changes in the number of incidences of health effects predicted as a result of reductions in particulate matter reductions can be estimated using these functions. Recent studies have identified concentration-response functions for sulfates, for all particulate matter, and for fine particulate

1 matter (U.S. EPA, 1996a and 1996b). Due to limited scientific literature addressing the health
2 impacts of nitrates, exposure currently is analyzed as exposure to fine particulate matter. Many
3 epidemiological studies suggest an association between both mortality and morbidity and exposure
4 to ambient fine particulate matter, but specific biological mechanisms have not yet been identified.
5 Studies in laboratory animals are beginning to provide preliminary evidence of plausible biological
6 mechanisms (Godleski et al., 1997). Definitive results, however, are not expected for several
7 years.

What are the economic benefits related to the effects areas?

- Economic methods for valuing the effects of pollution on marketed goods and services have been available for many years. The ability to estimate benefits for these effects is limited by the availability of economic and scientific data.
- In contrast, methods for estimating nonmarket values and passive-use values have only recently become widely used and accepted. The reliability of economic estimates for specific environmental effects from Title IV is greatest for the health benefits although significant uncertainties characterize the estimates. Estimation of recreational damages is less reliable because the economic studies are lacking, there are gaps in the science, and there is a lack of baseline recreation participation information specific to reference environments of interest.

The 1990 Clean Air Act Amendments directs NAPAP to coordinate federal agencies' efforts to analyze the costs and benefits of Title IV controls. Although NAPAP's *1990 Integrated Assessment Report* (NAPAP, 1991) identified a large number of potential economic benefits, three obstacles limited a quantitative assessment: (1) linkage to end points that directly affect welfare from an individual's perspective, that is, a determination of the quantitative effect of emission reductions on an item that is valued by individuals, was often poorly established; (2) inconsistent levels of rigor and reliability characterized the economic methods that were available for valuation; and (3) even where end points and methods for valuation were well developed, poor scientific or economic data often undermined the ability to estimate potential benefits. Since 1990, there have been advances in science and economics that make estimates of economic benefits more feasible. In some cases, even though estimates remain uncertain, more is known about the range of potential benefits and about the contribution that additional research could make to reducing uncertainty.

This assessment of economic benefits addresses the following areas: human mortality, human morbidity, visibility, materials and cultural resources, ecosystem health (nonuse values), aquatics (recreation), forests (recreation), agriculture and commercial forestry, and radiative forcing. For this assessment, NAPAP asked the following questions about each of the areas.

1. Does science provide the information needed for economic analysis? Are benefit end points well established?
2. Are economic methods adequately developed?
3. Are data available from science and economics for an assessment of benefits?
4. Are the expected benefits large?
5. With the goal of improving benefit estimates, what is the relative short-term return on investment for research?

A matrix was developed to display a qualitative, relative ranking for each benefit area with respect to these questions (**Table 7**). The major conclusions of this benefits assessment are summarized in the text below followed by a discussion of the rankings assigned each area.

Table 7. Qualitative, relative rankings for benefit areas with respect to the primary assessment questions of interest to NAPAP.

Categories ● high ◐ high-mid ◑ mid ◒ low-mid ○ low	<u>1. Link Between Science and Economics:</u> Are benefit endpoints well established? Does science provide information needed for economic analysis?	<u>2. Economic Methods:</u> Are economic methods adequately developed?	<u>3. Data Availability:</u> Is data available from science and from economics for an assessment of benefits?	<u>4. Expected Benefit:</u> Are expected benefits large?	<u>5. Value of Additional Information:</u> With the goal of improving benefit estimates, what is the relative short-term return on investment?
Health: Mortality	◐	◐	◐	●	●
Health: Morbidity	◐	◐	◐	◐	◐
Visibility	◐	◑	◒	◐	◐
Materials and Cultural Resources	◒	◑	○	◐	◐
Nonuse Values: Ecosystem Health	◒	◒	◒	●	◑
Aquatics: Recreation	◐	●	◒	◒	◑
Forests: Recreation	◒	◐	○	◒	◑
Agriculture and Commercial Forestry	◐	●	◑	◑	◒
Radiative Forcing	◒	○	○	◒	○

Benefits Estimation: Approaches and Conclusions

Benefits estimation provides a means to assess the values people place on the expected effects of Title IV. From an economic perspective, values are defined in relative terms and are measured by how much of one asset or service people are willing to sacrifice to obtain or preserve another. Economists refer to this as an *opportunity cost approach* to valuation.

Values are expressed in monetary terms, although in principle they can be expressed in other metrics. The methodologies used to obtain a monetary measure of value depend on the type of information available. For example, market goods have observable market prices and quantities from which estimates of value can be obtained. However, nonmarket goods and services and nonuse values must employ alternative techniques.

Table 7 displays a qualitative, relative ranking for each benefit area with respect to the primary assessment questions of interest to NAPAP. The qualitative rankings are ordinal; for instance, the highest ranking is better than the second highest, but should not be interpreted as twice as high. The rankings are primarily of interest relative to each other. They should be viewed both as rankings among benefit areas for a given question (within a column), and among questions for a given benefit area (within a row). Even in cases where the highest ranking is indicated, there still remains considerable room for improvement.

Market Goods and Services

The value or opportunity cost of goods and services that are readily traded in markets is reflected in their prices. Benefits from emission reductions to agriculture and commercial forestry fall in this category. However, the benefit in these areas is not simply the change in crop yield multiplied by crop price. Rather, it is the value of the change in yield compared with the next best use of the resources used to produce the crop. The opportunity to substitute among crops if the yield of one crop falls substantially provides a bound on the magnitude of the economic impact on agriculture. Also, if effects are widespread product prices will change as will consumer demand, which also must be taken into account.

The effect on materials is another benefit area that includes marketed goods and services. Again, the benefit in this area is not, for example, the fractional change in the physical lifetime of a material multiplied by its original cost. Rather, it is the lost economic use of the asset. Material assets often have shorter economic lives than physical lives, as is evident when a structure is torn down while it is still fully functional in order to construct a new one. The economic depreciation of material assets tends to lower the estimate of benefits to material resources from that which would occur if those resources were valued at their original cost less their physical depreciation. In other cases, the useful service life of an asset, such as an automobile, may be affected only slightly by damage to paint finish, but its economic value at each age may be greatly affected. The economic methods for valuing the effects of Title IV on marketed goods and services have been available for many years. The ability to estimate benefits is limited by the availability of economic and scientific data.

Nonmarket Goods and Services

The valuation of goods and services not traded in markets can be sensitive to the geographic location of impacts and cannot be observed directly from prices. Because the effects of Title IV are so widespread, it is prohibitively expensive to measure values with individual studies targeted at all the specific locations and effects. One approach to comprehensively measuring value in these situations is termed *benefit transfer*. Information from studies that value changes at specific geographic locations is "transferred" to estimate benefits at other locations.

The rigor and validity of benefit transfer vary with the types of benefits under consideration. They are central features in the assessment of the reliability of benefit estimates. A simplistic approach is the use of a uniform value for a day of recreational activity (unit-day value) calculated at a specific site and applied at various other sites with a wide range of characteristics. But estimates are unreliable because sites vary greatly, and uniform values for a day of recreation activity cannot be used to value a change in the quality of a recreational experience. A more sophisticated and preferable approach would be to identify the attributes of the sites that have been studied previously and to estimate the individuals' *willingness to pay* for these attributes at these and other locations.

Of the various effects of Title IV, impacts on aquatic and forest recreation are the most problematic with regard to transferring benefit estimates. The value of environmental assets has generally been calculated with unit-day values, but numerous values exist for all types of uses and environments (Walsh et al., 1990). Accounting for regional factors (such as the range and quality of substitute sites) and site-specific factors (such as congestion) is difficult, and there are no acceptable procedures for determining the spatial extent of the market. The methods for determining the size of the population that would be or is affected by a change in recreation quality or quantity are still being debated.

The most reliable valuation of environmental assets affected by Title IV is for aquatic recreation in the geographically specific but important Adirondack Park (New York). The seminal paper in this area (Englin et al., 1991) contains estimates for the value changes in the catch rate for trout and other fish species, given their relative sensitivity to lake acidity. The study also addresses the availability of substitute sites for aquatic recreation and an individual's willingness to pay to avoid degradation at sites affected by Title IV. However, the literature in general is weak on the issue of participation rates, assuming effectively that the number of anglers as a percentage of the population remains constant. Further, the valuation of benefits to lakes outside the northeastern region will require sophisticated benefit transfer.

Valuation of the benefits of Title IV for forest recreation poses the same potential problems as valuation of benefits for aquatic recreation. Forest recreation depends on the availability of substitute recreational activities, which has not been successfully modeled previously. Furthermore, there is no individual setting where credible estimates have been developed.

The area that affords the most credible benefit transfers is human morbidity. Once atmospheric or other natural processes are taken into account (e.g., in estimating the effect of reduced emissions on ambient air quality), the health effects and the values people place on avoiding them are assumed to be reasonably similar across locations in the United States, especially if the transfer occurs between populations with similar health status and life-styles. Applying unit values for health effects at all sites affected by Title IV is less problematic than applying unit-day values to specific environmental changes, because of the presumption that willingness to pay to avoid health effects is less influenced by region and site variables than willingness to pay for recreation. This is one reason why relatively good progress has been made in health benefits analysis.

The valuation of risks of human mortality depends on estimates of the willingness to pay of individuals for small changes in the risk of premature death. Most original studies in this area address accidental deaths in prime-age adults—a setting inappropriate for most types of environmental mortality risks, except perhaps accidental toxic waste releases and similar catastrophes. Hence, they are not appropriate to transfer to the valuation of health benefits from environmental changes. One study (Mitchell and Carson, 1986) addresses the latency issue that is important for valuing deaths due to cancer, but it does not consider the effect of prior health status and age on valuation, which influences rates of environment-related deaths, such as those from heart disease and respiratory failure. Few reliable, peer-reviewed studies are available to value life-years saved, and none do so for individuals with impaired health, even though this health end point can be estimated by health scientists. Nonetheless, the weight of evidence is so great regarding the

1 presence of mortality effects that in sensitivity analysis using a wide range of parameters for the
2 value of a statistical life, relatively large benefit estimates are consistently obtained.

3 Benefit transfer for valuing visibility presents formidable challenges because of the sensitivity of
4 values to regional, site, and personal characteristics. While visual range can be characterized in a
5 relatively straightforward way, the vista being affected is difficult to characterize in generic or
6 transferable terms beyond such labels as "urban," "rural," and "recreational area," which are not
7 likely to be sufficient. Despite these challenges, the literature on visibility benefits is fairly
8 conducive to benefit transfer. Several studies of visibility values in multiple cities (McClelland et
9 al., 1991; Tolley et al., 1986; Brookeshire et al., 1979) permit examination of city-specific factors
10 affecting values and derivation of functional relationships to predict willingness to pay given the
11 baseline visual range and the size of the change (NAPAP, 1989). There is a somewhat thinner
12 base upon which to construct estimates of benefits for changes in visibility at recreational sites
13 (Chestnut and Rowe, 1990; Electric Power Research Institute and Decision Focus, 1991). A
14 recent study estimates a significant annual value of total benefits of Title IV in 2010 for both (1)
15 residents of the eastern United States and (2) national parks (Chestnut and Dennis, 1997).

16 The major problem with calculating benefits for visibility is not the reliability of transferring
17 estimates from original studies to other study sites. Rather, it is the reliability of the original
18 studies. Often the valuation of visibility has relied on *stated preference methods*, or *contingent*
19 *valuation*, specifically. Contingent valuation techniques rely on information revealed through
20 choices made by individuals in a hypothetical setting, rather than information revealed through
21 actions in a real economic setting. Significant debates surround protocols for eliciting values in
22 contingent valuation studies. For example, the size of photographs shown to respondents appears
23 to influence willingness to pay. There are also concerns about joint valuation of visibility and
24 health (i.e., that reported values for visibility improvements are confounded by concerns about
25 health effects) and embedding (i.e., identification of the proper geographic scope for valuation).

26 Research on recreational values has concentrated on national parks in the Southwest. This focus
27 has the advantage that health and visibility may not be confounded, but provides little information
28 about changes in visibility at eastern national parks affected by Title IV. Nonetheless, these
29 existing studies indicate that benefits from improvements in visibility at recreational sites can be
30 substantial. In addition, previous studies indicate the valuation of visibility effects in residential
31 areas is likely to be as important as changes in visibility at national parks because these are use
32 values centered where people spend their daily lives.

33 Nonuse Values

34 Up to this point, *commodity values* have been used to describe the valuation of benefits to
35 marketed goods and services, and *use values* have described the value of nonmarketed goods and
36 services that individuals use or enjoy directly. Another potentially important category of values is
37 designated *nonuse values*. This value stems from an individual's willingness to pay to avoid
38 damages to environmental assets, even if the individual has no expectation of directly enjoying or
39 using that asset. Nonuse values are now widely accepted as a legitimate category of benefits
40 estimation, but estimating such values is highly controversial.

41 The most important area of potential benefits that would be characterized by nonuse values is
42 ecosystem health. By its nature, this area describes effects that occur over long time frames and
43 broad geographic areas, and that are the product of interactions between the effects of Title IV and
44 other impacts on the environment. People may hold significant nonuse values for the stability of
45 ecological systems, in addition to the use values for aquatic and forest recreation in these systems.
46 Some progress has been made in the last few years, in particular, in the valuation of estuaries.
47 These studies indicate economically significant nonuse values for environmental assets, but a range
48 of potential benefits from Title IV cannot yet be determined. Clearly, the literature on nonuse

values for environmental assets cannot yet support benefit transfers, because most of the studies concern nonmarginal changes in unique environments (e.g., species extinction, loss of an ecosystem).

Another area that would be characterized in large part by nonuse values is the potential benefits of Title IV emission reductions to cultural resources, including historic monuments and buildings, gravestones, and statues. Various studies in Europe have found very high estimates in this benefit area. A recent study indicates that important economic benefits measured by willingness to pay may result from the impacts of Title IV on national monuments (Morrey et al., 1997). However, the available scientific and economic data are insufficient even to place bounds on these potential benefits. Valuation of cultural materials typically involves summing estimates of an individual's willingness to pay to preserve the attribute of the cultural resource (e.g., a monument) that is being degraded by emissions. This will include the values of those that actually visit the cultural resource as well as the value that nonusers (those that may never visit the monument) may place on preserving the resource (existence value).

In summary, the reliability of economic estimates for specific environmental effects from Title IV is greatest for the human health benefits although significant uncertainties characterize the estimates. Recreation damage estimation is less reliable because the economic studies are lacking, there are gaps in the science, and there is a lack of baseline recreation participation information specific to reference environments of interest. The valuation of aquatic effects for one important region, the Adirondack Park, is fairly reliable. However, estimation of aquatics benefits in other regions will be difficult because of differences in water temperature, chemistry, availability of substitute sites, and the mix of species at other locations. Estimates of visibility damages are weak, partly because of weak methodology (e.g., some are limited to annual averages only) and partly because data on changes in visual range for the majority of affected recreational and residential vistas are of low quality. Methods for valuation of agricultural and commercial forestry and materials are well understood, but the availability of estimates is limited by scientific and economic data. Non-use value estimation studies for ecological systems and cultural resources suggest the value of changes in these assets may be relatively large, but there are almost no studies of marginal changes in the quality or quantity of these assets such as what may likely result from Title IV.

Expected Benefits and Associated Uncertainties

In general, quantifiable benefits are relatively large in the areas of health and visibility – areas where relatively large benefits may not have been anticipated, or that were not the primary motivation for Title IV originally. The magnitude of benefits in these areas compares favorably with the costs of Title IV, independent of benefit estimates for other areas.

While estimates of health and visibility benefits remain uncertain, the cost of reducing this uncertainty appears to be relatively less than reducing uncertainty for many other areas. Benefits to materials and cultural resources may also be sizable; an incremental research effort could provide substantial information about the likely extent of these benefits.

To evaluate Title IV on the basis of a comparison of benefits and costs, it may be sufficient to focus efforts on assessing benefits from health and visibility, because these benefits alone appear to outweigh the costs. Environmental areas, such as aquatics and forests, will also benefit, although quantification of ecosystem benefits is somewhat more uncertain. If future research reveals this to be an erroneous conclusion, then NAPAP would be justified in looking further into potential benefit areas where results may be more elusive. However, if NAPAP research is to be used to determine whether further controls on SO₂ emissions are warranted, then a comparison involving the additional benefits from all benefit categories and the additional costs would be necessary.

-----Text Box Begins-----

Relative Magnitude of Expected Benefits

Two characteristics help to explain the relative magnitude of expected benefits and the level of uncertainty in these estimates, as illustrated under Questions 4 and 5 in Table 7.

1. Expected benefits of Title IV tend to be greatest where opportunities for behavioral change to avoid damages are least.

Human health and visibility share the characteristic that individuals have few options to avoid detrimental changes in end points. An example where individuals may be able to do so is through increased use of air conditioning to avoid poor air quality. Such opportunities are much more evident in the recreational choices people make. For example, angling activity at one lake may be affected by diminished catch per unit of effort, but anglers are likely to respond by moving to other lakes. The willingness to pay is the opportunity cost of the substitution of the first lake, not the total value of activity at a given lake. Similarly, reduced crop yield in agriculture is mitigated by farmers' decisions to substitute different crops. The benefit measure is the opportunity cost of this substitution, not the entire value of reduced crop yield for a given crop.

2. Links between science and economics are strongest where changes in atmospheric concentrations directly affect end points relevant to individuals; they are weakest where ecological systems play a more important role.

In the cases of health, visibility, and agriculture, atmospheric modeling links directly to concentration-response functions with end points (human health, visibility, crop yield) of immediate relevance to individuals. In other benefit areas, the science is more complicated because ecological systems intervene between changes in atmospheric concentrations of pollutants and environmental consequences. One exception to this general rule is materials, where the linkage is weak due to a relative lack of research data.

-----Text Box Ends-----

A somewhat different perspective would be suggested if NAPAP were to consider a longer time frame and a more sustained research effort, with attention focused on possible amendments or extensions to the goals set forth in Title IV. In this case the assessment of nonuse values for ecosystem health should be afforded higher priority. However, a research emphasis in this area would require sustained levels of funding over several years to yield results that would be reliable. Also, agriculture and commercial forestry would receive a somewhat higher ranking were a sustained level of funding to be committed. One reason is that agriculture is undergoing structural change, due to 1996 congressional reforms, which may not be fully attained until the next decade. In addition, the cost of research in this area stems from the need for modeling of changes in ozone, which may be costly and time consuming, though such modeling would also contribute to an understanding of human health benefits and forest recreation.

Qualitative Ranking Summary

Assessment of economic benefits has significant room for improvement in each benefit area. The most pressing needs within each area are identified for the purpose of reducing uncertainty in benefit estimates. Brief explanations are provided for the qualitative rankings given for Questions 1–3 in Table 7. The rankings illustrate the relative strength at each step along the pathway linking environmental impacts and economic benefits. These rankings, coupled with a sense of the relative cost of improving information at each step (which is not considered formally in this document),

provide evidence for prioritizing incremental research to narrow the uncertainty associated with an overall assessment of benefits (Question 5).

Human Mortality Evidence of mortality effects from exposure to air pollutants is convincing, and their magnitude may be large, but major uncertainties remain about the particle sizes and composition affecting death risks. In addition, the traditional approach to estimating such risks and valuing them is problematic. Because health benefits will largely accrue to the older population, economists increasingly recognize that statistical years of life gained is the critical end point, rather than a “body count” of statistical lives lost. However, estimates of changes in life years have not been in standard use in their valuation of health effects (Question 3).

From an economic standpoint, health research is providing adequate concentration-response functions (Question 1), particularly from prospective studies linking cumulative exposures of health populations to premature mortality – through the increased probability of developing and exacerbating chronic illness and the vulnerability of old age. The added role of acute exposures, if any, is less clear. Just as important, the valuation of a statistical life for environmental risks draws on inappropriate estimates from accidental death and occupational risks (Question 2). The methods to calculate more appropriate measures of the willingness to pay for increased life expectancy are being developed, but only one study has been conducted (Johannesson and Johansson, 1996). A life-expectancy approach is likely to yield lower benefits than the conventional approach of valuing statistical lives.

The highest priority for reducing uncertainty in benefit estimates should be placed on human mortality. An improved basis is needed for the valuation of small risks to mortality due to environmental changes. Also, economists need to develop estimates for willingness to pay to avoid these risks that depend on the age and health status of the affected individual. Better estimates of the concentration-response functions and life years saved conditional on the age and health status of individuals are required from the health science and epidemiology disciplines.

Human Morbidity There is strong evidence of health morbidity effects from exposure to air pollutants (Question 1). From an economic perspective, the health science provides adequate concentration-response functions linking air concentrations to acute health effects, in the sense they are consistent and biologically plausible. Good functional relationships are available for medical and other indicators of distress, such as hospitalization and work-loss days for acute illness, and there are reliable, though partial, measures of willingness to pay to avoid these effects (Question 3). Also, literature exists that employs averting-behavior studies to value health morbidity, but they are less convincing (Question 2). The uncertainties in this area could be reduced substantially through additional research. Although, contingent valuation approaches for valuing acute health effects are reasonably well established, there are no recent studies in this area using such approaches. The literature is more ambiguous on the possible link between exposure and some chronic respiratory diseases and heart disease. Also, the willingness-to-pay estimates for chronic illness are less developed.

Visibility A comprehensive analysis of visibility benefits requires better monitoring data and atmospheric modeling. Monitoring data are important because visibility effects are nonlinear (Question 3). The economic methods for valuation include hedonic property valuation and stated preference for residential visibility, and travel cost and stated preference for recreational visibility. However, the economic research has failed to identify end points with adequate precision (Question 2). There is evidence that people care about characteristics of the distribution of visibility effects over seasons, weather conditions, and time of day, but valuation other than mean values has not been explored. Similarly, the atmospheric modeling has not addressed important distributions in time and space (Question 1), and the level of importance of this detail should be determined before additional resources are spent on monitoring and modeling. For moderate resources, additional valuation studies could be implemented to enhance the precision of the end

points that are important for assessing benefits. Particular attention should be paid to the nature of preferences for changes in visibility. This information would provide guidance for modeling of changes in visibility at a more decentralized level in more areas of the country.

Materials and Cultural Resources The main problem in both materials and cultural resource valuation is the lack of a complete inventory of affected assets (Question 3). A second problem is the lack of data about the economic lives of the assets. Further, information on behavioral responses is inadequate (e.g., changes in maintenance or replacement in response to physical deterioration). From an economic perspective, the methods for estimating general materials damage are more straightforward than those for estimating damage to cultural resources (Question 2). The latter requires stated preference methods, which have only recently been applied to cultural resources. Some dose-response functions of the effect of pollution on materials exist for cultural (e.g., monuments), commercial (e.g., buildings, clothes, rubber, automobile paint) and public (e.g., bridges, fences) assets, but there is uncertainty about how well these data describe the exposure and effects outside an experimental setting and how Title IV's controls contribute to reduced damages (Question 1).

Ecosystem Health (Nonuse Values) Ecosystem changes associated with Title IV cannot yet be determined (Question 1). Almost as significant is the lack of a valuation framework for assessing benefits from improvements in ecological indicators, especially given the temporal aspects of ecological dynamics (Question 2). Hence, there are only illustrative studies to serve as an indication of potential benefits (Question 3). Valuation studies should identify the attributes of ecosystems that affect willingness to pay most significantly. This information should then become the focus of ecosystem modeling.

Aquatics (Recreation) With respect to angler activities, the linkage between water chemistry and survival of fish fry is well understood in a laboratory setting, using dose-response methodology (Question 1). Translation of these results to wild populations may be imperfect because mortality as well as reproductive rates may be affected. Other possible contributing factors are not controlled for in laboratory tests. Results – expressed in terms of "acid stress indices" – are species-specific but may not have been carried out for all recreationally targeted species.

Three issues affect angler activities: effects of acid stress indices on angler catch-per-unit-effort; angler valuation for changes in catch per unit of effort; and effect of catch per unit of effort on angler participation rates. Only the first factor is relatively straightforward. The link between science and economics (acid stress indices to catch per unit of effort) is tenuous and limited to a few species (fewer than the number for which there are acid stress indices data). Links between acid stress indices and catch per unit of effort have been estimated only for Adirondack Park. Catch per unit of effort is valued using travel cost data and either of two competing methods: hedonics or random utility. Neither currently accounts for effects of changing fishing conditions on individual decisions to fish, which should be estimated jointly with valuation of catch per unit of effort and site choice. Such models have not yet been applied to freshwater fishing (Question 2). Nonetheless, among areas with nonmarket benefits, the techniques that have been developed in aquatic recreation are the most sophisticated.

With respect to boating and swimming, scientific models of eutrophication with predictions of turbidity (especially related to nitrogen) are being developed. Acidification may improve clarity of the water, thereby benefiting swimming and boating, however, nitrogen deposition may contribute to eutrophication, thereby worsening conditions for swimming and boating. Economic models of the benefits of boating and swimming involve valuation and participation rates. The methods share common characteristics with angler activities. As with angling, there are significant data on swimming and boating behavior, but little data on recruitment (Question 3). The science-to-economics links for affected regions outside the Northeast should be developed more fully. Within

economics, a model of participation decisions is needed, to be estimated jointly with angler valuation.

Forests (Recreation) There is ample evidence of the effects of sulfur, nitrogen and ozone on forests. A key concern is the decline of forest resources, particularly high-elevation trees. However, the link between primary pollutants and effects that people may care most about, such as foliage intensity and range, is not established (Question 1). A variety of confounding factors, such as drought and introduced pests, has made it difficult for forest researchers to quantify the relationship between air pollution levels and forest decline.

Economic methods for valuation of forest recreation draw essentially on the same techniques used in aquatics recreation (Question 2). However, the forest recreation literature is focused too narrowly to be of immediate value, does not provide evidence on the effect of Title IV's controls on forests, and typically has not incorporated substitution opportunities. Science has not established robust links to valuation end points, so data on changes do not exist (Question 3). Valuation data are also not available. Priority should be given to developing a regional model of forest recreation, to account for the range of recreation opportunities, and to develop estimates of willingness to pay to avoid damage to individual, or classes of recreationally related, forest resources. This economic research would help to focus the contribution of additional work in natural science and, especially, to identify end points that would help link science and economics.

Agriculture and Commercial Forestry Of all air pollutants, ambient ozone is likely to cause the most significant crop damage. Dose-response functions for ozone are available for most major agronomic crops and some specialty crops (Question 1). A number of studies have used ozone dose-response functions and standard economic data and models to estimate the monetary losses. However, most studies have failed to adequately account for behavioral responses, such as adaptation of crop varieties or adjustment of input mixes (e.g., using more fertilizer) although economic methods for doing so exist (Question 2). Moreover, data on changes in ozone resulting from Title IV are not currently available (Question 3). NO_x reductions under Title IV are believed to decrease the creation of ozone and damage to agriculture. Regional models of ozone are needed to predict its effects on agriculture and commercial forestry. (These models would also improve health benefit estimates for Title IV.) In addition, industry-level models are needed to predict changes in cropping patterns in response to changes in crop yields, which would mitigate the economic cost of reduced yields.

Radiative Forcing. Atmospheric models predict changes in particulates and their effect on radiative forcing (Question 1). Predictions of local changes in temperature are also available. However, changes in variability and the effect on general circulation in the atmosphere are not known. Economic methods for modeling damages of climate change are very uncertain, so measuring the benefits of avoided change are also uncertain (Question 2). Data for valuation of local effects are not available.

Valuation of Emission Reduction Benefits

- The largest expected benefits stem from human health and visibility, and potentially from ecosystem health. Significant potential benefits also arise from materials, agriculture (through the NO_x/ozone link), and commercial forestry.

EPA has estimated the benefits from reduced risk of human mortality resulting from SO₂ emission reductions under Title IV to be approximately \$1,806 per ton of SO₂ reduced in 1997 to \$3,438 per ton reduced in 2010 (in 1990 dollars) (U.S. EPA, 1995). The difference reflects, in part, population growth and a shift in emission reductions to areas upwind of larger population centers. (Pennsylvania and the Carolinas have larger shares of the emission reductions in 2010 than in 1997

while Ohio and Missouri have smaller shares.) The sensitivity of the EPA estimate has been explored for NAPAP by Resources for the Future (RFF, 1997) using the recently peer-reviewed Tracking and Analysis Framework (TAF) modeling system, which was developed to support NAPAP assessments.

One estimate that was varied in the RFF analysis was EPA's willingness-to-pay estimates of the expected "value of a statistical life" of \$3.2 million per statistical life with a range of \$1.7-7.9 million (1990 dollars). This is an age-weighted distribution, placing greater weight on values relevant to the cohort over 65 years of age than for the general population. The willingness-to-pay approach may overstate the value of excess premature mortality or the death of an extremely ill individual who is expected to die within days or weeks. As noted above, it may also overstate benefits in contrast to a life-expectancy approach.

In the RFF case, the mean of \$3.1 million per statistical life (in 1990 dollars) was used, along with a 90 percent confidence interval of \$1.6-6.0 million. This distribution generally accords with the valuation literature, but is somewhat on the conservative side because less weight was given to the labor market studies relative to the contingent valuation studies, the latter being marginally more appropriate for valuing mortality risks in the environmental health context and also captures age effects (as does the EPA value), based on Jones-Lee et al. (1985). The Jones-Lee study finds that the value of a statistical life for the 65 years and older group is about 75 percent of that of the average (40 year old) participant in their study. Even using the conservative \$3.1 million estimate, the EPA numbers appear to be quite robust. Examples of the value of a statistical life and the value of other health end points in these two studies are reported in **Table 8**. Ozone-related benefits due to Title IV's NO_x controls would be in addition to those health benefits of Title IV's SO₂ controls.

Table 8. Value of a statistical life and other health end points.

Health Effect	Value (in 1990 dollars per case)		
	Section 812 Study (1)	Health Benefits Report (2)	RFF (3)
Mortality	4,800,000*	3,200,000	3,100,000
Chronic Bronchitis	260,000	211,490	223,300
Respiratory Hospital Admission	6,100	12,340	6,650
Cardiac Hospital Admission	8,300	12,340	12,350
Work Loss Day	83	53	other measures of reduced activity days used
Asthma Attack	32	36	31

* Or \$293,000 per statistical life year. Note the Health Benefits Report (U.S. EPA, 1995) and RFF (1997) studies use an age-weighted expected value of statistical life while the Section 812 study is a best estimate not explicitly considering age.

Sources: (1) U.S. EPA, 1997 Table I-2.

(2) U.S. EPA, 1995 p. 5-24. The values are converted from 1994 dollars to 1990 dollars using the Consumer Price Index.

(3) RFF, 1997. Also see Appendix A for information on the Tracking and Analysis Framework (TAF).

The Health Benefits Report (U.S. EPA, 1995) and the RFF analysis (RFF, 1997) estimate the benefits from reduced risk of human health effects resulting from SO₂ emission reductions under Title IV. The 1990 Clean Air Act Amendments also require EPA to estimate the total benefits, including human health, achieved as part of the entire Clean Air Act (1970-1990, with a separate ongoing effort to estimate benefits achieved from 1990 to 2010). The valuation estimates of these health benefits are also listed in Table 8 under the Section 812 Study column. This analysis includes the benefits of reducing many pollutants, in addition to SO₂ emissions.

Using the TAF modeling system with various assumptions, RFF developed median estimates of health mortality benefits that range from \$2,845 to \$3,840 per ton of SO₂ emission reduction in 2010 (RFF, 1997). The lower estimate stems from the use of the Regional Acid Deposition Model to model atmospheric transport and pollutant concentrations in the EPA's 1995 Health Benefits Report, coupled with health epidemiology and valuation coefficients from the EPA's 1996 Regulatory Impact Assessment for the new ozone and particulate matter standard, which were equivalent to those used in the Section 812 Study. The higher estimate is based on the use of the Advanced Statistical Trajectory Regional Air Pollution model for atmospheric modeling and the health epidemiology and valuation coefficients cited above. The Regional Acid Deposition Model yields smaller health effects than the Advanced Statistical Trajectory Regional Air Pollution model; this is partly offset by the Regulatory Impact Assessment valuation numbers, which yield higher values than those used by RFF. Means are above the medians in all cases.

TAF incorporates sophisticated analysis of uncertainty by propagating the measures of uncertainty in demographic, climatological, dose-response, and valuation parameters and functions for all the modeled benefit pathways, resulting in a wide range of possible benefit estimates around the medians that are reported. The 90% confidence interval around TAF's reference case estimate for mortality benefits ranges from \$1,075 to \$15,020 per ton. This reflects the tremendous uncertainty in this estimate. However, the main result from this comparison is to confirm the probable magnitude of benefits relative to cost estimates reported elsewhere in this study. The median value of benefits from reduced risk of human morbidity effects estimated by TAF for 2010 is an additional \$475 per ton of SO₂ emission reduction.

The median estimates of benefits resulting from changes in NO_x emissions in 2010 are predicted by the TAF model to be \$476 per ton for reduced risk of human mortality and \$259 per ton for human morbidity. These estimates include effects from reduced secondary particulate formation, assuming nitrates are as potent as the average particle of particulate matter in affecting human health, but do not include the health effects from changes in ozone concentrations.

As a result of emission reductions, potentially significant benefits may be achieved in the areas of visibility, materials, and agriculture. As preliminary evidence discussed earlier indicates, reasonable estimates of benefits to materials and cultural resources are not available. With regard to visibility, in one study standard visual range with and without Title IV was compared to assess economic benefits of improvements in visibility. Drawing on several previous survey studies to value changes in visibility, substantial monetary benefits were obtained for residential areas in 31 eastern states and for national parks in the southeastern United States. Benefits to this region were estimated to be \$3.4 billion (1994 dollars) in 2010, or about \$377 per ton of SO₂ emission reduction (Chestnut and Dennis, 1997). The sensitivity of this estimate has been explored using TAF, which predicts median estimates of visibility benefits for improvements at recreational sites that range from \$118 to \$224 per ton of SO₂ reduction under alternative scenarios. In part, the range reflects uncertainty about the baseline against which emission changes under Title IV should be measured. Visibility changes vary in a nonlinear fashion with emission changes, resulting in the variation in benefits per ton. Benefits at residential sites were found to be of substantial magnitude.

1 Reasonable estimates of potential ecosystem (nonuse) benefits are not attainable at this time.
2 Ecosystem health benefits are expected to be large in part because they encompass broad changes
3 that affect many environmental end points, perhaps to a small degree, but that taken together could
4 alter large-scale systems. Aquatic and terrestrial effects are likely to have significant benefits
5 through nonuse values, but uncertainties around these values remain among the largest. The link
6 between changes in emissions and changes in ecosystems is even more tenuous than is the
7 valuation based on these changes. Nonetheless, the evidence based on a small number of relatively
8 narrow studies suggests these values may be significant.

- 9 • Expected benefits (for use values) were found to be low in areas that were the primary
10 focus of benefits assessment in the 1980s--namely, ecological effects (aquatics, forests
11 and agriculture).

12 Public attention in the 1980s to air pollution from SO₂ and NO_x emissions largely centered on the
13 problem of acidification ("acid rain"), with particular concern for effects on water and soil
14 chemistry and ultimately ecological systems. Given that, some find it surprising that relatively low
15 benefit values have been estimated for effects on aquatics, forests, and agriculture. These low
16 values stem from an assessment of use values, or commodity values in the case of agriculture.
17 Agricultural benefits result in large part from changes in ozone concentrations, but modeling of
18 ozone changes resulting from Title IV is not complete. Also, these low values result from the
19 availability of substitutes that ameliorate the economic losses from emissions. The lack of
20 substitutes for increased life or health makes morbidity and mortality a more important contributor
21 to the benefits of emission reductions.

REDUCTIONS TO PREVENT ADVERSE ECOLOGICAL EFFECTS

Congress asked NAPAP to determine a "threshold" value for deposition below which adverse ecological effects will not occur. Critical deposition levels that correlate with a "threshold" of adversity are scientifically complex. NAPAP's 1990 State of Science and Technology Report No. 13 states that "biological responses to changes in acid-base chemistry occur along a continuum. There is not a single value or set of chemical concentrations that represents a 'threshold' for 'significant adverse biological effects'." (Baker et al., 1991). Therefore, NAPAP determined that the logical path was to describe the ecosystem responses along a continuum, thereby allowing decision makers to determine the level of acceptable risk.

What are "adverse ecological effects"?

A working definition of "adverse ecological effects" has been derived for use in this report based on the intent of Congress as expressed in the 1990 Clean Air Act Amendments and shaped by other relevant environmental statutes (Comprehensive Environmental Response, Compensation and Liability Act [CERCLA] and the Clean Water Act) and associated regulations. (See **Appendix C** for the analysis supporting this interpretation.)

Adverse ecological effects: any injury (i.e., loss of chemical or physical quality or viability), to any ecological or ecosystem component, up to and including at the regional level, over both long and short terms.

Similarly, other areas addressed by NAPAP (i.e., materials, visibility, and human health) would follow the same definition as that above, but are not considered here as ecological effects.

What are the dose-response relationships for sulfur and nitrogen deposition?

NAPAP has undertaken as its long-term goal the explanation, to the extent possible, of dose-response relationships along a continuum for ecological and welfare effects. As the dose of air pollution (specifically sulfur and nitrogen emissions and its subsequent formation of acid deposition) is reduced, how do aquatic and terrestrial ecosystems respond?

Generally, quantification of dose-response relationships between acid deposition and ecosystems, materials, visibility, and human health are complex. Some interactions are well understood, while those that are not are often extremely complicated relationships involving other chemicals, climate, geography, and time. For future reports, NAPAP will attempt to better understand and quantify dose-response relationships assuming appropriate research is undertaken. To the extent that they are known, dose-response relationships have been identified and are used to answer the following question.

How might ecological effects be reduced if deposition were reduced further?

Answering this question depends on dose-response functions used in a predictive mode. Such models were used extensively to answer a similar question posed by Congress to EPA under Title IV (Appendix B, Section 404 of the 1990 Clean Air Act Amendments). EPA's report focused on surface waters in a few specific regions; the findings are reviewed in this report. There has been no parallel review of soils and forest effects, but this report attempts to at least qualitatively address them.

1 Surface Waters

2 The feasibility and effectiveness of an acid deposition standard to protect sensitive aquatic and
3 terrestrial resources were examined in the Acid Deposition Standard Feasibility Study - Report to
4 Congress (U.S. EPA, 1995). In the study, EPA projects the impacts of implementation of the
5 Clean Air Act Amendments on surface waters for three case study regions and projects a range of
6 responses if deposition were reduced further.

- 7 • Both nitrogen and sulfur deposition are important contributors to chronic and episodic
8 acidification of surface waters. Further reductions in nitrogen as well as sulfur deposition
9 may be necessary to fully protect targeted sensitive systems.

10 The feasibility study concluded that establishing standards for sulfur and nitrogen deposition in the
11 United States is technically feasible, but that two critical areas of uncertainty advised against taking
12 such action at that time. First, policy decisions regarding appropriate or desired goals for
13 protecting sensitive systems were needed, especially with respect to the level of protection desired
14 and the costs and benefits of such protection. Second, key scientific unknowns, particularly
15 regarding watershed processes that govern nitrogen dynamics, limited EPA's ability to recommend
16 specific deposition standards at that time.

17 The Nitrogen Bounding Study was developed as a primary component of the Acid Deposition
18 Standard Feasibility Study. This study illustrates the modeled results of scenarios of potential
19 future nitrogen and sulfur deposition rates and different watershed nitrogen retention conditions
20 and their combined effects on surface water chemistry at regional scales. The study evaluated
21 target populations of surface waters in three regions: lakes in the Adirondack region of New York,
22 and stream reaches in the mid-Appalachian and southern Blue Ridge Mountains.

23 The Nitrogen Bounding Study illustrates that, for the near term, sulfur deposition is likely to
24 remain the primary acidification problem in the most sensitive areas of eastern North America, with
25 the possible exception of the Adirondacks. Model projections indicate that at certain times and
26 under deposition scenarios tested, sulfur and nitrogen are projected to have approximately equal
27 roles in surface water acidification. The modeling shows that nitrogen deposition effects
28 approximate sulfur deposition effects in the mid-Appalachians and southern Blue Ridge. For most
29 areas where current or near term needs for additional controls are projected, and where watershed
30 nitrogen saturation is not likely imminent, the greatest potential benefits will come primarily from
31 control of sulfur emissions and deposition. In regions where nitrogen deposition is now or would
32 likely become a more direct cause of chronically acidic conditions in sensitive waters, with
33 potential effects of sulfur and nitrogen deposition becoming approximately equal and directly
34 additive, further limits on nitrogen deposition could produce a twofold impact by both reducing
35 acid deposition rates and lengthening average times to watershed nitrogen saturation.

36 The Acid Deposition Standard Feasibility Study concluded that scientific uncertainties regarding
37 varying regional rates and differences in processes affecting watershed assimilation of acid forming
38 sulfur and nitrogen compounds preclude quantifying the reduction in deposition of either chemical
39 below which there would be no significant adverse impact. Available information does indicate
40 that additional decreases in deposition reductions would reduce regional proportions of chronically
41 acidic surface waters or proportions of surface waters most sensitive to episodic effects. The
42 magnitude of these potential benefits to each group of surface waters varies considerably by
43 region. The study projections indicate for some surface waters in the Adirondacks, mid-
44 Appalachians, and southern Blue Ridge that potential benefits may amount to a shift of a few
45 percentage points in proportions of acidic or sensitive surface waters. For other groupings in other
46 regions, deposition reductions could benefit 20% or more of the acidic or sensitive waters.
47 However, even a few percentage points may mean that many lakes or miles of stream reaches are
48 benefiting.

1 **Soils and Forests**

- 2 • Decreasing deposition of sulfur and nitrogen will lead to decreasing concentrations of
3 sulfate, nitrate, and cations in soil water and in surface water draining forest soils. Some
4 change will occur rapidly in soils that have lost their ability to retain nitrogen – such as
5 nitrogen saturated, high-elevation, spruce-fir forests – or in soils that do not have
6 appreciable sulfate sorbed to soils.
- 7 • Water in soils with low supplies of base cations will most likely remain acidic and will
8 contain aluminum, while water in soils with moderate to high supplies of base cations will
9 most likely experience an increase in pH.
- 10 • Reduced nitrogen and sulfur deposition will decrease leaching of base cations from forest
11 soils. The time to replenish base cation supplies in soils will be decades to centuries,
12 depending on rates of weathering and cation cycling, which are typically slow.
13 Consequently, an important portion of the effects on forests of low levels of base cations
14 (in particular calcium) will remain until base cation supplies can be replenished.

15 The acid-base chemistry of the soils and the water draining forest soils will depend on the
16 characteristics and sensitivity of the soils. In general, decreasing the input of sulfur and nitrogen
17 from deposition will decrease sulfate and nitrate levels in soils and soil water. The timing of this
18 decrease in soil and soil water sulfate depends on the amount of sulfur stored. Soils high in sulfur
19 will most likely experience a slower decrease in soil water sulfate as the soil slowly releases stored
20 sulfur. For nitrogen, the timing of the decrease will depend on the amount of nitrogen stored and
21 the growth needs of the forest. A forest that has high demand for nitrogen, relative to nitrogen
22 storage and throughput, will experience smaller decreases in nitrogen in soil water in response to
23 decreased nitrogen deposition compared to a forest that has a low demand for nitrogen. A forest's
24 need for nitrogen is strongly dependent upon the age of the forest. Thus, young, actively growing
25 forests are far less likely to experience nitrogen saturation than older stands that have reached
26 steady state conditions.

EFFECTIVENESS OF TITLE IV: A MARKET-BASED APPROACH

As part of the iterative process of establishing and refining air quality management choices it is essential that the choices are evaluated for how effective they have been in achieving the desired results. In the case of Title IV, the desired result was to reduce the adverse effects of acid deposition by reducing emissions at a cost lower than could be achieved under traditional methods of regulatory control by using market-based incentives.

Looking across the causal chain from emissions to effects, there are several points at which one can evaluate the effectiveness of reducing emissions. This section will review some of those measures such as, the costs of compliance for utilities, the administrative costs for the government, the reduction in emissions and acid deposition, and its subsequent minimizing effects on human health, ecosystems, visibility, and materials and cultural resources.

Did the market-based approach reduce costs?

There are 3 aspects to the evaluation of the cost effectiveness of Title IV. First, did simply the choice of a market-based approach to control over the traditional command and control approach reduce compliance costs by utilities? This is a basic economic question that, when answered, may have a significant influence on the development of future pollution control options. Second, is the management of the control program by the government cost effective? Third, given that the market-based approach was selected as the management strategy for Title IV, were the initial projections of compliance costs accurate and are we getting better at making these economic projections? It is necessary that uncertainties in projections be reduced to the point that they can be confidently used in developing policy. These three aspects are discussed below.

- The market-based approach in Title IV has achieved significant savings when compared to a command-and-control regulatory approach.

As was predicted at the time of enactment of the 1990 Amendments, the competition between different compliance options and the integration of the allowance and fuel markets have reduced compliance costs. Several recent analyses have estimated that the cost savings from the market-based approach of Title IV amount to 50% or more of the predicted cost of compliance. Several studies have been conducted since 1990 of the cost savings in 1995 from emissions trading. Although these studies used different assumptions, the projected cost savings from trading alone have ranged from \$230 million in 1992 dollars (GAO, 1994), to \$400-600 million in 1990 dollars (ICF, 1992) to about \$225-375 million in 1994 dollars (Ellerman et al., 1997). In 1994, the General Accounting Office estimated the annualized cost in 2010 to be less than \$2 billion with full inter-utility trading, compared to \$4.9 billion without trading.

The costs for the government to administer Title IV are less than the administrative costs associated with more traditional approaches to regulation. Title IV's performance-based approach eliminates the need to devise source-specific emission limits and to review control technologies and detailed compliance schedules. In addition, eliminating case-by-case review and approval of each trade greatly reduces administrative and transaction costs associated with emissions trading programs (McLean, 1996). The program's administrative costs of roughly \$12 million per year translate into a cost of about \$1.50 per ton of pollution reduced (McLean, 1997).

- Emission reductions have been achieved at a lower cost than previously estimated for Title IV.

Since 1990, estimates of total costs of Title IV continue to be revised downward. Most recently, a retrospective analysis shows compliance costs in 1995 to be \$726 million. Projections for annualized costs of SO₂ reduction by the year 2010 have declined from \$3.7 billion (ICF, 1992) to

\$2 billion (GAO, 1994). Lower costs have been linked to cost-reduction efforts and improved performance of scrubbers and changes in fuel markets. Some of these revisions can be attributed to the difficulty in estimating future technological improvements, the more efficient use of existing technologies, and future economic conditions. These technologies are being developed in a competitive environment. Anecdotal evidence indicates that technological innovation is leading to cost savings. However, without a flexible approach to compliance that allows different technologies and fuels to compete against each other, firms would not have any incentive to finding and developing cost-effective measures that exceed emission reduction targets.

The flexibility of the Title IV program has continued to apply downward pressure on compliance costs by allowing electric utilities to take advantage of market forces and the changes in the relative prices of compliance options. Allowance prices are much lower than experts had predicted and are an indicator of the success of the control program by reacting to the developments in the low sulfur coal market, reduced rail rates for delivering western low sulfur coal to some midwestern utilities, lower than expected future costs of scrubbers for Phase II of Title IV, and new efficiencies in the operation of existing scrubbers that reduce emissions at lower costs. The price of allowances has reflected the declines in control costs, dropping from an estimated \$500-600 per ton when the 1990 Amendments were passed to about \$90 per ton at the end of 1996.

How did emission and deposition changes compare to projections?

- In 1995, the first year of compliance with the provisions of Title IV, electric utility emissions of SO₂ were below the projected levels.

In 1995, total electric utility emissions of SO₂ were 12 million tons. Actual 1995 utility emissions were less than the projections available in 1990 (see Table 4 for the various projections). Only Phase I with its prescribed emission limits was in effect in 1995.

Utilities participating in Phase I were about 39% (3.4 million tons) below their 1995 allowable emissions level of 8.7 million tons. This overcompliance was beyond any projections or expectations. There is a general consensus that these reductions were a direct result of the market-based mechanism of Title IV. Of significance is the fact that most of the emission reductions were in the Midwest, the highest emitting area of the country. This is consistent with the fundamental premise of Title IV's market-based approach, that the highest emitting plants have an incentive to make deep reductions in SO₂ emissions because they usually face a lower cost per ton of SO₂ reduced.

By the only emissions goal available at this time--the allowable level for Phase I units in 1995--Title IV has been effective in reducing emissions of SO₂. The market-based approach contributed to a significant overcompliance in 1995. Care should be taken in extrapolating this performance into the future.

It is too early to quantify the effectiveness of Title IV in reducing NO_x emissions.

The Phase I requirements for NO_x reductions under Title IV only took effect in January 1996. Even so, there has been a steady decline in NO_x emissions from utilities between 1980 and 1995. The largest year-to-year decline occurred between 1994 and 1995 (1.3 million tons, or 19%). Reductions in NO_x emissions are also required under Title I and Title V.

- Emission reductions imposed by Title IV have decreased acid deposition in the eastern United States.

Analysis of the 1995 deposition monitoring data demonstrate that the 1995 reduction in SO₂ emissions in the eastern United States substantially decreased the acidity and sulfate concentration

1 in precipitation. The largest reductions occurred in the Midwest and Northeast. These deposition
2 reductions were geographically and climatologically consistent with the large emission reductions
3 in the Midwest.

4 On the other hand, nitrate concentrations in 1995 were about 5% greater than predicted in the
5 eastern part of the country. Unlike sulfate concentrations, higher nitrate concentrations were
6 recorded at only 61% of the monitoring sites. Even though utility emissions of NO_x declined in
7 1995, they account for only 29% of all NO_x emissions (vehicles contribute the largest fraction).
8 Combined with the annual variability in meteorology, the record is too short to be able to attribute
9 changes in nitrate concentration in precipitation to NO_x reductions by utilities.

10 Qualitatively, Title IV has been effective in reducing acid deposition. However, the geographic
11 distribution and the quantitative measure of the changes in total deposition resulting from emission
12 reductions require a longer monitoring record and better estimates of dry deposition.

13 **Has reduced acid deposition had an effect on sensitive receptors?**

14 It is still too early to make a definitive statement on the impacts of deposition changes on sensitive
15 receptors. First, the magnitude, spatial distribution, and timing of changes in sulfur and nitrogen
16 emissions and deposition -- before and after implementation of Title IV -- need to be established.
17 The location of changes in emissions and deposition is important because the distribution of
18 sensitive receptors is not uniform and responses depend on specific exposure factors. For most
19 effects areas, changes attributable to Title IV have not yet been observed. Some systems,
20 particularly aquatics, forest and materials, can take many years to respond to chemical changes as a
21 result of reductions in atmospheric deposition. Two years since initiation of emissions reductions
22 under of Title IV is too short of a time period to detect changes. In addition, for most effects areas
23 there has been no systematic attempt to monitor the changes in sensitive receptors during this time
24 period.

OUTLOOK TO 2000

NAPAP recognized with the passage of the 1990 Clean Air Act Amendments that a complete assessment of Title IV in 1996 would be premature because emission reductions would not begin until 1995. Furthermore, due to scientific uncertainties, weather variability, and the inherently slow response times of many ecosystems, a quantification of human and ecosystem responses to any changes in emissions could not be made with reasonable confidence in 1996. Hence, a limited assessment was planned for 1996, with a more comprehensive assessment in 2000.

The goals of the 1996 assessment were to (1) provide information about acid deposition in a format that facilitates communication with policymakers and the public, (2) validate the assessment methodology, and (3) identify the monitoring, research and modeling needs for future assessments. This section is devoted to a discussion of the last two goals.

In accordance with its congressional mandate, NAPAP will conduct another integrated assessment of the costs, benefits, and effectiveness of Title IV in 2000 and will document the results in its 2000 Report to Congress. Given the scope of the 1996 assessment and the advice and comments of participants and stakeholders in the assessment process, NAPAP will modify the scope of the assessment over the next three years, based on the availability of resources and relevant scientific and economic information.

The 2000 assessment will include more case studies and will improve the depth of regional assessments. The middle and southern Appalachian mountains, northern New England, and the high-elevation West are currently being considered as case studies. An issue expected to receive more attention in 2000 is characterizing and quantifying (both air concentrations and dry deposition) sulfate and nitrate fine particles, and quantifying their causal relationships to human health and materials. Improvements in predictive modeling of air chemistry and ecosystem responses and in valuation methodologies for health, cultural resources, and visibility will enhance our ability to assess the "lifetime" benefits of emission reductions in most of the effects areas.

Many of these needs are crosscutting; they serve to meet the goals and objectives of individual agency missions and improve the knowledge base for other national issues. Acid rain is a crosscutting issue itself, with linkages to several national and international environmental issues, such as fine particles, radiative forcing, and NAFTA, the North American Free Trade Agreement (these issues are discussed under the Linkages subheading toward the end of this section).

Research, Monitoring, Modeling, and Data Needs

Despite the concerted research effort on the effects of acid deposition during 1980-1995, many important scientific and socioeconomic areas require further investigation before a more quantitative and comprehensive assessment can be made. Fundamental monitoring and scientific inquiry must be continued and reinforced to provide a foundation upon which we can build better assessments. Near-term research, monitoring, and development needs must be addressed to improve the knowledge base for the 2000 NAPAP assessment, and long-term needs must be addressed for subsequent NAPAP assessments. The research, monitoring, modeling, and data needs listed below are intended to be general descriptions of the types of activities that will improve the knowledge base for future assessments. For a general overview of research and monitoring needs see **Table 9**. Specific projects will be conducted, depending on their individual merit and level of impact. The needs were identified by the contributors to this report, including reviewers and other stakeholders.

The fundamental needs are presented below, followed by the near-term and long-term needs, organized according to subject area. Each section is broken down by the four elements of a successful assessment: monitoring, research, modeling, and data availability.

TABLE 9. RESEARCH AND MONITORING NEEDS FOR FUTURE NAPAP ASSESSMENTS

NEED	ACTION	BENEFIT
Timely utility and nonutility emissions data.	Ensure timely and open access to data, including industrial, Mexican, and Canadian emissions.	Improved ability for timely attribution of changes in deposition and effects to changes in emissions due to Title IV.
Quantification of status and trends in deposition and visibility.	Continue wet and dry deposition and visibility monitoring and analysis.	Long-term record ensures a better way to measure the effectiveness of Title IV.
Regional assessment of nitrogen impacts on aquatic and terrestrial ecosystems.	Conduct regional nitrogen modeling and nitrogen field studies.	Provide projections of nitrogen impacts on aquatic and terrestrial ecosystems and actual data on nitrogen cycling to facilitate regional modeling
Improved and sustained ecosystem monitoring; data on ecosystem health.	Expand and enhance long-term ecosystem monitoring relevant to acid deposition.	Ability to associate trends in ecosystem health with emission trends.
Quantification of the relative role of sulfate and nitrate aerosols in particulate health effects.	(1) Improve sulfate and nitrate aerosol concentration estimates; and (2) determine causative agent(s) of health effects.	Improved understanding of the health benefits from Title IV.
Decreased uncertainty in mortality valuation.	Reduce variation in "value of statistical life" estimates.	More accurate estimate of the valuation of health benefits.
Quantification of trends in water quality.	Continue long-term monitoring of lakes and streams.	Long-term record required to evaluate the effectiveness of Title IV and improve understanding of watershed processes.
Reduced uncertainty in benefit estimates in the areas of visibility and materials.	Investigate preferences for changes in visibility and the possible magnitude of benefits from materials preservation.	More accurate assessment of the benefits and costs of Title IV for areas of greatest concern.

1 **Fundamental Needs**

2 An assessment of the effectiveness of acid rain controls cannot be undertaken without the most
 3 basic scientific and technical information. The 1996 assessment depended on a set of elements that
 4 are essential, but not sufficient, for future NAPAP assessments. Primary examples are long-term
 5 environmental monitoring data and analysis, research and process studies leading to evolutionary
 6 model improvements, and data access. In most cases, the information sources are longstanding
 7 monitoring, research, and modeling programs within the NAPAP agencies. The following
 8 fundamental programs and activities listed below require annual resources, attention, and
 9 evaluation. Although they are often under the programmatic control of one of the NAPAP
 10 agencies, they contribute to the mission of other agencies and are essential to interagency programs
 11 like NAPAP. For that reason, relevant agencies are also identified, where appropriate.

12 Monitoring

13 Monitoring forms the basis of assessment. It provides the status and trends of the state of the
 14 environment, and the data are often used in *post facto* statistical inquiry into the form and
 15 magnitude of causal relationships (e.g., epidemiological studies). Existing long-term
 16 environmental records show considerable year-to-year variability due to meteorology, changes in
 17 source emissions, and other factors. A longer period of data is required before environmental
 18 trends can be established with confidence and reasonable inferences can be made into the causal
 19 relationships between emissions and deposition and deposition and effects. Monitoring consists of
 20 regular data collection, quality control and assurance, routine analysis, and presentation of peer-
 21 reviewed findings. The following monitoring efforts need to be continued to ensure a credible and
 22 comprehensive assessment of atmospheric deposition and its effects in 2000.

- 23 • *Emission Inventories.* A comparison and benchmarking of continuous emissions
 24 monitoring data versus historical emission estimates will ensure a continuous, long-term
 25 data record. Such a record is the essential first step in linking changes in emissions to
 26 changes in deposition.
- 27 • *Air Concentration and Deposition.* Air chemistry and composition data represent the most
 28 direct indication of whether emission reductions produce the desired results. Long-term
 29 monitoring of ambient air concentrations and deposition is needed to determine whether the
 30 air quality benefits that appear to be accruing in terms of emission reductions will translate
 31 into reductions in the rate of atmospheric deposition, or of the loadings to sensitive
 32 ecosystems that the legislation is attempting to protect. A commitment to continuing
 33 monitoring of the kind reported here at key sites is needed. The current monitoring
 34 networks--EPA's CASTNet, the multiagency IMPROVE, and NOAA's AIRMoN--have
 35 had benefits. These networks are quite different in their design purposes, and each has
 36 proved successful in meeting its original expectations.

37 Continued monitoring and interpretation of the nation's atmospheric deposition are needed
 38 because they provide the only physically based, nationwide data set to determine whether
 39 the regulations reducing atmospheric deposition and its associated adverse environmental
 40 impacts. Continued acquisition and analysis of the deposition data from the federal, state,
 41 and private sector sponsored National Atmospheric Deposition Program/National Trends
 42 Network (NADP/NTN) will provide the information necessary to evaluate the effectiveness
 43 of Title IV in protecting the environment. Continued monitoring from all these networks
 44 will also provide information to evaluate other air quality issues, such as utility
 45 restructuring and expected changes in NO_x emissions from mobile and stationary sources
 46 of air pollution.

- 1 • *Soil and surface water chemistry.* Continued monitoring of lakes and streams in EPA's
2 Long Term Monitoring program, Temporally Integrated Monitoring Ecosystems program,
3 and other state and federal programs is necessary for quantifying the responses of surface
4 waters to changing levels of sulfur and nitrogen deposition. For the most part, the Long
5 Term Monitoring program represents the longest continuous record of surface water
6 chemistry in the United States and is often used to study watershed (soil) chemistry, an
7 essential element in both terrestrial and aquatic ecosystems.
- 8 • *Forest health.* Continued monitoring of sites in sensitive ecosystems with prior strong
9 databases on forest condition and soil chemistry is necessary to quantify the responses of
10 terrestrial ecosystems to changing levels of sulfur and nitrogen deposition.
- 11 • *Visibility.* Visibility is the most direct, well understood indicator of changes in emissions.
12 Aerosol composition and visibility monitoring, such as that provided by the multiagency
13 IMPROVE network, should be continued consistently at sites with a record of such
14 monitoring. Although the relationship between emissions and air concentrations is highly
15 variable because of the weather and other factors, the relationship between concentrations
16 and visibility is well understood.

17 Research and Analysis

- 18 • *Comparison of actual to predicted visibility trends.* Extensive periodic assessments (at least
19 every five years) of the fine particle sulfate and visibility trends, and their relationships to
20 SO₂ emission trends and variations in meteorology, need to be conducted to determine
21 whether expected benefits of SO₂ emission reductions are being realized and to test the
22 credibility of modeled predictions.

23 Modeling

- 24 • *Integrated assessment tool.* NAPAP has developed a prototype integrated assessment tool
25 called the Tracking and Analysis Framework (see Appendix A). TAF has been peer
26 reviewed and possesses unique capabilities to provide the kind of information that NAPAP
27 needs for its assessments. Continued maintenance and development of TAF, especially of
28 those components that would benefit all effects areas (e.g., emissions, air transport and
29 diffusion, and economic assumptions), would provide an analytical tool specifically
30 designed for an integrated assessment of Title IV.
- 31 • *Regional acid deposition modeling.* Regional atmospheric deposition models, such as the
32 Regional Acid Deposition Model (RADM), are necessary to study the geographic origins of
33 deposited pollutants and the anticipated effectiveness of future emission reductions.
34 Continuous evolutionary improvements of model physics and chemistry, such as
35 parameterizations of internal nitrogen cycling, are necessary to develop an improved
36 analytical tool. Consistent with NAPAP's principles, formal model evaluation and peer
37 review should be a part of the overall model development process.
- 38 • *Economic/emissions predictive models.* Predictions of air quality and atmospheric
39 deposition rely on reliable economic and emissions prediction models. The improvement
40 of current models and the development of new models are necessary if new insight are to
41 be expressed in future assessments.

42 Data Access and Database Development

43 The free and open exchange of data ensures the broadest participation in scientific and economic
44 inquiry. Basic to this are the availability and access to quality-assured data. In some cases, data

exist but have not been analyzed and/or quality-assured, access is neither timely nor inexpensive, or the data need to be placed in a useable form. In other cases, needed data have not been collected or observed.

- *Emission inventories.* Monthly, up-to-date inventories of utility and nonutility SO₂ and NO_x emissions are required at the substate level.
- *Deposition data.* Quality-assured wet deposition data are already available for downloading via the World Wide Web. Dry deposition estimates from EPA and NOAA monitoring networks also need to be made available in a timely manner.
- *Geographic Information Systems (GIS).* GIS are already used by NAPAP agencies to support their mission. These GIS should be screened and explored as acid rain analysis and display tools, and geographic measures of the effectiveness of emissions reductions should be developed.
- *Model assumptions.* Much of an assessment relies on modeling. NAPAP aims to use only peer reviewed, nonproprietary models and information in its assessments. To that end, models that are potential contributors to NAPAP assessments, including their assumptions and science, must be peer reviewed and accessible.

Near- and Long-term Needs

Near-term needs are defined as new activities that must be completed and peer reviewed in time to be useful to the 2000 assessment--nominally, the second quarter of fiscal year 2000. These needs can be divided into two groups: large benefit areas (where even small uncertainties could have large valuation implications or acceptable valuation methodologies are not yet developed), and reductions in scientific and economic uncertainties (unknown or large uncertainties in scientific and economic understanding prevent any improved analysis of costs and benefits, either physical or economic). The degree to which the expectations for the 2000 assessment can be met depends on which near-term needs are addressed by 2000. A summary of the near-term needs can be found in **Table 9**.

Long-term needs are defined as those that cannot be met in time for the 2000 assessment but which are needed to contribute to future assessments.

Emissions and Compliance

Most of the emissions and compliance data and information requirements for an assessment of Title IV are satisfied by EPA operational activities or will be satisfied if the relevant fundamental needs specified above are met. However, there is a need to address the issues of transboundary pollution, as well as the technological and state regulatory impacts on the effectiveness and costs of Title IV.

Near-Term Needs

- *Research.* The following investigations must be completed in time to be useful to the 2000 assessment.
 - Analysis of industrial SO₂ and NO_x emission inventories.
 - Analysis of Canadian and Mexican emission inventories.
 - Retrospective analysis of causes of emission changes in the late 1980s and early 1990s.

- Analysis of the impacts of state Public Utility Commission regulations and policies on the allowance market.
- Analysis of the relationship between Title IV requirements and technology innovations (dynamic efficiency).
- Analysis of the dynamics of allowance trading and its effects on economics and the spatial and temporal distributions of emissions.

Concentration and Deposition

Acid deposition rates and ambient air concentrations of acid rain precursors will continue to be the most direct measures of the effectiveness of acid rain controls. The space and time variability of the data presents a challenge to documenting regional deposition trends. It is also a challenge to integrate the information from many different networks, each having its own (often mission-oriented) objectives and purpose of design. Projections of future deposition rates will be necessary in developing or modifying emission controls. This is only possible through the use of validated, peer-reviewed models.

Near-Term Needs

- *Monitoring.* Additional monitoring and interpretation from the concentration and deposition monitoring networks will be necessary to determine what effect various changes in nitrogen emissions are having on nitrates and nitrogen deposition. The data can be used to assess if increases in nitrates from mobile sources in some areas are offsetting reductions achieved by the Clean Air Act Amendments.
- *Research.* There is a continuing need to assess rates of dry deposition from the atmosphere, especially to areas where ecosystems are sensitive to sulfate, nitrate, and other chemicals that together make up acid deposition. Although dry deposition constitutes a significant fraction of total deposition, the scientific understanding of the processes is incomplete, and measurement technology is inadequate. The questions of deposition to landscapes and spatial heterogeneity remain largely unanswered. Research is modeled to develop methods for extrapolating from areas with dry deposition measurements to nearby areas with different terrain, vegetation, and chemistry.
- *Modeling.* Learning how to link the site-specific deposition values made by monitoring programs with the spatial average deposition predictions yielded by regional atmospheric deposition models remains a challenge.

Long-Term Needs

- Increased attention to the processes that modify acid-generating compounds in the air are needed, in particular insofar as these processes include mixes of natural and anthropogenic chemicals.

Aquatic Ecosystems

Additional work is required to improve the scientific understanding of the effects of acid deposition on surface water chemistry. A variety of organizations and agencies have called for a coordinated, integrated assessment of the environmental effects of anthropogenic emissions. The occurrences of coastal eutrophication are concurrent with acid deposition. Continued model development and application will provide an adequate basis for preparing an assessment of aquatic effects in the year 2000. The principal needs relate to monitoring and modeling activities.

Near-Term Needs

- *Monitoring.* The monitoring of lakes and streams in EPA's Long Term Monitoring program must be continued at least through 1999 to allow quantification of the responses of surface waters to changing levels of sulfur and nitrogen deposition due to continued implementation of Title IV. This information will be needed to provide an adequate basis for the 2000 assessment.

- *Modeling.* Modeling needs fall into three general categories: further regional sulfur modeling; implementation of regional nitrogen modeling; and continued model testing and verification, especially of nitrogen models. The effect of seasonal variations in NO_x emissions and deposition on aquatic systems also needs to be studied. Testing and improvement of the MAGIC sulfur model since 1990 have substantially modified forecasts of the pH of Adirondack lakes as sulfur deposition declines in response to Title IV, as well as forecasts of ranges of watershed responses to simultaneous changes in sulfur and nitrogen deposition. Model revision and reapplication efforts should be pursued in the following priority order: mid-Appalachian Mountains, northern New England, and southern Appalachian Mountains. This effort will not require new data collection.

It is now evident that nitrogen may be critical in chronic and episodic acidification. Models to simulate the terrestrial and aquatic effects of nitrogen deposition are now available. One or more such models should be implemented in areas of the United States thought to be highly sensitive to potential nitrogen effects. Candidate regions, in priority order, include the Adirondack Mountains, mid-Appalachian Mountains, southern Appalachian Mountains, portions of the West, and northern New England. This effort would require some field data collection to quantify nitrogen and carbon storage in important ecosystem compartments, mainly soils and foliage. Field work would have to be conducted at the latest during the summer and fall of 1998 to be useful for the 2000 assessment. Testing of the nitrogen models should continue in 1997 and 1998. Available data sets, with which to further test the models, include ecosystem manipulation data sets from Maine and Europe and long-term monitoring data sets from New York, New Hampshire, Norway, and Germany.

Model development efforts are necessary to improve the characterization of short term events (episodic acidification) as they relate to surface water chemistry, adverse effects to aquatic biota, coastal eutrophication, and harmful algal blooms.

Long-Term Needs

- *Monitoring.* Long-term monitoring is the only approach that can be used to determine the ultimate effectiveness of implementing any policy or management option. Because model projections will always entail considerable uncertainty, it is critical that real monitoring data continue to be collected in the long term within all regions of interest.

- *Research.* Research is needed on the effects of atmospheric nitrogen deposition on estuarine and near-coastal marine ecosystems. Atmospheric nitrogen is known to contribute to the eutrophication of such systems (excessive productivity, which can deplete dissolved oxygen and harm aquatic biota). The quantitative importance of atmospheric inputs, in addition to other sources of nitrogen, such as agriculture, sewage, and industry, is not well known.

Forest Ecosystems

Forest ecosystems are exposed at varying rates to a multitude of stressors. Stresses can be delivered by atmospheric process or can be related to weather or soil condition. This multiple-stressor paradigm makes attribution of damages and the quantification of causal relationships difficult.

Near-Term Needs

- *Monitoring.* An updated analysis of the health of forest ecosystems should be undertaken. The analysis would include a designation of ecosystems that are and are not sensitive to acid deposition.
- *Research.* A focus on the degree to which sensitive ecosystems are affected by current and reduced deposition rates. In particular, studies on older trees in acid soils are needed to evaluate the long-term, interactive effects of acid deposition and ozone.
- *Modeling.* Development and refinement of regional scale, spatially explicit models needs to be accelerated, and data for running and testing the models (PnET, MAGIC, NuCUM) need to be acquired. These models are required to predict responses of soil and soil water chemistry to increases or decreases in sulfur and nitrogen deposition and to assess potential responses to changes in management practices.

Long-Term Needs

- *Monitoring.* Because of the often long lag time involved in ecosystem responses to changes in deposition, future monitoring is necessary to elucidate responses attributable to emission reductions already realized. Periodic monitoring of terrestrial resources, including forest and soil conditions, should be initiated for a suite of carefully selected forest research sites. The effects of changes from both sulfur and nitrogen deposition should be documented.
- *Research.* Mechanistic studies are needed to evaluate the consequences of changes in soil chemistry (e.g., calcium depletion) on essential plant processes ranging from physiological processes to ecosystem function. These studies should include the synergistic relationship of soil chemistry with changes in sensitivity of vegetation to natural stresses, such as cold damage; decreased resistance to disease; and decreased structural resistance to physical stresses, such as wind and ice (i.e., multiple stressors). There are seven specific areas.
 - Research on the effects of changes in soil chemistry on forest health in northern hardwoods, especially sugar maple.
 - Long-term studies on the role of cation depletion and nitrogen addition on nutrient cycling and tree growth for southern pine forests.
 - Studies on the relative importance of cation leaching from foliage and cation losses from soils on whole plant function (short-term versus longer-term effects).
 - Studies to explore the role of acid deposition in limiting cation availability in southern pine forests growing on poorly buffered soils.
 - Studies on the role of cation depletion in soil profiles on the distribution and function of roots.

- Evaluation of the role of acid deposition in contributing to nitrogen saturation of forest soils through impaired root uptake of soil nitrogen.
- Development of reliable indicators of ecosystem condition and potential susceptibility to acid deposition. These indicators are needed to assess potential responses to altered nitrogen and sulfur deposition rates, and the spatial variability that is evident within a region. The indicators must be applicable at the site and watershed scales.
- *Data Access and Database Development.* Determine the distribution of soils that are susceptible to leaching of base cations and mobilization of aluminum. This information is critical to understanding dose-effect relationships and to projecting future impacts and improvements in response to changing deposition.

Materials and Cultural Resources

There are particularly significant benefits arising from materials and cultural resources....

Near-Term Needs

- *Research.* Applied research is needed to develop and improve treatments to prevent or minimize injury of cultural resources. Effective preservation of cultural resources affects our ability to assess the economic impact of Title IV. Costs for long-term maintenance and corrective treatment of injury to cultural resources must be considered in the overall economic valuation. Innovative treatments built upon fundamental research and accurately tested through applied studies may lead to more cost-effective strategies that will ultimately influence economic assessments.
- Continued emphasis is needed on determining ways to value cultural resources and estimate damage. One of the most promising approaches for economic evaluation of the effects of Title IV is the study of improving the "health" of a group of cultural resources by retarding the process of deterioration through treatments. This results in the extension of the average lifetime of cultural resources. In the proposed approach, historically documented changes in deterioration rates can be related to the prolonged life of the cultural asset.

Long-Term Needs

- *Research.* A better understanding of the factors and variables influencing pollution-induced stone deterioration is needed to improve diagnoses of damage to cultural resources caused by individual pollutants. In particular, it is important to further resolve effects associated with anthropogenic activities from the natural background weathering effects. A systematic approach to identifying and studying the individual dominant interactions contributing to the overall weathering of stone is needed to elucidate the deterioration processes.
- *Data access and database development.* The main obstacle to valuing the benefits of both materials and cultural resources is the lack of a complete inventory of affected assets. Combined with the above, the potentially large database could be used to quantify the particularly significant materials and cultural resources benefits

Visibility

Near-Term Needs

- *Data Access and Database Development.* Sulfur dioxide emissions data must be developed with at least monthly averaged temporal resolution and at least county spatial resolution to

be able to assess relationships between fine particle sulfate and sulfur dioxide emission trends.

Long-Term Needs

- *Monitoring.* Visibility is a local phenomenon. However, its impairment is a national issue involving both urban and rural settings. Additional collocated visibility and ambient air concentration monitoring sites should be initiated to provide a more complete spatial coverage.

Human Health

Near-Term Needs

- *Monitoring.* Adequate monitoring data are needed for epidemiological research efforts and the characterization of exposure risks. Separate monitoring of fine and coarse particulate matter is essential. Further characterization of particulate matter is necessary to determine the quantitative relationship of the different atmospheric components, including transition metals, ultrafine fraction, particle number, hydrogen ion, gases associated with particulate matter (e.g., ozone and carbon monoxide), and sulfur and nitrogen oxides. The ambient conditions that potentially confound epidemiological studies – weather and other pollutants (e.g., biological particles, organic compounds) – must also be monitored. The monitoring must be consistent across time and space, and must be frequent enough to determine temporal and spatial variability, especially of an episodic and seasonal nature; annual levels are not adequate.

- *Research.* Further epidemiological research is necessary to improve our understanding of the individual health parameters affected by particulate matter and co-pollutants (i.e., lung function parameters, causes of death) and the relationships between co-pollutants and health effects. Improved characterization of long-term health effects is needed and, in particular, the extent of life-shortening that can be attributed to particulate matter in general and sulfates and nitrates in particular. Studies need to follow large healthy populations living in areas with elevated particulate matter, sulfate, nitrate, or ozone for an extended number of years, as well as during episodic pollution events. In addition, more studies should be focused on populations at risk and on the issue of confounders (e.g., weather, the age of those adversely affected). Further characterization of individual exposure is needed to adequately account for the large portion of time most individuals spend indoors. Improved personal exposure assessments for particulate matter, especially for the most vulnerable members of the population, are necessary for better risk assessments.

Studies need to continue to identify the underlying biological causes of the relationship observed in epidemiological studies between particulate matter and human health to better understand the aspects of particulate matter pollution that need to be reduced or controlled and how certain members of sensitive populations can be better protected. It is important to understand how particulate matter could lead to death immediately associated with exposure and how particulate matter could contribute to shortening the lives of the sick and healthy, old and young. Further evaluation is needed for specific components of particulate matter (especially different forms in which acids can exist), as well as their fates and their effects on the environment and on human health. However, the absence of specific biological mechanisms should not be allowed to diminish the policy implications of the relationship between particulate matter (and other pollutants) and human health observed in epidemiological studies.

- *Modeling.* Development, refinement, and more experience are needed with models that associate pollution and health, including air quality models, as well as models that associate

health effects with changes in air quality, and models that value those changes in health effects. Also, the models used in clinical and toxicological studies need to be improved, for example, in the pollutants used to model ambient pollution and in the subjects exposed to air pollutants in the lab.

Benefits Valuation

Near-Term Needs

The greatest value of incremental, near-term information to reducing uncertainty in benefit estimates is in the areas of health, visibility, and materials. These priorities are illustrated under Question 5 in Table 7.

- *Research.* Near-term research efforts (by 2000) would make a substantial contribution to reducing uncertainty in estimates of economic benefits. The areas most deserving of additional research are not necessarily the same as those where expected benefits are largest (see Table 7). For instance, though ecosystem health has potentially large benefits and there is tremendous uncertainty surrounding those benefits, the cost of reducing this uncertainty through additional research is great, and the time frame for such a research project is long. On the other hand, the costs of reducing uncertainty around human health mortality are probably much less. Research should be conducted to improve our basis for valuing life years lost or for adjusting the value of statistical life for age of individuals and various risks and health status. Also, the empirical basis for valuing morbidity effects is weak.
- *Modeling.* Economic models are incomplete in almost every benefit area, but they are adequate to provide some illustrative estimates and to reveal the relative magnitudes of potential benefits in many cases. Development of economic models requires both theoretical underpinnings and readiness for implementation. In aquatics recreation, agriculture, and human health, economics has the capability of providing reliable, replicable estimation of benefits. In the areas of visibility and materials, economics provides at least a sufficient foundation for investigating the relative magnitudes of potential benefits and the likely direction of potential bias resulting from factors left out of the model or flaws in the data.
- *Data Access and Database Development.* Limitations on environmental and/or economic data are the weakest link in estimating benefits, especially in areas that are suspected to have the largest potential benefits. Typically, the cost of research in economics and the collection of economic and environmental data are less than the cost of scientific research. This is reflected in the priorities for research aimed at improving benefit estimates. In the areas of health, visibility, and materials, the greatest gaps are in economics and data collection rather than science. Hence, the biggest "bang for the buck" in the near term can be bought through investments in these areas.

In some cases, weaknesses in the scientific understanding – as in the case of forest recreation or ecosystem health – preclude collection of data linking changes in primary pollutants with economic end points. In other cases, such as visibility, monitoring has been inadequate to calibrate links with economics. Where benefit end points are not well established, data will be inadequate.

Linkages to Other Issues

In the early years of NAPAP, it became clear that acid deposition could not be easily isolated from other air pollution issues. Acid rain is but one of many related environmental issues and is a focus

for several international activities. Research has established linkages between acid deposition and other environmental issues that are not within the scope of this report and are adequately addressed by other programs. However, from a policy perspective, it is important to understand the science linkages so that the public policy linkages can be made. The long-range transport of air pollution has made acid deposition an international concern and it is being addressed in several fora.

Related Science Issues

The environmental science issues related to acid deposition include radiative forcing, tropospheric ozone, fine particles, atmospheric deposition, and coastal ecosystem health.

Radiative Forcing

In addition to reducing visibility, interference with light transmission can lead to changes in climate on local, regional, or global scales. Climatic variables influenced by aerosols include temperature, relative humidity, amount of clouds or fog, amount and type of precipitation, and albedo (reflectivity) of both clouds and clear air.

Aerosol particles affect climate by (1) scattering and absorbing solar radiation, (2) absorbing infrared radiation, and (3) acting as nuclei for the formation of cloud droplets and ice particles, thereby increasing cloud albedo. The first two effects are referred to collectively as direct effects, whereas the third is referred to as an indirect effect. The term *indirect* is not an indication that cloud effects are less important than direct effects. Indeed, the highest sensitivity of regional and global heat balance to aerosols may involve interaction with clouds.

The average total anthropogenic decrease in solar radiation at the ground due to reflection and absorption is about 10W/m² (about 7%) over an area of approximately 10,000,000 km² in eastern North America (and evidently over similarly sized areas in Europe and eastern Asia). This effect acts in a manner opposite to the mechanism for greenhouse warming. Direct heat can occur through absorption of infrared radiation from the earth by coarse dust particles (as is done by greenhouse gases). This latter direct effect is not well characterized. Indirect effects are also highly uncertain and not well understood.

Sulfates are assumed to contribute about half of the total direct solar reflection in the East (a 7% reduction in solar radiation, compared with natural background levels). The role of sulfur and nitrogen oxides with respect to indirect cloud effects is not well understood (NAPAP, 1991).

Tropospheric Ozone

Nitrogen oxides and volatile organic compounds are the primary precursors of tropospheric ozone. As a result of oxidation reactions in the atmosphere these compounds form ozone. These same oxidation reactions convert sulfur oxides and nitrogen dioxides to their sulfate and nitrate forms, leading to acid deposition. Research contributing to the understanding of ozone chemistry, transport, and fate (now being conducted as part of the North American Research Strategy for Tropospheric Ozone - NARSTO) will be critical in interpreting the results of sulfur dioxide and nitrogen oxide controls to reduce acid deposition. It will be important to know how controls of these compounds have interacted to produce beneficial results in environmentally sensitive regions, and to be able to sort out their respective contributions.

Both issues have nitrogen oxides in common. Understanding their sources and sinks will be important in resolving the acid deposition and ozone issues. Nitrogen oxide control is emerging as an essential feature of the eastern ozone control strategy and is also a significant part of the national acid deposition reduction program. An important question to address in the future will be the degree to which ozone-related nitrogen oxide controls (which are predominately summertime controls) also mitigate acid deposition and coastal eutrophication problems.

Fine Particles

The fine particulate matter (PM) and acid deposition issues are closely linked in the eastern United States. Both share the same dominance of a large sulfate fraction in the chemical composition of collected samples. Sulfates are a significant (30-40%) component of fine particles in the East. However, where sulfates have been linked directly to ecosystem damage, they have yet to be identified as a principal causal agent in human health effects correlated to elevated concentrations of particulate matter. The characteristics of particulate matter, including sulfates responsible for morbidity and mortality effects, are the subject of ongoing research.

Both direct emissions of particulate matter and secondary particle formation caused by oxidation of sulfur dioxide, nitrogen dioxide, and aerosol organic carbon species contribute to overall levels of airborne particles. With sulfates being the focus of the nation's acid deposition control strategy, and an expectation of a 40% reduction in sulfur oxide emissions from 1980 levels by 2010, a dramatic decrease in eastern fine particles is expected, particularly in the sulfate fraction. This should also have a beneficial impact on health effects related to particulate matter. Ongoing health and exposure research and future environmental monitoring will provide the answers and bear watching closely for this synergy.

Atmospheric Deposition

The deposition of sulfates and nitrates originating from utilities is but one facet of a larger science and policy issue, atmospheric deposition. Many other pollutants find their way to the earth's surface by wet and dry deposition processes, including trace metals, hazardous air pollutants, and organic compounds. Prominent examples are pesticides and mercury. The atmosphere is now recognized as a major pathway by which these pollutants can be transported and deposited in areas sometimes far removed from their sources. Recent research suggests that mercury is more mobile than previously thought because of volatilization and atmospheric transport. Although actual ambient air concentrations of these pollutants are generally small and below air quality levels, it is the long term accumulation in the ecosystem via deposition that is most harmful.

The overabundance of certain nutrients, primarily nitrogen (nitrate and ammonia) and phosphorus, can cause adverse ecological effects. More than 3.2 million tons of nitrogen are deposited from the atmosphere each year in the United States. This nitrogen falls upon both land and water contributing to serious problems, such as eutrophication.

Commonalities often arise in studying the causes and effects of acid deposition. There are several significant sources of nitrogen deposition besides utilities, namely, other combustion sources (primarily automobiles) and ammonia (often a result of agricultural practices). Combustion sources (including utilities) also emit, albeit in much smaller amounts, trace metals and carbonaceous aerosols. Many of these pollutants (SO₂, NO_x, ozone, fine particles, etc.) are subject to the same (or similar) atmospheric transport and diffusion processes.

Coastal Ecosystem Health

It is now obvious that ammonium and nitrate deposition are central concerns to the health of coastal ecosystems. Although these species are major contributors to acid deposition, their main environmental consequence is eutrophication of coastal waters. The problem is not just the deposition to the water bodies themselves, but the transport of airborne nitrogen species through surrounding watersheds, streams and ground water into the water bodies that become overenriched with nutrients. Depending on the water body in question, atmospheric deposition is likely to account for as much as 30-40% of the total nutrient loading received. Reducing deposition is not simply a case of each state or region controlling its own emissions. The chemicals of concern

come from sources that are sometimes quite distant and outside the regulatory reach of any such local authority.

Along the eastern seaboard of the United States, the importance of long-range transport and atmospheric deposition in the regulatory process has been evidenced by the success of a series of "Shared Resources" workshops. Conducted by the Chesapeake Bay Program (EPA and NOAA), these workshops are focusing on the need to recognize that the emission decreases imposed by the Clean Air Act Amendments will benefit all eastern estuaries, but to an extent that is not yet well determined.

Related International Activities

United States / Canada Air Quality Agreement

The benefits of the SO₂ emission reduction program under the 1990 Clean Air Act Amendments are extensive and are not limited to the United States. Improvements in Canadian air quality that can be attributed to Title IV's reduction of U.S. emissions – and the associated benefits – should be expected. The United States and Canada continue to address transboundary air pollution issues under the bilateral Air Quality Agreement signed in March 1991. The third biennial report published in 1996, focuses on the substantial progress the United States and Canada have made in achieving emission reduction goals and in carrying out long-term programs to reduce the effects of acid rain in both countries. The report also discusses new areas of concern, such as ground-level ozone (smog), inhalable particles, and air toxics. Developments in scientific and technical cooperation are reported, including emission inventories and data trends from 1980 to 2020 for SO₂, NO_x, and volatile organic compounds, which along with NO_x emissions contribute to the formation of ground-level ozone.

The progress report also contains the first five-year review and assessment of the Air Quality Agreement by the two governments. While the Agreement does not currently focus on transboundary air pollutants other than acid rain, both governments have begun to study regional ozone management and are evaluating the role the Air Quality Committee may play regarding air toxics. The review concludes that, overall, both governments have been successful in fulfilling their obligations under the Agreement, particularly the acid rain control programs in both countries.

U.N. Convention on Long-Range Transboundary Air Pollution

The 1979 United Nations Convention on Long-Range Transboundary Air Pollution develops various protocols for air pollution control commitments. The United States participation in the Convention consists of providing expertise and advice on a number of issues, including compliance monitoring, enforcement, emission inventory development, and market-based mechanisms.

NAFTA

At the continental level, the issue of atmospheric transport and deposition is receiving considerable attention. The North American Agreement on Environmental Cooperation (a side agreement of the North American Free Trade Agreement) is engaging in a process to enlighten regulatory authorities in Mexico and Canada, as well as in the United States, on the need to consider local environmental problems related to atmospheric deposition (including acid rain and nutrient overenrichment) as problems that disregard political and geographic boundaries. The Council for Environmental Cooperation of the North American Agreement on Environmental Cooperation is currently finalizing documentation relating to "continental pollutant pathways" that will propose new multilateral steps toward protecting the continental environment from threats related to chemical deposition from the air.

Global Atmospheric Watch

At the global scale, the leadership of the United States science developed under the auspices of NAPAP is now well recognized. Three of the contributing agencies of NAPAP (EPA, DOE, and NOAA) have joined forces to set up a Quality Assurance/Science Activity Center for the Global Atmosphere Watch under the auspices of the World Meteorological Organization. A major contribution of this Center is related to precipitation chemistry. The Center leads the global quality assurance program for precipitation chemistry, including a major focus on acid precipitation. This activity positions United States' science in a leadership role globally, and ensures that future decisions regarding long-range transport of pollutants from one continent to another will have a sound data base to guide them.

REFERENCES

Emissions and Compliance Costs

Bailey, E.M. 1996. *Allowance Trading Activity and State Regulatory Rulings: Evidence from the U.S. Acid Rain Program*. Center for Energy and Environmental Policy Research, Massachusetts Institute of Technology, Cambridge, MA.

Burtraw, D. 1996. The SO₂ emissions trading program: Cost savings without allowance trades. *Contemporary Economic Policy* (XIV April): 79-94.

Cason, T.N. 1993. Seller incentive properties of EPA's emission trading auction. *Journal of Environmental Economics and Management* 25:177-195.

Cason, T.N. 1994. An experimental investigation of the seller incentives in the EPA's emission trading auction. *American Economic Review* 85:4, 905-922.

Dean, M., and J. Kruger. 1997. Using EPA's Allowance Tracking System to Assess the Allowance Market. Presented at AWMA Acid Rain II Specialty Conference. Scottsdale, AZ.

Ellerman, A.D., and J.P. Montero. 1996. *Why are allowance prices so low?* Working Paper 96-001, Center for Energy and Environmental Policy Research, Massachusetts Institute of Technology, Cambridge, MA.

Ellerman, A.D. 1997 (forthcoming). *Calculating cost savings from emissions trading*. Center for Energy and Environmental Policy Research, Massachusetts Institute of Technology, Cambridge, MA.

Ellerman, A.D., R.L. Schmalensee, P. Joskow, J.P. Montero, and E. Bailey. 1997 (in preparation). *SO₂ Emissions Trading Under Title IV of the 1990 Clean Air Act Amendments: Evaluation of Compliance Costs and Allowance Market Performance*. Center for Energy and Environmental Policy Research, Massachusetts Institute of Technology, Cambridge, MA.

EPRI. 1993. *Integrated Analysis of Fuel, Technology and Emission Allowance Markets: Electric Utility Responses to the Clean Air Act Amendments of 1990*. Electric Power Research Institute, Palo Alto, CA. Table C-1.

EPRI. 1995. *The Emission Allowance Market and Electric Utility SO₂ Compliance in a Competitive and Uncertain Future*. Electric Power Research Institute, Palo Alto, CA. Table A-3.

EPRI. 1996. *Coal Supply and Transportation Markets During Phase One: Change Risk and Opportunity*. Electric Power Research Institute, Palo Alto, CA. TR-105916.

EPRI. 1997. *SO₂ compliance and allowance trading: Developments and outlook*. Electric Power Research Institute, Palo Alto, CA. TR-107897.

GAO. 1994. *Allowance Trading Offers an Opportunity to Reduce Emissions at Less Cost*. U.S. General Accounting Office, Washington, DC. pp. 36, 74.

- 1 Hahn, R.H., and C. May. 1994. The behavior of the allowance market: Theory and evidence.
2 *Electricity Journal* 7(2):28-37.
- 3 ICF Resources Incorporated. 1989. Economic Analysis of Title V (Acid Rain Provisions) of the
4 Administration's Proposed Clean Air Act Amendments (H.R. 3030/S. 1490). Based on "Flexible"
5 implementation case in Tables A1, A5, B1 and B5, plus mid-point estimate for banking from page
6 24.
- 7 ICF Resources Incorporated. 1990. Comparison of the Economic Impacts of the Acid Rain
8 Provisions of the Senate Bill (S.1630) and the House Bill (H.R.1630[sic]). Draft report prepared
9 for the U.S. Environmental Protection Agency, Washington, DC.
- 10 ICF Resources Incorporated. 1992. *Regulatory Impact Analysis of the Final Acid Rain*
11 *Implementation Regulations*. Prepared for the U.S. Environmental Protection Agency,
12 Washington, DC.
- 13 Joskow, P.L., R.L. Schmalensee, and E. Bailey. 1996. *Auction Design and the Market for*
14 *Sulfur Dioxide Allowances*. Center for Energy and Environmental Policy Research,
15 Massachusetts Institute of Technology, Cambridge, MA.
- 16 Kruger, J., and M. Dean. 1997. Looking back on SO₂ trading: What's good for the environment
17 is good for the market. *Public Utilities Fortnightly* 135(15):30-37.
- 18 Kruger, J., K. Jestin, and L. Wright. 1995. Catalyzing the alternative energy market: Incentives
19 from emissions trading. In: *Proceedings of the Air and Waste Management Association Annual*
20 *Meeting*. San Antonio, TX.
- 21 McLean, B.J. 1996. *The Sulfur Dioxide (SO₂) Allowance Trading Program: The First Five*
22 *Years*. U.S. Environmental Protection Agency, Washington, DC.
- 23 McLean, B.J. 1997. Evolution of marketable permits: The U.S. experience with sulfur dioxide
24 trading. In: *Proceedings from Controlling Carbon and Sulphur, International Trading Initiatives*.
25 Royal Institute of International Affairs, London, England.
- 26 NAPAP. 1991a. *1990 Integrated Assessment Report*. U.S. National Acid Precipitation
27 Assessment Program, Washington, DC.
- 28 NAPAP. 1991b. *Acidic Deposition: State of Science and Technology*. Volumes I-IV. U.S.
29 National Acid Precipitation Assessment Program, Washington, DC.
- 30 Placet, M., R.E. Battye, F.C. Fehsenfeld, G.W. Bassett, et al. 1991. Emissions involved in
31 acidic deposition. In: *Acidic Deposition: State of the Science and Technology*. Report 1. U.S.
32 National Acid Precipitation Assessment Program, Washington, DC.
- 33 U.S. EPA. 1994. *Energy Efficiency and Renewable Energy Opportunities from Title IV of the*
34 *Clean Air Act Amendments*. U.S. Environmental Protection Agency, Washington, DC. EPA-
35 430/R-94-001.
- 36 U.S. EPA. 1995a. *National Air Quality Trends Report*. U.S. Environmental Protection Agency,
37 Washington, DC.

- 1 U.S. EPA. 1995b. *National Air Pollutant Emission Trends, 1900-1994*. U.S. Environmental
2 Protection Agency, Washington, DC. EPA-454/R-95-011.
- 3 U.S. EPA. 1996a. *National Air Pollutant Emission Trends, 1900-1995*. U.S. Environmental
4 Protection Agency, Washington, DC. EPA-454/R-96-007.
- 5 U.S. EPA. 1996b. *1995 Emissions Scorecard*. U.S. Environmental Protection Agency,
6 Washington, DC.
- 7 U.S. EPA. 1997. *Acid Rain Program Emissions Scorecard 1995*. U.S. Environmental Protection
8 Agency, Washington, DC. EPA-430/R-97-009.
- 9 U.S. EPA. 1996c. *1995 Compliance Results*. U.S. Environmental Protection Agency,
10 Washington, DC. EPA-430/R-96-012.

11 Concentration and Deposition

- 12 Cahill, T.A., R.A. Eldred, and P.H. Wakabayashi. 1996. Trends in fine particle concentrations
13 at Great Smoky Mountains National Park. Presented at the 89th Annual Meeting of the Air and
14 Waste Management Association, Nashville, TN. Paper No. 96-MP1A.05
- 15 Hicks, B.B., R.P. Hosker, Jr., T.P. Meyers, and J.D. Womack. 1991. Dry deposition
16 inferential techniques -- I. Design and tests of a prototype meteorological and air chemical system
17 for determining dry deposition. *Atmospheric Environment* 25A:2345-2370.
- 18 Lynch, J.A., V.C. Bowersox, and J.W. Grimm. 1996. *Trends in Precipitation Chemistry in the*
19 *United States, 1983-94 -- An Analysis of the Effects in 1995 of Phase I of the Clean Air Act*
20 *Amendments of 1990, Title IV*. U.S. Geological Survey, Washington, DC. Open-File Report 96-
21 0346.
- 22 Malm, W.C., J. F. Sisler, D. Huffman, R.A. Eldred, and T.A. Cahill. 1994. Spatial and
23 seasonal trends in particle concentration and optical extinction in the United States. *Journal of*
24 *Geophysical Research* 99(D1):1347-1370.
- 25 Meyers, T.P., P. Finkelstein, J. Clarke, T.G. Ellestad, and P.F. Williams. 1997. Description and
26 evaluation of a multi-layer model for inferring dry deposition using standard meteorological
27 measurement. National Oceanic and Atmospheric Administration/ARL/ATDD, Oak Ridge, TN.
28 Report Number 97/2.
- 29 NADP. 1996. *NADP/NTN Wet Deposition in the United States 1995*. National Atmospheric
30 Deposition Program, Natural Resources Ecology Laboratory, Colorado State University, Fort
31 Collins, CO.
- 32
33 Sickles, J.E. II, D. Holland, P. Principe, R. Baumgardner, G. Evans, P. Finkelstein, J. Suggs,
34 B. Eder, S. LeDuc, E. Cooter, J. Pleim, J. Clarke, D. Schwede, R. McMillen, L.A. Byrd, T.
35 Ellestad, D. Filmore, L. Truppi, J. Shreffler, B. Bloomer, K. Endres, and R. Dennis. 1997.
36 Examination of CASTNet: Data, results, costs, and implications. (Preliminary Results). U.S.
37 Environmental Protection Agency, Research Triangle Park, NC.

REFERENCES

- 1 Sisterson, D.L., V.C. Bowersox, T.P. Meyers, A.R. Olsen, and R.J. Vong. 1991. Deposition
- 2 monitoring: Methods and results. In: *Acidic Deposition: State of Science and Technology*. Report
- 3 6. U.S. National Acid Precipitation Assessment Program, Washington, DC. pp. 6-104, 6-105, 6-
- 4 145 – 6-174, 6-220.

1 **Aquatic Ecosystems**

- 2 Aber, J.D., K.J. Nadelhoffer, P. Steudler, and J.M. Melillo. 1989. Nitrogen saturation in
3 northern forest ecosystems: Excess nitrogen from fossil fuel combustion may stress the biosphere.
4 *BioScience* 39(6):378-386.
- 5 Baker, J.P., D.P. Bernard, S.W. Christensen, M.J. Sale, J. Freda, K. Heltcher, D. Marmorek,
6 L. Rose, P. Scanlon, G. Suter, W. Warren-Hicks, and P. Welbourn. 1990. Biological effects of
7 changes in surface water acid-base chemistry. In: *Acidic Deposition: State of Science and*
8 *Technology*. Report 13. U.S. National Acid Precipitation Assessment Program, Washington, DC.
- 9 Baker, J.P., J. Van Sickle, C.J. Gagen, D.R. DeWalle, W.E. Sharpe, R.F. Carline, B.P.
10 Baldigo, P.S. Murdoch, D.W. Bath, W.A. Kretser, H.A. Simonin, and P.J. Wigington, Jr.
11 1996. Episodic acidification of small streams in the northeastern United States: Effects on fish
12 populations. *Ecological Applications* 6:422-437.
- 13 Baron, J.S., D.S. Ojima, E.A. Holland, and W.J. Parton. 1994. Analysis of nitrogen saturation
14 potential in Rocky Mountain tundra and forest: Implications for aquatic systems. *Biogeochemistry*
15 27:61-82.
- 16 Bulger, A.J., C.A. Dolloff, B.J. Cosby, K.N. Eshleman, J.R. Webb, and J.N. Galloway. 1995.
17 The "Shenandoah National Park: Fish in Sensitive Habitats" (SNP:FISH) Project. An integrated
18 assessment of fish community responses to stream acidification. *Water, Air, and Soil Pollution*
19 85:309-314.
- 20 Cook, R.B., J.W. Elwood, R.R. Turner, M.A. Bogle, P.J. Mulholland, and A.V. Palumbo.
21 1994. Acid-base chemistry of high-elevation streams in the Great Smoky Mountains. *Water, Air,*
22 *and Soil Pollution* 72:331-356.
- 23 Cosby, B.J., R.F. Wright, G.M. Hornberger, and J.N. Galloway. 1985. Modeling the effects of
24 acid deposition: Assessment of a lumped-parameter model of soil water and streamwater
25 chemistry. *Water Resources Research* 21:51-63.
- 26 Cosby, B.J., S.A. Norton, and J.S. Kahl. 1996. Using a paired-watershed manipulation
27 experiment to evaluate a catchment-scale biogeochemical model. *The Science of Total*
28 *Environment* 183:49-66.
- 29 Dennis, T.E., S.E. MacAvoy, M.B. Steg, and A.J. Bulger. 1995. The association of water
30 chemistry variables and fish condition in streams of Shenandoah National Park (USA). *Water,*
31 *Air, and Soil Pollution* 85:365-370.
- 32 Driscoll, C.T., M.D. Lehtinen, and T.J. Sullivan. 1994. Modeling the acid-base chemistry of
33 organic solutes in Adirondack New York lakes. *Water Resources Research* 30:297-306.
- 34 Driscoll, C.T., K.M. Postek, W. Kretser, and D.J. Raynal. 1995. Long-term trends in the
35 chemistry of precipitation and lake water in the Adirondack region of New York, U.S.A. *Water,*
36 *Air, and Soil Pollution* 85:583-588.
- 37 Fenn, M.E., M.A. Poth, and D.W. Johnson. 1996. Evidence for nitrogen saturation in the San
38 Bernardino Mountains in southern California. *Forest Ecology Management* 82:211-230.

- 1 Ferrier, R.C., A. Jenkins, B.J. Cosby, R.C. Helliwell, R.F. Wright, and A.J. Bulger. 1995.
2 Effects of future N deposition scenarios on the Galloway region of Scotland using a coupled
3 sulphur & nitrogen model (MAGIC-WAND). *Water, Air, and Soil Pollution* 85:707-712.
- 4 Gerritsen, J., J.M. Dietz, and H.T. Wilson, Jr. 1996. Episodic acidification of coastal plain
5 streams: An estimation of risk to fish. *Ecological Applications* 6:438-448.
- 6 Gilliam, F.S., M.B. Adams, and B.M. Yurish. 1995. Ecosystem nutrient responses to chronic
7 nitrogen inputs at Fernow Experimental Forest, West Virginia. *Canadian Journal of Forest*
8 *Research* 26:196-205.
- 9 Gjessing, E. 1992. The HUMEX Project: Experimental acidification of a catchment and its
10 humic lake. *Environment International* 18:535-543.
- 11 Gunn, J.M., ed. 1995. *Restoration and Recovery of an Industrial Region*. Springer-Verlag,
12 New York, NY.
- 13 Havas, M., D.G. Woodfine, P. Lutz, K. Yung, H.J. MacIsaac, and T.C. Hutchinson. 1995.
14 Biological recovery of two previously acidified, metal-contaminated lakes near Sudbury Ontario,
15 Canada. *Water, Air, and Soil Pollution* 85:791-796.
- 16 Hedin, L.O., L. Granat, G.E. Likens, T.A. Bulshand, J.N. Galloway, T.J. Butler, and H.
17 Rodhe. 1994. Steep declines in atmospheric base cations in regions of Europe and North
18 America. *Nature* 367:351-354.
- 19 Husar, R.B., T.J. Sullivan, and D.F. Charles. 1991. Historical trends in atmospheric sulfur
20 deposition and methods for assessing long-term trends in surface water chemistry. In: *Acidic*
21 *Deposition and Aquatic Ecosystems. Regional Case Studies*. D.F. Charles, ed. Springer-Verlag,
22 New York, NY. pp. 65-82.
- 23 Jackson, D.A., and H.H. Harvey. 1995. Gradual reduction and extinction of fish populations in
24 acid lakes. *Water, Air, and Soil Pollution* 85:389-394.
- 25 Jenkins, A., R.F. Wright, N. Van Breemen, E.D. Schulze, F. Berendse, F.I. Woodward, and L.
26 Brussard. 1992. The CLIMEX Project - Climate change experiment. International Symposium
27 on Experimental Manipulations of Biota and Biogeochemical Cycling in Ecosystems - Approach,
28 Methodologies, Findings. Copenhagen, Denmark.
- 29 Kahl, J.S., S.A. Norton, I.J. Fernandez, K.J. Nadelhoffer, C.T. Driscoll, and J.D. Aber. 1993.
30 Experimental inducement of nitrogen saturation at the watershed scale. *Environmental Science and*
31 *Technology* 27:565-568.
- 32 Kauppi, P.E., K. Mielikäinen, and K. Kuusela. 1992. Biomass and carbon budget of European
33 forests, 1971 to 1990. *Science* 256:70-74.
- 34 Kirchner, J.W., and E. Lydersen. 1995. Base cation depletion and potential long-term
35 acidification of Norwegian catchments. *Environmental Science and Technology* 29:1953-1960.
- 36 Likens, G.E., C.T. Driscoll, and D.C. Buso. 1995. Long-term effects of acid rain: Response and
37 recovery of a forest ecosystem. *Science* 272:244-246.

- 1 MacAvoy, S.E., and A.J. Bulger. 1995. Survival of brook trout (*Salvelinus fontinalis*) embryos
2 and fry in streams of different acid sensitivity in Shenandoah National Park, USA. *Water, Air,
3 and Soil Pollution* 85:445-450.
- 4 Marcus, M.D., and D.R. Mount, eds. 1991. Physiological, toxicological, and population
5 responses of rainbow trout to acidification. Interim Report of the Lake Acidification and Fisheries
6 Project. Electric Power Research Institute, Palo Alto, CA. EPRI EN-7114.
- 7 McNicol, D.K., B.E. Bendell, and M.L. Mallory. 1995a. Evaluating macroinvertebrate
8 responses to recovery from acidification in small lakes in Ontario, Canada. *Water, Air, and Soil
9 Pollution* 85:451-456.
- 10 McNicol, D.K., M.L. Mallory, and H.S. Vogel. 1995b. Using volunteers to monitor the effects
11 of acid precipitation on common loon (*Gavia immer*) reproduction in Canada: The Canadian Lakes
12 Loon Survey. *Water, Air, and Soil Pollution* 85:463-468.
- 13 Mitchell, M.J., C.T. Driscoll, J.S. Kahl, G.E. Likens, P.S. Murdoch, and L.H. Pardo. 1996.
14 Climatic control of nitrate loss from forested watersheds in the northeast United States.
15 *Environmental Science and Technology* 30:2609-2612.
- 16 Murdoch, P.S., and J.L. Stoddard. 1992. The role of nitrate in the acidification of streams in the
17 Catskill Mountains of New York. *Water Resources Research* 28:2707-2720.
- 18 NAPAP. 1991. *1990 Integrated Assessment Report*. U.S. National Acid Precipitation
19 Assessment Program, Washington, DC.
- 20 Norton, S.A., R.F. Wright, J.S. Kahl, and J.P. Scofield. 1992. The MAGIC simulation of
21 surface water at, and first year results from, the Bear Brook Watershed Manipulation, Maine,
22 USA. *Environmental Pollution* 77:279-286.
- 23 Oreskes, N., K. Shrader-Frechette, and K. Belitz. 1994. Verification, validation, and
24 confirmation of numerical models in the earth sciences. *Science* 263:641-646.
- 25 Peterjohn, W.T., M.B. Adams, and F.S. Gilliam. 1996. Symptoms of nitrogen saturation in two
26 central Appalachian hardwood forest ecosystems. *Biogeochemistry* 35:507-522.
- 27 Schofield, C.L., and C. Keleher. 1996. Comparison of brook trout reproductive success and
28 recruitment in an acidic Adirondack lake following whole lake liming and watershed liming.
29 *Biogeochemistry* 32:323-337.
- 30 Sisterson, D.L., V.C. Bowersox, T.P. Meyers, A.R. Olsen, and R.J. Vong. 1991. Deposition
31 monitoring: Methods and results. In: *Acidic Deposition: State of Science and Technology*. Report
32 6. U.S. National Acid Precipitation Assessment Program, Washington, DC. pp. 6-104, 6-105, 6-
33 145 – 6-174, 6-220.
- 34 Stoddard, J.L. 1994. Long-term changes in watershed retention of nitrogen: Its causes and
35 aquatic consequences. In: *Environmental Chemistry of Lakes and Reservoirs*. L.A. Baker, ed.
36 Advances in Chemistry. Series, No. 237. American Chemical Society, Washington, DC. pp. 223-
37 284.

- 1 Stoddard, J.L., C.T. Driscoll, S. Kahl, and J. Kellogg. 1997. Can site-specific trends be
2 extrapolated to the regional level? A lake acidification example for the northeastern U.S.
3 *Ecological Applications*.
- 4 Sullivan, T.J., and B.J. Cosby. 1995. Testing, improvement, and confirmation of a watershed
5 model of acid-base chemistry. *Water, Air, and Soil Pollution* 85:2607-2612.

- 1 Sullivan, T.J., and J.M. Eilers. 1994. Assessment of Deposition Levels of Sulfur and Nitrogen
2 Required to Protect Aquatic Resources in Selected Sensitive Regions of North America. Final
3 report prepared for Technical Resources, Inc., Rockville, MD under Contract to U.S.
4 Environmental Protection Agency, Environmental Research Laboratory-Corvallis, Corvallis, OR.
- 5 Sullivan, T.J., B. McMartin, and D.F. Charles. 1996. Re-examination of the role of landscape
6 change in the acidification of lakes in the Adirondack Mountains, New York. *Scientific Total*
7 *Environment* 183(3):231-248.
- 8 Sullivan, T.J., J.A. Bernert, E.A. Jenne, J.M. Eilers, B.J. Cosby, D.F. Charles, and A.R. Selle.
9 1991. *Comparison of MAGIC and Diatom Paleolimnological Model Hindcasts of Lakewater*
10 *Acidification in the Adirondack Region of New York*. U.S. Department of Energy, Pacific
11 Northwest Laboratory, Richland, WA.
- 12 Sullivan, T.J., B.J. Cosby, C.T. Driscoll, H.F. Hemond, D.F. Charles, S.A. Norton, H.M.
13 Seip, and G. Taugbøl. 1994. *Confirmation of the MAGIC Model Using Independent Data:*
14 *Influence of Organic Acids on Model Estimates of Lakewater Acidification*. U.S. Department of
15 Energy, Washington, DC. DOE/ER/30196-4
- 16 Sullivan, T.J., B.J. Cosby, C.T. Driscoll, D.F. Charles, and H.F. Hemond. 1996. Influence of
17 organic acids on model projections of lake acidification. *Water, Air, and Soil Pollution* 91:271-
18 282.
- 19 Tietema, A., and C. Beier. 1995. A correlative evaluation of nitrogen cycling in the forest
20 ecosystems of the EC projects NITREX and EXMAN. *Forest Ecology and Management* 71:143-
21 151.
- 22 Webb, J.R., F.A. Devincy, J.N. Galloway, C.A. Rinehart, P.A. Thompson, and S. Wilson.
23 1994. The acid-base status of native brook trout streams in the mountains of Virginia. University
24 of Virginia, Charlottesville, VA.
- 25 Wigington, P.J. Jr., T.D. Davies, M. Tranter, and K.N. Eshleman. 1991. Episodic Acidification
26 of Surface Waters Due to Acidic Deposition. In: *Acidic Deposition: State of Science and*
27 *Technology*. Report 12. U.S. National Acid Precipitation Program, Washington, DC.
- 28 Wigington, P.J. Jr., J.P. Baker, D.R. DeWalle, W.A. Kretser, P.S. Murdoch, H.A. Simonin, J.
29 Van Sickle, M.K. McDowell, D.V. Peck, and W.R. Barchet. 1993. *Episodic Acidification of*
30 *Streams in the Northeastern United States: Chemical and Biological Results of the Episodic*
31 *Response Project*. U. S. Environmental Protection Agency, Corvallis, OR. EPA/600/R-93/190.
- 32 Wigington, P.J. Jr., D.R. DeWalle, P.S. Murdoch, W.A. Kretser, H.A. Simonin, J. Van Sickle,
33 and J.P. Baker. 1996. Episodic acidification of small streams in the northeastern United States:
34 Ionic controls of episodes. *Ecological Applications* 6(2):389-407.
- 35 Williams, M.W., J.S. Baron, N. Caine, R. Sommerfeld, and R. Sanford, Jr. 1996. Nitrogen
36 saturation in the Rocky Mountains. *Environmental Science and Technology* 30(2):640-646.

- 1 Wright, R.F., E. Lotse, and A. Semb. 1993. RAIN Project: Results after 8 years of
2 experimentally reduced acid deposition to a whole catchment. *Canadian Journal of Fish and*
3 *Aquatic Science* 50:258-268.
- 4 Yan, N.D., W. Keller, K.M. Somers, T.W. Pawson, and R.E. Girard. 1996a. Recovery of
5 crustacean zooplankton communities from acid and metal contamination: Comparing manipulated
6 and reference lakes. *Canadian Journal of Fish and Aquatic Science* 53:1301-1327.
- 7 Yan, N.D., P.G. Welsh, H. Lin, D.J. Taylor, and J.M. Fillion. 1996b. Demographic and genetic
8 evidence of the long-term recovery of *Daphnia galeata mendotae* (Crustacea: Daphniidae) in
9 Sudbury lakes following additions of base: The role of metal toxicity. *Canadian Journal of Fish*
10 *and Aquatic Science* 53:1328-1344.
- 11 **Forest Ecosystems**
- 12 Adams, M.B., T.R. Angradi, and J.N. Kochenderfer. 1997. Stream water and soil solution
13 responses to five years of nitrogen and sulfur additions at the Fernow experimental forest, West
14 Virginia. *Forest Ecology and Management* 95(1):79-91.
- 15 Allen, H.L. 1987. Forest fertilizers, nutrient amendment, stand productivity and environmental
16 impact. *Journal of Forestry* 85:37-46.
- 17 Amundson, R.G., J.L. Hadley, J.F. Fincher, S. Fellows, and R.G. Alscher. 1992.
18 Comparisons of seasonal changes in photosynthetic capacity, pigments, and carbohydrates of
19 healthy sapling and mature red spruce and/or declining and healthy red spruce. *Canadian Journal*
20 *of Forest Research* 22:1605-1616.
- 21 Anderson, C.P., McLaughlin, S.B., and Roy, W.K. 1991. Foliar injury symptoms and pigment
22 concentrations in red spruce saplings in the southern Appalachians. *Canadian Journal of Forest*
23 *Research* 21:1119-1123.
- 24 Binkley, D., C.T. Driscoll, H.L. Allen, P. Schoeneberger, and D. McAvoy. 1989. *Acidic*
25 *Deposition and Forest Soils*. Ecological Studies 72. Springer-Verlag, New York, NY.
- 26 Bytnerowicz, A., and M.E. Fenn. 1996. Nitrogen deposition in California forests: A review.
27 *Environmental Pollution* 92:127-146.
- 28 Cape, J.N., I.D. Leith, D. Fowler, M.B. Murray, L.J. Sheppard, D. Eamus, R.H.F. Wilson.
29 1991. Sulphate and ammonium in mist impair the frost hardiness of red spruce seedlings. *New*
30 *Phytology* 118:119-126.
- 31 Craig, B.W., and A.J. Friedland. 1991. Spatial patterns in forest composition and standing dead
32 red spruce in montane forests of the Adirondacks and northern Appalachians. *Environmental*
33 *Monitoring and Assessment* 18:129-143.
- 34 Cronan, C.S., and D.F. Grigal. 1995. Use of calcium/aluminum ratios as indicators of stress in
35 forest ecosystems. *Journal of Environmental Quality* 24:209-226.

- 1 DeHayes, D.H. 1992. Winter injury and developmental cold tolerance of red spruce. In: *Ecology*
2 *and Decline of Red Spruce in the Eastern United States*. C. Eager and M.B. Adams, eds.
3 Springer-Verlag, New York, NY. pp 295-337.
- 4 DeHayes, D.H., F.C. Thornton, C.E. Waite, and M.A. Ingle. 1991. Ambient cloud deposition
5 reduces cold tolerance of red spruce seedlings. *Canadian Journal of Forest Research* 21:1292-
6 1295.
- 7 DeHayes, D.H., C.E. Waite, M.A. Ingle, and M.A. Williams. 1990. Winter injury susceptibility
8 and cold tolerance of current and year old red spruce trees from several provenances. *Forest*
9 *Science* 36:982-994.
- 10 Desanker, P.V., D.D. Reed, and E.A. Jones. 1994. Evaluating forest stress factors using various
11 forest growth modeling approaches. *Forest Ecology and Management* 69:269-282.
- 12 Dougherty, P.M. 1996. Response of loblolly pine to moisture and nutrient stress. In: *Impact of*
13 *Air Pollutants on Southern Pine Forests*. C. Eager and M. B. Adams, eds. Ecological Studies
14 118. Springer-Verlag, New York, NY. pp. 173-195.
- 15 Eagar, C., and M.B. Adams, eds. 1992. *Ecology and Decline of Red Spruce in the eastern U.S.*
16 Springer-Verlag, New York, N.Y.
- 17 Fenn, M. 1991. Increased site fertility and litter decomposition rate in high-pollution sites in the
18 San Bernardino Mountains. *Forest Science* 37:1163-1181.
- 19 Fenn, M.E., and A. Bytnerowicz. 1993. Dry deposition of nitrogen and sulfur to ponderosa pine
20 and Jeffrey pine in the San Bernardino National Forest in southern California. *Environmental*
21 *Pollution* 81:277-285.
- 22 Fernandez, I.J. 1992. Characterization of eastern U.S. spruce-fir soils. In: *Ecology and Decline*
23 *of Red Spruce in the Eastern United States*. C. Eagar and M.B. Adams, eds. Springer-Verlag,
24 New York. pp. 40-63.
- 25 Flagler, R.B., and A.H. Chappelka. 1996. Growth responses of southern pines to acidic
26 deposition and ozone. In: *Impact of Air Pollutants on Southern Pine Forests*. S.A. Fox and R.A.
27 Mickler, eds. Springer-Verlag, New York, NY. pp 388-424.
- 28 Fowler, D., J.N. Cape, J.D. Deans, I.D. Leath, et al. 1989. Effects of acid mist on frost
29 hardiness of red spruce seedlings. *New Phytology* 113:321-335.
- 30 Gilliam, F.S., M.B. Adams, and B.M. Yrish. 1996. Ecosystem nutrient responses to chronic
31 nitrogen inputs at Fernow Experimental Forest, West Virginia. *Canadian Journal of Forest*
32 *Research* 26:196-205.
- 33 Johnson, A.H., S.B. McLaughlin, M.B. Adams, E.R. Cook, D.H. DeHayes, C. Eagar, I.J.
34 Fernandez, D.W. Johnson, R.J. Kohut, V.A. Mohnen, N.S. Nicholas, D.R. Peart, G.A. Schier,
35 and P.S. White. 1992. Synthesis and conclusions from epidemiological and mechanistic studies
36 of red spruce decline. In: *Ecology and Decline of Red Spruce in the Eastern United States*. C. Eagar
37 and M.B. Adams, eds. Springer-Verlag, New York, NY.

- 1 Johnson, D.W., H. Van Miegroet, S.E. Lindberg, R.B. Harrison, and D.E. Todd. 1991.
2 Nutrient cycling in red spruce forests of the Great Smoky Mountains. *Canadian Journal of Forest*
3 *Research* 21:767-787.
- 4 Johnson, D.W., W.T. Swank, and J.M. Vose. 1993. Simulated effects of atmospheric sulfur
5 deposition on nutrient cycling in a mixed deciduous forest. *Biogeochemistry* 23:169-196.
- 6 Johnson, D.W., R.B. Susfalk, and P.F. Brewer. 1996. Simulated responses of red spruce forest
7 soils to reduced sulfur and nitrogen deposition. *Journal of Environmental Quality* 25:1300-1309.
- 8 Joslin, J.D., and M.H. Wolfe. 1993. Temperature increase accelerates nitrate release from high-
9 elevation red spruce soils. *Canadian Journal of Forest Research* 23:756-759.
- 10 Joslin, J.D., and M.H. Wolfe. 1994. Foliar deficiencies of mature southern Appalachian red
11 spruce determined from fertilization trials. *Soil Science Society of America Journal* 58:1572-1579.
- 12 Joslin, J.D., C.M. McDuffie, and P.F. Brewer. 1988. Acidic cloudwater and cation loss from red
13 spruce foliage. *Water, Air, and Soil Pollution* 39:355-363.
- 14 Joslin, J.D., J.M. Kelly, and H. Van Miegroet. 1992. Soil chemistry and nutrition of North
15 American spruce-fir stands: Evidence for recent change. *Journal of Environmental Quality* 21:12-
16 30.
- 17 Kiefer, J.W., and M.E. Fenn. 1997. Using vector analysis to assess nitrogen status of ponderosa
18 and Jeffrey pine along deposition gradients in forests of southern California. *Forest Ecology and*
19 *Management* 94(1-3):47-59.
- 20 Lawrence, G.B., M.B. David, and W.C. Shortle. 1995. A new mechanism for calcium loss in
21 forest-floor soils. *Nature* 378:162-165.
- 22 Lawrence, G.B., M.B. David, S.W. Bailey, and W.C. Shortle. 1997. Assessment of calcium
23 status in soils of red spruce forests in the northeastern United States. *Biogeochemistry* 38:19-39.
- 24 Likens, G.E., C.T. Driscoll, and D.C. Buso. 1996. Long-term effects of acid rain: Response and
25 recovery of a forest ecosystem. *Science* 272:244-246.
- 26 MacDonald, N.W., A.J. Burton, H.O. Liechty, J.A. Witter, K.S. Pregitzer, G.D. Mroz, and
27 D.D. Richter. 1992. Ion leaching in forest ecosystems along a Great Lakes air pollution gradient.
28 *Journal of Environmental Quality* 21:614-623.
- 29 McLaughlin, S.B., and D.J. Downing. 1995. Interactive effects of ambient ozone and climate
30 measured on growth of mature trees. *Nature* 374:252-254.
- 31 McLaughlin, S.B., and D.J. Downing. 1996. Interactive effects of ambient ozone and climate
32 measured on growth of mature loblolly pine trees. *Canadian Journal of Forest Research* 26:670-
33 681.
- 34 McLaughlin, S.B., and R. Kohut. 1992. The effects of atmospheric deposition on carbon
35 allocation and associated physiological processes in red spruce. In: *Ecology and Decline of Red*

- 1 *Spruce in the Eastern United States*. C. Eagar and B. Adams, eds. Springer-Verlag, New York,
2 NY. pp. 338-384.
- 3 McLaughlin, S.B., M.G. Tjoelker, and W.K. Roy. 1993. Acid deposition alters red spruce
4 physiology: Laboratory studies support field observations. *Canadian Journal of Forest Research*
5 23:380-386.
- 6 McLaughlin, S.B., D.J. Downing, T.J. Blasing, E.R. Cook, and H.S. Adams. 1987. An analysis
7 of climate and competition as contributors to decline of red spruce in high-elevation Appalachian
8 forests of the eastern United States. *Oecologia* 72:487-501.
- 9 McLaughlin, S.B., C.P. Andersen, P.J. Hanson, M.G. Tjoelker, and W.K. Roy, 1991.
10 Increased dark respiration and calcium deficiency of red spruce in relation to acidic deposition at
11 high-elevation southern Appalachian Mountain sites. *Canadian Journal of Forest Research*
12 21:1234-1244.
- 13 McLaughlin, S.B., P.A. Layton, M.B. Adams, N.T. Edwards, P.J. Hanson, E.G. O'Neill, and
14 W.K. Roy. 1994. Growth responses of 53 open pollinated loblolly pine families to ozone and
15 acid rain. *Journal of Environmental Quality* 23:247-257.
- 16 McLaughlin, S.B., J.D. Joslin, A. Stone, R. Wimmer, and S. Wullschlegel. 1997 (in press).
17 The impact of global change on high-elevation spruce/fir forests. In: *The Productivity and*
18 *Sustainability of Southern Forest Ecosystems in a Changing Environment*. R.A. Mickler and S.
19 Fox, eds. Springer-Verlag, New York, NY.
- 20 Mickler, R.A., and S. Fox, eds. 1997. *The Productivity and Sustainability of Southern Forest*
21 *Ecosystems in a Changing Environment*. Springer-Verlag, New York, NY.
- 22 Miller, P.R. 1992. Mixed conifer forests of the San Bernardino Mountains, California. In: *The*
23 *Response of Western Forests to Air Pollution*. R.K. Olson, D. Binkley, and M. Bohm, eds.
24 Springer-Verlag, New York, NY.
- 25 Minocha, R., W.C. Shortle, D.J. Coughlin, Jr., and S.C. Minocha. 1996. Effects of aluminum
26 on growth, polyamine metabolism, and inorganic ions in suspension cultures of red spruce (*Picea*
27 *rubens*). *Canadian Journal for Forest Research* 26:550-559.
- 28 Mitchell, M.J., C.T. Driscoll, J.H. Porter, D.J. Raynal, D. Schaefer, and E.H. White. 1994.
29 The Adirondack manipulation and modeling project (AMMP): Design and preliminary results.
30 *Forest Ecology and Management* 68:87-100.
- 31 Mitchell, M.J., D.J. Raynal, and C.T. Driscoll. 1996a. Biogeochemistry of a forested watershed
32 in the central Adirondack Mountains: Temporal changes and mass balances. *Water, Air, and Soil*
33 *Pollution* 88:355-369.
- 34 Mitchell, M.J., C.T. Driscoll, J.S. Kahl, G.E. Likens, P.S. Murdoch, and L.H. Pardo. 1996b.
35 Climate control on nitrate loss from forested watersheds in the Northeast United States.
36 *Environmental Science and Technology* 30: 2609-2612.

- 1 Neff, J.C., W.D. Bowman, E.A. Holland, M.C. Fisk, and S.K. Schmidt. 1994. Fluxes of
2 nitrous oxide and methane from nitrogen-amended soils in a Colorado alpine ecosystem.
3 *Biogeochemistry* 27:23-33.
- 4 Nicholas, N.S., and S.M. Zedaker. 1989. Ice damage in spruce-fir forests of the Black
5 Mountains, North Carolina. *Canadian Journal of Forest Research* 19:1487-1491.
- 6 Norton, S.A., J.S. Kahl, I.J. Fernandez, L.E. Rustad, J.P. Scofield, and T.A. Haines. 1994.
7 Response of the West Bear Brook Watershed, Maine, USA, to the addition of $(\text{NH}_4)_2\text{SO}_4$: 3-year
8 results. *Forest Ecology and Management* 68:61-73.
- 9 Peart, D.R., N.S. Nicholas, S.M. Zedaker, M.M. Miller-Weeks, and T.G. Siccama. 1992.
10 Condition and recent trends in high-elevation red spruce populations. In: *Ecology and Decline of*
11 *Red Spruce in the Eastern United States*. C. Eagar and M.B. Adams, eds. Springer-Verlag, New
12 York, NY. pp. 125-191.
- 13 Peterjohn, W.T., M.B. Adams, and F.S. Gilliam. 1996. Symptoms of nitrogen saturation in two
14 central Appalachian hardwood forest ecosystems. *Biogeochemistry* 35:507-522.
- 15 Richter, D.D., and D. Markewitz. 1995. Atmospheric deposition and forest resources of the
16 Southern Pine Forest. In: *Impact of Air Pollutants on Southern Pine Forests*. S.A. Fox and R.A.
17 Mickler, eds. Springer-Verlag, New York, NY. pp 315-336
- 18 Robarge, W.P., and D.W. Johnson. 1992. The effects of acidic deposition on forested soils.
19 *Advances in Agronomy* 47:1-81.
- 20 Rustad, L.E., I.J. Fernandez, M.B. David, M.J. Mitchell, K.J. Nadelhoffer, and R.B. Fuller.
21 1996. Experimental soil acidification and recovery at the Bear Brook Watershed in Maine. *Soil*
22 *Science Society of America Journal* 60:1933-1943.
- 23 Sasser, C.L., and D. Binkley. 1989. Nitrogen mineralization in high elevation of the
24 Appalachians. II. Patterns with stand development in fir waves. *Biogeochemistry* 7:147-156.
- 25 Sheppard, L.J. 1994. Causal mechanisms by which sulphate, nitrate and acidity influence frost
26 hardiness in red spruce: Review and hypothesis. *New Phytology* 127:69-82.
- 27 Sheppard, L.J., J.N. Cape, and I.D. Leith. 1993a. Influence of acidic mist on frost hardiness
28 and nutrient concentrations in red spruce seedlings, Part I: Exposure of the foliage and rooting
29 environment. *New Phytology* 124:595-605.
- 30 Sheppard, L.J., J.N. Cape, and I.D. Leith. 1993b. Influence of acidic mist on frost hardiness
31 and nutrient concentrations in red spruce seedlings, Part II: Frequency and dose. *New Phytology*
32 124:607-615.
- 33 Shortle, W.C., K.T. Smith, R. Minocha, G.B. Lawrence, and M.B. David. 1997. Acidic
34 deposition, cation mobilization, and biochemical indicators of stress in healthy red spruce. *Journal*
35 *of Environmental Quality* 26(3):871-876.

- 1 Sievering, H., D. Rusch, and L. Marquez. 1996. Nitric acid, particulate nitrate and ammonium in
2 the continental free troposphere: Nitrogen deposition to an alpine tundra ecosystem. *Atmospheric*
3 *Environment* 30:2527-2537.
- 4 Teskey, R.O. 1995. Synthesis and conclusions from studies of southern pines. In: *Impact of Air*
5 *Pollutants on Southern Pine Forests*. S.A. Fox and R.A. Mickler, eds. Springer-Verlag, New
6 York, NY. pp. 467-490.
- 7 Thornton, F.C., J.D. Joslin, P.A. Pier, H. Neufeld, J.R. Seiler, and J.D. Hutcherson. 1994.
8 Cloudwater and ozone effects upon high-elevation red spruce: A summary of study results from
9 Whitetop Mountain, Virginia. *Journal of Environmental Quality* 23:1158-1167.
- 10 Turner, R.S., R.J. Olson, and C.C. Brandt. 1986. Areas Having Soil Characteristics That May
11 Indicate Sensitivity to Acidic Deposition Under Alternate Forest Damage H. U.S. Department of
12 Energy, Oak Ridge National Laboratory, Oak Ridge, TN. TM 9917.
- 13 Van Miegroet, H., D.W. Cole, and N.W. Foster. 1992. Nitrogen distribution and cycling. In:
14 *Atmospheric Deposition and Forest Nutrient Cycling*. D.W. Johnson and S.E. Lindberg, eds.
15 Springer-Verlag, New York, NY. pp. 178-196.
- 16 Van Miegroet, H.V., D.W. Johnson, and D.E. Todd. 1993. Foliar response of red spruce
17 saplings to fertilization with Ca and Mg in the Great Smoky Mountains National Park. *Canadian*
18 *Journal of Forest Research* 23:89-95.
- 19 Vann, D.R., G.R. Strimbeck, and A.H. Johnson. 1992. Effects of ambient levels of airborne
20 chemicals on freezing resistance of red spruce foliage. *Forest Ecology and Management* 51:69-
21 79.
- 22 Williams, M.W., J.S. Baron, N. Caine, R. Sommerfield, and R. Sanford, Jr. 1996a. Nitrogen
23 saturation in the Rocky Mountains. *Environmental Science and Technology* 30:640-646.
- 24 Williams, M.W., P.D. Brooks, A. Mosier, and K.A. Tonnessen. 1996b. Mineral nitrogen
25 transformations in and under seasonal snow in a high-elevation catchment in the Rocky Mountains,
26 United States. *Water Resources Research* 32:3161-3171.
- 27 Williams, M.W., M. Losleben, N. Caine, and D. Greenland. 1996c. Changes in climate and
28 hydrochemical responses in a high-elevation catchment in the Rocky Mountains, USA. *Limnology*
29 *and Oceanography* 45:939-946.
- 30 **Materials and Cultural Resources**
- 31 Haynie, F.H., and T.J. Lemmons. 1990. *Aerosol Science and Technology* 13:356-367.
- 32 McGee, E.S. 1996. *Development of Gypsum Alteration on Marble and Limestone*. American
33 Society for Testing and Materials, West Conshohocken, PA. Standard Technical Publication
34 1258.
- 35 Meierding, T.C. 1993. Marble tombstone weathering and air pollution in North America. *Annals*
36 *of the Association of American Geographers* 83(4):568-588.

- 1 Mossotti, V.G., A.R. Eldeeb, T.L. Fries, V.N.Naudé, L. Soderberg, and G.S. Wheeler. 1997
2 (in preparation). Effect of cleaning techniques on Berkshire Lee marble. Case study: Philadelphia
3 City Hall, East Center Pavilion Facade. U.S. Geological Survey, Washington DC. Digital Data
4 Series DDS-xx-87 (CD-ROM).
- 5 Reimann, K.J. 1994. Color measurements on marble and limestone briquettes exposed to
6 outdoor environment in the eastern United States. Volume I: Results of exposure 1984-1990.
7 Argonne National Laboratory, Argonne, IL. Open file report ANL-94/27.
- 8 Reimann, K.J. 1995a. Weight losses of marble and limestone briquettes exposed to outdoor
9 environment in the eastern United States: Results of exposure 1998-1992. Argonne National
10 Laboratory, Argonne, IL. Open file report ANL-95/22.
- 11 Reimann, K.J. 1995b. Chemical alteration of limestone and marble samples exposed to acid rain
12 and weathering in the eastern United States, 1988-1992. Argonne National Laboratory, Argonne,
13 IL. Open file report ANL-95/xx.
- 14 Spence, J.W., F.H. Haynie, F.W. Lipfert, S.D. Cramer, and L.G. McDonald. 1992. *Corrosion*
15 48:1009-1019.
- 16 Spence, J.W., and J.N. McHenry. 1994. *Atmospheric Environment* 28 (18):3033-3046.
- 17 Spiker, E., R.P. Hosker Jr., V.J. Comer, J.R. White, R.W. Werre Jr., F.L. Harmon, G.D.
18 Dandy, and S.I. Sherwood. 1992. Environmental chamber for study of the deposition flux of
19 gaseous pollutants to material surfaces. *Atmospheric Environment* 26a(16):2885-2892.
- 20 Spiker, E., R.P. Hosker Jr., V.C. Weintraub, and S.I. Sherwood. 1995. Laboratory study of
21 sulfur dioxide dry deposition on limestone and marble: Effects of humidity and surface variables.
22 *Water, Air, and Soil Pollution*.
- 23 **Visibility**
- 24 Cahill, T.A., R.A. Eldred, and P.H. Wakabayashi. 1996. Trends in fine particle concentrations at
25 Great Smoky Mountains National Park. Presented at the 89th Air and Waste Management
26 Association Annual Meeting in Nashville, TN. Paper No. 96-MP1A.05.
- 27 Eldred, R.A., and T.A. Cahill. 1994. Trends in elemental concentration of fine particles at remote
28 sites in the United States. *Atmospheric Environment* 28(5):1009-1019.
- 29 Malm, W.C., J.F. Sisler, D. Huffman, R.A. Eldred, and T.A. Cahill. 1994. Spatial and seasonal
30 trends in particle concentration and optical extinction in the United States. *Journal of Geophysical*
31 *Research* 99(D1):1347-1370.
- 32 Pitchford, M.L., and W.C. Malm. 1994. Development and application of a standard visual index.
33 *Atmospheric Environment* 28(5):1049-1054.
- 34 Sisler, J.F., W.C. Malm, K.A. Gebhart, and M.L. Pitchford. 1996. Spatial and seasonal
35 patterns and long-term variability of the composition of the haze in the United States: An analysis
36 of data from the IMPROVE Network. Report by the Cooperative Institute for Research in the
37 Atmosphere, Colorado State University, Fort Collins, CO.

- 1 and Technology. Report 27. U.S. National Acid Precipitation Assessment Program, Washington,
2 DC.
- 3 Chestnut, L.G., and R.L. Dennis. 1997. Economic benefits of improvements in visibility: Acid
4 rain provisions of the 1990 Clean Air Act Amendments. *Journal of Air and Waste Management*
5 *Association* 47:395-402.
- 6 Chestnut, L.G., and R.D. Rowe, 1990. *Preservation Values for Visibility Protection at the*
7 *National Parks*. U.S. Environmental Protection Agency, Research Triangle Park, NC.
- 8 Electric Power Research Institute and Decision Focus, Inc. 1991. *Analysis of Alternative SO₂*
9 *Reduction Strategies*. Electric Power Research Institute, Palo Alto, CA. EN/GS-7132.
- 10 Englin, J.E., et al. 1991. Valuation of Damages to Recreational Trout Fishing in the Upper
11 Northeast Due to Acidic Deposition. Pacific Northwest Laboratory, Seattle, WA.
- 12 Johannesson, M., and P. Johannesson. 1996. To be or not to be, that is the question: An empirical
13 study of the willingness-to-pay for an increased life expectancy at an advanced age. *Journal of*
14 *Risk and Uncertainty* 13(2):163-174.
- 15 Jones-Lee, M.W., M. Hammerton, and P.R. Philips. 1985. The value of safety: Results of a
16 national sample survey. *The Economic Journal* 95(March):49-72.
- 17 McClelland, G., et al. 1991. Valuing eastern visibility: A field test of the contingent valuation.
18 U.S. Environmental Protection Agency, Washington, DC. Draft.
- 19 Mitchell, R.C., and R.T. Carson. 1986. *Valuing Drinking Water Risk Reductions Using the*
20 *Contingent Valuation Method: A Methodological Study of Risks from THM and Giardia*. U.S.
21 Environmental Protection Agency, Washington, DC.
- 22 Morey, E., K. Rossman, L. Chestnut, and S. Ragland. 1997. Valuing Acid Deposition Injuries
23 to Cultural Resources. U.S. National Acid Precipitation Assessment Program, Washington DC.
- 24 RFF. 1997. The costs and benefits of Title IV: An integrated assessment using TAF. Resources
25 for the Future, Washington, DC. Discussion Paper 97-31.
- 26 Tolley, G.A., et al. 1986. *Establishing and Valuing the Effects of Improved Visibility in the*
27 *Eastern United States*. U.S. Environmental Protection Agency, Washington, DC.
- 28 U.S. EPA. 1995. *Human Health Benefits from Sulfate Reductions Under Title IV of the 1990*
29 *Clean Air Act Amendments*. U.S. Environmental Protection Agency, Office of Air and Radiation,
30 Washington, DC.
- 31 U.S. EPA. 1997. *The Benefits and Costs of the Clean Air Act 1970 to 1990: Report to Congress*
32 (also known as Section 812 Retrospective Study). U.S. Environmental Protection Agency,
33 Washington, DC.
- 34 Walsh, R.G., et al. 1990. Nonmarket values from two decades of research on recreation demand.
35 *Advances in Applied Micro-Economics* 5:167-193.

APPENDICES

- A. Integrated Models (TAF, RAISON, RAINS)
Process Models (MAGIC)**
- B. Design and Performance of Pollution Trading Programs**
- C. Interpretation of "Adverse" Effects for NAPAP Report to
Congress Advances in Relevant Modeling**

APPENDIX A: ADVANCES IN RELEVANT MODELING

A successful assessment relies on modeling. Issue-specific models are used to understand such processes as pollution transport and diffusion or visibility impairment and, in some cases, to make predictions. In performing an integrated assessment of a science-policy issue that involves many disciplines, it is important to be able to understand and assess the relationships and dependencies between the disciplines. Integrated models that link issue-specific models are tools that scientists and economists can use to provide the understanding necessary for an integrated assessment. This appendix provides descriptions of a few issue-specific and integrated models that are relevant to the acid rain issue.

INTEGRATED MODELS

Integrated models link issue-specific models to enhance understanding of a complex issue that is broad in scope. This section describes three integrated models of the acid deposition issue. They all include emissions, transport and diffusion, deposition, and economics in their frameworks but have different modeling approaches that match the needs of their users. The Tracking and Analysis Framework was developed in the U.S. while RAINS and RAISON were developed by the European community and Canada, respectively.

Tracking and Analysis Framework: TAF

With the passage of the 1990 Clean Air Act Amendments, the United States embarked on an acid deposition control policy that has been estimated to cost billions of dollars. The Amendments created a major innovation in environmental regulation by introducing market-based incentives specifically, trading among electric utility companies in allowances to emit sulfur dioxide. NAPAP is charged with evaluating the status of implementation, the effectiveness, and the costs and benefits of the acid deposition control program created by Title IV of this Act, and to determine whether additional reductions in deposition are necessary to prevent adverse ecological effects. To help NAPAP face this challenge, the U.S. Department of Energy, with support from other federal agencies, has sponsored the development of an integrated assessment model, known as the Tracking and Analysis Framework (TAF).

TAF has been developed in less than two years with relatively modest resources for such a comprehensive model. This rate of progress has been made possible by an innovative combination of methods for integrated assessment. An overview of TAF follows with an outline of these methods, including references to companion documents that provide more details on selected methods and modules.

General Objectives

The following are the general objectives of TAF.

A framework for integrated assessment: TAF is designed to provide a comprehensive framework to address the major issues of concern, from end to end — that is, from the effects of the Clean Air Act Amendments on reducing emissions of pollutants, atmospheric transport, deposition, and environmental effects, all the way to economic valuation of the environmental benefits of emission reductions. A variety of modules can be slotted into this framework. At present, the areas of environmental effects addressed by TAF include visibility, aquatic ecosystems, soils, and human health. Modules for forests and terrestrial ecosystems, crops, and materials remain to be added.

1 *Complete integration:* TAF is designed to include all components within a unified computing
2 environment so that they can be examined and evaluated together, including exploration of the
3 interactions among components.

5 *Agility and flexibility:* TAF is designed to be run on a personal computer in a few minutes, and to
6 allow easy modification of input assumptions and reconfiguration to assess alternative policy
7 scenarios, as new policy issues arise and new data and science become available. It is designed to
8 allow analysts to address new questions in hours or days, rather than weeks or months as is
9 needed with many models.

11 *Transparency:* TAF is designed to provide the models in a form whose structure, relationships, and
12 assumptions can easily be inspected and reviewed. It is designed as a “glass box,” rather than a
13 “black box,” model.

15 *Scientific credibility:* TAF is based on the best available peer-reviewed science and data.

17 *Explicit treatment of uncertainty:* TAF provides explicit representation of the uncertainties due to
18 limitations in scientific understanding, lack of data, and model precision.

20 Modeling Methods

21 The following set of methods as adopted to achieve these objectives:

23 *Influence diagrams:* Influence diagrams provide a graphical representation for display of the
24 qualitative structure of models.

26 *Modular structure:* The model is organized throughout in a hierarchy of modules so that each
27 module is simple enough to be easily understood.

29 *Integrated documentation:* Documentation is integrated, explaining the variables and their role in the
30 computer representation.

32 *Reduced form models:* Most modules are “reduced-form” models — that is, simplified models
33 fitted to more detailed, scientific full-form models. They derive their scientific credibility from the
34 quality of their fit to the detailed models. Both the full-form and the reduced-form models used in
35 TAF are peer reviewed.

37 *Probabilistic analysis of uncertainty:* Probability distributions are used to represent variability,
38 uncertainty due to lack of scientific knowledge or data, and imprecision due to model
39 approximations. Monte Carlo and related methods are used to propagate and combine these
40 distributions to assess the implied uncertainty in the results, and to compare the importance of the
41 various sources of uncertainty.

43 *Progressive refinement:* The team has developed TAF as a series of prototypes of increasing
44 sophistication and refinement, progressively reviewing and refining each to create the next version.
45 Several of these methods have been used in the development of other integrated assessment
46 models. In adopting and refining the entire set of methods, the team found significant synergies
47 among them, leading to what is believed to comprise some important innovations in integrated
48 assessment methodology. The use of many of these methods has been facilitated by the use of
49 Analytica, general modeling software for quantitative modeling and integrated assessment,
50 developed by Lumina. Analytica (Henrion et al., 1996) provides a variety of features used in TAF,
51 including influence diagrams, hierarchies of modules, integrated documentation, and Monte Carlo
52 simulation.

Overview of TAF Modules

A comprehensive assessment of the costs, benefits, and effectiveness of Title IV of the Clean Air Act Amendments requires consideration of many issues. Figure 1 shows the top level of TAF as an influence diagram. Each of the nodes on the diagram represents a module currently in TAF.




Figure 1: A computer screen image shows a top-level influence diagram from the TAF prototype .

The scenario selector on the top left allows the model user to select one or more scenarios for projecting future emissions and to assess and compare effects of those emissions. Users can specify their own scenarios, making assumptions about future growth rates in emissions, by pollutant type (SO_x and NO_x), and source region. Alternatively, users can select a predefined scenario from recent EPA projections, or estimated projections from one of 16 scenarios defined by TAF's emissions module. These 16 scenarios are based on combinations of Phase I caps only and Phase II caps, with and without trading in emission allowances, and with alternative assumptions about future electricity demand growth rates and power plant retirement ages.

TAF currently contains 10 modules developed by over 30 people at 10 different sites, including four consulting firms, three national laboratories, two universities, and a nonprofit foundation.

Model Transparency and Organization

A common complaint about computer models – be they scientific or policy models – is that they are too complicated and too poorly documented to be understood, verified, or trusted. Typically, model documentation is created and updated separately from the computer model, with the result

1 being inconsistent with the model it is supposed to document. In some cases, models are
2 proprietary, and their developer wishes to keep their internal structure secret. Since a major
3 objective of TAF is to support communication and coordination among scientists and policy
4 analysts, an essential requirement for TAF is that the models be documented clearly and
5 consistently.

7 *The Module Hierarchy*

8 TAF employs features of Analytica to display the model as a hierarchy of influence diagrams and to
9 integrate model documentation in the same computer representation used for computation. Figure 1
10 shows the top-level influence diagram, including the key modules and arrows indicating the
11 dependencies among these modules.

12 Each module consists of a diagram, showing the key inputs and outputs, and submodules
13 containing the details of the model. These submodules are themselves arranged hierarchically, as
14 illustrated in Figure 2. Clicking the mouse on one of these nodes in the diagram opens up the
15 diagram for the model it contains. This model hierarchy in TAF extends down to six levels in parts
16 of TAF.




Figure 2: This example of the module hierarchy in TAF shows that double clicking the mouse on a module node (thick outline) opens up the diagram for that module. We repeat to see the third level down in the hierarchy. Parts of TAF contain up to six levels.

Each variable in a model is represented in a diagram by a node with a thin outline. Variables that are defined as uncertain, using a probability distribution are represented by oval nodes. Other variables are represented as rounded rectangles. Index variables are represented by parallelogram nodes.

1 Integrated documentation

2 Each variable in TAF is documented by a card (Object window), containing a set of attributes
3 describing the variable, as illustrated in Figure 3. The card shows the variable class, name, units of
4 measurement, description, definition (mathematical relationship for calculation), list of inputs and
5 outputs, and, optionally, a reference to the publication or authority on which the definition is
6 based. The card also lists the inputs and outputs of the variable. When the definition of a variable is
7 specified or modified, Analytica automatically updates the lists of inputs and outputs and the
8 arrows in the parent diagram to reflect any changes in the dependency relationships.




Figure 3: Each variable is documented internally with an object window. The card shows key information about the variable, including its mathematical definition, and a source or citation.

14 Scientific Credibility and Reduced-Form Models

15 Previous attempts to develop integrated assessment models have sometimes been criticized as
16 lacking sound scientific foundations due to the degree of simplification (Balson & North, 1982;
17 Alcamo et al., 1987). The challenge is to reconcile the need for models to be based on the best
18 available scientific data and models, and to be small, agile, flexible, and comprehensible. TAF
19 meets this challenge by building most modules as reduced-form models based directly on the best
20 available detailed scientific model or data.

22 *Reduced-form models* are simplified models, intended to approximate the behavior of larger, more
23 complicated *full-form models* or data sets. Reduced-form models are simplified in containing
24 fewer variables, less causal detail, or higher levels of aggregation. Their performance is calibrated
25 against or fitted to the performance of the detailed models. In practice, the approximation
26 uncertainty introduced by the simplification for the reduced-form models in TAF is usually
27 dwarfed by the inherent uncertainty in the full-form model. In these cases, the loss in precision
28 from the reduced-form model is negligible.

1 In integrated assessments, it is generally necessary to link several models together – the outputs of
2 one are matched to the inputs of the next. Typically, problems arise because the detailed models
3 are at different levels of aggregation. For example, emission projections may be by season for each
4 power plant; but the atmospheric transport model may need emissions on a daily basis aggregated
5 by a 20-kilometer grid square. Also, the file formats and platforms are often incompatible.
6 Moreover, the models are so large that it is too expensive and time consuming to run them for
7 many different scenarios, especially to handle uncertainty using Monte Carlo or other techniques. It
8 is often impractical to reconfigure and rerun them every time a new policy problem arises.
9 Reduced-form models can obviate these problems, provided they are designed explicitly to use
10 compatible levels of aggregation and file formats.

12 Reduced-form models may be developed or formulated in a wide variety of ways. Following is a
13 description of the approach employed for the atmospheric transport module.

15 *Reduced -Form Models for Atmospheric Transport Module*

16 The atmospheric pathways module of TAF is a reduced-form model based on results from the
17 Advanced Statistical Trajectory Regional Air Pollution (ASTRAP), the detailed long-range
18 atmospheric transport model developed at Argonne National Lab (Shannon, 1981). The reduced-
19 form models consist of source-receptor matrices, normalized to unit emissions at each source. The
20 normalization allows the model to be applied to any emission scenario. Since ASTRAP generates
21 ambient concentrations and deposition rates that are linear in emission rates, this normalization
22 involves no additional approximation. The 60 sources are centroids of the United States, Canadian
23 provinces, and northern Mexico. Temporal aggregation is by season and year. Transport matrices
24 are provided for dry and wet deposition for SO_x and NO_x . Specific receptors have been selected for
25 the visibility, aquatics, crops, and human health effects. Figure 4 shows the top left corner of a
26 source-receptor matrix for ambient SO_2 in winter. (See the documentation on the Atmospheric
27 Pathways module for more details.)




Figure 4: The lower window shows part of the normalized transport matrix by source plant and receptor region, as a detail of the diagram in the upper window. The two-dimensional transport matrix displayed is for winter and ambient SO₂, and is a slice from a four-dimensional array, indexed by four seasons and ambient species.

(Shannon et. al., 1997) has compared the performance of ASTRAP with a nonlinear transport model called RADM (Regional Acid Deposition Model), and actual observations for annual average atmospheric concentration at selected receptor sites in the eastern United States Figure 5 shows an example comparison with regression lines fitting the observations to predictions for each model. Both models appear to underestimate the observations on average. Both models show a similar quality of fit to the data. Since ASTRAP is a linear model, it generates ambient concentrations and deposition at each receptor that are proportional to the emissions at the sources. Therefore, representing it by normalized transport matrices, as in TAF, introduces no additional approximation imprecision for a given time period (seasons). In other words, there is no approximation uncertainty introduced by the reduced-form model beyond the uncertainty inherent in the detailed model on which it is based. Moreover, the uncertainty of the latter appears to be no more at the selected levels of aggregation over space and time than RADM, which is a significantly more complex model than ASTRAP.




Figure 5: A comparison of predictions by ASTRAP (the TAF module) and RADM (another more detailed transport model) with actual observations of annual average concentrations of atmospheric sulfate ($\mu\text{g}/\text{m}^3$) in the eastern United States in 1990.

Progressive Refinement

Model development is, or should be, a learning process. It requires many decisions about the level of detail and aggregation of each variable, making compromises between accuracy and practicality, between detail and computer time and memory, between the policy questions of concern and the pragmatic limitations on what questions the model can address. Finding the best trade-offs is a major challenge, even for the most experienced modelers. The most satisfactory results are obtained when the modelers can revisit decisions in the light of experience with early versions of the model — expanding, simplifying, and refocusing models as the process continues. This process is called *progressive refinement*.

Progressive refinement was adopted as the approach to TAF from the start, beginning with an earlier model named ADAM, developed for NAPAP in the mid 1980s at Carnegie Mellon University. In 1993, NAPAP commissioned a revised prototype integrated assessment model, based on ADAM, that came to be known as TAF. And so began the multiple cycles of progressive refinement, which were essential in obtaining a fully integrated model.

2 TAF Peer Review

3 An intensive peer review of TAF by twelve scientists in December 1995 concluded that TAF was
4 generally successful in meeting its objectives. The development team provided considerable
5 refinement in the following year to address remaining concerns and to improve the analysis.

7 Hierarchical influence diagrams have proved valuable as a visual tool to support transparency for
8 organizing and communicating complex models. Analytica's tools for integrated documentation
9 and array abstraction have also proved helpful. Members of the TAF team and reviewers have been
10 able to scrutinize model structure and assumptions using the built-in model diagrams and
11 documentation.

13 TAF is small enough to run in a few minutes, allowing multiple Monte Carlo runs for
14 comprehensive uncertainty analysis. It is also flexible enough to be rapidly reconfigured to address
15 new policy issues. Yet it is derived from credible, detailed scientific models. The key to reconciling
16 these apparently conflicting goals has been the development of reduced-form models for key
17 modules. Thus, the relatively small size and simplicity of TAF imposes no important loss of
18 precision in the results that it generates.

20 The general approach has been one of progressive refinement, in which each module and the
21 integrated model are developed as a series of versions, starting with module specifications, being
22 progressively refined in response to review and critique by other members of the team. The current
23 version of TAF is the result of four major cycles of refinement, each comprised of a number of
24 minor cycles.

26 The methods and tools that have been developed and the experience gained in developing TAF
27 could be of value to other teams involved in the collaborative development of models for integrated
28 assessment. Other domains of application might include integrated assessments for regional or
29 local air pollution policy, and for international environmental problems, especially for global
30 climate change.

32 Exercises with Historical Data and the Aquatics Module

34 In this section, the transport and aquatics modules in TAF are exercised using historical emissions
35 data from the period 1980 through 1995. Some of the intermediate ambient concentration and
36 deposition results are presented, and the aquatics module is used to estimate the changes in
37 alkalinity for a series of lakes in the Adirondack Mountain region of New York. The acid stress
38 index for three fish species within these lakes is also estimated. Finally, some of the sensitivity
39 and uncertainty analyses possible using the TAF framework are presented.

41 *Historical Emission Trajectories*

42 The utility SO_x and NO_x emission trajectories used in this exercise are derived from the
43 Environmental Protection Agency's emissions databases.¹ Additional non-utility emissions (from
44 commercial, industrial, and transportation sources) are from the emission databases at Argonne
45 National Laboratory.² Mexican and Canadian emission trajectories are also included because of
46 their contributions to ambient pollutant concentrations within United States borders.

1 Acid Rain Division, Environmental Protection Agency, Washington DC

2 Argonne National Laboratory.

4 The emissions are aggregated to the state level in TAF. Figure 6 includes utility and nonutility
5 sources of U.S. emissions during 1980-1995.




Figure 6. National SO_x and NO_x emissions.

9 TAF also contains emission projections based on models run by ICF for EPA and models run by
10 Argonne National Labs for the Department of Energy. Other emission projections can be input by
11 TAF users, and propagated through the entire model.

13 *Estimating Ambient Concentrations and Deposition in the Pathways Module*

15 The pathways module uses linear source-receptor matrices to calculate seasonal ambient pollutant
16 concentrations and deposition estimates integrated over states and at a few selected point receptors,
17 based on state level data from the emissions module. Because the TAF module is primarily
18 concerned with annual averages of deposition and ambient pollutant concentration levels (a few
19 exceptions are handled downstream in the assessment), a linear approximation of transport
20 processes is appropriate.

22 The source-receptor matrices are from Advanced Source Trajectory Regional Air Pollution model
23 (ASTRAP). Using historical emissions data, the ASTRAP matrices have been validated against
24 ambient concentration/deposition data. Eleven years of wind and precipitation data have been used
25 in the model to estimate the variability of model results based on climatological variability. The
26 resulting variability in ambient concentration and deposition estimates was then incorporated into

the module to represent climatological variability. Normal distributions representing the annual variability of the source/receptor relationship are multiplied by the concentrations and depositions estimated at each receptor site.

The variability in ambient concentrations based on climatic fluctuations is illustrated in Figure 7, for micrograms per cubic meter of SO₂ in the Grand Canyon area. Confidence bands of 5, 25, 50 (median), 75, and 95 percent are used to represent the annual variability of average ambient pollutant concentration.




Figure 7. Annual average species concentration of SO₂ in the Grand Canyon area.

This variability is significant when examining the baseline or Title IV pollutant concentrations alone, but when the Title IV concentrations are subtracted from the baseline concentrations to obtain an estimate of concentration reductions under Title IV, much of the year-to-year variability due to climatological differences is canceled out, resulting in estimates of reduced ambient concentrations as shown in Figure 5. The climatological variability factored into the transport of pollutants has a measurable effect on reductions in pollutant concentrations, as demonstrated by the confidence interval surrounding the mean estimate of ambient pollutant concentrations. The next section compares this variability to other sources of variability and uncertainty that contribute to the aquatic effects of acid deposition.

Aquatics Effects: Using TAF to Rank Effects

The aquatics module is a reduced-form version of the Model of Acidification of Groundwater in Catchments (MAGIC). Using deposition data from the pathways module and Adirondack lake background data, the module calculates lake pH, acid neutralizing capacity, base saturation, fish species richness, and fish acid stress indexes for 33 Adirondack lakes. The module has been calibrated to data and results from the MAGIC model, and performs comparably, despite its much more modest computational requirements.

1 This discussion is limited to the acid stress index, also known as the conditional mortality rate.
2 The acid stress index is a common estimate of the increased likelihood that a fish of a given life
3 stage will die when exposed to the specified water quality conditions, over and above the mortality
4 expected in a circumneutral reference water. Higher numbers indicate higher stress and increased
5 likelihood of death. The benefits module in TAF uses the acid stress index computed at the
6 Adirondack lake sites, for three fish species, to estimate the catch per unit of effort expended by
7 recreational fishermen. Figure 8 contains the median brook trout acid stress index results for a
8 single Adirondack lake. Figure 8 also shows 25 % and 75 % probability intervals. The confidence
9 intervals are not symmetric, indicating that the distribution of possible acid stress index values is
10 itself asymmetric, with a right-hand tail.




Figure 8. Acid Stress Index and confidence intervals for an Adirondack lake.

Aquatics Effects: Using TAF to Rank Sensitivities and Uncertainties

The uncertainty around the acid stress index includes a fraction above zero. This indicates that, when the uncertainty in the aquatics modeling and natural climatological variability is taken into account, a reduction in the acid stress index cannot be guaranteed. That said, the chance of a nonzero, favorable change in acid stress index (i.e., a reduction) is quite large. An importance analysis can be used to compare the relative contributions of the uncertainties in the model to the acid stress index results. The uncertainties affecting the acid stress index include:

- *Uncertainty in deposition from the pathways module.* This is similar to the climatological variability in the visibility module, except it is expressed as cumulative acid deposition instead of annual ambient concentration.
- *Uncertainties in the fit between the MAGIC model and empirical data.* There are four components to this uncertainty: uncertainty in the estimation of lake calcium concentrations, uncertainty in the estimation of acid neutralizing capacity, uncertainty in the estimation of lake pH from acid neutralizing capacity, and uncertainty in the estimation of acid stress index from lake pH (described with four parameters).
- *Uncertainties in the fit between the reduced-form model version in TAF and the full-form*

MAGIC. There are two components to this uncertainty: uncertainty in the estimation of lake calcium concentrations, and uncertainty in the estimation of lake acid neutralizing capacity.

The reduced-form model and *MAGIC* uncertainties were quantified from the results of linear and nonlinear regressions. Climatic variability was quantified by measuring variability in *ASTRAP* deposition results using historical wind trajectory data from eleven separate years. These sources of uncertainty are ranked using an importance analysis. The results of the analysis are shown in Figure 9.




Figure 9. Importance of uncertain inputs on uncertainty in ASI output for brook trout.

The uncertainty in the relationship that translates pH to acid stress index (ASI parameters 1, 2, and 4) dominates the result. The conversion from acid neutralizing capacity to pH also provides a significant amount of uncertainty in the result. These uncertainties swamp the other sources of uncertainty in the model, including the imprecision in the reduced-form model and the variability caused by year-to-year changes in deposition. Because the overarching uncertainties in *MAGIC* dominate the uncertainty in the result, we conclude that the reduced-form version of *MAGIC* within TAF performs comparably to *MAGIC*.

Note also that the climatological variability is not large compared to some of the other uncertainties. This is true in part because much of the climatological uncertainty is canceled out when the difference of the baseline and comparison scenario results is taken. The climatological uncertainty is the same across the two scenarios, so it is reduced when the difference of the two scenarios is taken.

This analysis identifies the conversion of pH to acid stress index and of acid neutralizing capacity to pH as critical sources of uncertainty in the aquatics module. Acid neutralizing capacity is converted to pH by using a four-parameter nonlinear equation based on work by Small and Sutton (1986), calibrated to data for the 33 Adirondack lakes considered in TAF. Whether this source of uncertainty should be refined and reduced in future versions of the aquatics module depends on the effect of this uncertainty in calculation of aquatics benefits.

Future TAF Analyses

The analyses described here are just a small sample of the potential of an integrated assessment. Future analyses in TAF can compare results not only across effect modules, but also across unmodeled effects using back-of-the envelope scoping analyses. These analyses will permit prioritization of additional modules to be added to the TAF framework.

As additional information on the costs of Title IV regulations on utilities is integrated, utility costs can be compared to the benefits calculated in TAF to determine whether the subset of benefits calculated are sufficient to suggest that Title IV is cost-effective. Also, the capability exists to compare the geographic distribution of costs with the distribution of benefits, because TAF calculates both costs and benefits on a state level.

The model is able to compare both uncertainties that propagate through several modules, and uncertainties across different effects and benefits. It enabled comprehensive identification of those inputs and model forms sensitive to change and most influential in their effects on output uncertainty. These abilities allow TAF to provide important information on future research priorities and the confidence in current estimates of acid rain damages and Title IV benefits.

In a further effort to share TAF-related research, information on the TAF project, including draft models and the Analytica modeling software, is being made available over the World Wide Web via Internet (<http://www.lumina.com/taflist>). In addition, examples of TAF used as an analytical tool will appear in the following forthcoming publications:

Palmer, K.L., and D. Burtraw. 1997. Electricity Restructuring and Regional Air Pollution. *Resource and Energy Economics* , 19:139-174.

Shannon, J.D., E.C. Trexler, and R.S. Sonnenblick. 1997. Modeling Visibility for Assessment. *Atmospheric Environment* , 31(22):3719-3727.

References

Alcamo, J., M. Amann, J.P. Hettelingh, N. Holmber, L. Hordjik, et al. 1987. "Acidification in Europe: A simulation model for evaluating control strategies", *Ambio* 16(5) 232-245.

Balson, W.E. and D.W. North. 1982. " Acid deposition: Decision Framework", EA-2540, prepared by Decision Focus, Inc for the Electric Power Research Institute, Palo Alto, CA, 1982.

Henrion, M. et al. 1996. *Analytica User Guide*, Lumina Decision Systems, Los Altos, Ca.

Shannon, J.D. 1981. A model of long term average sulfur atmospheric pollution, surface removal, and net horizontal flux. *Atmospheric Environment* 15(5):689-701.

Small, M.J. and M.C. Sutton. 1986. A regional pH-alkalinity relationship. *Water Resources* 20:335-343.

1 RAISON

2 The National Water Research Institute of Environment Canada has developed a decision model to
3 support environmental decision makers. The RAISON system is designed for a teamwork
4 approach to developing decision-support systems for various environmental problems. The
5 teamwork approach considers not just the software but also the scientists and end users who must
6 be involved in the development from an early stage. The RAISON prototype is sufficiently robust
7 to interact with scientists and policy advisors, first to overcome any communication problems
8 among themselves and with the computer, and then to iteratively improve the system toward
9 creating a useful final product. While RAISON has some similarities to geographic information
10 systems (GIS), it differs significantly insofar as it emphasizes decision support and analysis that
11 are difficult or impossible in a typical GIS.

12 RAISON offers a generic framework to integrate data, text, maps, satellite images, pictures, video,
13 and other knowledge input. The system provides the user with a library of software functions and
14 tools—including algorithms, models, optimization procedures, expert systems, neural network and
15 other information technologies—to produce customized interfaces and output, including
16 interpretation, advice, scenario tests, strategic analysis, and policy recommendations. For
17 example, data can be entered to RAISON through conversion interfaces available for many off-the-
18 shelf databases. GIS maps can be entered in vector and raster formats. Models can be
19 incorporated into the system by: (1) using the codes as given if compatible with the programming
20 languages used in RAISON (Visual Basic, Visual C, C++), (2) building an interface that intercepts
21 the input and output to connect to the database in RAISON, or (3) executing the model off-line but
22 writing and reading the input and output. By using the various modules in RAISON, since all
23 components are linked via graphical interfaces, different applications can be adapted with
24 customized interface using optimization procedures or expert system techniques to direct control of
25 the information and knowledge.

26 The RAISON system is property of the Government of Canada. It is distributed under license by
27 NWRI Software.

28 For further details, please consult the references or contact:
29 Dr. David Lam
30 Phone: (905) 336-4916
31 Fax: (905) 336-4582
32 Email: david.lam@cciw.ca

33 *References:*

34 Lam, D.C.L., C.I. Mayfield, C.I. Swayne, and K. Hopkins. 1994. A prototype information
35 system for watershed management and planning, Vol. 2(4):499-517.

36 Lam, D.C.L., D.A. Swayne, G.S. Bowen, and D. Kay. 1995. A teamwork approach to decision
37 support system development for watershed management problems. Proceedings of the ICCI
38 Conference on Computer Modeling and Application, Peterborough, June 1995.

39 RAINS-Asia

40 Expanded energy use in Asia, combined with utilization of indigenous coal, will result in an
41 increase in emissions of acidifying compounds and greenhouse gases. By the year 2010, sulfur
42 dioxide emissions for Asia will most likely exceed the emissions of North America and Europe
43 combined.

In recent years, integrated assessment models have been used for international negotiations on acid rain in Europe and North America. These models provide negotiators and regulators with a full regional picture of the problems associated with the entire causal process, from energy systems and emissions to the ultimate impact on the natural and man-made environment. The model user can analyze the regional and national implications of various scenarios, which include options for energy use, control strategies, and mitigation policies.

The RAINS-ASIA model is a tool for integrated analysis of air pollution. It consists of three modules: (1) Resource and Energy Scenario Generator (RESGEN), (2) Energy Emissions (ENEM), and (3) Deposition and Critical Loads Assessment (DEP). Each module describes various aspects of acidifying emissions and dry and wet deposition. A set of menus and options guides users through the modules.

RAINS-ASIA covers East, South and Southeast Asia with particular emphasis on Japan, India, China, Indonesia, Thailand, and South Korea. It contains databases on energy consumption for 23 countries, 94 subregions, and 250 large point sources of deposition and estimates acid deposition carrying capacity of 31 types of ecosystems. Values for sulfur depositions are based on a yearly average and are calculated at a 1 x 1 degree of resolution. Aggregate results of source/receptor information on acid deposition patterns for each subregion or country can be obtained, and local impacts can be estimated as well. The temporal range of the model is 1990-2020.

RESGEN. Based on best-estimate assumptions of economic development and population growth, RESGEN makes possible the generation of scenarios of energy consumption pathways. Energy consumption is disaggregated into industrial, transportation, residential, commercial, and other sectors. The supply side of the module identifies different technologies for energy generation and sources of emissions, such as electricity generation, oil refining, and other energy sector operations with combustion of fossil fuels. This framework makes it possible to generate rough estimates of future energy demand and supply trends under a variety of socioeconomic and technical assumptions.

ENEM. The ENEM module takes the energy consumption at the sectoral and regional levels as given by the RESGEN module and estimates the corresponding SO₂ emissions and costs of various emission control options. Sulfur emissions from combustion of fossil fuels are calculated based on fuel characteristics, combustion technology, and emission control assumptions. Emissions are characterized as low-level area sources and high-level large point sources. The module considers a number of options to reduce sulfur emissions, including fuel desulfurization and flue-gas desulfurization. To explore integrated abatement strategies, the users can apply specific control policies to selected countries or regions within countries. The results of energy conservation measures and fuel substitution can be explored by analyzing alternative energy pathways, either by selecting one of the several preset energy-use scenarios or by creating a new scenario based on expectations of fuel use.

Emission control and associated costs are based on the most commonly used emission control technologies. Cost evaluation is based on international operating experiences of pollution control equipment, and extrapolating them to country context. It is assumed that a relatively free and competitive market exists for control technology. This module also computes national cost curves that rank abatement measures by their cost-effectiveness.

DEP. This module estimates ambient levels of acid rain precursors and acid deposition throughout the region and compares them with data on environmental sensitivities that are presented in the form of critical-load maps. This module consists of two submodules: Atmospheric Transportation and Deposition (ATMOS), and Ecosystems Impact (IMPACT).

The *ATMOS* submodule is based on the National Oceanic and Atmospheric Administration's Branching Atmospheric Trajectory (BAT) model, which calculates wet and dry depositions of SO₂ and sulfates from a particular source as the pollutant is transported by meteorological fields. If these trajectories are run for an entire year, then the submodule estimates the amount of annual deposition on the entire region from a particular source. If the calculations are repeated for all sources, then the total annual deposition in the region can be estimated. Inputs for this submodule include SO₂ emission rates, winds and temperature, precipitation rates, and estimates of dispersion coefficients, dry deposition velocities, and wet-scavenging coefficients. The module is run for each large point source and area source estimated by ENEM, and sulfur deposition is calculated on a 1 x 1 grid. The results are aggregated to provide source/receptor information on acid deposition patterns for each of the subregions in the study region, and further aggregated to provide country-by-country source/receptor information.

The *IMPACT* submodule estimates critical loads (the maximum long-term deposition levels that could be tolerated without damage) for fifteen different ecosystems.³ The critical loads are compared with the estimates of sulfur deposition from the *ATMOS* submodule to determine which ecosystems may be at risk for different emission scenarios. This assessment is based on complex dynamics of processes in key ecosystems, such as soils, surface waters, and vegetation systems. These models include the computation of the depletion of acid buffer capacity of ecosystems under the influence of precipitation, evaporation, water flows, and budgets of chemical ecosystem constituents.

The critical-load calculation involves a two-step process. The first step applies a qualitative relative-sensitivity approach to distinguish an ecosystem's sensitivity to acidification. In this method, weights are assigned to four indicators of ecosystem sensitivity -- bedrock lithology, soil type, land use, and annual rainfall. In the second step, based on the Steady-State-Mass Balance Method, computations are performed to assign critical loads to all areas distinguished on the map of relative sensitivities. This method assumes steady state equilibrium between soil solid phase and soil solution, and computes the maximum acid input to the system that will not cause an excess of the critical alkalinity value, which is computed from average thresholds for chemical values, such as pH, aluminum, and aluminum-calcium ratios.

Weathering rate, which is a key input to the process, is determined by estimates of soil mineralogy, which are then modified by climate and soil attributes. The resulting map of weathering rates and ratios of land use and precipitation to potential evapotranspiration are used in the critical-load calculations.

The three modules together permit users to operate the PC-based RAINS-ASIA policy model in a scenario-analysis mode. They can estimate current costs and impacts of alternative emission control strategies on a country, or regional basis. Emissions can be tracked through the deposition process to assess their potential impacts on critical ecosystems. Control strategies can be specified for specific fuel types, economic sectors, emission-generating power plants, emission control technologies, regions or countries under study, and they can be applied individually or in any combination. The model can also be used to identify potential maximum impact locations for establishing monitoring sites, which in turn could assist in model validation.

3 Desert/semi-desert, mangrove woods, irrigated land/paddy, semi-arid and thorn woods, dry tropical/subtropical/savanna, agriculture, cool scrub/grassland, temperate broadleaf wood, tropical montane forest, other conifer trees, bog/mire/moor, wet tropical forest, Tibetan cold grass, northern/main/southern taiga/tundra.

1 ISSUE-SPECIFIC MODELS: MAGIC

2 A need has arisen to test the veracity of model projections, especially in cases where policy and/or
3 economic interests are at stake. As Oreskes et al. (1994) pointed out, however, verification and
4 validation of mathematical models of natural systems are impossible, because natural systems are
5 never closed, and model results are not unique. Model confirmation is possible and entails
6 demonstration of agreement between prediction and observation. Because such confirmation is
7 inherently partial, it is critical that policy-relevant models be tested in a variety of settings and under
8 a variety of conditions.

9 Since 1990, the Model of Acidification of Groundwater in Catchments (MAGIC) (Cosby et al.,
10 1985) has been widely used throughout North America and Europe, and has been the principal
11 model used by NAPAP to project the response of surface waters to changing levels of sulfur
12 deposition. MAGIC projections of the effects on surface water chemistry of various sulfur
13 emission scenarios formed the technical foundation for a large part of NAPAP's *1990 Integrated*
14 *Assessment Report* (NAPAP, 1991). Subsequently, a research effort was conducted from 1990 to
15 1996 to improve the performance of MAGIC and to test the model and confirm its results at
16 multiple sites. Model evaluations have included hindcast comparisons with diatom
17 reconstructions⁴ of preindustrial lake-water chemistry in the Adirondack Mountains of New York,
18 and tests of the veracity of model forecasts using the results of whole-catchment acidification
19 experiments in Maine (Norton et al., 1992) and Norway (Gjessing, 1992) and whole-catchment
20 acid-exclusion experiments in Norway (Wright et al., 1993).

21 Based on the results of this testing, it appears that MAGIC provides reasonably accurate forecasts
22 of changes in surface water acid-base chemistry in response to changing levels of acidic
23 deposition. Although some uncertainties remain, particularly with respect to watershed nitrogen
24 dynamics, MAGIC provides a generally accurate and well-tested tool for integrated assessment
25 modeling.

26 The testing of MAGIC over the last six years has elucidated several potentially important
27 deficiencies in structure and method of application, and has resulted in changes to the model and its
28 calibration procedures. The work has included in-depth evaluation of issues related to regional
29 aggregation of soils data, background sulfur deposition, natural organic acidity, nitrogen, and
30 aluminum mobilization. The result has been an improved and more thoroughly tested version of
31 MAGIC, which yields forecasts different from the version that served as the technical foundation
32 for NAPAP's *1990 Integrated Assessment Report*.

33 Changes to the Model

34 Background Sulfate and Subregional Calibration

35 Subsequent to the regional MAGIC modeling that was conducted for NAPAP (1991), there was
36 concern that (subregional) Adirondack soils might differ in their chemical properties from related
37 (regional) soils in other areas of the Northeast, and that MAGIC projections for Adirondack
38 watersheds might be biased because they were based on soil attributes that actually reflected
39 conditions in the Northeast other than those in the Adirondacks. Therefore, the model input data
40 were reaggregated to use only soil collected from Adirondack sites.

41 Modeling for the *1990 Integrated Assessment Report* also assumed that the deposition of sulfur in

4 Diatoms are microscopic algae, the remains of that are incorporated into lake sediments which accumulate over time. The species composition and relative abundance of diatoms at different levels in the sediment can be used to estimate the pH of lake water using sophisticated mathematical relationships.

preindustrial times was limited to sea salt contributions. Based on analyses presented by Husar et al. (1991), this assumption was modified so that preindustrial deposition of sulfate was assumed to be equal to 13% of current values (Sullivan et al., 1991).

Recalibration of MAGIC to the Adirondack lakes database using the regionally corrected soil and background sulfate data resulted in approximately $10 \mu\text{eq L}^{-1}$ lower model estimates of current acid neutralizing capacity. A substantial downward shift was also observed in predicted preindustrial and current lake water pH (~ 0.25 pH units) for lakes having current pH greater than about 5.5. These differences were attributed to lower calibrated values for lake water sulfate concentrations and higher values for the partial pressure of carbon dioxide estimated for Adirondack lakes, compared with the Northeast as a whole (Sullivan et al., 1991).

Organic Acids

Concern was raised subsequent to the *1990 Integrated Assessment* regarding potential bias from the failure to include organic acids in the MAGIC formulations used by NAPAP (1991). MAGIC hindcasts of preindustrial lake water pH showed poor agreement with diatom-inferences of preindustrial pH, and preliminary analyses suggested that these differences could be partly due to the presence of naturally occurring organic acids in Adirondack lake waters.

An organic acid model was developed by Driscoll et al. (1994) using data collected by the Adirondack Lakes Survey Corporation (Kretser et al., 1989) for 1,400 lakes located in the Adirondack region. This model was coupled with MAGIC. Model hindcasts using the unmodified MAGIC yielded preindustrial pH values that were substantially higher than diatom-based estimates, and the discrepancy was greatest for those lakes in the most biologically sensitive portion of the pH range (pH of 5.0 to 6.0). Furthermore, MAGIC hindcast pH estimates were greater than 6.0 for all lakes investigated, whereas diatom estimates of preindustrial pH ranged from as low as 5.2 to above 7.0. When the organic acid model was incorporated into MAGIC and simulated preindustrial pH values from the new model were compared with diatom-inferred pH, the comparison yielded considerably closer agreement between model estimates of preindustrial pH than did the simulations that did not consider the effects of organic acids (Sullivan et al., 1996).

The lakes of greatest relevance with respect to potential biological effects of acidification, especially those having a pH of less than 5.5, exhibited increasingly larger discrepancies with decreasing pH between diatom and MAGIC model estimates of preindustrial pH when organic acids were omitted from the analysis. Including an organic acid representation in the MAGIC simulations greatly improves the agreement between these two modeling approaches.

The results of these analyses of Adirondack lakes demonstrated that: (1) organic acids must be considered in modeling the response of lake waters in the Adirondack Mountains (and possibly other regions) to acidic deposition; and (2) once organic acids are included in the modeling approach, reasonable agreement is obtained in hindcast comparisons with diatom-inferred pH. It should be emphasized, however, that this test included only two points in time and involved only pH. Even though the model adjustment with organic acids improved agreement for pH, other variables in the model may have been poorly represented. MAGIC and other process models require further testing and confirmation. Many potentially important geochemical processes are not well represented in the model or the input data, and it was not clear how inclusion of such processes might affect model results.

Aluminum

Aluminum mobilization is now widely believed to be one of the most important ecological effects of surface water acidification. Potential effects of aluminum mobilization from soils to surface and

soil waters include alterations in nutrient cycling, pH buffering effects, and toxicity to aquatic biota and terrestrial vegetation. MAGIC simulates aluminum solubility based on an assumed equilibrium with the mineral gibbsite. The model first calculates the total concentration of acidic cations (e.g., hydrogen plus aluminum) on the basis of simulated concentrations of base cations and mineral acid anions (e.g., sulfate, nitrate, chloride) using mass balance and electroneutrality constraints. The acidic cations are then partitioned between hydrogen and aluminum, using the gibbsite mineral equilibrium, thermodynamic equations, the partial pressure of carbon dioxide, and the organic acid formulation. This partitioning is important because inorganic aluminum in solution can be highly toxic to aquatic biota, even at low concentrations (Baker and Schofield, 1982). Model simulations often overpredict the change in aluminum concentration. The aluminum formulation in the MAGIC model has recently been modified to better reflect empirical relationships between aluminum and hydrogen ion. The revised formulation was used to predict aluminum concentrations in runoff at experimental ecosystem manipulation sites in Maine and Norway, and in both cases yielded closer agreement with measured values than the original MAGIC predictions (Sullivan and Cosby, in review).

Nitrogen

MAGIC contains an extremely simplified representation of nitrogen dynamics within catchment soils. There are no processes controlling the details of nitrogen cycling in the model. The version of MAGIC used for the *1990 Integrated Assessment Report* (NAPAP, 1991) was not appropriate for simulation of changes in atmospheric deposition of nitrogen. In light of the increasing concern about nitrogen saturation in forested ecosystems, this was a serious shortcoming in the model. A new, coupled sulfur and nitrogen model, MAGIC-WAND, was developed by extending the MAGIC model to incorporate the major ecosystem nitrogen fluxes and their changes through time (Ferrier et al., 1995). MAGIC-WAND is perhaps the most generalized model, but several more detailed nitrogen models are also available, including MERLIN, NuCM, and PNET-CN. MAGIC-WAND has been applied regionally to simulate the response of lakes in the Galloway region of southwestern Scotland to changing deposition of sulfur and nitrogen from 1988 to 1993. The model is currently being further evaluated for watersheds in the southern Appalachian, Cascade, and Rocky Mountain regions of the United States.

Cumulative Impacts to Changes to MAGIC

- The improved MAGIC predicts that sensitive lakes and watersheds in the Adirondack Mountains are less responsive (in terms of change in acid neutralizing capacity, pH, and inorganic aluminum) than was predicted by the earlier version of MAGIC used for the *1990 Integrated Assessment*.

To evaluate the incremental and cumulative impact of the modifications to MAGIC, Sullivan and Cosby (1995) conducted a suite of model simulations for the Adirondack Direct/Delayed Response Project (DDRP) lakes. They used the baseline model structure from the Direct/Delayed Response Project and the NAPAP *1990 Integrated Assessment Report* (NAPAP, 1991). The changes to the model they examined included modifying the assumption regarding background sulfur deposition, reaggregating the soils data, recalibrating the model specifically for the Adirondack subregion, adding the organic acid model to the surface water compartment, and changing the aluminum/hydrogen ion relationship from cubic to quadratic. However, these analyses did not examine the effects on model output of including nitrogen dynamics in the model simulations.

A suite of simulations was conducted based on the application of an assumed deposition scenario to derive a fifty year forecast using each model structure. The deposition scenario assumed constant sulfur deposition from 1984 (the calibration year) to 1994, followed by a 30% decrease in sulfur deposition from 1995 to 2009, with constant deposition thereafter until 2034 and considered the modeled responses of thirty-three Adirondack lakes to this scenario. The impacts of the

changes were illustrated by tabulating the percentage of lakes predicted to have pH, acid neutralizing capacity, or aluminum values in excess of commonly accepted thresholds of potential biological effects.

The overall effect of the various changes to the model structure and application procedures was an increase in the percentage of lakes exceeding various biological thresholds with respect to pH, aluminum, and acid neutralizing capacity subsequent to an hypothesized 30% decrease in sulfur deposition (Table 1). The largest changes were observed for pH and aluminum; acid neutralizing capacity projections were less affected. The modifications to the model that caused the greatest changes in projected output were the recalibration of the model to the Adirondack subregion, modification of the assumption regarding background sulfate, and the incorporation of the organic acid model into MAGIC. The modification of the aluminum algorithm caused fewer lakes to be projected to exceed aluminum threshold values in response to the reduced deposition scenario; this change was quantitatively less important than the previous changes.

The magnitude of effect of the cumulative modifications to the model was considerable (Table 1). For example, 32% of the lakes had measured pH less than 5.5 in 1984, whereas only 8% were projected to still have pH less than 5.5 after the reduction in sulfur deposition, using the original

Table 1. Cumulative effects of recent (post-1990) changes to the structure and method of application of the MAGIC model. MAGIC predictions of the percentage of Adirondack DDRP lakes having pH, acid neutralizing capacity (ANC), and Al above or below threshold values in the year 2034 subsequent to an hypothesized 30% decrease in sulfur deposition (Source: Sullivan and Cosby 1995)									
Data Type	Percentage of Lakes Having pH Below Value			Percentage of Lakes Having ANC Below Value ($\mu\text{eq L}^{-1}$)			Percentage of lakes Having Al Above Value ($\mu\text{eq L}^{-1}$)		
	5	5.5	6	0	25	50	50	100	200
Measured 1984 Values	12	32	38	18	48	59	30	18	10
MAGIC Projection of 2034									
1990 Version of MAGIC used for NAPAP	0	8	20	6	34	44	4	0	0
Current Version of MAGIC*	8	32	44	14	40	44	30	10	4
* Does not include nitrogen dynamics, which are included in MAGIC-WAND.									

MAGIC application used for NAPAP. In contrast, the improved version of MAGIC projected that 32% of lakes would still have pH less than 5.5 in the year 2034. Similarly, of the 30% with measured inorganic aluminum concentrations greater than $50 \mu\text{g L}^{-1}$ in 1986, the original model structure projected only 4% would still have concentrations greater than $50 \mu\text{g L}^{-1}$ in 2034, compared to 30% projected to continue to have high inorganic aluminum by the improved version of MAGIC. Based on model projections using the improved version of MAGIC, little recovery of

Adirondack lakes would be expected subsequent to a 30% reduction in sulfur deposition. The number of lakes having pH lower than 6.0 was actually projected to increase, and the number of lakes projected to have acid neutralizing capacity lower than 0 only decreased slightly in response to lower deposition. These estimates were independent of any possible increases in nitrate leaching that might occur. The lack of recovery suggested by these revised model projections is attributable partly to a decrease in the modeled base saturation of watershed soils. These results may affect expectations of recovery in response to sulfur emission controls mandated by Title IV.

The future response of lakes and streams to acid deposition is also highly dependent on the extent to which watersheds in acid-sensitive regions become nitrogen-saturated. EPA scientists conducted MAGIC model simulations for fifty years into the future that effectively bounded the range of possible water chemistry responses—ranging from no watersheds reaching nitrogen saturation to all simulated watersheds reaching nitrogen saturation during the simulation period. The model projections for Adirondack lakes, for example, suggested that the percent of chronically acidic lakes in the target population in fifty years could range from 11% to 43%, depending on the number of watersheds that become nitrogen saturated (U.S. EPA, 1995). Similarly, for mid-Appalachian streams, the modeled percent of streams acidic in fifty years ranged from 0% to 9%, depending on the extent of nitrogen saturation (U.S. EPA, 1995).

Magic Model Confirmation

MAGIC has been tested after inclusion of many of the model modifications discussed in the preceding sections. The revised model with Driscoll et al.'s (1994) organic acid model yielded reasonable agreement between model hindcast pH and diatom-inferred pH for the data set of thirty-three Adirondack lakes. Differences between diatom and MAGIC estimates of preindustrial pH of Adirondack lakes, based on the version of MAGIC that includes an organic acid representation, were well within the range of expected differences due to annual and seasonal variability and uncertainties in the model algorithms.

However, "successful" comparison of MAGIC with diatom hindcasts in one region does not constitute a sufficient verification to impart complete confidence in using MAGIC, or any process model, for predicting the response of surface water chemistry to changes in acidic inputs. Additional model confirmation in the form of comparison of model output with *measured* data, is required. This has been the focus of modeling efforts at the experimental manipulation site at Bear Brook Lake in Maine and at two sites in Norway.

Initial modeling efforts at Bear Brook (Norton et al., 1992; Sullivan et al., 1994; Cosby et al., 1996) predicted a much larger increase in stream water sulfate concentration than was observed in the treated stream. Although there is considerable uncertainty regarding the lag in sulfate release/adsorption in soils, it appears that MAGIC overpredicted the increase in stream water sulfate concentrations at Bear Brook by nearly a factor of two. This overprediction was due to the high value assumed for the half saturation of sulfur adsorption, which was based on laboratory measurements. As a consequence, other key variables (especially acid neutralizing capacity and aluminum) were also predicted to increase to a greater degree in response to the experimental acidification than was actually observed.

The original calibration of MAGIC for the Bear Brook forecast was based on four years of data from the reference stream, East Bear Brook. To assess the degree to which discrepancies between predicted and observed stream water chemistry at Bear Brook could be improved by correcting the error in predicting sulfur dynamics and *a priori* differences between treatment and control catchments, a revised calibration was conducted. The revised calibration corrected for the obvious large bias in effective sulfur adsorption in watershed soils and also corrected for *a priori* differences between the treatment and reference catchments. In essence, in the latter case, expert judgment was substituted for strictly laboratory-derived information. The resulting simulations

206 matched measured values in West Bear Brook to a substantially greater degree than the earlier
207 forecasts.

208 Projected stream water sulfate concentrations closely agreed with measured values in West Bear
209 Brook for the first three years of manipulation in the revised model simulation (Cosby et al.,
210 1996). The model simulation also showed much better agreement with measured values for the
211 sum of base cations and acid neutralizing capacity, than the initial MAGIC simulation. Although
212 the effects of a drought year (1992) on base cation concentrations and acid neutralizing capacity
213 were still not captured by the simulation, the overall agreement between predicted and observed
214 base cation concentrations and acid neutralizing capacity was much improved. Slight
215 underestimation of pH decrease and overestimation of aluminum increase were still evident in the
216 revised projections, although the magnitudes of these biases were reduced dramatically because of
217 the improvement in predicted sulfate concentration and acid neutralizing capacity.

218 Results of modeling efforts at Bear Brook, as well as measured chemical changes at Bear Brook,
219 illustrate that a remaining major weakness of MAGIC (and other process models) relative to the
220 needs of NAPAP is the failure to include algorithms to simulate nitrogen cycling and nitrogen
221 retention in watershed soils and vegetation. The success of the nitrogen component of the
222 modeling effort at Bear Brook was totally dependent on adjusting the nitrogen inputs to the model
223 to match measured outputs in stream water. Nitrogen dynamics were extremely important at this
224 site (Kahl et al., 1993), although this had not been anticipated at the inception of the Watershed
225 Manipulation Project at Bear Brook.

226 The process of evaluating and improving MAGIC is iterative. It has now been shown that the
227 inclusion of organic acids in the model is important and that MAGIC often yields acceptable model
228 simulations of past and future change. It has also been shown that further improvements are
229 needed, particularly with respect to nitrogen, which is the focus of the extended version of the
230 model MAGIC-WAND. The model simulations at Bear Brook also revealed important weaknesses
231 and uncertainties in several aspects of the model structure and/or the manner in which the model is
232 applied to a given catchment. Results at Bear Brook verified that key remaining uncertainties relate
233 to the modeling of aluminum dissolution, sulfur retention in soils, and the dependence of runoff
234 chemistry on hydrological variations that are difficult to simulate.

APPENDIX B

Design and Performance of Pollution Trading Programs

Control of SO₂ emissions under the 1990 Clean Air Act Amendments instituted two important innovations in U.S. environmental policy. The more widely acknowledged of these is the SO₂ emissions trading program. Less acknowledged is the average annual cap on aggregate emissions by electric utilities, which guarantees that nationwide emissions will not increase as economic growth occurs in the future.

The two innovations are designed to work together. Firms are allocated annual SO₂ emission "allowances" in proportion to their historic emissions, which they may transfer among facilities or "bank" for future use. Under this approach, the environmental goal emissions cap is established in the statute, but the means for accomplishing that goal is left to the ingenuity of the interested parties. In addition, each affected plant must continue to meet all other applicable state and federal emission standards.

The main attraction of a permit trading program or, more generally, of what is known as an incentive-based approach to environmental regulation, is the promise that it can achieve an environmental goal at a lower cost than regulatory approaches that dictate specific actions for individual facilities or groups of facilities. Because the cost of reducing emissions often varies tremendously among facilities, trading programs can help ensure that the least expensive means are pursued before undertaking more costly efforts. The savings can be a boon for consumers and industry by reducing the cost of regulation, and, a boon for the environment by allowing society to purchase greater environmental protection at the same cost. The evidence to date for the SO₂ emission trading program indicates that the cost savings have been substantial.

A second type of cost savings from incentive-based regulation are those that are expected to be achieved over time as firms find ways to lower the cost of reducing their emissions. Emission trading provides incentives for firms to innovate because firms can expect to keep the cost savings. At this juncture it is premature to say whether significant innovation has resulted from the SO₂ program, but there are many anecdotes of process changes and efficiency improvements that have contributed to the low cost of emission reductions to date. It can be said at this point that competition between different methods of compliance has lowered the cost of compliance.

With the apparent success of the SO₂ trading program, the question arises whether this approach should be used to guide other environmental protection efforts. To be sure, there are other environmental problems that would seem to lend themselves to the use of one or another incentive-based approach such as tradable permits, emission fees or deposit-refund systems. However, there are also a variety of problems that are less well suited. This appendix considers the characteristics of environmental problems in general that may or may not be amenable to this approach, especially in the context of controlling air pollution. It evaluates why the SO₂ problem appears well suited to the use of tradable permits, and describes new applications of incentive-based approaches for controlling CO₂ and NO_x.

Characteristics That Contribute to the Success of Permit Trading Programs

Emission permits are similar in many ways to other goods traded in markets. The success of the market for emission permits depends on the breadth of supply and demand. The larger the number of firms that can trade pollution permits the more successful a trading program will be in reducing control costs.

1 Paradoxically, while the participation of many firms makes it easier for buyers and sellers to find
2 each other, it also makes it easier for them to remain anonymous. Firms may want to remain
3 anonymous when it comes to trading permits because they do not want to signal to competitors
4 their plans for the future. In a small permit market, the actions of one firm are easily recognized by
5 its competitors. In a large permit market it is less likely that any one firm can dominate the market.
6 Most important, the greater the number and diversity of firms, the greater is the likelihood that
7 there will be differences among firms in the cost of reducing emissions and, hence, greater
8 potential gains from trading.

10 While a greater number of participants may offer greater potential cost savings, it also makes
11 monitoring and enforcement more difficult. In cases where regulators cannot measure emissions
12 accurately -- either because of technological limitations or because there are a very large number of
13 emitters (e.g., small sources and automobiles) -- they are likely to prefer specific technology
14 controls to incentive-based approaches. Technology standards ensure that the concentrations of
15 pollution from a facility will meet the design value of the controls in place. The drawback,
16 however, is that since the use utilization of the facility or vehicle may vary, this approach will not
17 allow regulators to achieve a cap on the total volume of emissions.

19 The environmental consequences of pollution may depend on where and when emissions occur.
20 For some pollutants, a ton of emission will have basically the same effect on the environment,
21 regardless of its location or source. These type of pollutants are called "uniformly mixing," and
22 their homogeneity broadens the potential market by expanding the realm for trades among greater
23 numbers of emitters. Ozone-depleting substances and greenhouse gases are good examples of
24 uniformly mixing air pollutants, because their effects on the environment do not depend on the
25 individual sources of emissions.

27 On the other hand many pollution problems have important local attributes. For example, NO_x
28 emissions contribute to ground level ozone, which is a local and, sometimes, a regional problem.
29 The role of NO_x in ozone creation depends not only on where emissions occur, but also on the
30 presence of other pollutants and on the season and the time of day when emissions occur. At the
31 same time, NO_x emissions contribute to other environmental problems--such as particulate
32 pollution and nitrogen deposition--that have distinct spatial characteristics.

34 Incentive-based approaches, such as tradable permits, can be designed to overcome local concerns
35 in a variety of ways. A trading program may restrict trading between geographic zones, or may
36 allow trading among zones at ratios that reflect the relative environmental damage that results from
37 a unit of emissions at each location. Similarly, concerns about the timing of emissions can be
38 addressed through trading rules that restrict increases in emissions during certain time periods. In
39 some cases, these approaches have been applied successfully. However, one lesson from previous
40 experience is that restrictions on trading programs detract from their likely success because they
41 limit the breadth or scope of the market and raise transaction costs.

43 In considering the applicability of trading programs to various pollution problems, there appears to
44 be a simple trade-off. The more localized over space or time is the environmental effect of concern
45 is, the smaller the potential market will be, and the less likely trading programs will offer
46 significant potential cost savings.

48 On the other hand, experience suggests that it is not always necessary to allow local concerns about
49 the spatial and temporal effects of emissions to narrowly determine the ultimate design of a trading
50 program. In some cases, the cost savings from a broader-scale, less encumbered trading program
51 outweigh local environmental concerns and ultimately lead to net gains for environmental
52 protection. 2) Moreover, in some cases very simple safeguards are sufficient to protect against
53 most adverse local effects, while not significantly restricting the scope for trading. For example,

1 although the local effects of some pollutants vary greatly, the lion's share of this variation may be
2 controlled by a simple system of trading zones that generally limits the pattern of trading.

4 **Elements That Led to Success of the SO₂ Program**

6 Several features of the SO₂ problem make it a good candidate for a permit trading program. Among
7 these is the large number of electricity-generation facilities that are responsible for a significant
8 majority of all SO₂ emissions and that provide ample potential buyers and sellers of pollution
9 permits. The physical differences among these facilities and their fuels correspond to the great
10 variation in the cost--and cost-saving opportunities of reducing emissions.

12 An interesting feature of the SO₂ program is that it is a national trading program, absent restrictions
13 on trading between or among facilities in different locations. However, the environmental
14 problems caused by SO₂ are regional in nature and relate to the geographic source of emissions.
15 For instance, emissions from a coal-fired power plant in the Ohio Valley make a greater
16 contribution to acidification of the Adirondacks than do emissions from a plant in Mississippi.

18 By treating all emissions equally, the program has extended the breadth of the market and has
19 opened up greater opportunities for trading. The presumption is that opportunities to reduce costs
20 and to apply those savings to achieve greater emissions reductions outweigh the possibility that
21 pollution control will be greater or less in some locations than in others. Any differentiation in the
22 benefits in local areas should be measured against this substantial reduction overall.

24 The program also affords significant potential cost savings by allowing sources of SO₂ emissions
25 to bank their emission reductions in one period to offset necessary future reductions.
26 Environmental problems associated with SO₂ emissions are not particularly sensitive to the timing
27 of emissions. For example, the ecological impacts of acidification are primarily the result of
28 accumulated deposition of sulfur. The secondary particulates that cause health effects are evident
29 in the atmosphere for time periods that extend over several days, which also helps to mitigate the
30 effects of changes in emissions at a particular time.

32 One further important ingredient in the successful formula for SO₂ trading is the availability of
33 continuous emissions monitoring systems, which provide a means to ensure compliance when
34 regulators are not sure what actions should be taken by individual firms. Indeed, compliance has
35 been achieved by 100% of facilities affected under the first two years of the Title IV program.

37 **Closely Related Experiments**

39 Several closely related experiments on a somewhat smaller scale also provide lessons about the
40 potential viability of permit trading programs in different settings. One of the earliest and most
41 successful was the program to phase out lead in gasoline. In the 1980s, EPA set a schedule to
42 virtually eliminate lead in gasoline. To achieve this schedule, it allowed trading among refineries in
43 order to obtain, on average, an average lead content in gasoline specified in their 1983-1987 phase
44 out time schedule. The program also allowed refineries to bank their lead rights, so that a refiner
45 could lower the average lead content of its fuels ahead of schedule in order to fall behind schedule
46 at a later point in time.

48 The primary success in the case of the lead phase out, as in the case of any environmental program,
49 rests in achieving its environmental goal. However, the trading aspect of the phase-out program
50 contributed to this success because it helped to reduce the cost of achieving the environmental goal
51 and, in so doing, helped build consensus among industry and consumers for the viability of that
52 goal. Environmental concerns were not significantly affected by the program, and monitoring lead
53 content and measuring of performance were straight forward. Further, although some variation in

1 the timing and geography of emissions inevitably resulted under the program, there is no evidence
2 that it was significant.

4 Another closely related experiment is the phase down of chlorofluorocarbons and halons--
5 collectively described as ozone-depleting chemicals--under international agreements to which the
6 United States has been a signatory. The ultimate goal is to phase out or severely limit the use of
7 these chemicals. To achieve this goal, EPA issued regulations to control both the production and
8 consumption of ozone-depleting chemicals through a quota system that allocated tradable quotas to
9 producers and consumers in proportion to their historic levels of use. Producers need to use both
10 types of allowances, while importers only need to use consumption allowances. Allowances may
11 be traded domestically or internationally among signatory countries.

13 An interesting aspect of this phase out was concern that the decreasing availability of ozone-
14 depleting chemicals would raise their profitability. In response, Congress imposed a tax on these
15 chemicals in part to capture some of the windfalls that would result from their increased scarcity,
16 and also to promote development of substitutes.

18 The ozone-depleting chemical program is widely viewed as a success. The cost of the phase-out
19 has not been exorbitant, due in part to the large number of substitutes that have been brought to the
20 market, allowing the phase-out schedule to be moved forward in some cases. One class of ozone-
21 depleting chemicals was eliminated in 1996, except for very limited uses. A second class of
22 chemicals is to be phased out in the early decades of the next century.

24 A third experiment that has been repeated in a number of local settings is the use of emission-
25 reduction credits for criteria air pollutants. These experiments have taken a number of forms.
26 "Offset" programs allow a new source of emissions locating in an urban areas in violation of the
27 national ambient quality standards to obtain "offsets" for new emissions by reducing emissions at
28 older sources. The "bubble" program allows new or modified emission sources to avoid stricter
29 new source performance standards as long as total emissions from an entire industrial facility do
30 not increase as a result of the changes. In some cases, of emission reduction credits may be
31 "banked" for subsequent use.

33 Most analyses of the numerous efforts to promote local emission reduction credit programs have
34 concluded that the programs have fallen short of their expectations. One reason is the programs
35 have had to accommodate the geographic and time-sensitive nature of emissions of criteria air
36 pollutants in urban nonattainment areas. Also, there has been concern that offsets may be generated
37 from reduced activity at facilities that were about to shut down anyway, and that trading a credit to
38 a new facility effectively increased emissions in an urban area. The result of these concerns has
39 been a variety of controls on the nature of trading that limit the scope of the market, effectively
40 raising transaction costs and reducing the volume of trading.

42 Programs other than emission-reduction credit programs are geared toward controlling the total
43 quality of emissions. For instance, both the SO₂ and ozone-depleting chemicals programs cap
44 overall emissions. The lead phase-out program comes close to doing the same thing, because there
45 was expected to be relatively little variation in refinery production over the brief period when the
46 phase-out was achieved. However, the emission-reduction credit programs are calibrated with
47 emission rates (tons of pollutant per volume of output), rather than emission quantities (tons of
48 pollutant per year). This provides no guarantee that emissions will not increase with intensified
49 economic activity.

51 That possibility meant the program had to impose additional constraints to make sure emissions
52 remain stable or decrease. These constraints undermined the performance of the emission-
53 reduction credit programs to some extent. This approach was taken in many cases, since it was not
54 possible for environmental agencies to consider the alternative approach of capping emission

quantities. For many types of emission sources, monitoring of emissions has not been established, and a historic emissions profile is not known with which a baseline can be established, making an overall cap unattainable.

In general, as noted previously one finds the greater number and type of sources emitting a pollutant, the more difficult it is to monitor emissions. This makes it difficult to design an incentive-based approach to regulate emissions, and more likely that a traditional command and control approach will be necessary. Given the inability to monitor total emissions, regulators have relied on the use of specific technologies to ensure that emission rates are controlled, even though total emissions are not known or controlled with certainty.

New Applications of Emission Trading Programs

Two air pollution problems have recently attracted growing attention as potential new applications of emission trading programs. These two problems—CO₂ and NO_x emissions—are aligned at opposite ends of the spectrum with respect to the breadth of their environmental impacts. CO₂ emissions contribute to global climate change, regardless of their location. However, the contribution of NO_x emissions to pollution problems depends strategically on the timing and location of those emissions. Also, CO₂ emissions cannot readily be controlled through post-combustion abatement technologies, so emission reductions must be achieved through efficiency improvements and fuel switching. However, NO_x emissions can be controlled through abatement technologies, although significant opportunities also exist for efficiency improvements and fuel switching.

In other ways, the two pollutants are similar. A significant portion of national emissions of both pollutants comes from large electricity-generating facilities and industrial facilities, which are easily monitored. However, a significant portion of both also comes from smaller sources, including vehicles, which are not easily monitored. Furthermore, in both cases there is tremendous variation in the costs of reducing emissions among the various sources, which provides considerable motivation to find ways to design trading programs that can overcome these obstacles.

The prospects for a CO₂ trading program were significantly bolstered by the Draft Protocol Framework for an international agreement proposed by the United States in January 1997. The protocol promotes the use of permit trading for CO₂ reductions among so-called Annex A and Annex B countries, which roughly correspond to the more developed economies. Furthermore, the draft protocol calls for expanded use of "joint implementation" between Annex A/B and other countries, allowing more developed countries to invest in projects in less developed countries to generate requisite CO₂ emission reductions. The prospect for the proposed CO₂ trading program is very uncertain at the time of this report's publication, as well as is the potential design for such a program. However, the differences in the cost of emission reductions among potentially affected sources and different countries is enormous. These differences argue strongly for designing a program that will allow the international community to reduce emissions more cost-effectively than uniform national approaches would allow.

At the other end of the spectrum are local experiments to reduce NO_x emissions. Previously established emission reduction credit programs for NO_x and other pollutants are delivering important cost savings. However, some of their conditions have prevented them from achieving widespread success. And, as with the use of traditional technology standards, they cannot necessarily contain the level of emissions that result.

Increasingly, regulators are considering the use of cap and trade programs to explicitly limit the total quantity of emissions and to allow flexibility in attaining this goal. However, these trading programs must consider the local nature of the NO_x pollution--hence, calling for local or regional

1 markets, sometimes with restrictions on the direction or timing of trading. Consequently, an
2 emission-trading program will produce less savings than a market covering a larger geographic
3 area with more potential traders. However, there are such a large number of potential sources of
4 NO_x emission reductions that cost savings could still be great. Regulators are striving to design
5 programs that maintain environmental safeguards while providing incentives to capture these
6 potential savings.

8 Table A-1 provides a status report on a number of efforts to establish markets for NO_x trading. In
9 some of these cases, regional markets have been established that are significantly large enough to
10 overcome this geographic limitation. For instance, the RECLAIM program in Los Angeles has
11 over 500 participants. Others, such as the Texas market, have been so small as to all but prevent
12 trading between firms, though some trading among facilities within the same firm has occurred.

14 The regional nature of the environmental impacts of NO_x emissions also has caused other
15 restrictions to be placed on NO_x trading. The southern California RECLAIM and emerging
16 Ontario, Canada markets can only trade in directions that reflect regional wind patterns. The Texas
17 market has regional boundaries, which trades are not allowed to cross, and the proposed
18 northeastern market may have a similar form. These restrictions limit the amount and the
19 corresponding potential gains from trading.

21 As mentioned, NO_x emissions come from a wide variety of sources that include mobile and
22 stationary emitters, small and large sources, and facilities that may be on the geographic fringe of
23 the emission trading market. When many small and mobile sources exist in the emission-trading
24 market, it is difficult to include all of them because of difficulties in monitoring and administering
25 the program. Nonetheless, in some cases, the small sources can contribute significant emissions to
26 the local airshed. Hence, regulators tend to rely on conventional approaches to control emissions
27 from numerous, small sources, which typically control emission rates but not the overall level of
28 emissions from these sources. Their exclusion from the trading program misses not only
29 significant opportunities for cost savings, but also significant opportunities for emission reductions
30 and, hence is an issue attracting increasing attention.

32 Episodic constraints are also an important issue in NO_x trading markets. In contrast to the SO₂
33 emission trading market, some experts believe that the potential of emission "spikes" from NO_x are
34 a key issue. Some markets have restrictions on the amount of trading or banking between seasons
35 of the year. Additional episodic restrictions placed on NO_x trades include limitations on when
36 banked credits may be used.

APPENDIX C

Interpretation of "Adverse" Effects for NAPAP Report to Congress

Summary

The 1990 Clean Air Act Amendments require that the National Acid Precipitation Assessment Program (NAPAP) prepare biennial reports to Congress, and that "every four years . . . the report . . . shall include the reduction in deposition rates that must be achieved in order to prevent adverse ecological effects." Public Law 101-549, Title IX, Section 903 (j)(3)(F)(i), [codified as amended at 42 USC §7403(j)(3)(F)(I)]. This 1996 report is the first report to address this requirement.

Although the term "adverse ecological effects" is not specifically defined in the Clean Air Act Amendments, a working definition can be derived from relevant statements at various locations in the statute. Congress expresses its concern with ecological components (the scope is broad and inclusive, since ecology encompasses the interrelationships of organisms and their environment) in the preceding subsection (E) of the statute. It requires reporting on "the status of ecosystems (including forest and surface waters) . . . affected by acid deposition . . . including changes in surface water quality and forest and soil conditions . . . [and] high elevation watersheds 42 USC §7403(j)(3)(E)(iii-v). The adverse effects of concern to Congress, as evidenced in its findings and declaration of purpose (Clean Air Act (CAA) 42 USC §7401(a)), are the "dangers to the public health and welfare . . . including injury . . . damage . . . and . . . deterioration."

Conclusions

Based on the intent of Congress as expressed above and elsewhere in the Clean Air Act Amendments, and shaped by indications of intent expressed in other relevant environmental statutes and regulations, the following working definition of "adverse ecological effects" has been derived and is used in the preparation of this report: any injury (i.e., loss of chemical or physical quality or viability) to any ecological or ecosystem component, up to and including at the regional level, over both long and short terms. Similarly, adverse effects for other areas of concern addressed in this report--i.e., visibility, materials, and human health --consist of loss of quality up to and including at the regional level, over both long and short terms.

Background

Ecological components of concern to Congress are addressed in the Clean Air Act Amendments section titled "Research, investigation, training, and other activities" 42 USC §7403. In the subsection that includes the provision for the continuation of NAPAP, the ecological components mentioned include ecosystems, forests, surface waters, soil, and high-elevation watersheds; ecological effects that could be adverse include changes in surface-water quality, changes in forest and soil conditions, and occurrence of episodic acidification (especially in high-elevation watersheds) 42 USC §7403(j)(3)(E)(iii-v). Additional ecological components and attributes of interest to Congress with respect to adverse effects are named in a preceding Clean Air Act Amendment subsection titled "Ecosystem research" (42 USC § 7403(e)), and include "regionally representative and critical ecosystems . . . crops, biological diversity, wetlands, estuaries, groundwater, other terrestrial systems, and other aquatic systems"; effects named that could be adverse include those that are "short-term and long-term . . . [show] trends of ecosystem damage . . . [are due to] chronic and episodic exposures . . . [and] multiple environmental stresses." Also, "sensitive and critically sensitive aquatic and terrestrial resources" are the subject of specific congressional protection to be achieved through the adoption of an acid deposition standard(s) (Appendix B of § 404 (42 USC § 7651c) of Title IV of the 1990 Clean Air Act Amendments).

Nowhere in the Clean Air Act Amendments, or associated case law, is the specific type of damage or injury that would constitute an adverse ecological effect specified. Other environmental statutes, however, deal with similar concepts. The Comprehensive Environmental Response,

Compensation, and Liability Act (CERCLA) assigns liability for damage to natural resources from releases of hazardous substances, a broad concept encompassing adverse ecological effects. The term *damages* is defined as "injury or loss of natural resources." Natural resources are "land, fish, wildlife, biota, air, water, groundwater, drinking water supplies, and other such resources" 42 USC §9601(6,16). The regulations delineating how natural resource damage assessments may be carried out for CERCLA (and also the Clean Water Act (CWA)), are at 43 CFR §11.10 et seq., and define injury as:

a measurable adverse change, either long- or short-term, in the chemical or physical quality or the viability of a natural resource resulting either directly or indirectly from exposure to . . . a hazardous substance, or . . . to a product of reactions resulting from . . . a hazardous substance. 43 CFR §11.14(v).

There are, in turn, detailed injury definitions applicable to specific resources. Portions of these definitions potentially relevant to ecological effects of acid deposition are at 43 CFR §11.62 and include:

(e) Geologic resources. An injury . . . has resulted . . . if one or more of the following changes . . . is measured:

(4) Concentrations of substances sufficient to decrease the water holding capacity such that plant, microbial, or invertebrate populations are affected;

(5) Concentrations of substances sufficient to impede soil microbial respiration to an extent that plant and microbial growth have been inhibited;

(6) Concentrations in the soil of substances sufficient to inhibit mineralization resulting from a reduction in soil microbial populations;

(7) Concentrations of substances sufficient to restrict the ability to access, develop, or use mineral resources within or beneath the geologic resources

(9) Concentrations in the soil of substances sufficient to cause a toxic response to soil invertebrates;

(10) Concentrations in the soil of substances sufficient to cause a phytotoxic response such as retardation of plant growth.

(f) Biological resources. (1) An injury . . . has resulted . . . if concentration of the substance is sufficient to:

(i) Cause the biological resource [fish and wildlife and other biota] or its offspring to have undergone at least one of the following adverse changes in viability: death, disease, behavioral abnormalities, cancer, genetic mutations, physiological malfunctions (including malfunctions in reproduction), or physical deformations.

Since natural resources are ecological components, and the injuries are adverse effects, these definitions give a further indication of adverse ecological effects that lie within the intent of Congress within the environmental statutes of CERCLA and the CWA.

Incorporating the content and concepts above, the adverse ecological effects of acid deposition that could lie within the scope of this NAPAP report are those effects that cause--

injury, damage, or deterioration

72 *consisting of*
73 a measurable adverse change, either long- or short-term, in the chemical or physical quality
74 or the viability of

75 ecosystems (sometimes causing trends), regionally representative ecosystems, critical
76 ecosystems, other terrestrial systems, other aquatic systems, sensitive and critically
77 sensitive aquatic and terrestrial resources, forests, surface waters, wetlands, estuaries,
78 groundwater, high elevation watersheds, soil, crops, biological diversity, land, fish,
79 wildlife, biota, air, water, drinking water supplies, and other such resources

80 *due to*

81 occurrence of episodic and chronic exposures, short- and long-term exposure, multiple
82 environmental stresses; and resulting either directly or indirectly from exposure to acid
83 deposition, or exposure to a product of reactions resulting from acid deposition,

84 *and including for* geological resources

85 decreases in the water holding capacity such that plant, microbial, or invertebrate
86 populations are affected; impedance of soil microbial respiration to an extent that plant and
87 microbial growth have been inhibited; inhibition of mineralization resulting from a
88 reduction in soil microbial populations; restrictions in the ability to access, develop, or use
89 mineral resources within or beneath the geologic resources; toxic responses to soil
90 invertebrates; and phytotoxic responses, such as retardation of plant growth;

91 *and including for* biological resources

92 changes sufficient to cause the biological resource or its offspring to have undergone at
93 least one of the following adverse changes in viability: death, disease, behavioral
94 abnormalities, cancer, genetic mutations, physiological malfunctions (including
95 malfunctions in reproduction), or physical deformations.

96 From this detailed, descriptive, lengthy, and sometimes redundant definition the working definition
97 of adverse ecological effects in the text was derived.

[Image][Image]

[Image] Overview | Effects | SO₂ Program | NO_x Program | Emissions | Conservation

Acid Rain Program Information: Allowance Tracking System

EPA is responsible for recording the transfer of allowances that are used for compliance and for confirming that utilities hold at least as many allowances as tons of SO₂ emitted by the end of the year. To fulfil this role, EPA designed and maintains the Allowance Tracking System (ATS), a computer program that is the official record of allowance holdings and transfers.

File Descriptions

Information contained in ATS is provided here in several files, with the specific file name in parentheses:

Allowances Held by Each Account (ALLOW.EXE):

Account number, account name, allowance use year, quantity of allowances held for the use year, starting serial number, and ending serial number (290 kB)

Allowance Transactions (TRANNEW.TXT):

Transaction number, transaction type, transferror account number, transferror name, transferee name, indication of allowances transferred in perpetuity, date recorded, total number of allowances transferred, allowance use year, starting serial number, ending serial number, and number of allowances within the specified block. For allowance transactions since June 6, 1997 (250 kB)

Allowance Transactions (TRANOLD.EXE):

The same categories as found in TRANNEW.TXT, but for allowance transactions prior to June 6, 1997 (1,300 kB)

Allowance Accounts (ACCOUNT.TXT):

Account number, account name, state, account representative ID, and alternate representative ID (120 kB)

Account Representatives (AAR.TXT):

Representative ID, representative name, address, phone, and fax number (150 kB)

Binding Parties (OWNER.TXT):

Account name, account number, binding party (owner) number, and binding party name (225 kB)

Another file, UPDATE.txt, provides information on changes in ATS

since the previous weekly update.

PURPOSE

As EPA's official record of allowance accounts, representatives, transactions, and allowance holdings, ATS information is critical to the effective operation of the allowance market and the Acid Rain Program. Because the Program is a testing ground for market-based regulation, this information is of keen interest to public groups, especially the environmental community. This information is also integral to the missions of other Federal and State regulatory agencies that regulate the generation and dispatch of electric power.

ORIGINATOR: Acid Rain Division

PUBLICATION DATE: Not applicable

ACCESS CONSTRAINTS: None

[Image]

AVAILABILITY:

- a. Distributor: Hardcopy information is available from EPA's Acid Rain Division.
- b. Order process: Information can be requested by calling the Acid Rain Hotline, (202) 564-9620.
- c. Technical Prerequisites: Files are provided as both text (in comma-delimited format) and compressed (.exe) format. The usefulness of these files is enhanced by using a spreadsheet or database program.
- d. Automated Linkage: Not applicable

COVERAGE: Regulated entities are located within the continental United States, most densely in the Midwest and Mid-Atlantic. Initially, no query capability is available, but will be provided eventually.

TIME PERIOD OF COVERAGE: Information is cumulative since the start of the Program.

Point of Contact for Further Information:

Melanie Dean (6204J)

U.S. Environmental Protection Agency

Office of Air and Radiation

Office of Atmospheric Programs

Acid Rain Division

Permits and Allowance Market Branch

401 M Street, SW
Washington, DC 20460

Phone: 202-564-9189

Email: dean.melanie@epamail.epa.gov

Cataloging Source:
Alexander Salpeter (6204J)
U.S. Environmental Protection Agency
Office of Air and Radiation
Office of Atmospheric Programs
Acid Rain Division
Permits and Allowance Market Branch
401 M Street, SW
Washington, DC 20460

Phone: 202-564-9157

Email: salpeter.alex@epamail.epa.gov

DATE OF CREATION: April 21, 1995

DATE OF LAST MODIFICATION: December 10, 1997

ELEMENTS FOR INFORMATION SYSTEMS

- a. Agency program: Acid Rain Program (Title IV, Clean Air Act)
- b. Sources of Data: Regulated entities, mainly power plants

AGENCY SUPPLEMENTAL INFORMATION: Initially, the information will be updated weekly, and eventually will be revised daily.

EXPIRATION DATE: Not applicable

Acid Rain Program Home | ATS Data | Contact Us
<http://www.epa.gov/acidrain/ats/atsdetl.html>