

FOIA MARKER

This is not a textual record. This is used as an administrative marker by the William J. Clinton Presidential Library Staff.


Collection/Record Group: Clinton Presidential Records
Subgroup/Office of Origin: National AIDS Policy Office
Series/Staff Member: Patsy Fleming
Subseries: National AIDS Strategy

OA/ID Number: 9003
FolderID:


Folder Title:
CDC [Centers for Disease Control] Charts

Stack:	Row:	Section:	Shelf:	Position:
S	66	5	11	2


Death Rates from Leading Causes of Death in White Men Aged 25-44 Years, USA, 1982-93




Death Rates from Leading Causes of Death in White Women Aged 25-44 Years, USA, 1982-1993




Death Rates from Leading Causes of Death in Black Men Aged 25-44 Years, USA, 1982-1993



Death Rates from Leading Causes of Death in Black Women Aged 25-44 Years, USA, 1982-1993




Death Rates from Leading Causes of Death in Children 1-4 Years of Age, United States, 1983-1992




Source: National vital statistics

Death Rates from Leading Causes of Death in Men Aged 25-44 Years, USA, 1982-1993



Death Rates from Leading Causes of Death in Women Aged 25-44 Years, USA, 1982-93



Women with AIDS

Mode of transmission by Age at Diagnosis


United States, Reported in 1994

Mode of Transmission	Age at Diagnosis			
	13-19 (n=191)	20-29 (n=3935)	30-49 (n=11,311)	50+ (n=1351)
Injection drug use	11%	37%	55%	25%
Heterosexual contact	66%	49%	33%	45%
Transfusion recipient	5%	2%	2%	13%
No identified risk*	18%	12%	9%	17%

*Includes patients under investigation; patients who died, were lost to follow-up, or refused interview; and patients whose mode of exposure to HIV remains undetermined after investigation


Women with AIDS

Incidence and Percent of Adult AIDS Cases by Year, United States, 1981-1994*



*Data adjusted for reporting delay and the 1993 change in the AIDS case definition

AIDS Incidence, and Estimated AIDS-opportunistic Illness Incidence, Adjusted for Delays in Reporting, by Quarter-year of Diagnosis, January 1986 through June 1994, United States



Adult Male AIDS Cases and Annual Rates per 100,000 Population by Race/Ethnicity, Reported in 1994, United States

<u>Race/Ethnicity</u>	<u>Cases</u>	<u>Rate</u>
White, not Hispanic	29,910	38.9
Black, not Hispanic	22,838	208.0
Hispanic	12,016	109.8
Asian/Pacific Islander	518	15.3
American Indian/ Alaska Native	184	26.8
<u>All men*</u>	<u>65,591</u>	<u>63.7</u>


* Includes 125 men whose race/ethnicity is unknown.

Adult Female AIDS Cases and Annual Rates per 100,000 Population by Race/Ethnicity, Reported in 1994, United States

<u>Race/Ethnicity</u>	<u>Cases</u>	<u>Rate</u>
White, not Hispanic	3,148	3.8
Black, not Hispanic	8,016	62.7
Hispanic	2,814	26.0
Asian/Pacific Islander	49	1.3
American Indian/ Alaska Native	42	5.8
<u>All women*</u>	<u>14,081</u>	<u>12.8</u>

* Includes 12 women whose race/ethnicity is unknown.

Adult Cases of AIDS by Transmission Category Reported in 1994, United States




- Homosexual/bisexual men (44%)
- Injecting drug user (27%)
- Homosexual and injecting drug user (5%)
- Hemophilia cases (1%)
- Heterosexual contact (10%)
- Transfusion recipient (1%)
- Under investigation (12%)*
- Other/undetermined (<1%)


*Most patients have a risk identified after follow-up

† Includes patients who declined interview, died, or were lost to follow-up, and whose HIV exposure remains undetermined

Pediatric AIDS Cases Reported in 1994, United States



Cases of AIDS by Age and Gender, Reported 1981 through 1994, United States




AIDS-indicator Conditions in Adult Patients Reported in 1994, United States

<u>Most Frequently Reported Conditions</u>	<u>Percent of Cases</u>
Immunosuppression, severe HIV-related	81
<i>Pneumocystis carinii</i> pneumonia	19
HIV wasting syndrome	10
Candidiasis of esophagus	7
<i>Mycobacterium tuberculosis</i> infection	5
Kaposi's sarcoma	4
<i>M. avium</i> or <i>M. kansasii</i> infection	3

Many patients have more than one condition.

HIV Infections (not AIDS) Reported in 1994, from States with Confidential HIV Infection Reporting



* Includes 623 persons reported from states with confidential HIV reporting who were residents of other states.

Adult/Adolescent AIDS Annual Rates per 100,000 Population, for Cases Reported in 1994, United States

