



WORKING PAPER NO. 46
November 2015



Occupancy-related metrics for wildlife status assessment

Dr. Timothy O'Brien, Dr. Samantha Strindberg, Dr. Robert Wallace

We Stand for Wildlife

WCS envisions a world where wildlife thrives in healthy lands and seas, valued by societies that embrace and benefit from the diversity and integrity of life on earth.

WCS uses science to discover and understand the natural world. This knowledge helps us engage and inspire decision-makers, communities and millions of supporters to take action with us to protect the wildlife and wild places we all care about.

SAVING WILDLIFE

WCS scientists study what wildlife need to thrive. With this knowledge we invest in abating threats to wildlife within their most important strongholds and the corridors that connect them. We target large, iconic, wide-ranging wildlife because they are vital to ecosystem health and, by saving them, we protect all other biodiversity that shelters under their conservation canopy.

SAVING WILD PLACES

Over the past century, WCS has established long-term conservation presence in the last wild places across the Americas, Africa, Asia and Oceania, built strong and trusting partnerships, and acquired a depth of knowledge that ensures effective conservation action. We protect these last wild places because they are intact, biodiverse, most resilient to climate change, and bastions for large, iconic wildlife.

WCS WORKING PAPER No. 46
NOVEMBER 2015

Occupancy-related metrics for wildlife status assessment

By Tim O'Brien, Samantha Strindberg, Robert Wallace

Dr. Timothy O'Brien
Wildlife Conservation Society
Conservation Measures
Email: tobrien@wcs.org

Dr. Samantha Strindberg
Wildlife Conservation Society
Conservation Measures
Email: sstrindberg@wcs.org

Dr. Robert Wallace
Wildlife Conservation Society
Bolivia Program
Email: rwallace@wcs.org

WCS Working Papers:

Print ISSN 1530-4426

Online ISSN 1534-7389

Copies of the WCS Working Papers are available at
http://ielc.libguides.com/wcs/library_wps

Suggested citation:

O'Brien, T., Srindberg, S., and R. Wallace. 2015. Measuring conservation effectiveness: occupancy-related metrics for wildlife. WCS Working Paper No. 46 New York: Wildlife Conservation Society.

Cover photograph: Bolivian portion of Greater Madidi-Tambopata Landscape with 81 km² Occupancy Analysis Cell Design for Jaguar (*Panthera onca*)



Copyright:

This work is licensed under a [Creative Commons Attribution-NonCommercial-ShareAlike 4.0 International License](https://creativecommons.org/licenses/by-nc-sa/4.0/)

Table of Contents

Executive Summary	iv
Introduction	1
Results-based management	1
The five measures of our impact	2
The workshop	3
Occupancy and abundance for conservation monitoring	6
Flavors of occupancy	9
WCS monitoring and decisions making in the face of uncertainty	10
Examples of occupancy-based monitoring at WCS	11
Tigers forever: baselines for a large-scale occupancy by tigers and tiger prey	11
Conservation effectiveness of patrolling and law enforcement in BBSNP	12
Livestock management and large mammals in Laikipia County, Kenya	14
Modeling wolverine occurrence using aerial surveys of tracks in snow	15
Monitoring rare species in the Andean Patagonian Steppe landscapes	16
Occupancy methods to assess changes in species fish diversity on coral reefs	17
Migratory species	18
Design considerations for occupancy methods	19
The way forward	23
Literatures cited	24
Appendix 1: A brief overview of occupancy methods	26
Appendix 2: A short bibliography of occupancy methods	33
Methods and general	33
Amphibians	39
Birds	42
Mammals	48
Reptiles	54
Invertebrates	55
Plants	55
Other journals with recent papers on occupancy	56
Acknowledgements	58

Executive summary

WCS Global Conservation Program is exploring the development of measures of conservation effectiveness focusing on how WCS management interventions affect wildlife populations, wildlife habitat, threats to wildlife and law enforcement effectiveness, natural resource governance, and livelihoods. Here we report on a two-day internal workshop in August 2012, where we invited WCS staff and friends with expertise in occupancy analysis and monitoring to examine the use of occupancy as a metric for within and cross-site comparisons of conservation effectiveness in WCS programs. Occupancy methods are potentially efficient in terms of their application across large areas and the ability to combine multiple sources of information, while providing a scientifically defensible metric of the proportion of a landscape of interest that is occupied. Occupancy-based analyses also can yield information on species richness, relative abundance, and dynamics of species and communities; explanatory variables for detectability and occupancy can be included during analysis. In addition, the results of occupancy analyses can be displayed in a spatially explicit manner, which permits conservation decision makers or donors to easily visualize how conservation management is influencing the status of conservation targets in space and time.

This document highlights some of the key results of the meeting, as well as provides a background to the uses of occupancy analysis in a variety of landscape contexts. In the appendices we give an overview of occupancy methods and a bibliography of recent uses of occupancy models for a variety of taxa.

- ◆ We agreed that WCS should use unbiased estimates of occupancy, abundance, density or relative abundance whenever possible.
- ◆ We reached a broad consensus that occupancy will be a useful metric for comparing landscape/seascape level wildlife trends between sites and over time.

- ◆ We also recognized the limitations of occupancy to provide information on certain kinds of species that are important to our landscapes and seascapes.
- ◆ We did not suggest that occupancy be the only metric of conservation effectiveness or that current best practices be abandoned in favor of occupancy.
- ◆ We recognized that many interventions are indirect and often we are not the management authority in our Scapes. Under these circumstances, it may be hard to attribute changes in target species abundance and occupancy to our intervention efforts.
- ◆ We concluded that occupancy would be a useful metric for the majority of target species in the WCS landscape and seascape portfolio.
- ◆ We suggested four activities to move forward on the use of occupancy as a WCS landscape monitoring metric:
 - ◆ Investigate retro-fitting occupancy analyses to current sampling designs and datasets.
 - ◆ Encourage collaboration within and among Regional Programs to increase the power of their monitoring programs by standardizing monitoring data (The development of the camera trap database is a start).
 - ◆ A series of workshops at NCEAS under the NCEAS/TNC/WCS NATURELAB program to investigate structured decision making, monitoring, and assessment of conservation effectiveness in WCS Scapes.
 - ◆ Development of test landscapes in each region (NA, LA, Asia, Africa, Marine) where we engage in occupancy analyses to establish the utility of the efforts.

Introduction

Results-based management

Within WCS, we advocate the practice of results-based management. This is the process by which, given a management action, we measure effectiveness and adapt project strategies as we learn what does and does not work in promoting conservation. The keys to results-based management are 1) Use of conceptual models to make explicit a results-chain that specifies causal connections between our measurable objective, the direct and indirect barriers that presently prevent us from achieving our objectives and the strategies that we will deploy to take down these barriers; and 2) monitoring how well we implement our chosen strategies, tracking whether threats and barriers are being abated and overcome, and assessing whether the status of our conservation and development targets (our objectives) is improving, declining or static.

By monitoring at three levels - Strategies (also called activities, management or interventions), Threats (and threat reduction) and Targets (wildlife populations response to strategies), we can assess the effectiveness of our investments and adapt our strategies as we learn what works and as the situation changes (Strindberg and O'Brien 2012). Thus, we need to be able to track the status of our principal conservation targets, the wildlife species we care about, habitats and threats, as well as additional information that is of perennial interest to major donors. At the same time we want to keep our measures to a minimum because there is a strong inverse relationship between the number of indicators to be measured and the probability that they will be measured well.

In selecting impact measures we need to ensure that: 1. they are perceived as transparent and credible (i.e., reported results will be believed); 2. they can be collected at modest additional cost; and 3. they are able to show change over a relatively short time frame.

Recently, WCS Conservation Support was asked to develop a set of impact measures to assess the conservation impact of WCS in its entirety.

Five measures of our conservation impact

We are proposing a five measure dashboard to aggregate information from our landscapes, seascapes, and species programs in an effort to assess and report the conservation impact of our organization as a whole. The dashboard will have the capacity to track changes in our conservation targets (wildlife), their habitat, level of threats, law enforcement effectiveness, natural resource governance, and livelihoods.

We propose to produce a graphical WCS Conservation Assessment on programs.wcs.org that provides a measure of our global impact. The Assessment will report a standard set of terrestrial and marine impact measures at each of our 74 landscapes and seascapes, as well as for our Species Programs (clearly Scapes or Species Programs may choose to measure a broader range of factors). The Assessment will be presented as simple, time-series graphs for each individual landscape and seascape (by clicking on a map) and provide a global assessment by aggregating data across landscapes and seascapes, or Species Program elements. Audiences for this Assessment include the Board of Trustees, all NY and field staff at WCS, donors, other conservation NGOs, and the general public.

The Conservation Assessment will do two important things for WCS:

- ◆ Provide a regular, publicly available, accounting of WCS conservation achievements, reinforcing our position as the most effective, science-based, global conservation NGO;
- ◆ Be the guiding impetus, within our organization, for putting in place the systems needed to more easily and

more precisely track, analyze and report our conservation effectiveness.

These measures will be built on the best science and practice available, based on what we are already doing, to ensure that we measure changes that result from our interventions.

Five teams have been put together to develop each of the measures. The remit of each team is to produce the following outputs by 31, December 2012: 1. An analysis of a method or comparable methods that would be deployed across all our Scapes and Species Programs to produce the measure (keeping in mind that the measure needs to be inexpensive but credible, with the premise that less is more because complicated systems are likely to be too costly to implement across appropriate time scales in order to establish and observe trends); 2. Technical guidance for deploying the method or methods within our Scapes and Species Programs; 3. Characterization of a data visualization approach that would make the results of our measures monitoring efforts readily understandable to both a lay and professional audience, and make clear the trends in the measures over time.

In this document, we focus on the progress made in developing a measure to track changes in our conservation targets (wildlife). We convened a workshop to examine the use of occupancy methods as a wildlife metric for within and cross site comparisons of conservation effectiveness in WCS programs. Occupancy methods were considered due to their statistical rigor, potential efficiency of application across large areas and the ability to combine multiple sources of information. They also provide a measure that is of interest for wildlife targets: their changing distribution across the landscape or seascape.

The Workshop

The Occupancy metrics workshop was convened on 2-3 August 2012 to examine the use of occupancy methods as

a metric for within and cross-site comparisons of conservation effectiveness in WCS programs. Occupancy may be used to ask questions about whether our interventions have an effect on the target species we care about. Because it explicitly addresses detectability of target species, occupancy provides an unbiased estimate of the proportion of a landscape that is occupied by a species of interest, and is thus an indicator of species distribution, which together with species abundance form the core of target population-level responses to conservation actions. Occupancy-based analyses also can yield information on species richness, relative abundance, meta-population dynamics, dynamics of species and communities. Explanatory variables, or covariates, can also be incorporated in the estimation of occupancy and detection probability allowing us to test hypotheses about interventions and to explore factors that affect occupancy.

Workshop participants were chosen for their expertise in statistics and occupancy analysis (Jim, Aaron, Arjun, Samantha, Tim) and for their interest and experience in landscape monitoring (David, Steve, Rob, Andrés).

David Wilkie: Director, Conservation Measures

Jim Nichols: Senior Scientist, Patuxent Wildlife Research Center, U.S. Geological Survey

Tim O'Brien: Senior Scientist, Conservation Support

Samantha Strindberg: Associate Conservation Statistician, Conservation Support

Steve Zack: Senior Conservationist, Coordinator Bird Conservation, Species Conservation Program

Rob Wallace: Conservation Scientist, Latin America Program

Andrés Novaro: Conservation Scientist, Latin America Program

Arjun Gopaldaswamy: PhD Student, Oxford University

Aaron MacNeil: Research Scientist, Australian Institute of Marine Science

We were tasked with reaching a conclusion on the following issues:

- ◆ Consensus on an occupancy metric(s) for within and cross-site comparisons;
- ◆ Agreement on methods for relating occupancy metrics to threats and interventions;
- ◆ A clear understanding of the use of covariates in occupancy modeling and how covariates might be standardized across sites;
- ◆ How changes in occupancy relate to changes in abundance and the strength of the relationship.

We agreed that abundance and distribution are the two most important state variables we can use to measure the state of wildlife populations and the impact of WCS interventions. We recognized the value (and challenge) of developing unbiased abundance or relative abundance estimates for parts of large landscapes. We reached a broad consensus that occupancy will be a useful metric for comparing landscape level trends between sites and over time, not just because it may be more feasible sometimes to measure distribution than abundance, but also because of the intrinsic value of having an indicator of the distribution of the target species across the landscape. We also recognized the limitations of occupancy to provide information on certain kinds of species that are important to our landscapes and seascapes. We puzzled over how occupancy relates to the status of extremely rare species, migratory species, nomadic species, and especially pelagic species that may range beyond the bounds of the Scapes or range across such vast areas that we might never be able to afford to achieve adequate coverage to make meaningful inferences. Finally, we recognized that many interventions are indirect and often we are not the management authority in our Scapes. Under these circumstances, it may be hard to attribute changes in target species abundance and occupancy to our intervention efforts. Nonetheless, we concluded that occupancy would be a useful metric for the majority of target species in the WCS landscape and seascape portfolio. We noted that the focus would be on monitoring with-

in Scapes, which might capture some elements of a more comprehensive monitoring program required for tracking the status of conservation targets within the Species Program. Below we summarize our discussions and present some ideas for moving forward with measuring the effectiveness of WCS interventions in our conservation Scapes in terms of a wildlife metric.

Occupancy and Abundance for Conservation Monitoring

The extent of area occupied by a species is an important state variable for conservation. IUCN uses changes in area of occurrence and distribution to guide the listing and change of status of species in the Red List and the USGS uses occupancy in several national and regional monitoring programs. A few definitions are useful to guide our thoughts about occupancy.

Occupancy is the proportion of a region (landscape, study area) that is occupied and can be thought of as the range of the species in the region. Abundance can be expressed as the sum of all patch-specific densities across a region. A species can be present and observed in a sampling unit, present and not observed in a sampling unit or absent from the sampling unit. The more individuals that occur in a sampling unit, the more likely the species will be detected during a sample survey. In IUCN-speak the range of a species is described by the Extent Of Occurrence (EOO – the minimum convex polygon that includes all of the area of normal (or likely) occurrence) and Area Of Occupancy (AOO – the area where the species actually occurs).

Most agree that the positive relationship between occupancy and abundance is real and not a statistical artifact. First, the relationship between AOO and abundance is always positive because zero abundance equals zero AOO and increases from there. Second, a number of studies have shown that species undergoing changes in abundance exhibit a concurrent change in AOO. This principle underlies

the use of occupancy as a surrogate for abundance in IUCN Red Lists, setting harvest rates in fisheries and tracking extinction risk and invasive species in conservation biology. Basically as a species' abundance declines, its AOO shrinks, and as a species' abundance increases, its AOO can be expected to increase.

The strength of AOO-abundance relationship, however, is not linear, is scale dependent, being affected by the size of the study area, the size of the sampling unit and in our case, can be species- and context- specific. The linearity of the relationship decreases as the size of the sampling unit increases; for a given abundance, the AOO often increases with sampling unit size; variance in AOO increases with size of sampling unit and size of the study area. We can envision why these relationships hold. Consider a rare exploited species that is suddenly well-protected. In a study area of s potential sampling units, the species occurs in some number of sampling units (s_o) out of the total study area. As it recovers abundance grows first in the occupied sampling units, then the species disperses into unoccupied sampling units. So a lag may occur between increases in abundance and the AOO of the species. As the species continues to increase in abundance, it fills the study area and AOO reaches an asymptote, even though the population may continue to increase. For a given species, as the sampling unit size increases, it may take longer to observe dispersal into unoccupied cells, but it may become easier to detect the species within a sampling unit, if more individuals result in a more even distribution across the sampling unit. Therefore, in any monitoring effort using occupancy as a metric, special attention must be paid to issues of scale of movements and home range size in relation to the size of the landscape. Clearly whether the species of interest is territorial or not plays a key role in the relationship between occupancy and abundance, as does the aggregating characteristics of the species with increasing group size weakening the relationship between occupancy and abundance.

Traditionally, ecologists considered the AOO as the proportion of sampling units in a study area where at least one

individual was present in a sampling unit. This ignored the state of present but not detected during the sampling effort (detection was assumed to be perfect), and led to estimates of AOO that were biased low because present but not detected cells were lumped with absent cells. To correct the bias, we must estimate the probability p that, if at least one individual is present, it will be detected. Replicated sampling (multiple observers, temporal replication, or spatial replication) allows us to estimate p and correct s_o to obtain an unbiased estimate of AOO, which is represented by the symbol ψ , the probability that a sampling unit is occupied or the proportion of the study area that is occupied.

Methods are available that allow us to estimate p and ψ simultaneously in a maximum likelihood or Bayesian framework. The basic sampling scheme involves multiple visits to all sampling units or a randomly selected subset of potential sampling units in a Scape. Detection/non-detection data are collected during each visit. The sampling is conducted during a time interval short enough that the occupancy state of each sampling unit remains unchanged (the population is closed to change). We also assume that the sampling units and detections are independent. This means that sampling in unit 1 does not affect sampling in unit 2 and that a detection in unit 1 does not change the likelihood of a detection in unit 2. It also means that a detection in unit 1 does not affect the likelihood of subsequent detections in unit 1 (though we have methods to handle with spatial autocorrelation). Furthermore, we assume that species are not misidentified and that occupancy and detection are constant across all sites or that heterogeneity in either detectability, occupancy or both can be modeled with covariates. Fortunately, models exist within the occupancy framework that allow us to relax many of these assumptions.

If we extend this idea to a monitoring framework we would revisit the sampling units over some time interval t . At t_0 and t_1 , we would estimate occupancy. During the secondary sampling period within each primary sampling period, occupancy states of each sampling unit remain constant. Between primary sampling periods, occupancy states may

change. Occupied sites may become unoccupied (local extinction or ε) and unoccupied sites may become occupied (local colonization or γ). Colonization and extinction are the dynamic parameters (also called vital rates) for the state variable occupancy. The change in occupancy between time 0 and time 1 can be described by the relative effect of γ and ε on ψ_0 : $\psi_1 = \psi_0(1 - \varepsilon) + \gamma(1 - \psi_0)$. Conservation interventions affect occupancy over time through their effects on local colonization and extinction.

The strength of occupancy as a monitoring metric is that we can model occupancy, colonization, extinction and detection as functions of covariates. For example, if we are estimating occupancy in a landscape that consists of 4 management regimes that might affect the presence of a target species, we can incorporate “management regimes” as a covariate to estimate the impact of management types on occupancy of the target species. Models incorporating different covariates represent competing hypotheses about factors believed to affect occupancy, colonization and extinction. Over time, multi-year models can be used to evaluate the effectiveness of conservation interventions.

Flavors of Occupancy

There is a wide range of questions and models that can be applied in the occupancy framework (see Appendix 1). In addition to the single season and multi-season occupancy models discussed above, we briefly describe four types of models that may be of interest to WCS for monitoring in wildlife in our Scapes. A recent bibliography of published research using occupancy analyses is included in Appendix 2. Most of the occupancy models can be analyzed in a maximum likelihood or a Bayesian framework.

Species richness models: If our conservation targets include communities of species, one metric of interest is how species richness or the number of species present in our Scape, is changing over time or remaining constant as the result of interventions.

Repeated count models: Occupancy based abundance estimation procedures used to estimate the number of individuals at a point when individuals cannot be identified or marked. Rather than using species presence-absence (detection-nondetection) data, these models are based on counts of individuals obtained at replicate visits. These can be very useful when the sample sites are discrete (i.e. ponds or woodlots) and where the area of sampling can be defined (i.e. fixed distance point count).

Abundance-induced heterogeneity models: Similar to repeated count models but are based on the idea that heterogeneity in abundance generates heterogeneity in detection probability across the Scape. Uses detection/non-detection data to estimate point abundance, and estimates occupancy as a function of point abundance.

Multi-state models: These models are used when we are interested in not only whether a site is occupied, but whether there are different states that the occupied sites might attain. Two examples might be occupancy with breeding and non-breeding birds, or relative abundance surveys that classify occupancy as none, rare, common, and abundant. These analyses move closer to the kind of monitoring that has relevance for traditional WCS monitoring programs. Multi-state models may be run for multiple seasons to assess questions of change in the state of the population over time.

WCS monitoring and decision making in the face of uncertainty

Jim Nichols gave a thoughtful talk on monitoring for structured decision making in the face of uncertainty. He pointed out that we always have uncertainty about how to manage our Scapes. This uncertainty arises from a lack of understanding of the ecological processes that affect our Scapes (energy flow for example), environmental variation (Climate change), partial control, either due to lack of management authority, or lags in response to intervention, and

imperfect observation of the state of nature. Jim argues that well-designed monitoring systems should be able to identify uncertainty and allow us to: 1. Act conservatively when uncertainty is high; and 2. Use knowledge gained in an iterative fashion to reduce uncertainty.

In order to implement a monitoring program using a structured decision making process, we need to follow a series of clear steps, similar to our approaches for Scape conservation planning. First, we need a clear set of objectives regarding what we expect to accomplish. We need a clear set of management alternatives regarding what we are able to accomplish (we are in control) and what we are not able to accomplish (someone else is in control). We need quantitative models to predict the responses to the potential interventions for use in selecting the one that is 'best' with respect to our objectives. We need a method for determining the credibility of the competing models and an algorithm to identify the optimal decision. Finally, we need a monitoring system to gather data to determine the state of the system and the other relevant variables that affect our conservation targets.

Examples of occupancy-based monitoring in WCS

Tigers Forever: Obtaining baselines for large-scale occupancy by tigers and tiger prey

Spurred by the successes of wild tiger recovery in Nagarhole, India and Sikhote-Alin, Russia, the Tigers Forever program was established by WCS in 2006 with support from Panthera. The goal of Tigers Forever is to increase wild tiger numbers by 50% in ten years in key WCS landscapes across Asia. Following a "source-sink" strategy, each selected landscape consisted of source/potential source sites within a larger sink landscape. The objective was to identify key threats and target interventions at

source sites, and document rigorously the recovery of wild tiger numbers. The monitoring strategy involved assessing tiger and prey densities at the source populations annually and assessing the hypothesized “ripple” effect in the larger landscapes using occupancy surveys once every 4-5 years.

The large cell occupancy surveys that employed the Hines et al. (2010) model with spatial replication were used to assess tiger occupancy in the Malenad-Mysore Tiger Landscape in India. The naïve occupancy was found to be 47% lower than the estimated occupancy of 0.66 in this landscape (Karanth et al 2011). The estimated area occupied by tigers was ~14,100 km² of the 22,000 km² of potential tiger habitat. A similar tiger occupancy survey in Indonesia (Wibisono et al. 2011), but confronted with different models, revealed an estimated occupancy of tigers of 0.70 in comparison to a naïve estimate of 0.52.

Assessment of tiger prey in most source sites in southeast Asia (apart from Huai Kha Kaeng, Thailand, and sites in MMTL, India) posed a major challenge because traditional line-transect surveys were difficult to implement in practice. As an alternative, small cell occupancy surveys were designed to best meet the assumptions of Royle and Nichols (2003) abundance-induced heterogeneity models to estimate large ungulate densities using the field survey protocols of a trial survey in Bhadra (Gopaldaswamy et al 2012). Large sample sizes for estimating ungulate prey densities in Malaysia, Laos and Myanmar yielded precise estimates of abundance and occupancy using these models.

Conservation effectiveness of patrolling and law enforcement in BBSNP

Bukit Barisan Selatan National Park (BBSNP) in southern Sumatra, Indonesia was a stronghold for Sumatran tiger, elephant and rhinoceros during the 1990's. In 1995, the International Rhino Program, in collaboration with WWF-Indonesia and PHKA/BBSNP initiated joint patrols in BBSNP. This work was expanded to four patrol units in

1997 to improve coverage of patrols and continues today with 11 patrol units. In 1999, WCS assisted with funding from Save the Tiger Fund. The goal of the armed patrols was to reduce poaching of rhinos, tigers and elephants. In 2003, WCS initiated the Wildlife Crime Unit (WCU) with the objective of filling a capacity gap in being able to respond to and follow up on information regarding poaching of tigers, elephant, rhino and birds at the local and regional level. WCU is an investigation unit that helps the government by supplying the best information on illegal poaching and hunting and also deals with informant networks and promoting public awareness. WCU activities continue today. Between 1998 and 2012, WCS has conducted camera trap surveys throughout the park (1998-2006) and in the southern part of the park (2010-2011), arguably the best zone for wildlife, especially elephant, rhinos and tigers.

To determine whether the interventions (patrols and WCU activity) were effective in reducing poaching and unsustainable hunting, we looked at wildlife trends using the occupancy-based Wildlife Picture Index (WPI: O'Brien et al 2010). We found a park-wide decline in the WPI between 1998 and 2006 suggesting an erosion in biodiversity of medium and large sized mammals. Tigers, elephants and rhinos declined faster than the rate of forest loss and faster than the rate of loss for species that were hunted as crop-raiders or for subsistence. Species hunted for subsistence and persecuted for crop-raiding declined at rates similar to forest loss. Species that had no economic value showed no consistent pattern: declines and increases were balanced.

A more recent analysis looked at trends between 1998 and 2011 in the south of BBSNP, a well-protected area due to patrols, a private concession in the south and the presence of the WCS Way Canguk Research Station. In this area, there was a significant increase in the WPI between 1998 and 2011 for 34 mammal species. Tigers, elephants and rhinos, however, declined over this time period. Thus we conclude that the patrol efforts and WCU efforts may not have been sufficient to stem the loss of the 3 most endan-

gered large mammals in the park.

Effect of livestock management on diversity, distribution and abundance of large mammals in Laikipia County, Kenya.

Successful conservation of large terrestrial mammals on private lands requires that landowners be able to manage wildlife to derive benefits that offset costs of wildlife. In Laikipia County, Kenya (9,666 km²), all of the wildlife are on private lands, and land use and tolerance attitudes play a major role in the fate of wildlife. Kinnaird and O'Brien (2012) used camera traps to sample large mammal communities to determine the impact of 4 different livestock management systems (rhinoceros sanctuaries: no livestock; conservancies: intermediate stocking rates; fenced ranches and pastoralist group ranches: highest stocking rates) to examine whether management and stocking rates affect wildlife communities. We deployed cameras across 8 properties and used the photographs to estimate species richness in an occupancy framework, species' occupancies, and relative abundances. Species richness was highest on conservancies and sanctuaries, and lowest on fenced and group ranches. Occupancy estimates were, on average, twice as high on sanctuaries and conservancies as on fenced ranches, and five times higher than on group ranches. The relative abundances of most species were highest or second-highest on sanctuaries or conservancies. We identified exclusion fencing and overstocking of sheep and goats as major impediments to wildlife distribution and abundance. But we also found that wildlife thrived under a moderate stocking level that allowed space for wildlife. Kenya's policy toward wildlife is government ownership and no consumptive use. The lack of landowner rights to manage wildlife is therefore a key policy issue to tackle for wildlife conservation in Laikipia. Current policies prevent many direct management options that might improve the situation. Alternatives that are available include applying a landscape-level approach to land use planning that aims to

increase the area under conservation by providing incentives for conservation on overstocked and fenced properties.

Modeling Wolverine Occurrence Using Aerial Surveys of Tracks in Snow

Justina Ray and colleagues (Magoun et al. 2007) designed a novel approach to determine the extent of distribution and area of occupancy for wolverines (*Gulo gulo*) by using aerial surveys of tracks in snow and hierarchical occupancy modeling. They used a small, fixed-wing aircraft with pilot and one observer to search 575 of 588 survey units for wolverine tracks in approximately 60,000 km² of boreal forest in northwestern Ontario, Canada. They used sinuous flight paths to scan open areas in the forest in the 100-km² survey units. They detected tracks in 138 (24%) of the 575 sampled units. There was strong evidence of occurrence (probability of occurrence > 0.80) in 30% of the 588 survey units, weak evidence of occurrence (0.50–0.80) in 12%, weak evidence of absence (0.20–0.50) in 15%, and strong evidence of absence (< 0.20) in 43%. Wolverine range comprised 59% of the study area and area of occupancy was 33,400 km². With information on probability of occurrence and core areas of occupation for wolverines in the study area, resource managers and others can examine factors that influence wolverine distribution patterns and use this information to formulate best management practices that will maintain wolverines on the landscape in the face of increasing resource development. Comparing future survey results with those of the 2005 survey will provide an objective way to assess the efficacy of management practices.

Monitoring rare species in the Andean Patagonian Steppe Landscape

The unique wildlife of the Andean Patagonian Steppe Landscape includes a large (25 kg), flightless bird, the Darwin's rhea, with populations that have been decimated by hunting, egg collection, and habitat degradation due to livestock grazing and hydrocarbon development. To assess effectiveness of conservation interventions and also learn about the factors that affect rhea distribution, WCS is developing a monitoring system based on an occupancy analysis with a single-season model. Unlike the other large-bodied herbivore in the landscape, the conspicuous guanaco, which is easy to observe and count on transects from vehicles in the open steppe, rheas are in lower densities and harder to observe, although feces can be easily detected in walking transects. A calibration by the WCS team also indicates that fecal counts accurately reflect rhea densities. Based on a pilot survey of 20 transects (where we did repeated surveys throughout the year) we estimated a very high detection rate for rhea feces (83%). Because the portion of the landscape for which we wished to estimate occupancy is very large (>2.5 million hectares) and many areas are difficult to reach, repeated visits in a single season would be prohibitively expensive in both time and money. Therefore, we decided to conduct multiple, 1-km transect surveys on a single visit within each 2 X 2 km area. To design the survey we performed simulation analysis using program PRESENCE, and determined that 100 cells sampled with 2 transects and a sub-sample of 20 cells sampled with 4 transects would be sufficient to estimate occupancy with high precision (SE = 0.04). We overlaid a 2 X 2 km grid over the area, and randomly selected 100 cells which we sampled with transects with randomly selected starting points and orientations within the cell. Sampling was carried out in six 10-13-day campaigns by 3-5 people at total cost of \$ 18,000. We estimated detection probability for rheas to be 0.78 (SE = 0.04), and the proportion of sites occupied in the landscape was 0.63 (SE = 0.05). Using logistic regression analysis we evaluated the relationship

between rhea presence and elevation, vegetation type, livestock density, road density (including seismic lines for oil exploration that are useable by 4-wheel-drive vehicles), and distance to rural houses (people collect eggs and hunt birds for food). We found that the presence of rheas was negatively affected by elevation and livestock density. We are currently developing a protocol for repeating this survey every 3-5 years to determine changes in rhea occupancy through time and assess success of conservation measures.

Feces of other species of conservation concern can be detected in the walking transects, so we are evaluating the use of this survey technique and design to simultaneously monitor occupancy by the assemblage of large (2-8 kg) hystricomorph rodents. These rodents have more restrictive habitat requirements than rheas, so some adaptation of the sampling scheme will need to be employed. In the case of the guanaco, occupancy surveys may be cheaper to use than line transects used to estimate abundance and sufficient for our monitoring purposes in some portions of the landscape and for evaluating the effects of some interventions or threats. A WCS radiotelemetry study has determined that ca 60% of the large northern guanaco population is migratory, and our transect data over the last 20 years show that social structure is also fluid, with group size and conformation changing seasonally. Therefore, if we decide to employ occupancy analysis as a monitoring technique for guanacos, surveys will need to be conducted at least twice during the year to determine distribution in summer and winter ranges.

Occupancy methods to assess changes in species fish diversity on coral reefs

Surveys for coral reef fishes are frequently conducted using underwater visual census (UVC) methods that use one of two common survey types, line transects or point counts, to take replicate samples from a given reef at a specific point in time. These surveys are often repeated annually, making conventional UVC monitoring a natural fit for species rich-

ness occupancy analysis, whereby within-year UVC replicates are capture occasions used to estimate detectability of individual species and the total occupancy of species across the assemblage provides an estimate of species richness. This approach is only just being applied to coral reef ecosystems but a recent application by Cheal et al. (2012) used a hierarchical occupancy model to infer relative functional redundancy of reef fish communities on the Great Barrier Reef (GBR). To do this, Cheal et al. estimated the average probability of occupancy for herbivorous fishes on reefs within management sectors of the GBR. These reef and sector-level estimates represented the average probability of presence for herbivorous fishes, a direct analogue of functional redundancy that was readily comparable among reefs and sectors. Cheal et al. found that inner-shelf reefs, particularly near the cities of Cairns and Townsville, had the lowest functional redundancy across the GBR, potentially an intrinsic pattern of the GBR ecosystem. Importantly the occupancy approach used matched the current long-term sampling scheme for the Australian Institute of Marine Science and should be equally applicable to most reef monitoring programs.

Migratory Species

For migratory species, with life cycle activities often widely separated by geography, the problem at first seems intractable by scale. In the case of migratory birds and many marine mammals, however, most species studied show remarkable site fidelity at both their breeding grounds and (with more spatial variation) in their wintering grounds. Thus there seems to be the opportunity to examine occupancy estimates for populations of such migrants at the ends of their migrations. Hypothetically, such estimates could provide information as to where interventions are most needed. Assessment in passage migration (moving latitudinally or altitudinally) is more problematic, unless the migratory species has few and well known stopover areas (as is true for red knots and their dependence on Delaware Bay in their northward migration from southern Argentina to Arctic

Canada). This perspective is all hypothetical, as no migratory species has been assessed by these means to date.

Design considerations for occupancy studies

An occupancy metric is not a panacea for all of the WCS species of interest. As species become more common, often the cost of estimating abundance goes down and the precision increases. At the same time, occupancy analyses are not useful (or interesting) when true occupancy approaches 100%. Occupancy methods perform best when detection is high, occupancy is moderate, and we have many sampling sites. In general assessment of occupancy states and their dynamics seem to work best when the occupancy states are in the range of 0.3 to 0.8. Low detection probabilities can cause problems for occupancy because the estimates become increasingly less accurate as detection probabilities become very small. Thus it is important to maximize detection probabilities, which is a challenge when the study area is largely unknown, although some general rules may apply (e.g., in dry forest pellets work well for ungulates, in rainy places tracks are better). Increasing the number of sample units and sampling occasions generally increases the accuracy and precision of occupancy estimates. In general, for a rare species it is more efficient to survey more sampling units with fewer replications because we are interested to identify populations of the rare species, while for a common species fewer sampling units should be surveyed with more replicates. In either case, we should always measure meaningful covariates to better understand what drives species occupancy.

Few wildlife species of interest to WCS are likely to be widely distributed on the landscape. Many species will have clumped distribution; some will be widespread but patchy for many reasons. Fortunately, there are a couple of modeling tools to help with designing and evaluating the

potential sensitivity of occupancy monitoring programs. Occupancy software PRESENCE includes a simulation routine that allows you to determine the degree of bias and precision likely to occur for a given level of sampling units and sampling effort when estimates or guesses of true occupancy and detection probability are provided. This allows one to quickly evaluate a number of scenarios of sampling effort. For more complex sampling designs, GENPRES software may be used to assist in developing sampling designs. We recommend using GENPRES when designing occupancy studies.

Choice of sites (sampling units, grid cells) is important both for interpretation of occupancy statistics and for cost of sampling design. At the spatial scale of a site, the intent is to infer if the target species is present or absent. By combining data from a number of sites, we calculate the probability of occupancy as a value between 0 and 1. The size of a site is dependent on many factors, including the management objectives for the Scape (and potentially the scale of the threat being mitigated). As discussed earlier, measures of occupancy can be scale dependent, especially for arbitrarily determined sites within contiguous habitat (i.e. sampling grids in forests). A larger site has a higher probability of occupancy than a smaller site. MacKenzie et al. (2006) suggest that a site should be large enough to have a reasonable probability of the species being present, but small enough that the measure of occupancy is meaningful and site can be surveyed in a cost-effective manner. In order to draw inferences about a landscape, sites should be selected using some probabilistic sampling scheme (simple random, stratified random, systematic with random starting site, etc.) unless the entire landscape is sampled. The closure assumption means that either the occupancy state does not change during the sampling period (true occupancy-site is permanently occupied) or that the changes in occupancy within a season are random (use-site is sometimes occupied during the season). This interpretation suggests that the closure assumption may be more easily met for large sites than for small sites.

When designing occupancy over time we need to consider the biological characteristics of the species of interest, as well as how our investments are likely to influence occupancy and over what time frame. This of course becomes more complicated with multiple target species because species-specific landscapes may vary.

Sampling designs are very flexible in occupancy analysis. Sample replication may be spatial or temporal or by means of independent observers. There are three basic sampling strategies: Standard design in which all sites are sampled K times; double sampling design in which a subset of sites is sampled K times and the remaining sites are sampled once; and a removal design in which all sites are sampled up to K times but sampling at a site stops when a target species is detected.

We discussed the problem of misidentification of species due to a variety of causes. Target species may be misidentified in a photograph, or a track or other indirect sign may be wrongly attributed to the target species. Interviews may also lead to misidentification if the interviewee is unsure of when and where he saw a species, or confuses the target species with another species. We discussed a number of methods to control for misidentification of target species, including additional questions to verify truthfulness and accuracy and independent sampling in a selection of sites covered in the context of the sign or interviews surveys. We also discussed the use of mixed methods to increase sample sizes. Mixed methods can be incorporated into a sampling design to yield unbiased estimates of occupancy. For example, we might define sampling units appropriate for jaguar and conduct structured interviews with communities to determine within which of these sampling units community members have observed jaguars within the last three months. We could then visit a subset of the same sampling units identified for the community interview (and possibly visit additional units) and survey for jaguars sign along rivers. As even our field teams might not correctly distinguish jaguar from puma sign, we could also potentially set up a

camera trap or DNA-based study to obtain further information to deal with potential misidentification of jaguar. Under either of these sampling schemes, the investigator would visit areas where jaguars were designated as being present as well as areas where jaguar were designated as being absent.

We advocate the balanced use of covariates to model heterogeneity in detection probability and occupancy, and to test hypotheses of interest to monitoring. Covariates may be measured in the field during sampling or may be assigned using global knowledge of the landscape (e.g. GIS information). If covariates are measured in the field, then the inferences using those covariates is strictly valid only for the area sampled because we cannot use the covariates to make predictions about areas not visited. If covariates are assigned based on some global knowledge (land-use, habitat, climate, deforestation), then we can make predictions about expected occupancy of locations within the surveyed area that were not sampled. It is especially important to include covariates that might affect detection probability because the models assume that all heterogeneity in detection is modeled and violation of this assumption can lead to biased inferences. Note that a modeling approach that involves finite mixture models provides an alternative for dealing with potential heterogeneity from an unknown source, i.e. when we do not have the appropriate covariate information. Ignoring covariates that might affect occupancy (e.g. habitat types) will reduce the precision of the estimates but should not induce bias. When evaluating management interventions or conservation effectiveness, we can use stratification of the landscape into treatments and calculate occupancy separately for each treatment then compare results. A more efficient alternative is to treat the interventions as covariates within a single analysis. The nature of the objectives and the data may lend itself to one strategy over the other. For example, if law enforcement is one of our interventions then we could stratify by law enforcement effort, e.g. none, low, medium, high, or alternatively assign a continuous variable value to each of the sampling units.

Way Forward

We found it quite productive to discuss the design and analytical issues for Madidi, Bolivia as a way to move the discussion from the abstract to the realm of possibility. Rob Wallace prepared a companion planning document (Wallace 2012) for implementing an occupancy-based monitoring program in the Madidi Landscape. In addition, we offered several ideas for moving forward with the investigation of the use of occupancy as a conservation effectiveness metric for wildlife:

- ◆ Investigate retro-fitting occupancy analyses to current sampling designs and datasets. Samantha and Boo Maisels are looking at the Central Africa elephant and ape data set to determine if it is appropriate to integrate the traditionally analyzed transect data with both the travel and guided reconnaissance data (usually not included in analyses due to known sampling biases). This would provide additional spatial coverage and increase sample sizes for species for which estimates can generally not be obtained from the transect data (e.g. forest buffalo, bongo, red river hog, leopard). Margaret Kinnaird, Samantha and Tim have done the same for aerial surveys in Laikipia. Aaron suggested further application of the species richness estimation methods to UVC data to WCS sites in Kenya and elsewhere. A key question that we need to answer is whether designs that maximize detection probability for a single (or few) target species can be used to assess occupancy of non-target species. This is a fundamental question for the utility of large cat monitoring programs for non-target species monitoring, i.e. placing camera traps to detect tigers you may be less likely to detect tiger prey species that avoids tiger trails. Ullas and Arjun are of the opinion that it is not appropriate for the India sites. However, we believe that the results may be biased, but if the bias is consistent, or there are covariates that can be used to address the sampling bias the data may still be useful. We need to explore this option further.

- ◆ We need to bring together programs that are interested in increasing the power of their monitoring programs to explore ways to get more from their data – turning by-catch into filets. A first step is to develop a standard database for WCS monitoring data. We recommend that WCS, as part of the TEAM data federation exercise, integrate camera trap and other monitoring data from WCS TEAM sites and two additional Latin America sites into a WCS camera trap database. Partial funding for this may be provided by TEAM.
- ◆ We recommend that WCS engage in a series of workshops at NCEAS under the NCEAS/TNC/WCS NATURELAB program to investigate structured decision making, monitoring, and assessing conservation effectiveness in WCS Scapes.
- ◆ We recommend the development of a test landscape in each region (NA, LA, Asia, Africa, Marine) where we engage in occupancy analyses (either retro-fit or design of new surveys) to establish the utility of the efforts.

Literature cited

- Cheal, AJ, M Emslie, A MacNeil, I Miller, and H Sweatmann. 2012. Spatial variation in the functional characteristics of herbivorous fish communities and the resilience of coral reefs. *Ecological Applications* In press.
- Gopalaswamy, AM, KU Karanth, NS Kumar, and D.W. McDonald. 2012. Estimating tropical forest ungulate densities from sign surveys using abundance models of occupancy. *Animal Conservation* DOI: 10.1111/j.1469-1795.2012.00565.x
- Hines, J.E., Nichols, J.D., Royle, J.A., MacKenzie, D.I., Gopalaswamy, A.M., Kumar, N.S., Karanth, K.U. 2010. Tigers on trails: occupancy modeling for cluster sampling. *Ecological Applications* 20:1456-1466.
- Karant, KU, AM Gopalaswamy, NS Kumar, S. Vaidyanathan, JD Nichols, JD, and DI MacKenzie. 2011.

- Monitoring carnivore populations at the landscape scale: occupancy modeling of tigers from sign surveys. *Journal of Applied Ecology* 48:1048-1056.
- Kinnaird, MF and TG O'Brien. 2012. The effects of private land-use, livestock management, and human tolerance on diversity, distribution and abundance of large African mammals. *Conservation Biology* In Press.
- MacKenzie, D.I., Nichols, J.D., Royle, J.A., Pollock, K.P., Bailey, L.L. and Hines, J.E. 2006. *Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species Occurrence*. Academic Press, New York.
- Magoun, AJ, JC Ray, DS Johnson, P Valkenburg, FN Dawson, and J Bowman, J. 2007. Modeling wolverine occurrence using aerial surveys of tracks in snow. *Journal of Wildlife Management* 71:2221-2229.
- O'Brien, TG, Baillie, JEM, Krueger, L, Cuke, M. 2010. The Wildlife Picture Index: monitoring top trophic levels. *Animal Conservation* 13:335-343.
- Royle, JA and JD Nichols. 2003. Estimating abundance from repeated presence-absence data or point counts. *Ecology* 84:777-790.
- Strindberg, S and T O'Brien. 2012. A decision tree for monitoring wildlife to assess the effectiveness of conservation interventions. WCS Working Papers No. 41.
- Wallace R. 2012 Landscape scale for species and threats: Jaguars in the greater Madidi-Tambopata landscape. Unpublished report to the Wildlife Conservation Society. 16 pp.
- Wibisono, HT, M Linkie, G, Guillera-Arroita, JA Smith, Sunarto, et al. 2011. Population status of a cryptic top predator: An island-wide assessment of tigers in Sumatran rainforests. *PLoS One* 6(11): e25931. doi:10.1371/journal.pone.0025931

Appendix 1: Brief Overview of Occupancy Methods

Compiled by Tim O'Brien and Samantha Strindberg March 2012

Occupancy methods estimate the proportion of a habitat or number of patches occupied when detection is incomplete (Mackenzie et al., 2002, Mackenzie et al., 2003, Mackenzie et al., 2006). When sampling units are scaled to the home range size of target species, the interpretation is proportion of area occupied. When sampling units are smaller than home ranges, the interpretation is intensity of use. The analysis recognizes three states: occupied and species detected, occupied and species not detected, and not occupied. It provides estimates of the probability that a sampling unit is occupied and the probability that at least one individual animal (or sign, if sign surveys are used) is detected. It requires replicated observations on each sampling unit and it allows for covariates that might affect occupancy or detection to be incorporated into the analysis. The basic method assumes demographic and spatial closure during a sampling period (referred to as a season) such that the occupancy status does not change and that sampling units states are independent. Additional assumptions include no errors in identifying species and that observations are independent. There are analysis options that relax most of these assumptions should this be needed. The fundamental reference for occupancy analysis is the MacKenzie et al. 2006 book *Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species Occurrence*. The University of Vermont Cooperative Fish and Wildlife Research Unit Spreadsheet Project has an excellent website for self-training in occupancy methods (<http://www.uvm.edu/rsenr/vtcfwru/spreadsheets/occupancy.htm>). The Patuxent Wildlife Research Center Software page (<http://www.mbr-pwrc.usgs.gov/software/doc/presence/presence.html>) also provides a useful user manual that outlines the different analyses available in PRESENCE. Key references for de-

signing occupancy studies include MacKenzie and Royle (2005) and Bailey et al. (2007). The methods are continually evolving and some of the analysis options currently available include:

Single Season models - this is the basic occupancy model, which allows for estimates of the proportion of the study area occupied and the detection probability. Occupancy and detection parameters may be constant across the sampling area or be estimated as a function of site and survey-specific covariates. Single season models based on mixture models to deal with unobservable heterogeneity can also be used. Substitution of species for samples permits estimation of species richness in a study area and exploration of the covariates that affect species richness. When covariates are used to estimate occupancy, predictive maps can be developed to include occupancy estimates for sites in which no detections were made and for sites that were not sampled (but fall within the study area and have covariate data) can be generated. Single season models can also be used for meta-population modeling.

Mackenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Royle, J.A. and Langtimm, C.A. 2002. Estimating site occupancy when detection probabilities are less than one. *Ecology* 83:2248-2255.

Multi-season models - are an extension of single season models and can be used for inferences about occupancy and meta-population dynamics. Sites can change between being occupied and unoccupied over time allowing for estimates of rates of local extinction and local colonization. Single and multi-season models are ideal for large scale surveys of single species, single populations, meta-populations and communities.

Mackenzie, D.I., Nichols, J.D., Hines, J.E., Knutson, M.G. and Franklin, A.B. 2003. Estimating site occupancy, colonization and local extinction probabilities when a species is not detected with certainty. *Ecology* 84:2200–2207.

Single season with false positive detections models - are useful when there is a good chance that sign, or aural cues or visual identifications are incorrect. For example, this happens with scat surveys, bird counts using calls, and blurry camera trap photos of duikers and viverrids.

Royle, J.A., Link, W.A. 2006. Generalized site occupancy models allowing for false positive and false negative errors. *Ecology* 87:835-841.

Miller, DA, Nichols, JD, McClintock, BT, Grant, EHC, Bailey, LL, Weir, LA. 2011. Improving occupancy estimation when two types of observational error occur: non-detection and species misidentification. *ECOLOGY* 92:1422-1428.

Multi-method models - allow estimation of occupancy when more than one method for detection is employed across sites, providing detection probabilities for each method used. This is useful for hybrid surveys or surveys using multiple cues (e.g., species richness estimation for bird communities using visual and aural cues).

Nichols, J.D., Bailey, L.L., O'Connell, A.F., Talancy, N.W., Grant, E.H.C., Gilbert, A.T., Annand, E.M., Husband, T.P., Hines, J.E. 2008. Multi-scale occupancy estimation and modeling using multiple detection methods. *Journal of Applied Ecology* 45:1321-1329.

Single season multi-state models - are used when we are interested in not only whether a site is occupied, but whether there are different states that the occupied site might attain (for example occupied with breeders, occupied with non-breeders, or occupied but with different classes of abundance). This can be very important for breeding bird surveys, and for meta-population analyses.

Nichols, J.D., Hines, J.E., MacKenzie, D.I., Seamans, M.E., Gutierrez, R.J. 2007. Occupancy estimation and modeling with multiple states and state uncertainty. *Ecology* 88:1395-1400.

MacKenzie, D.I., J.D. Nichols, M.E. Seamans, and R.J. Gutierrez. 2009. Modeling species occurrence dynamics with multiple states and imperfect detection. *Ecology* 90:823–835.

Multi-season multi-state - extend multi-state models to multiple seasons. For example, occupancy models can be used to estimate if a species is absent, rare, or abundant (i.e., 3 population states) or alternatively, if different life history stages are present such as: absent, juvenile, adults. When combined, models can be used to estimate meta-demographic rates such as colonization, extinction, reproduction and recruitment.

MacKenzie, D.I., J.D. Nichols, M.E. Seamans, and R.J. Gutierrez. 2009. Modeling species occurrence dynamics with multiple states and imperfect detection. *Ecology* 90:823–835.

MacKenzie, D.I., L.L. Bailey, J.E. Hines, and J.D. Nichols. 2011. An integrated model of habitat and species occurrence dynamics. *Methods in Ecology and Evolution* 2:612-622.

Multi-season integrated habitat occupancy - can be used to examine how habitat suitability and factors that affect habitat suitability can influence the distribution and relative abundance of organisms over time. (This has been used with elephants at water holes in Zimbabwe and would be a good candidate for comparing dung surveys over time).

MacKenzie, D.I., J.D. Nichols, M.E. Seamans, and R.J. Gutierrez. 2009. Modeling species occurrence dynamics with multiple states and imperfect detection. *Ecology* 90:823–835.

Martin, J, Chammille-Jammes, S, Nichols, J.D., Fritz, H., Hines, J.E., Fonnesebeck, C.J., MacKenzie, D.I., Bailey, L.L. 2010. Simultaneous modeling of habitat suitability, occupancy, and relative abundance: African elephants in Zimbabwe. *Ecological Applications* 20:1173-1182.

2 species co-occurrence models - are used when the goal is to determine if 2 species occupy a site, whether occupancy is affected by co-occurrence, and to assess whether they affect each other's detection probabilities. We can also test if the detection probability of one species changes in the presence of the other.

MacKenzie D.I., Bailey, L.L., Nichols, J.D. 2004. Investigating species co-occurrence patterns when species are detected imperfectly. *Journal of Animal Ecology* 73:546-555.

Repeated count models - are occupancy based abundance estimation procedures used to estimate the number of individuals at a point when individuals cannot be identified or marked. Rather than using species presence-absence (detection-nondetection) data, these models are based on counts of individuals obtained at replicate visits. These can be very useful when the sample sites are discrete (i.e. ponds or woodlots) and where the area of sampling can be defined (i.e. fixed distance point count).

Royle, J.A. 2004. N-Mixture Models for Estimating Population Size from Spatially Replicated Counts. *Biometrics* 60, 108-115.

Single season heterogeneity models - similar to above but is based on the idea that heterogeneity in abundance generates heterogeneity in detection probability. Uses presence/absence data to estimate point abundance, and occupancy as a function of point abundance.

Royle, J.A., Nichols, J.D. 2003. Estimating abundance from repeated presence-absence data or point counts. *Ecology* 84:777-790.

Single season staggered entry - are used when we cannot assume that the population is closed within a season. Instead, individuals of the species are assumed to arrive and depart from the study area. We estimate $P(\text{arrival})$, $P(\text{departure})$ and $P(\text{detection})$ to develop occupancy esti-

mates.

Kendall, W.L., Hines, J.E., Nichols, J.D., Grant, E.H. (in prep.) Relaxing the closure assumption in single-season occupancy models: staggered arrival and departure times.

Single season spatial/temporal autocorrelation - occupancy analysis and many other estimation methods assume that detections are independent in space and time. When conducting sign surveys along trails, or using camera traps we may encounter situations where observations are correlated in space or time. These models incorporate autocorrelation in detections to produce unbiased occupancy and detection estimates.

Hines, J.E., Nichols, J.D., Royle, J.A., MacKenzie, D.I., Gopalaswamy, A.M., Kumar, N.S., Karanth, K.U. 2010. Tigers on trails: occupancy modeling for cluster sampling. *Ecological Applications* 20:1456-1466.

Bled, F., Royle, J.A. and Cam, E. 2011. Hierarchical modeling of an invasive spread: The Eurasian collared-dove *Streptopelia decaocto* in the United States. *Ecological Applications* 21:290-302.

Occupancy analysis can be carried out in a maximum likelihood (frequentist) framework or a Bayesian framework. The PRESENCE software facilitates frequentist analysis of occupancy data and can be used for single species studies, community level studies and estimation of species richness. It is available as a free download from Patuxent Software Archive (www.mbr.pwrc.usgs.gov/software.html). Occupancy analysis can also be carried out in R using the Unmarked package (<http://github.com/rbchan/unmarked>).

References

Bailey, L.L., Hines, J.E., Nichols, J.D., MacKenzie, D.I. 2007. Sampling design trade-offs in occupancy studies with imperfect detection: Examples and software.

- Ecological Applications 17:281-290.
- Mackenzie, D.I., Nichols, J.D., Lachman, G.B., Droege, S., Royle, J.A., Langtimm, C.A. 2002. Estimating site occupancy when detection probabilities are less than one. *Ecology* 83:2248-2255.
- Mackenzie, D.I., Nichols, J.D., Hines, J.E., Knutson, M.G., Franklin, A.B. 2003. Estimating site occupancy, colonization and local extinction probabilities when a species is not detected with certainty. *Ecology* 84:2200-2207.
- Mackenzie, D.I., Royle, J.A. 2005. Designing occupancy studies: general advice and allocating survey effort. *Journal of Applied Ecology* 42:1105-1114.
- MacKenzie, D.I., Nichols, J.D., Royle, J.A., Pollock, K.P., Bailey, L.L. and Hines, J.E. 2006. *Occupancy Estimation and Modeling: Inferring Patterns and Dynamics of Species Occurrence*. Academic Press, New York.

Appendix 2: A Short Bibliography of Occupancy Methods Related Publications

Compiled by Tim O'Brien

As part of the WCS Conservation Support work on measures of conservation effectiveness, I compiled a partial list of publications that use occupancy methods. I surveyed *Animal Conservation*, *Auk*, *Biological Conservation*, *Bird Conservation International*, *Conservation Biology*, *Ecology*, *Ecological Applications*, *Herpetologica*, *Ibis*, *Journal of Applied Ecology*, *Journal of Herpetology*, *Journal of Wildlife Management*, and *Wildlife Society Bulletin* for articles that apply presence/absence and detection/nondetection methods, published between 2002 (first paper by MacKenzie et al. on occupancy analysis) and 2012. Most articles use occupancy-based methods, but allogistic regressions, presence only methods, and incidence functions. I focused on wildlife applications, but there is a wide array of tax-specific papers included.

Methods and general

- Aing, C, Halls, S, Oken, K, Dobrow, R, Fieberg, J. 2011. A Bayesian hierarchical occupancy model for track surveys conducted in a series of linear, spatially correlated, sites. *JOURNAL OF APPLIED ECOLOGY* 48:1508-1517.
- Bled, F, Royle, JA, Cam, E. 2011. Assessing hypotheses about nesting site occupancy dynamics. *ECOLOGY* 92:938-951.
- Christy, MT, Adams, AAY, Rodda, GH, Savidge, JA, Tyrrell, CL. 2010. Modelling detection probabilities to evaluate management and control tools for an invasive species. *JOURNAL OF APPLIED ECOLOGY* 47:106-113.
- Conroy, MJ, Runge, JP, Barker, RJ, Schofield, MR,

- Fonnesbeck, CJ. 2008. Efficient estimation of abundance for patchily distributed populations via two-phase, adaptive sampling. *ECOLOGY* 89:3362-3370.
- Delaney, DG, Leung, B. 2010. An empirical probability model of detecting species at low densities. *ECOLOGICAL APPLICATIONS* 20:1162-1172.
- Dorazio, RM. 2007. On the choice of statistical models for estimating occurrence and extinction from animal surveys. *ECOLOGY* 88:2773-2782.
- Dorazio, RM, Royle, JA, Soderstrom, B, Glimskar, A. 2006. Estimating species richness and accumulation by modeling species occurrence and detectability. *ECOLOGY* 87:842-854.
- Dorazio, RM, Kery, M, Royle, JA, Plattner, M. 2010. Models for inference in dynamic metacommunity systems. *ECOLOGY* 91:2466-2475.
- Drielsma, M, Ferrier, S. 2009. Rapid evaluation of metapopulation persistence in highly variegated landscapes. *BIOLOGICAL CONSERVATION* 142:529-540.
- Ferraz, G, Sberze, M, Cohn-Haft, M. 2010. Using occupancy estimates to fine-tune conservation concerns. *ANIMAL CONSERVATION* 13:19-20.
- Field, SA, Tyre, AJ, Possingham, HP. 2005. Optimizing allocation of monitoring effort under economic and observational constraints. *JOURNAL OF WILDLIFE MANAGEMENT* 69:473-482.
- Fitzpatrick, MC, Preisser, EL, Ellison, AM, Elkinton, JS. 2009. Observer bias and the detection of low-density populations. *ECOLOGICAL APPLICATIONS* 19:1673-1679.
- Griffin, SC, Taper, ML, Hoffman, R. 2010. Ranking Mahalanobis Distance Models for Predictions of Occupancy From Presence-Only Data. *JOURNAL OF WILDLIFE MANAGEMENT* 74:1112-1121.
- Grouios, CP, Manne, LL. 2009. Utility of Measuring Abundance versus Consistent Occupancy in Predicting Biodiversity Persistence. *CONSERVATION BIOLOGY* 23:1260-1269.
- Gu, WD, Swihart, RK. 2004. Absent or undetected? Effects

- of non-detection of species occurrence on wildlife-habitat models. *BIOLOGICAL CONSERVATION* 116:195-203.
- Halstead, BJ, Wylie, GD, Coates, PS, Casazza, ML. 2011. Bayesian Adaptive Survey Protocols for Resource Management. *JOURNAL OF WILDLIFE MANAGEMENT* 75:450-457.
- Hodgson, JA, Moilanen, A, Thomas, CD. 2009. Metapopulation responses to patch connectivity and quality are masked by successional habitat dynamics. *ECOLOGY* 90:1608-1619.
- Hui, C, McGeoch, MA, Reyers, B, le Roux, PC, Greve, M, Chown, SL. 2009. Extrapolating population size from the occupancy-abundance relationship and the scaling pattern of occupancy. *ECOLOGICAL APPLICATIONS* 19:2038-2048.
- Johnson, DM. 2005. Metapopulation models: An empirical test of model assumptions and evaluation methods. *ECOLOGY* 86:3088-3098.
- Jones, JE, Kroll, AJ, Giovanini, J, Duke, SD, Betts, MG. 2011. Estimating thresholds in occupancy when species detection is imperfect. *ECOLOGY* 92:2299-2309.
- Jones, JPG. 2011. Monitoring species abundance and distribution at the landscape scale. *JOURNAL OF APPLIED ECOLOGY* 48:9-13.
- Joseph, LN, Possingham, HP. 2008. Grid-based monitoring methods for detecting population declines: Sensitivity to spatial scale and consequences of scale correction. *BIOLOGICAL CONSERVATION* 141:1868-1875.
- Joseph, LN, Elkin, C, Martin, TG, Possingham, HP. 2009. Modeling abundance using N-mixture models: the importance of considering ecological mechanisms. *ECOLOGICAL APPLICATIONS* 19:631-642.
- Kery, M, Royle, JA, Plattner, M, Dorazio, RM. 2009. Species richness and occupancy estimation in communities subject to temporary emigration. *ECOLOGY* 90:1279-1290.
- Kery, M, Royle, JA, Schmid, H, Schaub, M, Volet, B, Hafliger, G, Zbinden, N. 2010. Site-Occupancy Dis-

- tribution Modeling to Correct Population-Trend Estimates Derived from Opportunistic Observations. *CONSERVATION BIOLOGY* 24:1388-1397.
- McClintock, BT, Bailey, LL, Pollock, KH, Simons, TR. 2010. Unmodeled observation error induces bias when inferring patterns and dynamics of species occurrence via aural detections. *ECOLOGY* 91:2446-2454.
- MacKenzie, DI. 2005. What are the issues with presence-absence data for wildlife managers? *JOURNAL OF WILDLIFE MANAGEMENT* 69:849-960.
- MacKenzie, DI, Nichols, JD, Lachman, GB, Droege, S, Royle, JA, Langtimm, CA. 2002. Estimating site occupancy rates when detection probabilities are less than one. *ECOLOGY* 83:2248-2255.
- MacKenzie, DI, Nichols, JD, Hines, JE, Knutson, MG, Franklin, AB. 2003. Estimating site occupancy, colonization, and local extinction when a species is detected imperfectly. *ECOLOGY* 84:2200-2207.
- MacKenzie, DI, Nichols, JD, Sutton, N, Kawanishi, K, Bailey, LL. 2005. Improving inferences in population studies of rare species that are detected imperfectly. *ECOLOGY* 86:1101-1113.
- MacKenzie, DI. 2006. Modeling the probability of resource use: The effect of, and dealing with, detecting a species imperfectly. *JOURNAL OF WILDLIFE MANAGEMENT* 70:367-374.
- Mackenzie, DI, Royle, JA. 2005. Designing occupancy studies: general advice and allocating survey effort. *JOURNAL OF APPLIED ECOLOGY* 42:1105-1114.
- Mackenzie, DI, Nichols, JD, Seamans, ME, Gutierrez, RJ. 2009. Modeling species occurrence dynamics with multiple states and imperfect detection. *ECOLOGY* 90:823-835.
- Manley, PN, Schlesinger, MD, Roth, JK, Van Horne, B. 2005. A field-based evaluation of a presence-absence protocol for monitoring ecoregional-scale biodiversity. *JOURNAL OF WILDLIFE MANAGEMENT* 69:950-966.
- Martin, J, Nichols, JD, McIntyre, CL, Ferraz, G, Hines, JE. 2009. Perturbation analysis for patch occupancy dynamics. *ECOLOGY* 90:10-16.

- Martin, J, Runge, MC, Nichols, JD, Lubow, BC, Kendall, WL. 2009. Structured decision making as a conceptual framework to identify thresholds for conservation and management. *ECOLOGICAL APPLICATIONS* 19:1079-1090.
- Mattfeldt, SD, Bailey, LL, Grant, EHC. 2009. Monitoring multiple species: Estimating state variables and exploring the efficacy of a monitoring program. *BIOLOGICAL CONSERVATION* 142:720-737.
- Miller, DA, Nichols, JD, McClintock, BT, Grant, EHC, Bailey, LL, Weir, LA. 2011. Improving occupancy estimation when two types of observational error occur: non-detection and species misidentification. *ECOLOGY* 92:1422-1428.
- Mordecai, RS, Mattsson, BJ, Tzilkowski, CJ, Cooper, RJ. 2011. Addressing challenges when studying mobile or episodic species: hierarchical Bayes estimation of occupancy and use. *JOURNAL OF APPLIED ECOLOGY* 48:56-66.
- Moreno, M, Lele, SR. 2010. Improved estimation of site occupancy using penalized likelihood. *ECOLOGY* 91:341-346.
- Nichols, JD, Hines, JE, MacKenzie, DI, Seamans, ME, Gutierrez, RJ. 2007. Occupancy estimation and modeling with multiple states and state uncertainty. *ECOLOGY* 88:1395-1400.
- Nichols, JD, Bailey, LL, O'Connell, AF, Talancy, NW, Grant, EHC, Gilbert, AT, Annand, EM, Husband, TP, Hines, JE. 2008. Multi-scale occupancy estimation and modelling using multiple detection methods. *JOURNAL OF APPLIED ECOLOGY* 45:1321-1329.
- O'Hara, RB, Arjas, E, Toivonen, H, Hanski, I. 2002. Bayesian analysis of metapopulation data. *ECOLOGY* 83:2408-2415.
- Otto, CRV, Roloff, GJ. 2011. Using Multiple Methods to Assess Detection Probabilities of Forest-Floor Wildlife. *JOURNAL OF WILDLIFE MANAGEMENT* 75:423-431.
- Pavlacky, DC, Blakesley, JA, White, GC, Hanni, DJ, Lukacs, PM. 2012. Hierarchical multi-scale occupancy

- estimation for monitoring wildlife populations. *JOURNAL OF WILDLIFE MANAGEMENT* 76:154-162.
- Patten, MA, Smith-Patten, BD. 2011. Predictors of Occupancy Trend Across Spatial Scale. *CONSERVATION BIOLOGY* 25:1203-1211.
- Pellet, J, Schmidt, BR. 2005. Monitoring distributions using call surveys: estimating site occupancy, detection probabilities and inferring absence. *BIOLOGICAL CONSERVATION* 123:27-35.
- Rhodes, JR, Tyre, AJ, Jonzen, N, McAlpine, CA, Possingham, HP. 2006. Optimizing presence-absence surveys for detecting population trends. *JOURNAL OF WILDLIFE MANAGEMENT* 70:8-18.
- Rhodes, JR, Callaghan, JG, McAlpine, CA, De Jong, C, Bowen, ME, Mitchell, DL, Lunney, D, Possingham, HP. 2008. Regional variation in habitat-occupancy thresholds: a warning for conservation planning. *JOURNAL OF APPLIED ECOLOGY* 45:549-557.
- Royle, JA, Nichols, JD. 2003. Estimating abundance from repeated presence-absence data or point counts. *ECOLOGY* 84:777-790.
- Royle, JA, Link, WA. 2005. A general class of multinomial mixture models for anuran calling survey data. *ECOLOGY* 86:2505-2512.
- Royle, JA, Link, WA. 2006. Generalized site occupancy models allowing for false positive and false negative errors. *ECOLOGY* 87:835-841.
- Royle, JA, Kery, M. 2007. A Bayesian state-space formulation of dynamic occupancy models. *ECOLOGY* 88:1813-1823.
- Stanley, TR, Royle, JA. 2005. Estimating site occupancy and abundance using indirect detection indices. *JOURNAL OF WILDLIFE MANAGEMENT* 69:874-883.
- Vojta, CD. 2005. Old dog, new tricks: Innovations with presence-absence information. *JOURNAL OF WILDLIFE MANAGEMENT* 69:845-848.
- Wenger, SJ, Freeman, MC. 2008. Estimating species occurrence, abundance, and detection probability using zero-inflated distributions. *ECOLOGY* 89:2953-2959.
- Wintle, BA, McCarthy, MA, Parris, KM, Burgman, MA.

2004. Precision and bias of methods for estimating point survey detection probabilities. *ECOLOGICAL APPLICATIONS* 14:703-712.
- Zillio, T, He, FL. 2010. Modeling spatial aggregation of finite populations. *ECOLOGY* 91:3698-3706.
- Zipkin, EF, Dewan, A, Royle, JA. 2009. Impacts of forest fragmentation on species richness: a hierarchical approach to community modeling. *JOURNAL OF APPLIED ECOLOGY* 46:815-822.
- Zipkin, EF, Royle, JA, Dawson, DK, Bates, S. 2010. Multi-species occurrence models to evaluate the effects of conservation and management actions. *BIOLOGICAL CONSERVATION* 143:479-484.

AMPHIBIANS

- Adams, MJ, Chelgren, ND, Reinitz, D, Cole, RA, Rachowicz, LJ, Galvan, S, McCreary, B, Pearl, CA, Bailey, LL, Bettaso, J, Bull, EL, Leu, M. 2010. Using occupancy models to understand the distribution of an amphibian pathogen, *Batrachochytrium dendrobatidis* *ECOLOGICAL APPLICATIONS* 20:289-302.
- Adams, MJ, Pearl, CA, Galvan, S, McCreary, B. 2011. Non-Native Species Impacts on Pond Occupancy by an Anuran. *JOURNAL OF WILDLIFE MANAGEMENT* 75:30-35.
- Bailey, LL, Simons, TR, Pollock, KH. 2004. Estimating site occupancy and species detection probability parameters for terrestrial salamanders. *ECOLOGICAL APPLICATIONS* 14:692-702.
- Bradford, DF, Neale, AC, Nash, MS, Sada, DW, Jaeger, JR. 2003. Habitat patch occupancy by toads (*Bufo punctatus*) in a naturally fragmented desert landscape. *ECOLOGY* 84:1012-1023.
- Brander, SM, Royle, JA, Eames, M. 2007. Evaluation of the status of anurans on a refuge in suburban Maryland. *JOURNAL OF HERPETOLOGY* 41:52-60.
- Chelgren, ND, Adams, MJ, Bailey, LL, Bury, RB. 2011. Using multilevel spatial models to understand salamander site occupancy patterns after wildfire. *ECOLOGY* 92:408-421.

- de Solla, SR, Shirose, LJ, Fernie, KJ, Barrett, GC, Brousseau, CS, Bishop, CA. 2005. Effect of sampling effort and species detectability on volunteer based anuran monitoring programs. *BIOLOGICAL CONSERVATION* 121:585-594.
- Dodd, CK, Dorazio, RM. 2004. Using counts to simultaneously estimate abundance and detection probabilities in a salamander community. *HERPETOLOGICA* 60:468-478.
- Hartel, T, Schweiger, O, Ollerer, K, Cogalniceanu, D, Arntzen, JW. 2010. Amphibian distribution in a traditionally managed rural landscape of Eastern Europe: Probing the effect of landscape composition. *BIOLOGICAL CONSERVATION* 143:1118-1124.
- Hayes, MP, Quinn, T, Dugger, DJ, Hicks, TL, Melchior, MA, Runde, DE. 2006. Dispersion of Coastal Tailed Frog (*Ascaphus truei*): An hypothesis relating occurrence of frogs in non-fish-bearing headwater basins to their seasonal movements. *JOURNAL OF HERPETOLOGY* 40:531-543.
- Jackson, JT, Weckerly, FW, Swannack, TM, Forstner, MRJ. 2006. Inferring absence of Houston toads given imperfect detection probabilities. *JOURNAL OF WILDLIFE MANAGEMENT* 70:1461-1463.
- Knapp, RA, Matthews, KR, Preisler, HK, Jellison, R. 2003. Developing probabilistic models to predict amphibian site occupancy in a patchy landscape. *ECOLOGICAL APPLICATIONS* 13:1069-1082
- Kroll, AJ, Maccracken, JG, McBride, TC, Bakke, J, Light, J, Peterson, P, Bach, J. 2010. Basin-Scale Surveys of Stream-Associated Amphibians in Intensively Managed Forests. *JOURNAL OF WILDLIFE MANAGEMENT* 74:1580-1587.
- McClintock, BT, Bailey, LL, Pollock, KH, Simons, TR. 2010. Experimental Investigation of Observation Error in Anuran Call Surveys. *JOURNAL OF WILDLIFE MANAGEMENT* 74:1882-1893.
- Mazerolle, MJ, Desrochers, A, Rochefort, L. 2005. Landscape characteristics influence pond occupancy by frogs after accounting for detectability. *ECOLOGICAL APPLICATIONS* 15:824-834.

- Russell, KR, Mabee, TJ, Cole, MB. 2004. Distribution and habitat of Columbia torrent salamanders at multiple spatial scales in managed forests of northwestern Oregon. *JOURNAL OF WILDLIFE MANAGEMENT* 68:405-417.
- Russell, KR, Mabee, TJ, Cole, MB, Rochelle, MJ. 2005. Evaluating biotic and abiotic influences on torrent salamanders in managed forests of western Oregon. *WILDLIFE SOCIETY BULLETIN* 33:1413-1424.
- Royle, JA. 2004. Modeling abundance index data from anuran calling surveys. *CONSERVATION BIOLOGY* 18:1378-1385.
- Schmidt, BR, Pellet, J. 2005. Relative importance of population processes and habitat characteristics in determining site occupancy of two anurans. *JOURNAL OF WILDLIFE MANAGEMENT* 69:884-893.
- Sepulveda, AJ, Lowe, WH. 2009. Local and Landscape-Scale Influences on the Occurrence and Density of *Dicamptodon aterrimus*, the Idaho Giant Salamander. *JOURNAL OF HERPETOLOGY* 43:469-484.
- Sewell, D, Beebee, TJC, Griffiths, RA. 2010. Optimising biodiversity assessments by volunteers: The application of occupancy modelling to large-scale amphibian surveys. *BIOLOGICAL CONSERVATION* 143:2102-2110.
- ter Braak, CJF, Etienne, RS. 2003. Improved Bayesian analysis of metapopulation data with an application to a tree frog metapopulation. *ECOLOGY* 84:231-240.
- Walls, SC, Waddle, JH, Dorazio, RM. 2011. Estimating Occupancy Dynamics in an Anuran Assemblage From Louisiana, USA. *JOURNAL OF WILDLIFE MANAGEMENT* 75:751-761.
- Weir, LA, Royle, JA, Nanjappa, P, Jung, RE. 2005. Modeling anuran detection and site occupancy on North American Amphibian Monitoring Program (NAAMP) routes in Maryland. *JOURNAL OF HERPETOLOGY* 39:627-639.
- Welsh, HH, Dunk, JR, Zielinski, WJ. 2006. Developing and applying habitat models using forest inventory data: An example using a terrestrial salamander. *JOURNAL*

OF WILDLIFE MANAGEMENT 70:671-681.

Welsh, HH, Pope, KL, Wheeler, CA. 2008. Using multiple metrics to assess the effects of forest succession on population status: A comparative study of two terrestrial salamanders in the US Pacific Northwest. *BIOLOGICAL CONSERVATION* 141:1149-1160.

BIRDS

Atwood, TC, Young, JK, Beckmann, JP, Breck, SW, Fike, J, Rhodes, OE, Bristow, KD. 2011. Modeling connectivity of black bears in a desert sky island archipelago. *BIOLOGICAL CONSERVATION* 144:2851-2862.

Bailey, LL, Reid, JA, Forsman, ED, Nichols, JD. 2009. Modeling co-occurrence of northern spotted and barred owls: Accounting for detection probability differences. *BIOLOGICAL CONSERVATION* 142:2983-2989.

Beaudry, F, Pidgeon, AM, Radeloff, VC, Howe, RW, Mladenoff, DJ, Bartelt, GA. 2010. Modeling regional-scale habitat of forest birds when land management guidelines are needed but information is limited. *BIOLOGICAL CONSERVATION* 143:1759-1769.

Beck, JL, Skorkowsky, RC, Hayward, GD. 2011. Estimating Occupancy to Monitor Northern Goshawk in the Central Rocky Mountains. *JOURNAL OF WILDLIFE MANAGEMENT* 75:513-524.

Beissinger, SR, Walters, JR, Catanzaro, DG, Smith, KG, Dunning, JB, Haig, SM, Noon, BR, Stith, BM. 2006. Modeling approaches in avian conservation and the role of field biologists. *AUK* 123:1-56.

Belthoff, JR, Smith, BW. 2003. Patterns of artificial burrow occupancy and reuse by burrowing owls in Idaho. *WILDLIFE SOCIETY BULLETIN* 31:138-144.

Betts, MG, Forbes, GJ, Diamond, AW. 2007. Thresholds in songbird occurrence in relation to landscape structure. *CONSERVATION BIOLOGY* 21:1046-1058.

Blakesley, JA, Noon, BR, Anderson, DR. 2005. Site occupancy, apparent survival, and reproduction of California spotted owls in relation to forest stand characteristics. *JOURNAL OF WILDLIFE MANAGEMENT* 69:1554-1564.

Bled, F, Royle, JA, Cam, E. 2011. Hierarchical modeling of

- an invasive spread: the Eurasian Collared-Dove *Streptopelia decaocto* in the United States. *ECOLOGICAL APPLICATIONS* 21:290-302.
- Burton, AC, Sam, MK, Kpelle, DG, Balangtaa, C, Buedi, EB, Brashares, JS. 2011. Evaluating persistence and its predictors in a West African carnivore community. *BIOLOGICAL CONSERVATION* 144:2344-2353.
- Cassey, P, Lockwood, JL, Fenn, KH. 2007. Using long-term occupancy information to inform the management of Cape Sable seaside sparrows in the Everglades. *BIOLOGICAL CONSERVATION* 139:139-149.
- Castellon, TD, Sieving, KE. 2006. Landscape history, fragmentation, and patch occupancy: Models for a forest bird with limited dispersal. *ECOLOGICAL APPLICATIONS* 16:2223-2234.
- Collier, BA, Morrison, ML, Farrell, SL, Campomizzi, AJ, Butcher, JA, Hays, KB, Mackenzie, DI, Wilkins, RN. 2010. Monitoring Golden-Cheeked Warblers on Private Lands in Texas. *JOURNAL OF WILDLIFE MANAGEMENT* 74:140-147.
- Darrah, AJ, Kremetz, DG. 2009. Distribution and Habitat Use of King Rails in the Illinois and Upper Mississippi River Valleys. *JOURNAL OF WILDLIFE MANAGEMENT* 73:1380-1386.
- De Wan, AA, Sullivan, PJ, Lembo, AJ, Smith, CR, Maerz, JC, Lassoie, JP, Richmond, ME. 2009. Using occupancy models of forest breeding birds to prioritize conservation planning. *BIOLOGICAL CONSERVATION* 142:982-991.
- Dewey, SR, Kennedy, PL, Stephens, RM. 2003. Are dawn vocalization surveys effective for monitoring goshawk nest-area occupancy? *JOURNAL OF WILDLIFE MANAGEMENT* 67:390-397.
- Donner, DM, Ribic, CA, Probst, JR. 2010. Patch dynamics and the timing of colonization-abandonment events by male Kirtland's Warblers in an early succession habitat. *BIOLOGICAL CONSERVATION* 143:1159-1167.
- Donovan, TM, Flather, CH. 2002. Relationships among North American songbird trends, habitat fragmentation, and landscape occupancy. *ECOLOGICAL APPLI-*

- CATIONS 12:364-374.
- dos Anjos, L, Collins, CD, Holt, RD, Volpato, GH, Mendonca, LB, Lopes, EV, Bocon, R, Bisheimer, MV, Serafini, PP, Carvalho, J. 2011. Bird species abundance-occupancy patterns and sensitivity to forest fragmentation: Implications for conservation in the Brazilian Atlantic forest. *BIOLOGICAL CONSERVATION* 144:2213-2222.
- Driscoll, MJL, Donovan, T, Mickey, R, Howard, A, Fleming, KK. 2005. Determinants of wood thrush nest success: A multi-scale, model selection approach. *JOURNAL OF WILDLIFE MANAGEMENT* 69:699-709.
- Dugger, KM, Anthony, RG, Andrews, LS. 2011. Transient dynamics of invasive competition: Barred Owls, Spotted Owls, habitat, and the demons of competition present. *ECOLOGICAL APPLICATIONS* 21:2459-2468.
- Duren, KR, Buler, JJ, Jones, W, Williams, CK. 2011. An Improved Multi-Scale Approach to Modeling Habitat Occupancy of Northern Bobwhite. *JOURNAL OF WILDLIFE MANAGEMENT* 75:1700-1709.
- Evans, KL, Newson, SE, Gaston, KJ. 2009. Habitat influences on urban avian assemblages. *IBIS* 151:19-39.
- Ferland, CL, Forsman, ED, Hargis, CD. 2006. A field test of the northern goshawk bioregional monitoring design: Is it cost effective? *WILDLIFE SOCIETY BULLETIN* 34:215-217.
- Goodwin, SE, Shriver, WG. 2011. Effects of Traffic Noise on Occupancy Patterns of Forest Birds. *CONSERVATION BIOLOGY* 25:406-411.
- Groce, JE, Morrison, ML, 2010. Habitat Use by Saw-Whet Owls in the Sierra Nevada. *JOURNAL OF WILDLIFE MANAGEMENT* 74:1523-1532.
- Hamer, TE, Varland, DE, McDonald, TL, Meekins, D. 2008. Predictive model of habitat suitability for the marbled murrelet in western Washington. *JOURNAL OF WILDLIFE MANAGEMENT* 72:983-993.
- Hansen, CP, Millsbaugh, JJ, Rumble, MA. 2011. Occupancy Modeling of Ruffed Grouse in the Black Hills National Forest. *JOURNAL OF WILDLIFE MANAGEMENT* 75:71-77.
- Henneman, C, Mcleod, MA, Andersen, DE. 2007. Red-

- shouldered hawk occupancy surveys in central Minnesota, USA. *JOURNAL OF WILDLIFE MANAGEMENT* 71:526-533.
- Henneman, C, Andersen, DE. 2009. Occupancy Models of Nesting-Season Habitat Associations of Red-Shouldered Hawks in Central Minnesota. *JOURNAL OF WILDLIFE MANAGEMENT* 73:1316-1324.
- Hollenbeck, JP, Saab, VA, Frenzel, RW. 2011. Habitat Suitability and Nest Survival of White-Headed Woodpeckers in Unburned Forests of Oregon. *JOURNAL*
- Howell, JE, Peterson, JT, Conroy, MJ. 2008. Building hierarchical models of avian distributions for the state of Georgia. *JOURNAL OF WILDLIFE MANAGEMENT* 72:168-178.
- Huste, A, Boulinier, T. 2011. Determinants of bird community composition on patches in the suburbs of Paris, France. *BIOLOGICAL CONSERVATION* 144:243-252.
- Kennedy, CM, Grant, EHC, Neel, MC, Fagan, WF, Marra, PP. 2011. Landscape matrix mediates occupancy dynamics of Neotropical avian insectivores. *ECOLOGICAL APPLICATIONS* 21:1837-1850.
- Kroll, AJ, Duke, SD, Runde, DE, Arnett, EB, Austin, KA. 2007. Modeling habitat occupancy of orange-crowned warblers in managed forests of Oregon and Washington, USA. *JOURNAL OF WILDLIFE MANAGEMENT* 71:1089-1097.
- Kroll, AJ, Fleming, TL, Irwin, LL. 2010. Site Occupancy Dynamics of Northern Spotted Owls in the Eastern Cascades, Washington, USA, 1990-2003. *JOURNAL OF WILDLIFE MANAGEMENT* 74:1264-1274
- Kudo, T, Ozaki, K, Takao, G, Sakai, T, Yonekawa, H, Ikeda, K. 2005. Landscape analysis of northern goshawk breeding home range in northern Japan. *JOURNAL OF WILDLIFE MANAGEMENT* 69:1229-1239.
- Lawes, MJ, Fly, S, Piper, SE. 2006. Gamebird vulnerability to forest fragmentation: patch occupancy of the crested guineafowl (*Guttera edouardi*) in Afromontane forests. *ANIMAL CONSERVATION* 9:67-74.
- McNew, LB, Prebyl, TJ, Sandercock, BK, 2012. Effects of

- rangeland management on the site occupancy dynamics of prairie-chickens in a protected prairie preserve. *JOURNAL OF WILDLIFE MANAGEMENT* 76:38-47.
- McShea, WJ, Stewart, C, Peterson, L, Erb, P, Stuebing, R, Gimán, B. 2009. The importance of secondary forest blocks for terrestrial mammals within an Acacia/secondary forest matrix in Sarawak, Malaysia. *BIOLOGICAL CONSERVATION* 142:3108-3119.
- Magness, DR, Wilkins, RN, Hejl, SJ. 2006. Quantitative relationships among golden-cheeked warbler occurrence and landscape size, composition, and structure. *WILDLIFE SOCIETY BULLETIN* 34:473-479.
- Martin, J, McIntyre, CL, Hines, JE, Nichols, JD, Schmutz, JA, MacCluskie, MC. 2009. Dynamic multistate site occupancy models to evaluate hypotheses relevant to conservation of Golden Eagles in Denali National Park, Alaska. *BIOLOGICAL CONSERVATION* 142:2726-2731.
- Martinez, JE, Pagan, I, Calvo, JF. 2006. Factors influencing territorial occupancy and reproductive output in the Booted Eagle *Hieraaetus pennatus*. *IBIS* 148:807-819.
- Montague-Drake, RM, Lindenmayer, DB, Cunningham, RB. 2009. Factors affecting site occupancy by woodland bird species of conservation concern. *BIOLOGICAL CONSERVATION* 142:2896-2903.
- O'Brien, TG, Kinnaird, MF. 2008. A picture is worth a thousand words: the application of camera trapping to the study of birds. *BIRD CONSERVATION INTERNATIONAL* 18:S144-S162.
- Olson, GS, Anthony, RG, Forsman, ED, Ackers, SH, Loschl, PJ, Reid, JA, Dugger, KM, Glenn, EM, Ripple, WJ. 2005. Modeling of site occupancy dynamics for northern spotted owls, with emphasis on the effects of barred owls. *JOURNAL OF WILDLIFE MANAGEMENT* 69:918-932.
- Parker, TH, Stansberry, BM, Becker, CD, Gipson, PS. 2005. Edge and area effects on the occurrence of migrant forest songbirds. *CONSERVATION BIOLOGY* 19:1157-1167.
- Popescu, VD, Gibbs, JP. 2009. Interactions between cli-

- mate, beaver activity, and pond occupancy by the cold-adapted mink frog in New York State, USA. *BIOLOGICAL CONSERVATION* 142:2059-2068.
- Radford, JQ, Bennett, AF. 2004. Thresholds in landscape parameters: occurrence of the white-browed treecreeper *Climacteris affinis* in Victoria, Australia. *BIOLOGICAL CONSERVATION* 117:375-391.
- Richmond, OMW, Hines, JE, Beissinger, SR. 2010. Two-species occupancy models: a new parameterization applied to co-occurrence of secretive rails *ECOLOGICAL APPLICATIONS* 20:2036-2046.
- Risk, BB, de Valpine, P, Beissinger, SR. 2011. A robust-design formulation of the incidence function model of metapopulation dynamics applied to two species of rails. *ECOLOGY* 92:462-474.
- Roberts, SL, van Wagendonk, JW, Miles, AK, Kelt, DA. 2011. Effects of fire on spotted owl site occupancy in a late-successional forest. *BIOLOGICAL CONSERVATION* 144:610-619.
- Ruiz-Gutierrez, V, Zipkin, EF, Dhondt, AA. 2010. Occupancy dynamics in a tropical bird community: unexpectedly high forest use by birds classified as non-forest species. *JOURNAL OF APPLIED ECOLOGY* 47:621-630.
- Studds, CE, Marra, PP. 2005. Nonbreeding habitat occupancy and population processes: An upgrade experiment with a migratory bird. *ECOLOGY* 86:2380-2385.
- Syartinilia, Tsuyuki, S. 2008. GIS-based modeling of Javan Hawk-Eagle distribution using logistic and autologistic regression models. *BIOLOGICAL CONSERVATION* 141:756-769.
- Szabo, JK, Davy, PJ, Hooper, MJ, Astheimer, LB. 2009. Predicting avian distributions to evaluate spatiotemporal overlap with locust control operations in eastern Australia. *ECOLOGICAL APPLICATIONS* 19:2026-2037.
- Tipton, HC, Dreitz, VJ, Doherty, PF. 2008. Occupancy of mountain plover and burrowing owl in Colorado. *JOURNAL OF WILDLIFE MANAGEMENT* 72:1001-

1006.

- Vogeli, M, Serrano, D, Pacios, F, Tella, JL. 2010. The relative importance of patch habitat quality and landscape attributes on a declining steppe-bird metapopulation. *BIOLOGICAL CONSERVATION* 143:1057-1067.
- Watson, CA, Weckerly, FW, Hatfield, JS, Farquhar, CC, Williamson, PS. 2008. Presence-nonpresence surveys of golden-cheeked warblers: detection, occupancy and survey effort. *ANIMAL CONSERVATION* 11:484-492.
- Wiens, JD, Anthony, RG, Forsman, ED. 2011. Barred Owl Occupancy Surveys Within the Range of the Northern Spotted Owl. *JOURNAL OF WILDLIFE MANAGEMENT* 75: 531-538.
- Winchell, CS, Doherty, PF. 2008. Using California gnatcatcher to test underlying models in habitat conservation plans. *JOURNAL OF WILDLIFE MANAGEMENT* 72:1322-1327.
- Winarni, NL, O'Brien, TG, Carroll, JP, Kinnaird, MF. 2009. Movements, distribution, and abundance of great argus pheasants (*Argusianus argus*) in a Sumatran rainforest, *AUK* 126:341-350.
- Zicus, MC, Rave, DP, Das, A, Riggs, MR, Buitenwerf, ML. 2006. Influence of land use on mallard nest-structure occupancy. *JOURNAL OF WILDLIFE MANAGEMENT* 70:1325-1333.

MAMMALS

- Andelt, WF, White, GC, Schnurr, PM, Navo, KW. 2009. Occupancy of Random Plots by White-Tailed and Gunnison's Prairie Dogs. *JOURNAL OF WILDLIFE MANAGEMENT* 73:35-44.
- Avila-Flores, R, Boyce, MS, Boutin, S. 2010. Habitat Selection by Prairie Dogs in a Disturbed Landscape at the Edge of Their Geographic Range. *JOURNAL OF WILDLIFE MANAGEMENT* 74:945-953.
- Ball, LC, Doherty, PF, McDonald, MW. 2005. An occupancy modeling approach to evaluating a Palm Springs ground squirrel habitat model. *JOURNAL OF WILDLIFE MANAGEMENT* 69:894-904.

- Ball, LC, Doherty, PF, Ostermann-Kelm, SD, McDonald, MW. 2010. Effects of Rain on Palm Springs Ground Squirrel Occupancy in the Sonoran Desert. *JOURNAL OF WILDLIFE MANAGEMENT* 74:954-962.
- Boulanger, J, White, GC, Proctor, M, Stenhouse, G, Machutchon, G, Himmer, S. 2008. Use of occupancy models to estimate the influence of previous live captures on DNA-based detection probabilities of grizzly bears. *JOURNAL OF WILDLIFE MANAGEMENT* 72:589-595.
- Buij, R, McShea, WJ, Campbell, P, Lee, ME, Dallmeier, F, Guimondou, S, Mackaga, L, Guisseougou, N, Mboumba, S, Hines, JE, Nichols, JD, Alonso, A. 2007. Patch-occupancy models indicate human activity as major determinant of forest elephant *Loxodonta cyclotis* seasonal distribution in an industrial corridor in Gabon. *BIOLOGICAL CONSERVATION* 135:189-201.
- Conde, DA, Colchero, F, Zarza, H, Christensen, NL, Sexton, JO, Manterola, C, Chavez, C, Rivera, A, Azuara, D, Ceballos, G. 2010. Sex matters: Modeling male and female habitat differences for jaguar conservation. *BIOLOGICAL CONSERVATION* 143:1980-1988.
- Cotner, LA, Schooley, RL. 2011. Habitat Occupancy by Riparian Muskrats Reveals Tolerance to Urbanization and Invasive Vegetation. *JOURNAL OF WILDLIFE MANAGEMENT* 75:1637-1645.
- Eaton, MJ, Hughes, PT, Nichols, JD, Morkill, A, Anderson, C. 2011. Spatial Patch Occupancy Patterns of the Lower Keys Marsh Rabbit. *JOURNAL OF WILDLIFE MANAGEMENT* 75:1186-1193.
- Finley, DJ, White, GC, Fitzgerald, JP. 2005. Estimation of swift fox population size and occupancy rates in eastern Colorado. *JOURNAL OF WILDLIFE MANAGEMENT* 69:861-873.
- Gardner, CL, Lawler, JP, Hoef, JMV, Magoun, AJ, Kellie, KA. 2010. Coarse-Scale Distribution Surveys and Occurrence Probability Modeling for Wolverine in Interior Alaska. *JOURNAL OF WILDLIFE MANAGEMENT*

74:1894-1903.

- Gopalaswamy, AM, Karanth, KU, Kumar, NS, Macdonald, DW. (2012) Estimating tropical forest densities using abundance models of occupancy. *ANIMAL CONSERVATION*
- Graves, TA, Kendall, KC, Royle, JA, Stetz, JB, Macleod, AC. 2011. Linking landscape characteristics to local grizzly bear abundance using multiple detection methods in a hierarchical model. *ANIMAL CONSERVATION* 14:652-664.
- Griffin, SC, Taper, ML, Hoffman, R, Mills, LS. 2008. The case of the missing marmots: Are metapopulation dynamics or range-wide declines responsible? *BIOLOGICAL CONSERVATION* 141:1293-1309.
- Hines, JE, Nichols, JD, Royle, JA, MacKenzie, DI, Gopalaswamy, AM, Kumar, NS, Karanth, KU. 2010. Tigers on trails: occupancy modeling for cluster sampling. *ECOLOGICAL APPLICATIONS* 20:1456-1466.
- Jaquiere, J, Guelat, J, Broquet, T, Berset-Brandli, L, Pellegrini, E, Moresi, R, Hirzel, AH, Perrin, N. 2008. Habitat-quality effects on metapopulation dynamics in greater white-toothed shrews, *crocidura russula*. *ECOLOGY* 89:2777-2785.
- Jeffress, MR, Paukert, CP, Sandercock, BK, Gipson, PS. 2011. Factors Affecting Detectability of River Otters During Sign Surveys. *JOURNAL OF WILDLIFE MANAGEMENT* 75:144-150.
- Karanth, KK, Nichols, JD, Hines, JE, Karanth, KU, Christensen, NL. 2009. Patterns and determinants of mammal species occurrence in India. *JOURNAL OF APPLIED ECOLOGY* 46:1189-2000.
- Karanth, KK, Nichols, JD, Hines, JE. 2010. Occurrence and distribution of Indian primates. *BIOLOGICAL CONSERVATION* 143:2891-2899.
- Karanth, KU, Gopalaswamy, AM, Kumar, NS, Vaidyanathan, S, Nichols, JD, MacKenzie, DI. 2011. Monitoring carnivore populations at the landscape scale: occupancy modelling of tigers from sign surveys. *JOURNAL OF APPLIED ECOLOGY*
- Kawanishi, K, Sunquist, ME. 2004. Conservation status of tigers in a primary rainforest of Peninsular Malaysia.

- BIOLOGICAL CONSERVATION 120:329-344.
- Licona, M, McCleery, R, Collier, B, Brightsmith, DJ, Lopez, R. 2011. Using ungulate occurrence to evaluate community-based conservation within a biosphere reserve model. ANIMAL CONSERVATION 14:206-214.
- Linkie, M, Dinata, Y, Nugroho, A, Haidir, IA. 2007. Estimating occupancy of a data deficient mammalian species living in tropical rainforests: Sun bears in the Kerinci Seblat region, Sumatra. BIOLOGICAL CONSERVATION 137:20-27.
- Litvaitis, JA, Tash, JP Litvaitis, MK Marchand, MN Kovach, Al Innes, R. 2006. A range-wide survey to determine the current distribution of New England cottontails. WILDLIFE SOCIETY BULLETIN 34:1190-1197.
- Magle, SB, Reyes, P, Zhu, J, Crooks, KR. 2010. Extirpation, colonization, and habitat dynamics of a keystone species along an urban gradient. BIOLOGICAL CONSERVATION 143:2146-2155.
- Magoun, AJ, Ray, JC, Johnson, DS, Valkenburg, P, Dawson, FN, Bowman, J. 2007. Modeling wolverine occurrence using aerial surveys of tracks in snow. JOURNAL OF WILDLIFE MANAGEMENT 71:2221-2229.
- Martin, J, Chamaille-Jammes, S, Nichols, JD, Fritz, H, Hines, JE, Fongesbeck, CJ, MacKenzie, DI, Bailey, LL. 2010. Simultaneous modeling of habitat suitability, occupancy, and relative abundance: African elephants in Zimbabwe. ECOLOGICAL APPLICATIONS 20:1173-1182.
- Miller, MW, Pearlstine, EV, Dorazio, RM, Mazzotti, FJ. 2011. Occupancy and Abundance of Wintering Birds in a Dynamic Agricultural Landscape. JOURNAL OF WILDLIFE MANAGEMENT 75:836-847.
- Moore, JE, Swihart, RK. 2005. Modeling patch occupancy by forest rodents: Incorporating detectability and spatial autocorrelation with hierarchically structured data. JOURNAL OF WILDLIFE MANAGEMENT 69:933-949.
- Moriarty, KM, Zielinski, WJ, Forsman, ED. 2011. Decline in American Marten Occupancy Rates at Sagehen Experimental Forest, California. JOURNAL OF WILDLIFE

- MANAGEMENT 75:1774-1787.
- Mortelliti, A, Amori, G, Capizzi, D, Cervone, C, Fagiani, S, Pollini, B, Boitani, L. 2011. Independent effects of habitat loss, habitat fragmentation and structural connectivity on the distribution of two arboreal rodents. *JOURNAL OF APPLIED ECOLOGY* 48:153-162.
- Nielsen, CLR, Wakamiya, SM, Nielsen, CK. 2008. Viability and patch occupancy of a swamp rabbit metapopulation at the northern edge of its distribution. *BIOLOGICAL CONSERVATION* 141:1043-1054.
- Nielsen, SE, McDermid, G, Stenhouse, GB, Boyce, MS. 2010. Dynamic wildlife habitat models: Seasonal foods and mortality risk predict occupancy-abundance and habitat selection in grizzly bears. *BIOLOGICAL CONSERVATION* 143:1623-1634.
- Norris, D, Rocha-Mendes, F, Ferraz, SFD, Villani, JP, Galetti, M. 2011. How to not inflate population estimates? Spatial density distribution of white-lipped peccaries in a continuous Atlantic forest. *ANIMAL CONSERVATION* 14:492-501.
- O'Brien, CM, Crowther, MS, Dickman, CR, Keating, J. 2008. Metapopulation dynamics and threatened species management: Why does the broad-toothed rat (*Mastacomys fuscus*) persist? *BIOLOGICAL CONSERVATION* 141:1962-1971.
- O'Connell, AF, Talancy, NW, Bailey, LL, Sauer, JR, Cook, R, Gilbert, AT. 2006. Estimating site occupancy and detection probability parameters for meso- and large mammals in a coastal ecosystem. *JOURNAL OF WILDLIFE MANAGEMENT* 70:1625-1633.
- Ober, HK, Hayes, JP. 2008. Influence of vegetation on bat use of riparian areas at multiple spatial scales. *JOURNAL OF WILDLIFE MANAGEMENT* 72:396-404.
- Pillay, R, Johnsingh, AJT, Raghunath, R, Madhusudan, MD. 2011. Patterns of spatiotemporal change in large mammal distribution and abundance in the southern Western Ghats, India. *BIOLOGICAL CONSERVATION* 144:1567-1576.
- Potvin, MJ, Drummer, TD, Vucetich, JA, Beyer, DE, Peterson, RO, Hammill, JH. 2005. Monitoring and habitat analysis for wolves in upper Michigan. *JOURNAL OF*

- WILDLIFE MANAGEMENT 69:1660-1669.
- Reunanen, P, Nikula, A, Monkkonen, M, Hurme, E, Nivala, V. 2002. Predicting occupancy for the Siberian flying squirrel in old-growth forest patches. ECOLOGICAL APPLICATIONS 12:1188-1198.
- Smith, JB, Jenks, JA, Klaver, RW. 2007. Evaluating detection probabilities for American marten in the Black Hills, South Dakota. JOURNAL OF WILDLIFE MANAGEMENT 71:2412-2416.
- Spencer, W, Rustigian-Romsos, H, Strittholt, J, Scheller, R, Zielinski, W, Truex, R. 2011. Using occupancy and population models to assess habitat conservation opportunities for an isolated carnivore population. BIOLOGICAL CONSERVATION 144:788-803.
- Thorn, M, Green, M, Bateman, PW, Waite, S, Scott, DM. 2011. Brown hyaenas on roads: Estimating carnivore occupancy and abundance using spatially auto-correlated sign survey replicates. BIOLOGICAL CONSERVATION 144:1799-1807.
- Tosh, CA, Reyers, B, van Jaarsveld, AS. 2004. Estimating the abundances of large herbivores in the Kruger National Park using presence-absence data. ANIMAL CONSERVATION 7:55-61.
- Trudeau, C, Imbeau, L, Drapeau, P, Mazerolle, MJ. 2011. Site Occupancy and Cavity Use by the Northern Flying Squirrel in the Boreal Forest. JOURNAL OF WILDLIFE MANAGEMENT 75:1646-1656.
- Vors, LS, Schaefer, JA, Pond, BA, Rodgers, AR, Patterson, BR. 2007. Woodland caribou extirpation and anthropogenic landscape disturbance in Ontario. JOURNAL OF WILDLIFE MANAGEMENT 71:1249-1256.
- Weller, TJ. 2008. Using occupancy estimation to assess the effectiveness of a regional multiple-species conservation plan: Bats in the Pacific Northwest. BIOLOGICAL CONSERVATION 141:2279-2289.
- Weir, RD, Corbould, FB, 2010. Factors Affecting Landscape Occupancy by Fishers in North-Central British Columbia. JOURNAL OF WILDLIFE MANAGEMENT 74:405-410.
- Wilson, TL, Odei, JB, Hooten, MB, Edwards, TC. 2010.

- Hierarchical spatial models for predicting pygmy rabbit distribution and relative abundance. *JOURNAL OF APPLIED ECOLOGY* 47:401-409.
- Winchester, C, Castleberry, SB, Mengak, MT. 2009. Evaluation of Factors Restricting Distribution of the Endangered Key Largo Woodrat. *JOURNAL OF WILDLIFE MANAGEMENT* 73:374-379.
- Wintle, BA, Kavanagh, RP, McCarthy, MA, Burgman, MA. 2005. Estimating and dealing with detectability in occupancy surveys for forest owls and arboreal marsupials. *JOURNAL OF WILDLIFE MANAGEMENT* 69:905-917.
- Yates, MD, Muzika, RM. 2006. Effect of forest structure and fragmentation on site occupancy of bat species in Missouri ozark forests. *JOURNAL OF WILDLIFE MANAGEMENT* 70:1238-1248.
- Zeller, KA, Nijhawan, S, Salom-Perez, R, Potosme, SH, Hines, JE. 2011. Integrating occupancy modeling and interview data for corridor identification: A case study for jaguars in Nicaragua. *BIOLOGICAL CONSERVATION* 144:892-901.
- Zielinski, WJ, Slauson, KM, Bowles, AE. 2008. Effects of off-highway vehicle use on the American marten. *JOURNAL OF WILDLIFE MANAGEMENT* 72:1558-1571.

REPTILES

- Durso, AM, Willson, JD, Winne, CT. 2011. Needles in haystacks: Estimating detection probability and occupancy of rare and cryptic snakes. *BIOLOGICAL CONSERVATION* 144:1508-1515.
- Kery, M. 2002. Inferring the absence of a species - A case study of snakes. *JOURNAL OF WILDLIFE MANAGEMENT* 66:330-338.
- Zylstra, ER, Steidl, RJ. 2009. Habitat Use by Sonoran Desert Tortoises. *JOURNAL OF WILDLIFE MANAGEMENT* 73:747-754.
- Zylstra, ER, Steidl, RJ, Swann, DE, 2010. Evaluating Survey Methods for Monitoring a Rare Vertebrate, the Sonoran Desert Tortoise. *JOURNAL OF WILDLIFE MANAGE-*

INVERTEBRATES

- Hodgson, JA, Moilanen, A, Bourn, NAD, Bulman, CR, Thomas, CD. 2009. Managing successional species: Modelling the dependence of heath fritillary populations on the spatial distribution of woodland management. *BIOLOGICAL CONSERVATION* 142:2743-2751.
- Grundel, R, Pavlovic, NB. 2007. Resource availability, matrix quality, microclimate, and spatial pattern as predictors of patch use by the Karner blue butterfly. *BIOLOGICAL CONSERVATION* 135:135-144.
- Krauss, J, Steffan-Dewenter, I, Tschardtke, T. 2004. Landscape occupancy and local population size depends on host plant distribution in the butterfly *Cupido minimus*. *BIOLOGICAL CONSERVATION* 120:355-361.
- Maes, D, Bonte, D. 2006. Using distribution patterns of five threatened invertebrates in a highly fragmented dune landscape to develop a multispecies conservation approach. *BIOLOGICAL CONSERVATION* 133:490-499.
- Nowicki, P, Pepkowska, A, Kudlek, J, Skorka, P, Witek, M, Settele, J, Woyciechowski, M. 2007. From metapopulation theory to conservation recommendations: Lessons from spatial occurrence and abundance patterns of *Maculinea* butterflies. *BIOLOGICAL CONSERVATION* 140:119-129.
- van Strien, AJ, van Swaay, CAM, Kery, M. 2011. Metapopulation dynamics in the butterfly *Hipparchia semele* changed decades before occupancy declined in The Netherlands. *ECOLOGICAL APPLICATIONS* 21:2510-2520.

PLANTS

- Verheyen, K, Vellend, M, Van Calster, H, Peterken, G, Hermy, M. 2004. Metapopulation dynamics in changing landscapes: A new spatially realistic model for forest plants. *ECOLOGY* 85:3302-3312.

OTHER JOURNALS WITH RECENT PAPERS ON OCCUPANCY

ACTA OECOLOGICA - INTERNATIONAL JOURNAL OF ECOLOGY
ADVANCES IN ECOLOGICAL RESEARCH
AMERICAN MIDLAND NATURALIST
AMERICAN NATURALIST
ANNALES ZOOLOGICI FENNICI
APPLIED VEGETATION SCIENCE
AUSTRAL ECOLOGY
BASIC AND APPLIED ECOLOGY
BEHAVIORAL ECOLOGY
BIODIVERSITY AND CONSERVATION
COMMUNITY ECOLOGY
DIVERSITY AND DISTRIBUTIONS
ECOGRAPHY
ECOLOGICAL COMPLEXITY
ECOLOGICAL ECONOMICS
ECOLOGICAL INFORMATICS
ECOLOGICAL MODELLING
ECOLOGICAL MONOGRAPHS
ECOLOGICAL RESEARCH
ECOLOGY LETTERS
ECOSCIENCE
ECOSYSTEMS
EKOLOJI
ENVIRONMENTAL BIOLOGY OF FISHES
EUROPEAN JOURNAL OF WILDLIFE RESEARCH
EVOLUTION
EVOLUTIONARY ECOLOGY RESEARCH
FUNGAL ECOLOGY
GLOBAL CHANGE BIOLOGY
GLOBAL ECOLOGY AND BIOGEOGRAPHY
INTERCIENCIA
ISRAEL JOURNAL OF ECOLOGY & EVOLUTION
JOURNAL OF ANIMAL ECOLOGY
JOURNAL OF BIOGEOGRAPHY
JOURNAL OF ECOLOGY

JOURNAL OF EXPERIMENTAL MARINE BIOLOGY AND
ECOLOGY
JOURNAL OF PLANT ECOLOGY-UK
JOURNAL OF THE NORTH AMERICAN BENTHOLOGICAL
SOCIETY
JOURNAL OF TROPICAL ECOLOGY
JOURNAL OF VEGETATION SCIENCE
LANDSCAPE AND URBAN PLANNING
LANDSCAPE ECOLOGY
MARINE ECOLOGY-PROGRESS SERIES
METHODS IN ECOLOGY AND EVOLUTION
MICROBIAL ECOLOGY
MOLECULAR ECOLOGY RESOURCES
NATURAL AREAS JOURNAL
NEW ZEALAND JOURNAL OF ECOLOGY
NORTHEASTERN NATURALIST
OECOLOGIA
OHIO JOURNAL OF SCIENCE
OIKOS
ORYX
PEDOBIOLOGIA
PLANT ECOLOGY
POPULATION ECOLOGY
PROCEEDINGS OF THE ROYAL SOCIETY B-BIOLOGICAL
SCIENCES
PROCEEDINGS OF THE ROYAL SOCIETY OF LONDON
SERIES B-BIOLOGICAL SCIENCES
RANGELAND ECOLOGY & MANAGEMENT
REVISTA CHILENA DE HISTORIA NATURAL
REVUE D ECOLOGIE-LA TERRE ET LA VIE
SOUTH AFRICAN JOURNAL OF WILDLIFE RESEARCH
SOUTHWESTERN NATURALIST
TEXAS JOURNAL OF SCIENCE
THEORETICAL ECOLOGY
THEORETICAL POPULATION BIOLOGY
WETLANDS
WILDLIFE BIOLOGY
WILDLIFE MONOGRAPHS
WILDLIFE RESEARCH

Wildlife Conservation Society
Global Program
2300 Southern Boulevard
Bronx, NY 10460
Tel: 718-220-5100
www.wcs.org

