

eAppendix Table. Distribution of Comorbidities by False-Positive Group

	Total (n = 3920)	Current False Positive (n = 500)	Previous False Positive (n = 594)	No False Positive (n = 2826)	<i>P</i>
Comorbidity Group	n (%)	n (%)	n (%)	n (%)	
Congestive heart failure	69 (1.8)	4 (0.80)	8 (1.35)	57 (2.0)	.11
Cardiac arrhythmias	96 (2.4)	11 (2.2)	14 (2.4)	71 (2.5)	.91
Valvular disease	50 (1.3)	3 (0.60)	4 (0.67)	43 (1.5)	.09
Pulmonary circulation disorders	40 (1.0)	3 (0.60)	8 (1.35)	29 (1.0)	.48
Peripheral vascular disorders	71 (1.8)	7 (1.40)	17 (2.9)	47 (1.7)	.10
Hypertension, uncomplicated	1801 (45.9)	182 (36.4)	314 (52.9)	1305 (46.2)	<.0001
Hypertension, complicated	17 (0.4)	2 (0.40)	1 (0.17)	14 (0.5)	.67
Paralysis	14 (0.4)	3 (0.60)	1 (0.17)	10 (0.4)	.51
Other neurological disorders	85 (2.2)	10 (2.0)	16 (2.7)	59 (2.1)	.63
Chronic pulmonary disease	562 (14.3)	73 (14.6)	109 (18.4)	380 (13.4)	.008
Diabetes, uncomplicated	152 (3.9)	19 (3.8)	31 (5.2)	102 (3.6)	.18
Diabetes, complicated	169 (4.3)	21 (4.2)	20 (3.4)	128 (4.5)	.44
Hypothyroidism	336 (8.6)	48 (9.6)	69 (11.6)	219 (7.7)	.006
Renal failure	68 (1.7)	6 (1.20)	12 (2.0)	50 (1.8)	.56
Liver disease	54 (1.4)	5 (1.00)	10 (1.7)	39 (1.4)	.63
Peptic ulcer disease, excluding bleeding	30 (0.8)	2 (0.40)	9 (1.52)	19 (0.7)	.08
Rheumatoid arthritis/collagen vascular disorders	88 (2.2)	9 (1.80)	17 (2.9)	62 (2.2)	.47
Coagulopathy	42 (1.1)	5 (1.00)	6 (1.01)	31 (1.1)	.97
Obesity	228 (5.8)	31 (6.2)	20 (3.4)	177 (6.3)	.022
Weight loss	131 (3.3)	15 (3.0)	21 (3.5)	95 (3.4)	.88
Fluid/electrolyte disorders	65 (1.7)	10 (2.0)	13 (2.2)	42 (1.5)	.39
Deficiency anemias	270 (6.9)	31 (6.2)	36 (6.1)	203 (7.2)	.50
Alcohol abuse	91 (2.3)	9 (1.80)	11 (1.9)	71 (2.5)	.44
Drug abuse	194 (4.9)	30 (6.0)	31 (5.2)	133 (4.7)	.44
Psychoses	58 (1.5)	6 (1.20)	10 (1.7)	42 (1.5)	.80
Depression	1140 (29.1)	133 (26.6)	195 (32.8)	812 (28.7)	.06
≥1 Elixhauser comorbidity	2971 (75.8)	357 (71.4)	477 (80.3)	2137 (75.6)	.003