



# Giving Chicks a Good Start

Mahmoud El-Begearmi, Extension Poultry Specialist  
H. Michael Opitz, Extension Veterinarian

Bulletin #2219

## Sources of Chicks

First and foremost, get your chicks from a reliable hatchery that is part of the National Poultry Improvement Plan (NPIP). A listing of NPIP participating hatcheries is available through your county Extension office. Request that egg or meat-type chickens be vaccinated at the hatchery against Marek's disease.

## Be Prepared Before Chicks Arrive

Clean the brooder house before the chicks arrive. Start by brushing down cobwebs and dust from ceilings and walls. Sweep and wash the floor thoroughly. Scrub the area with a good disinfectant, then air out the building. Cover the floor with two to four inches of clean, dry litter. Dry pine shavings or sawdust and other absorbent materials, such as shredded newspapers, make good litter. Hardwood shavings or hay are not suitable for litter. Make certain the brooder area is free of rats and mice and protected against predators. Start the brooder 24 hours before the chicks arrive to be sure everything is in a good working order.

## Brooding Temperature

The temperature at the chicks' level (two to three inches from the floor) should be 95 degrees F through the first week. Then reduce the temperature five degrees each week until you reach a temperature of 70 degrees F. You may turn off the brooder on warm days.

Observe the chicks around the brooder to see if they are comfortable and getting enough heat. If they huddle together, they are cold. If they are spread uniformly under the heat source, they are comfortable. If they stay away from the heat source, they are probably too warm. Chicks need fresh air but avoid draught.

## Floor, Feeder and Waterer Space

It is important to provide enough floor, feeder and waterer space. Table 1 (on back) lists the recommended guidelines for space for chicks at different ages. These guidelines work in most cases. Even so, observe your chicks to make sure that they have enough room and easy access to feed and water.

For the first week to 10 days, you can feed chicks in cut-down cardboard boxes. The edges should be about 1 <sup>1</sup>/<sub>4</sub> inches high on the sides. Place feed in piles. The chicks will scratch around. After 10 days, provide enough space (see Table 1) so they can all eat at once. Do not fill the feeders more than half full to avoid wasting feed.

**Table 1: Average Space Requirements per Bird**

Age (Weeks)	Floor (square foot)	Feeder (linear inches)	Waterer (linear inches)
0-4	0.5	1.0	0.5
4-8	1.0	2.0	0.75
8-12	2.0	2.5	1.0
Over 12	3.0 Light	3.0	1.0
	3.5 Heavy	3.0	1.5

**Table 2: Suggested Feeding Program**

Type of Bird	Age (weeks)	Type of Feed
Broilers/Fryers	0-6	Starter ..... 22-23% protein
	6-8	Finisher ..... 18-20% protein
Roasters/Capons	0-6	Starter ..... 22-24% protein
	6-14	Grower/Developer.... 15-17% protein
	To Market	Finisher ..... 16% protein
Pullets (to be layers)	0-6	Starter ..... 18-20% protein
	6-18	Grower ..... 15% protein
	18+	Layer ..... 17% protein

Also, if you use fountain-type waterers, will be used, provide a one-quart jar for every 16 chicks for the first week, a one-quart jar per 20 one- to four-week-old chicks and one-gallon waterer per 20 chicks at 12 weeks of age. Water is essential for proper growth and development. Fresh, clean water should be available at all times. Clean the waterers each day.

### **Suggested Feeding Program**

Start chicks on an all-mash starter for the first six weeks, at which time they can be changed to a finisher ration (meat-type birds) or to an all-mash grower/ developer (other types of birds). Table 2 lists a simple feeding program for different type of chickens at different ages. Always use fresh feed that has not been stored for more than four weeks.

The use of medication against coccidiosis is recommended in starter and grower feed only. Refer to "Biological Control of Coccidiosis in Small Poultry Flocks," bulletin #2259, if you prefer to raise chickens on non-medicated feed. Coccidiosis is a common killer of chickens. Either medicated feed or biological control of this disease is needed.

Other more comprehensive feeding programs are also available for small semi-commercial operations. For more information on these programs, please contact your county Extension office.