



Larder Beetle

Dermestes lardarius L.



larder beetle

Dermestes lardarius Linnaeus

Adult(s)

Photo by Joseph Berger

www.insectimages.org

Injury

In the past, home stored meats and raw hides were frequently damaged by larder beetles. Modern methods of meat storage and meat distribution have eliminated this food source for the beetle larvae. Presently, larder beetles are more of a nuisance pest, although they may attack some pantry products such as dried pet food.

The occurrence of larder beetles in the home may be associated with the presence of a dead rodent—mouse, rat, chipmunk or squirrel—between walls of the house or in an attic or crawl space. Accumulations of dead insects such as cluster flies in lamp globes, between walls or attic windows may also support large numbers of larder beetles. Fledgling birds or abandoned nests beneath the eaves or in an attic may attract larder beetles. Museum specimens, feathers, horn, hair, hides and beeswax along with dried meat or fish, biscuits or other dry pet food are susceptible to attack.

Description and Life History

Adult beetles are about 1/3 inch (8-10mm) long, dark brown in color for most of the body, interrupted by a broad, somewhat yellowish gray band across the front portion of the forewings. The band may show six darker spots. Adult beetles are sometimes observed outdoors where they have been feeding upon the pollen of flowers. See Fig. 1.

Adult beetles are attracted to larval food sources on which they lay their eggs. Eggs are deposited on suitable food sources and are seldom observed. In summer they hatch in 12 days or less.

Larvae are dark reddish brown to near black in color, hairy, and bear a pair of small backward curved spines (urogomphi) near the tip of the abdomen. (See Fig. 2). Larvae feed until the food source is consumed, after which they move about in search of further nourishment. It is this movement that frequently leads to their discovery. In a heavy infestation hundreds of individuals may creep from beneath baseboards, around windowsills or from electrical fixtures. Full-grown larvae are nearly 1/2 inch in length.

When larvae are full-grown, they search for a place to pupate and often bore into nearby structural components, especially wood or paper. The actual damage they do is slight, as they do not bore to eat, simply to have a sheltered place to pupate. Pupae (an inactive stage) are covered with the last larval skin and are found in sheltered places near larval food.

Helping You Put Knowledge to Work

Adults emerge from the pupae and the cycle begins again. A complete generation can occur in 40 to 50 days under ideal conditions (64-68 degrees F. and plenty of food) but, it is not unusual to have only one generation per year in many areas.

Management

Adult beetles can be picked up by hand and discarded when seen. When larger numbers of larder beetles are seen, an effort should be made to search out the food source and clean up any uneaten food. If pet food is the source, consider purchasing smaller quantities, or storing unused portions in an insect resistant container. Rodents should be removed from traps regularly. Birds' nests should be cleared away. Light globes should be emptied of insects regularly, and accumulations of dead insects on windowsills should be vacuumed or cleaned up regularly and discarded. Empty or dispose of vacuum bags- if live insects are left in a vacuum bag they may crawl back out into the storage area.

Screening attics and practicing vigorous rodent control will help prevent larder beetle infestations. Pesticides should only be used as a last resort. A thorough search and clean up is the best approach for managing these insect pests. However, if a pesticide is deemed necessary, the insecticide cyfluthrin is labeled for larder beetle control. Bayer Advanced Home: Home Pest Control Indoor & Outdoor Killer RTU and Real-Kill Home Insect Control Indoor & Outdoor Insect Killer are some products available to consumers for larder beetle control containing cyfluthrin.

4/96, Prepared by: Carolyn Klass

Sr. Extension Associate

Department of Entomology

Cornell University

6/04, Revised by: Carolyn Klass

This publication contains pesticide recommendations. Changes in pesticide regulations occur constantly and human errors are still possible. Some materials mentioned may no longer be available and some uses may no longer be legal. All pesticides distributed, sold or applied in New York State must be registered with the New York State Department of Environmental Conservation (DEC). Questions concerning the legality and/or registration status for pesticide use in New York State should be directed to the appropriate Cornell Cooperative Extension Specialist or your regional DEC office. ***READ THE LABEL BEFORE APPLYING ANY PESTICIDE.***