



**Restriction Map and Multiple Cloning Site (MCS) of pDsRed2.** Unique restriction sites are in bold.

### Description

pDsRed2 is a prokaryotic expression vector that encodes DsRed2, a DsRed variant engineered for faster maturation and lower non-specific aggregation. Derived from the *Discosoma sp.* red fluorescent protein (drFP583; 1), DsRed2, like its progenitor DsRed1, contains a series of silent base-pair changes corresponding to human codon-usage preferences for high expression in mammalian cells (2). In addition to these changes, DsRed2 contains six amino acid substitutions: A105V, I161T, and S197A, which result in the more rapid appearance of red fluorescence in transfected cell lines; and R2A, K5E, and K9T, which prevent the protein from aggregating. (DsRed2 may, however, form the same tetrameric structure as DsRed1 [3].) In mammalian cell cultures when DsRed2 is expressed constitutively, red-emitting cells can be detected by fluorescence microscopy within 24 hours of transfection. Large insoluble aggregates of protein, often observed in bacterial and mammalian cell systems expressing DsRed1, are dramatically reduced in cells expressing DsRed2. The faster-maturing, more soluble red fluorescent protein is also well tolerated by host cells; mammalian cell cultures transfected with DsRed2 show no obvious signs of reduced viability—in those cell lines tested, cells expressing DsRed2 display the same morphology (e.g., adherence, light-refraction) and growth characteristics as non-transfected controls.

In pDsRed2, the DsRed2 coding sequence is flanked by separate and distinct multiple cloning sites at the 5' and 3' ends so that it is easy to excise the gene for use in other applications. Alternatively, the DsRed2 coding sequence can be amplified by PCR. The DsRed 2 gene was inserted in frame with the *lacZ* initiation codon from pUC19 so that DsRed2 is expressed from the *lac* promoter ( $P_{lac}$ ) in *E. coli* host cells. A Kozak consensus sequence is located immediately upstream of the DsRed2 gene to enhance translational efficiency in eukaryotic systems (4). The entire DsRed2 expression cassette in pDsRed2 is supported by a pUC backbone, which contains a high-copy number origin of replication and an ampicillin resistance gene for propagation and selection in *E. coli*.

(PR063558; published 30 June 2010)



**Clontech**

United States/Canada  
800.662.2566

Asia Pacific  
+1.650.919.7300

Europe  
+33.(0)1.3904.6880

Japan  
+81.(0)77.543.6116

Clontech Laboratories, Inc.  
A Takara Bio Company  
1290 Terra Bella Ave.  
Mountain View, CA 94043  
Technical Support (US)  
E-mail: tech@clontech.com  
www.clontech.com

**Use**

pDsRed2 is primarily intended to serve as a source of DsRed2 cDNA. The flanking MCS regions make it possible to excise the DsRed2 coding sequence and insert it into other expression vectors of choice. The vector can also be used in bacteria to produce DsRed2 protein.

**Location of features**

- *lac* Promoter: 95–178
  - CAP binding site: 111–124
  - 35 region: 143–148; –10 region: 167–172
  - Transcription start point: 179
  - lac* operator: 179–199
- *lacZ*-DsRed2 fusion protein expressed in *E. coli*
  - Ribosome binding site: 206–209
  - Start codon (ATG): 217–219; Stop codon 964–966
- 5' Multiple Cloning Site: 240–281
- *Discosoma sp.* Red Fluorescent Protein (DsRed2) gene
  - Kozak consensus translation initiation site: 282–292
  - Start codon (ATG): 289–291; Stop codon: 964–966
- 3' Multiple cloning site: 968–1065
- Ampicillin resistance gene
  - Promoter
    - 35 region: 1441–1446; –10 region: 1464–1469
    - Transcription start point: 1476
  - Ribosome binding site: 1499–1503
  - β-lactamase coding sequences
    - Start codon (ATG): 1511–1513; Stop codon: 2369–2371
    - β-lactamase signal peptide: 1511–1579
    - β-lactamase mature protein: 1580–2368
- pUC plasmid replication origin: 2519–3162

**Sequencing Primer Location**

- DsRed1-C Sequencing Primer (Cat. No. 632388; 5'-AGCTGGACATCACCTCCCACAACG-3'): 881–904

**Propagation in *E. Coli***

- Recommended host strain: DH5α
- Selectable marker: plasmid confers resistance to ampicillin (50 µg/ml) to *E. coli* hosts.
- *E. coli* replication origin: pUC
- Copy number: ~500
- Plasmid incompatibility group: pMB1/ColE1

**Red Fluorescent Protein (DsRed2)**

- Excitation/Emission Maxima: 558 nm / 583 nm

**References**

1. Matz, M. V., *et al.* (1999) *Nature Biotech.* **17**:969–973.
2. Haas, J., *et al.* (1996) *Curr. Biol.* **6**:315–324.
3. Yarbrough, D., *et al.* (2001) *Proc. Natl. Acad. Sci. USA* **98**:462–467.
4. Kozak, M. (1987) *Nucleic Acids Res.* **15**:8125–8148.

**Notice to Purchaser**

Clontech products are to be used for research purposes only. They may not be used for any other purpose, including, but not limited to, use in drugs, *in vitro* diagnostic purposes, therapeutics, or in humans. Clontech products may not be transferred to third parties, resold, modified for resale, or used to manufacture commercial products or to provide a service to third parties without written approval of Clontech Laboratories, Inc.

The RCFP's (including DsRedExpress and DsRedExpress2) are covered by one or more of the following U.S. Patent Nos. 7,166,444; 7,157,565; 7,217,789; 7,338,784; 7,338,783; 7,537,915 6,969,597, 7,150,979 and 7,442,522.

**Living Colors Fluorescent Protein Products:**

**Not-For-Profit Entities:** Orders may be placed in the normal manner by contacting your local representative or Clontech Customer Service at 650.919.7300. At its discretion, Clontech grants Not-For-Profit Entities a non-exclusive, personal, limited license to use this product for non-commercial life science research use only. Such license specifically excludes the right to sell or otherwise transfer this product, its components or derivatives thereof to third parties. No modifications to the protein coding sequence may be made without express written permission from Clontech. Any other use of this product requires a license from Clontech. For license information, please contact a licensing representative by phone at 650.919.7320 or by e-mail at [licensing@clontech.com](mailto:licensing@clontech.com).

For-Profit Entities wishing to use this product are required to obtain a license from Clontech. For license information, please contact a licensing representative by phone at 650.919.7320 or by e-mail at [licensing@clontech.com](mailto:licensing@clontech.com).

Clontech, the Clontech logo and all other trademarks are the property of Clontech Laboratories, Inc., unless noted otherwise. Clontech is a Takara Bio Company. ©2010 Clontech Laboratories, Inc.