

# National Radiology Data Registry

---

QCDR Report, September 2016

Public Facility  
(Facility ID: 100853)



[nrdr.acr.org](http://nrdr.acr.org)



Breast Cancer Surveillance Consortium



Working together to advance  
breast cancer research

***The National Mammography Database (NMD) thanks the National Cancer Institute funded Breast Cancer Surveillance Consortium (BCSC) for providing benchmark data for comparison with measures for participating facilities.***

## ***NMD QCDR Measure Definitions***

<b>Measure</b>	<b>Definition</b>
All exams	Number of screening exams performed
ACRad 3: Cancers and CDR per 1000	Number of screening mammograms with a BI-RADS assessment category of 0, 4 or 5, total and per 1000, that had a tissue diagnosis of cancer within 12 months.
ACRad 4: Invasive CDR per 1000	Number of screening mammograms with a BI-RADS assessment category of 0, 4 or 5, total and per 1000, that had a tissue diagnosis of invasive cancer within 12 months. (Invasive cancer rate x CDR)
ACRad 5: Recall rate	Number and percent of screening exams with a BI-RADS assessment category of 0, 3,4 or 5.
ACRad 6: Cancers and PPV2	Number of screening exams and percent of screening exams where biopsy was recommended that result in a tissue diagnosis of cancer within 12 months.
Invasive cancer	Number and percent of cancers that were invasive
ACRad 7: Nodal status - negative	Number and percent of exams with nodal status indicated where status was negative
ACRad 8: Minimal cancer	Number and percent of cancers that were DCIS or had a tumor size of 10mm or less
PQRS 146: Inappropriate use of BI-RADS	Number and percent of screening exams with a BI-RADS assessment category of 3.

**Section 1: Outcome measures and demographic data for exams with at least 365 days follow-up (Includes uploaded follow up data through September 30, 2016)**

*(Facilities with no exams with at least 365 days follow-up receive only NMD aggregate and benchmark data in this section. Preliminary data for the facility are available in Section 2.)*

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**January 2015 - December 2015: Comparison to all NMD facilities and BCSC benchmarks**  
**Follow up available through September 30, 2016**

Measure	Facility 100853			All NMD facilities (N=274)			BCSC benchmark*		
	Rate	(Num-Den)		Rate	(Num-Den)		Rate	(Num-Den)	
All exams	.	.	10,214	.	.	2,359,219	.	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	. 11.02%	(1,126 /10,214 )		. 10.00%	(235,908 /2,359,219)		. 10.95%	(263,905 /2,410,932)	
ACRad 6: Cancers and PPV2	. 28.75%	(23 /80 )		. 21.30%	(7,868 /36,946 )		. 22.90%	(7,613 /33,239 )	
ACRad 3: Cancers and CDR per 1000	. 2.45	(25 /10,214 )		. 4.16	(9,805 /2,359,219)		. 4.18	(10,072 /2,410,932)	
ACRad 4: Invasive cancers and ICDR per 1000	. 2.45	(25 /10,214 )		. 3.13	(7,396 /2,359,219)		.		
Invasive cancer rate (Used to calculate ICDR)	. 100.00%	(25 /25 )		. 75.48%	(7,396 /9,798 )		. 76.42%	(7,697 /10,072 )	
ACRad 8: Minimal cancer rate	. .%	( / )		. 72.62%	(3,830 /5,274 )		. 52.40%	(4,980 /9,504 )	
ACRad 7: Nodal status negative rate	. 0.00%	(0 /21 )		. 87.60%	(3,023 /3,451 )		. 75.95%	(5,629 /7,411 )	
PQRS 146: Inappropriate use of BI-RADS 3	. 0.29%	(30 /10,214 )		. 0.29%	(6,775 /2,359,100)		.		

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006.

\*BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1013909175: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1013909175			Facility 100853			All NMD facilities (N=274)			BCSC benchmark*		
	Rate	(Num-Den)		Rate	(Num-Den)		Rate	(Num-Den)		Rate	(Num-Den)	
All exams	.	.	146	.	.	10,214	.	.	2,359,219	.	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	. 16.44%	(24	/146 )	. 11.02%	(1,126	/10,214 )	. 10.00%	(235,908	/2,359,219)	. 10.95%	(263,905	/2,410,932)
ACRad 6: Cancers and PPV2	. 0.00%	(0	/2 )	. 28.75%	(23	/80 )	. 21.30%	(7,868	/36,946 )	. 22.90%	(7,613	/33,239 )
ACRad 3: Cancers and CDR per 1000	. 0.00	(0	/146 )	. 2.45	(25	/10,214 )	. 4.16	(9,805	/2,359,219)	. 4.18	(10,072	/2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	. 0.00	(0	/146 )	. 2.45	(25	/10,214 )	. 3.13	(7,396	/2,359,219)	.		
Invasive cancer rate (Used to calculate ICDR)	. .%	(	/ )	. 100.00%	(25	/25 )	. 75.48%	(7,396	/9,798 )	. 76.42%	(7,697	/10,072 )
ACRad 8: Minimal cancer rate	. .%	(	/ )	. .%	(	/ )	. 72.62%	(3,830	/5,274 )	. 52.40%	(4,980	/9,504 )
ACRad 7: Nodal status negative rate	. .%	(	/ )	. 0.00%	(0	/21 )	. 87.60%	(3,023	/3,451 )	. 75.95%	(5,629	/7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	. 0.00%	(0	/146 )	. 0.29%	(30	/10,214 )	. 0.29%	(6,775	/2,359,100)	.		

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1104033950: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1104033950		Facility 100853		All NMD facilities (N=274)		BCSC benchmark*	
	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)
All exams	.	2,050	.	10,214	.	2,359,219	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	14.20%	(291 /2,050 )	11.02%	(1,126 /10,214 )	10.00%	(235,908 /2,359,219)	10.95%	(263,905 /2,410,932)
ACRad 6: Cancers and PPV2	13.33%	(2 /15 )	28.75%	(23 /80 )	21.30%	(7,868 /36,946 )	22.90%	(7,613 /33,239 )
ACRad 3: Cancers and CDR per 1000	0.98	(2 /2,050 )	2.45	(25 /10,214 )	4.16	(9,805 /2,359,219)	4.18	(10,072 /2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	0.98	(2 /2,050 )	2.45	(25 /10,214 )	3.13	(7,396 /2,359,219)	.	.
Invasive cancer rate (Used to calculate ICDR)	100.00%	(2 /2 )	100.00%	(25 /25 )	75.48%	(7,396 /9,798 )	76.42%	(7,697 /10,072 )
ACRad 8: Minimal cancer rate	.%	( / )	.%	( / )	72.62%	(3,830 /5,274 )	52.40%	(4,980 /9,504 )
ACRad 7: Nodal status negative rate	0.00%	(0 /2 )	0.00%	(0 /21 )	87.60%	(3,023 /3,451 )	75.95%	(5,629 /7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	0.54%	(11 /2,050 )	0.29%	(30 /10,214 )	0.29%	(6,775 /2,359,100)	.	.

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1235112152: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1235112152		Facility 100853		All NMD facilities (N=274)		BCSC benchmark*	
	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)
All exams	.	2,164	.	10,214	.	2,359,219	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	9.43%	(204 /2,164 )	11.02%	(1,126 /10,214 )	10.00%	(235,908 /2,359,219)	10.95%	(263,905 /2,410,932)
ACRad 6: Cancers and PPV2	35.29%	(6 /17 )	28.75%	(23 /80 )	21.30%	(7,868 /36,946 )	22.90%	(7,613 /33,239 )
ACRad 3: Cancers and CDR per 1000	3.23	(7 /2,164 )	2.45	(25 /10,214 )	4.16	(9,805 /2,359,219)	4.18	(10,072 /2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	3.23	(7 /2,164 )	2.45	(25 /10,214 )	3.13	(7,396 /2,359,219)	.	.
Invasive cancer rate (Used to calculate ICDR)	100.00%	(7 /7 )	100.00%	(25 /25 )	75.48%	(7,396 /9,798 )	76.42%	(7,697 /10,072 )
ACRad 8: Minimal cancer rate	0.00%	(0 /6 )	0.00%	(0 /21 )	87.60%	(3,023 /3,451 )	75.95%	(5,629 /7,411 )
ACRad 7: Nodal status negative rate	0.00%	(0 /6 )	0.00%	(0 /21 )	87.60%	(3,023 /3,451 )	75.95%	(5,629 /7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	0.23%	(5 /2,164 )	0.29%	(30 /10,214 )	0.29%	(6,775 /2,359,100)	.	.

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.



**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1619948387: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1619948387			Facility 100853			All NMD facilities (N=274)			BCSC benchmark*		
	Rate	(Num-Den)		Rate	(Num-Den)		Rate	(Num-Den)		Rate	(Num-Den)	
All exams	.	.	1,245	.	.	10,214	.	.	2,359,219	.	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	12.05%	(150	/1,245 )	11.02%	(1,126	/10,214 )	10.00%	(235,908	/2,359,219)	10.95%	(263,905	/2,410,932)
ACRad 6: Cancers and PPV2	0.00%	(0	/13 )	28.75%	(23	/80 )	21.30%	(7,868	/36,946 )	22.90%	(7,613	/33,239 )
ACRad 3: Cancers and CDR per 1000	0.00	(0	/1,245 )	2.45	(25	/10,214 )	4.16	(9,805	/2,359,219)	4.18	(10,072	/2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	0.00	(0	/1,245 )	2.45	(25	/10,214 )	3.13	(7,396	/2,359,219)	.	.	.
Invasive cancer rate (Used to calculate ICDR)	.	.	( / )	100.00%	(25	/25 )	75.48%	(7,396	/9,798 )	76.42%	(7,697	/10,072 )
ACRad 8: Minimal cancer rate	.	.	( / )	.	.	( / )	72.62%	(3,830	/5,274 )	52.40%	(4,980	/9,504 )
ACRad 7: Nodal status negative rate	.	.	( / )	0.00%	(0	/21 )	87.60%	(3,023	/3,451 )	75.95%	(5,629	/7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	0.32%	(4	/1,245 )	0.29%	(30	/10,214 )	0.29%	(6,775	/2,359,100)	.	.	.

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1841243599: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1841243599		Facility 100853		All NMD facilities (N=274)		BCSC benchmark*	
	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)
All exams	.	2,178	.	10,214	.	2,359,219	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	7.94%	(173 /2,178 )	11.02%	(1,126 /10,214 )	10.00%	(235,908 /2,359,219)	10.95%	(263,905 /2,410,932)
ACRad 6: Cancers and PPV2	55.00%	(11 /20 )	28.75%	(23 /80 )	21.30%	(7,868 /36,946 )	22.90%	(7,613 /33,239 )
ACRad 3: Cancers and CDR per 1000	5.05	(11 /2,178 )	2.45	(25 /10,214 )	4.16	(9,805 /2,359,219)	4.18	(10,072 /2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	5.05	(11 /2,178 )	2.45	(25 /10,214 )	3.13	(7,396 /2,359,219)	.	.
Invasive cancer rate (Used to calculate ICDR)	100.00%	(11 /11 )	100.00%	(25 /25 )	75.48%	(7,396 /9,798 )	76.42%	(7,697 /10,072 )
ACRad 8: Minimal cancer rate	.	( / )	.	( / )	72.62%	(3,830 /5,274 )	52.40%	(4,980 /9,504 )
ACRad 7: Nodal status negative rate	0.00%	(0 /9 )	0.00%	(0 /21 )	87.60%	(3,023 /3,451 )	75.95%	(5,629 /7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	0.09%	(2 /2,178 )	0.29%	(30 /10,214 )	0.29%	(6,775 /2,359,100)	.	.

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1912950429: January 2015 - December 2015**  
**Follow up available through September 30, 2016**

Measure	1912950429		Facility 100853		All NMD facilities (N=274)		BCSC benchmark*	
	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)	Rate	(Num-Den)
All exams	.	2,431	.	10,214	.	2,359,219	.	2,410,932
ACRad 5: Abnormal interpretation and recall rate	11.68%	(284 /2,431 )	11.02%	(1,126 /10,214 )	10.00%	(235,908 /2,359,219)	10.95%	(263,905 /2,410,932)
ACRad 6: Cancers and PPV2	30.77%	(4 /13 )	28.75%	(23 /80 )	21.30%	(7,868 /36,946 )	22.90%	(7,613 /33,239 )
ACRad 3: Cancers and CDR per 1000	2.06	(5 /2,431 )	2.45	(25 /10,214 )	4.16	(9,805 /2,359,219)	4.18	(10,072 /2,410,932)
ACRad 4: Invasive cancers and ICDR per 1000	2.06	(5 /2,431 )	2.45	(25 /10,214 )	3.13	(7,396 /2,359,219)	.	.
Invasive cancer rate (Used to calculate ICDR)	100.00%	(5 /5 )	100.00%	(25 /25 )	75.48%	(7,396 /9,798 )	76.42%	(7,697 /10,072 )
ACRad 8: Minimal cancer rate	.%	( / )	.%	( / )	72.62%	(3,830 /5,274 )	52.40%	(4,980 /9,504 )
ACRad 7: Nodal status negative rate	0.00%	(0 /4 )	0.00%	(0 /21 )	87.60%	(3,023 /3,451 )	75.95%	(5,629 /7,411 )
PQRS 146: Inappropriate use of BI-RADS 3	0.33%	(8 /2,431 )	0.29%	(30 /10,214 )	0.29%	(6,775 /2,359,100)	.	.

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below

<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

NCI-funded Breast Cancer Surveillance Consortium (BCSC) data on screening mammograms performed 2002-2006. BCSC benchmark data are calculated using screen-detected cancers; NMD data may include some interval cancers.

***Section 2: Preliminary data on exams performed in the most recent 12 months.  
(Includes uploaded data through September 30, 2016)***

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**January 2016 - September 2016: Comparison to all NMD facilities and BCSC benchmarks**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1013909175: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1104033950: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1235112152: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>



**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1619948387: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1841243599: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

**PRELIMINARY OUTCOME DATA; FINAL DATA WILL BE AVAILABLE 12 MONTHS AFTER EXAM DATE**  
**QCDR Measures: Abnormal Interpretations and Cancers for Screening Mammography**  
**Physician 1912950429: January, 2016 - September 2016**

Measure	All NMD facilities (N=274)	
	Rate	(Num-Den)
All exams	.	1,225,921
ACRad 5: Abnormal interpretation and recall rate	10.00%	(122,547 /1,225,921)
ACRad 6: Cancers and PPV2	20.73%	(3,661 /17,658 )
ACRad 3: Cancers and CDR per 1000	3.87	(4,745 /1,225,921)
ACRad 4: Invasive cancers and ICDR per 1000	3.00	(3,673 /1,225,921)
Invasive cancer rate (Used to calculate ICDR)	77.42%	(3,673 /4,744 )
ACRad 8: Minimal cancer rate	74.82%	(1,780 /2,379 )
ACRad 7: Nodal status negative rate	85.01%	(1,293 /1,521 )
PQRS 146: Inappropriate use of BI-RADS 3	0.25%	(3,012 /1,219,812)

To learn more about the Qualified Clinical Data Registry (QCDR) option please click below  
<http://www.acr.org/Quality-Safety/National-Radiology-Data-Registry/Qualified-Clinical-Data-Registry>

Promote your participation in NRDR™!  
Download your marketing toolkit today.

---

[acr.org/NMDtoolkit](http://acr.org/NMDtoolkit)

**CONTACT US**

1.800.227.5463, ext. 3535  
[nrd@acr.org](mailto:nrd@acr.org) | [nrd.acr.org](http://nrd.acr.org)

