

Fifteen Things Physicians and Patients Should Question

1

Don't do an inherited thrombophilia evaluation for women with histories of pregnancy loss, intrauterine growth restriction (IUGR), preeclampsia and abruption.

Scientific data supporting a causal association between either methylenetetrahydrofolate reductase (MTHFR) polymorphisms or other common inherited thrombophilias and adverse pregnancy outcomes, such as recurrent pregnancy loss, severe preeclampsia and IUGR, are lacking. Specific testing for antiphospholipid antibodies, when clinically indicated, should be limited to lupus anticoagulant, anticardiolipin antibodies and beta 2 glycoprotein antibodies.

2

Don't place a cerclage in women with short cervix who are pregnant with twins.

Women with a short cervical length who are pregnant with twins are at very high risk for delivering preterm, but the scientific data, including a meta-analysis of data published on this issue, shows that cerclage in this clinical situation not only is not beneficial, but may in fact be harmful, i.e., associated with an increase in preterm births.

3

Don't offer noninvasive prenatal testing (NIPT) to low-risk patients or make irreversible decisions based on the results of this screening test.

NIPT has only been adequately evaluated in singleton pregnancies at high risk for chromosomal abnormalities (maternal age >35, positive screening, sonographic findings suggestive of aneuploidy, translocation carrier at increased risk for trisomy 13, 18 or 21, or prior pregnancy with a trisomy 13, 18 or 21). Its utility in low-risk pregnancies remains unclear. False positive and false negative results occur with NIPT, particularly for trisomy 13 and 18. Any positive NIPT result should be confirmed with invasive diagnostic testing prior to a termination of pregnancy. If NIPT is performed, adequate pretest counseling must be provided to explain the benefits and limitations.

4

Don't screen for intrauterine growth restriction (IUGR) with Doppler blood flow studies.

Studies that have attempted to screen pregnancies for the subsequent occurrence of IUGR have produced inconsistent results. Furthermore, no standards have been established for the optimal definition of an abnormal test, best gestational age for the performance of the test or the technique for its performance. However, once the diagnosis of IUGR is suspected, the use of antenatal fetal surveillance, including umbilical artery Doppler flow studies, is beneficial.

5

Don't use progestogens for preterm birth prevention in uncomplicated multifetal gestations.

The use of progestogens has not been shown to reduce the incidence of preterm birth in women with uncomplicated multifetal gestations.

Fifteen Things Physicians and Patients Should Question

6

Don't perform routine cervical length screening for preterm birth risk assessment in asymptomatic women before 16 weeks of gestation or beyond 24 weeks of gestation.

The predictive ability of cervical length measurement prior to 16 weeks of gestation for preterm birth risk assessment is limited. It should be performed, when indicated, between 16 and 24 weeks of gestation. Routine cervical length screening for preterm birth risk assessment in asymptomatic women beyond 24 weeks of gestation has not been proven to be effective.

7

Don't perform antenatal testing on women with the diagnosis of gestational diabetes who are well controlled by diet alone and without other indications for testing.

Monitoring of glucose levels and maintaining adequate glycemic control for gestational diabetes are paramount to decreasing adverse outcomes, including stillbirth. If nutritional modification and glucose monitoring alone control maternal glycemic status such that pharmacological therapy is not required, the risk of stillbirth due to uteroplacental insufficiency is not increased. Thus, the use of routine antepartum testing (e.g. biophysical profile (BPP) or nonstress test (NST)) in the absence of other co-morbidities is not indicated.

8

Don't place women, even those at high-risk, on activity restriction to prevent preterm birth.

There are no studies documenting an improvement in outcomes in women at risk for preterm birth who are placed on activity restriction, including bed rest. There are multiple studies documenting untoward effects of routine activity restriction on the mother and family, including negative psychosocial effects. Therefore, activity restriction should not be routinely prescribed as a treatment to reduce preterm birth.

9

Don't order serum aneuploidy screening after cfDNA aneuploidy screening has already been performed.

Serum biochemistry and cell free DNA (cfDNA) are both screening tests for fetal aneuploidy. When low-risk results have been reported on either test, there is limited clinical value of also performing the other screen. While serum screening may identify some aneuploidies not detected by cfDNA, the yield is too low to justify this test if cfDNA screening has already been performed.

10

Don't perform maternal serologic studies for cytomegalovirus and toxoplasma as part of routine prenatal laboratory studies.

Routine serologic screening of pregnant women for CMV and toxoplasmosis is not recommended due to poor predictive value of these tests and potential for harm due to false positive results. Serologic screening during pregnancy for both diseases should be reserved for situations in which there is clinical or ultrasound suspicion of maternal or fetal infection.

Fifteen Things Physicians and Patients Should Question

11

Don't recommend diagnostic testing following sonographic identification of an isolated echogenic intracardiac focus (EIF) or choroid plexus cyst (CPC) in women with low-risk aneuploidy screening results.

The concept of using ultrasonographic soft markers for aneuploidy, such as EIF and CPC, was introduced in an era that predated screening for Down syndrome based on factors other than maternal age. Because the sensitivity of cell free (cfDNA) screening for Down syndrome approaches 99%, the residual risk for Down syndrome is very low in patients who have a negative cfDNA screening test result. Given the low a priori risk, the presence of an isolated EIF or CPC is unlikely to increase the detection rate for aneuploidy to any measurable degree. In addition, for a woman with an isolated EIF or CPC on a second-trimester ultrasound in the setting of any negative first- or second-trimester aneuploidy screening test result, a reasonable approach is to consider the presence of the isolated finding as a normal variant. Recent guidelines from the Society for Maternal-Fetal Medicine state that diagnostic testing should not be recommended to patients solely for the indication of an isolated EIF or CPC in the setting of a negative cfDNA screening test result or a negative first- or second-trimester screening test result.

12

Don't perform serial cervical length measurement following cerclage placement.

Although progressive cervical shortening after cerclage placement increases the risk of preterm birth, neither overall cervical length nor the length below the stitch correlates well with outcomes. Most importantly, there are currently no additional treatment options for a short cervix after cerclage (e.g., reinforcement suture does not improve outcomes). Although there may be theoretical psychological benefits to the patient and provider to visualize the stitch, there are insufficient data to suggest a clinical benefit of routine post-cerclage serial cervical length measurement.

13

Don't test women for MTHFR mutations.

MTHFR is responsible for the conversion of 5,10-methylenetetrahydrofolate to 5-methyltetrahydrofolate. Genetic variant C677T and A1286C have been associated with a mild decrease in enzymatic activity, which in the setting of reduced folate levels has been found to be a risk factor for hyperhomocysteinemia. Although hyperhomocysteinemia is a risk factor for cardiovascular disease and venous thrombosis, its cause is multifactorial and independent of the MTHFR genotype, even in homozygotic individuals. Despite earlier (mostly case control) studies that found an association between the MTHFR genotype and adverse outcomes, recent studies of more robust design have not replicated these findings. Due to the lack of evidence associating genotype independently with thrombosis, recurrent pregnancy loss, or other adverse pregnancy outcomes, MTHFR genotyping should not be ordered as part of a workup for thrombophilia.

14

Don't screen asymptomatic pregnant women for subclinical hypothyroidism.

Subclinical hypothyroidism (SCH) is defined as an elevated serum TSH level in the presence of a normal free T4 level and is found in 2% to 5% of otherwise healthy pregnant women. SCH is unlikely to progress to overt hypothyroidism during pregnancy. While some authorities and organizations have recommended routine screening for all pregnant women and subsequent treatment with levothyroxine, two recent, large (>100,000 women) prospective randomized clinical trials of screening and treatment for SCH demonstrated no effect of treatment on offspring IQ at age 5 years. Because treatment for SCH has not resulted in a beneficial effect on outcomes, routine screening for SCH is not currently recommended. Targeted screening for women at risk for overt hypothyroidism is still appropriate.

15

Don't use amniotic fluid index to make a diagnosis of oligohydramnios (in the third trimester).

Amniotic fluid volume can be measured using either the amniotic fluid index (AFI) or the deepest vertical pocket (DVP). Diagnosis of oligohydramnios based on an AFI of <5 cm has been found to lead to a greater number of obstetric interventions without a significant benefit in improving perinatal outcomes when compared to use of a DVP of <2 cm for diagnosis.

How This List Was Created

As a national medical specialty society, the Society for Maternal-Fetal Medicine relies on the input of any number of its committees in the development of various documents. In the case of the items included in this list, the Publications Committee reviewed the literature and evidence from SMFM's published documents for possible topics. For SMFM's first set of five recommendations a sub-group of the Committee initially developed a list of 10 items that the Committee then ranked for the top five with input and suggestions by the Society's Executive Committee. For SMFM's second set of recommendations, the sub-group of the Committee developed a list of 12 items that the Committee then ranked for the top five, again soliciting input and suggestions by the Society's Executive Committee. For SMFM's third set of five recommendations, the sub-group of the Publications Committee developed a list of 10 items that the Committee ranked for the top five, again soliciting input and suggestions by the Society's Executive Committee. The final lists have been reviewed and approved by the Society's Document Review Committee and Executive Committee.

SMFM's disclosure and conflict of interest policy can be found at www.smfm.org.

Sources

- Dizon-Townson D, Miller C, Sibai B, Spong CY, Thom E, Wendel G Jr, Wenstrom K, Samuels P, Cotroneo MA, Moawad A, Sorokin Y, Meis P, Miodovnik M, O'Sullivan MJ, Conway D, Wapner RJ, Gabbe SG; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network (NICHD MFMU). The relationship of the factor V Leiden mutation and pregnancy outcomes for mother and fetus. *Obstet Gynecol*. 2005 Sep;106(3):517–24.

Silver RM, Zhao Y, Spong CY, Sibai B, Wendel G Jr, Wenstrom K, Samuels P, Caritis SN, Sorokin Y, Miodovnik M, O'Sullivan MJ, Conway D, Wapner RJ; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units (NICHD MFMU) Network. Prothrombin gene G20210A mutation and obstetric complications. *Obstet Gynecol*. 2010 Jan;115(1):14–20.

Kupferminc MJ, Eldor A, Steinman N, Many A, Bar-Am A, Jaffa A, Fait G, Lessing JB. Increased frequency of genetic thrombophilia in women with complications of pregnancy. *N Engl J Med*. 1999 Jan;340(1):9–13. [published erratum appears in *N Engl J Med* 1999 Jul 29;341(5):384].
- Durnwald CP, Momirova V, Rouse DJ, Caritis SN, Peaceman AM, Sciscione A, Varner MW, Malone FD, Mercer BM, Thorp JM Jr, Sorokin Y, Carpenter MW, Lo J, Ramin SM, Harper M, Spong CY; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network (NICHD MFMU). Second trimester cervical length and risk of preterm birth in women with twin gestations treated with 17-alpha hydroxyprogesterone caproate. *J Matern Fetal Neonatal Med*. 2010 Dec;23(12):1360–4.

Berghella V, Odibo AO, To MS, Rust OA, Althuisius SM. Cerclage for short cervix on ultrasonography: meta-analysis of trials using individual patient-level data. *Obstet Gynecol*. 2005;106:181–9.
- American College of Obstetricians and Gynecologists Committee on Genetics. Noninvasive prenatal testing for fetal aneuploidy. Committee Opinion No. 545. *Obstet Gynecol*. 2012 Dec;120(6):1532–4.
- Society for Maternal-Fetal Medicine Publications Committee, Berkley E, Chauhan SP, Abuhamad A. Doppler assessment of the fetus with intrauterine growth restriction. *Am J Obstet Gynecol*. 2012 Apr;206(4):300–8.
- Society for Maternal-Fetal Medicine Publications Committee, Berghella V. Progesterone and preterm birth prevention: translating clinical trials data into clinical practice. *Am J Obstet Gynecol* 2012 May;206(5):376–86.

Combs CA, Garite T, Maurel K, Das A, Porto M; Obstetrix Collaborative Research Network. 17-hydroxyprogesterone caproate for twin pregnancy: a double-blind, randomized clinical trial. *Am J Obstet Gynecol*. 2011 Mar;204(3):221.e1–8.

Combs CA, Garite T, Maurel K, Das A, Porto M; Obstetrix Collaboration Research Network. Failure of 17-hydroxyprogesterone to reduce neonatal morbidity or prolong triplet pregnancy: a double-blind, randomized clinical trial. *Am J Obstet Gynecol*. 2010 Sep;203(3):248.e1–9.

Caritis SN, Rouse DJ, Peaceman AM, Sciscione A, Momirova V, Spong CY, Iams JD, Wapner RJ, Varner M, Carpenter M, Lo J, Thorp J, Mercer BM, Sorokin Y, Harper M, Ramin S, Anderson G; Eunice Kennedy Shriver National Institute of Child Health and Human Development Maternal-Fetal Medicine Units Network (NICHD MFMU). Prevention of preterm birth in triplets using 17 alpha-hydroxyprogesterone caproate: a randomized controlled trial. *Obstet Gynecol*. 2009 Feb;113(2 Pt 1):285–92.
- Iams JD, Goldenberg RL, Meis PJ, Mercer BM, Moawad A, Das A, Thom E, McNellis D, Copper RL, Johnson F, Roberts JM. The length of the cervix and the risk of spontaneous premature delivery. National Institute of Child Health and Human Development Maternal Fetal Medicine Unit Network. *N Engl J Med*. 1996 Feb 29;334(9):567-72.

Conoscenti G, Meir YJ, D'Ottavio G, Rustico MA, Pinzano R, Fischer-Tamaro L, Stampalija T, Natale R, Maso G, Mandruzzato G. Does cervical length at 13–15 weeks' gestation predict preterm delivery in an unselected population? *Ultrasound Obstet Gynecol*. 2003 Feb;21(2):128-34.

Ozdemir I, Demirci F, Yucel O, Erkorkmaz U. Ultrasonographic cervical length measurement at 10-14 and 20-24 weeks gestation and the risk of preterm delivery. *Eur J Obstet Gynecol Reprod Biol*. 2007 Feb;130(2):176-9.

Berghella V, Talucci M, Desai A. Does transvaginal sonographic measurement of cervical length before 14 weeks predict preterm delivery in high-risk pregnancies? *Ultrasound Obstet Gynecol*. 2003 Feb;21(2):140-4.
- Rosenstein MG, Cheng YW, Snowden JM, Nicholson JM, Doss AE, Caughey AB. The risk of stillbirth and infant death stratified by gestational age in women with gestational diabetes. *Am J Obstet Gynecol*. 2012;206:309.e1-7.

- 8 Society for Maternal-Fetal Medicine (SMFM), Habebber E, Sciscione A. SMFM Consult Activity Restriction in Pregnancy. *Contemp Ob Gyn*. 2014.
- 9 Society for Maternal-Fetal Medicine (SMFM) Publications Committee. Society for Maternal-Fetal Medicine Consult Series #36: Prenatal aneuploidy screening using cell-free DNA. *Am J Obstet Gynecol*. 2015 Jun;212(6):711-6.
Committee Opinion No. 640: Cell-Free DNA Screening For Fetal Aneuploidy. *Obstet Gynecol*. 2015;126(3):e31-7.
- 10 Society for Maternal-Fetal Medicine (SMFM), Hughes BL, Gyamfi-Bannerman C. Society for Maternal-Fetal Medicine Consult Series #39: Diagnosis and antenatal management of congenital cytomegalovirus (CMV) infection. *Am J Obstet Gynecol*. 2016 (in press).
American College of Obstetricians and Gynecologists. Practice Bulletin #151: Cytomegalovirus, Parvovirus B19, varicella zoster, and toxoplasmosis in pregnancy. *Obstet Gynecol*. 2015 Jun;125(6):1510-25.
- 11 Society for Maternal-Fetal Medicine (SMFM). Norton ME, Biggio JR, Kuller JA, Blackwell SC. The role of ultrasound in women who undergo cell-free DNA screening. *Am J Obstet Gynecol*. 2017 Mar;216(3):B2-B7.
Benn P, Cuckle H, Pergament E. Non-invasive prenatal testing for aneuploidy: current status and future prospects. *Ultrasound Obstet Gynecol* 2013;42:15-33.
- 12 Society for Maternal-Fetal Medicine (SMFM). McIntosh J, Feltovich H, Berghella V, Manuck T. The role of routine cervical length screening in selected high- and low-risk women for preterm birth prevention. *Am J Obstet Gynecol*. 2016 Sep;215(3):B2-7.
Baxter JK, Airoidi J, Berghella V. Short cervical length after history indicated cerclage: Is a reinforcing cerclage beneficial? *Am J Obstet Gynecol* 2005;193:1204-7.
- 13 Inherited thrombophilias in pregnancy. ACOG Practice Bulletin No. 197. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2018;132:e18–34.
Hickey SE, Curry CJ, Toriello HV. ACMG Practice Guideline: lack of evidence for MTHFR polymorphism testing. *Genet Med*. 2013 Feb;15(2):153-6.
Bagliin T, Gray E, Greaves M, et al. Clinical guidelines for testing for heritable thrombophilia. *Br J Haematol* 2010;149:209–220.
- 14 Thyroid disease in pregnancy. Practice Bulletin No. 148. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2015; 125:996–1005.
Lazarus JH, Bestwick JP, Channon S, Paradise R, Maina A, Rees R, et al. Antenatal thyroid screening and childhood cognitive function [published erratum appears in *N Engl J Med* 2012;366:1650]. *N Engl J Med* 2012;366:493–501.
Society for Maternal-Fetal Medicine. Screening for thyroid disease during pregnancy. *Contemporary OB/GYN*; August 2012. <https://www.smfm.org/publications/88-screening-for-thyroid-disease-in-pregnancy>
- 15 Kehl S, Schelkle A, Thomas A, Puhl A, Meqdad K, Tuschy B, et al. Single deepest vertical pocket or amniotic fluid index as evaluation test for predicting adverse pregnancy outcome (SAFE trial): a multicenter, open-label, randomized controlled trial. *Ultrasound Obstet Gynecol*. 2016;47(6):674-9.
Ultrasound in pregnancy. Practice Bulletin No. 175. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2016;128:e241–56.
Antepartum fetal surveillance. Practice Bulletin No. 145. American College of Obstetricians and Gynecologists. *Obstet Gynecol* 2014;124:182–92.

About the ABIM Foundation

The mission of the ABIM Foundation is to advance medical professionalism to improve the health care system. We achieve this by collaborating with physicians and physician leaders, medical trainees, health care delivery systems, payers, policymakers, consumer organizations and patients to foster a shared understanding of professionalism and how they can adopt the tenets of professionalism in practice.



To learn more about the ABIM Foundation, visit www.abimfoundation.org.

About the Society for Maternal-Fetal Medicine

The Society for Maternal-Fetal Medicine (SMFM) is a society of physicians and scientists who are dedicated to the optimization of pregnancy and perinatal outcomes. SMFM was established in 1977 and is the membership organization for obstetricians/gynecologists who have additional formal education and training in maternal-fetal medicine. There are currently about 2,000 active members of SMFM. The Society hosts an annual scientific meeting in which new ideas and research in the area of maternal-fetal medicine are presented. The Society is also an advocate for improving public policy and expanding research funding and opportunities in the area of maternal-fetal medicine.



For more information about SMFM, visit www.smfm.org.