

Research Publications Featuring the MA900 Multi-Application Cell Sorter

Immunology

Levin N, Paria BC, Vale NR, et al. Identification and validation of T-cell receptors targeting RAS hotspot mutations in human cancers for use in cell-based immunotherapy. *Clin Cancer Res.* 2021;27:5084-5095. [PubMed](#)

Denstaedt SJ, Bustamante AC, Newstead MW, et al. Long-term survivors of murine sepsis are predisposed to enhanced LPS-induced lung injury and proinflammatory immune reprogramming. *Am J Physiol Lung Cell Mol Physiol.* 2021;321:L451-L465. [PubMed](#)

Snyder GA, Eliachar S, Connelly MT, et al. Functional characterization of hexacorallia phagocytic cells. *Front Immunol.* 2021;12:662803. [PubMed](#)

Calcagno DM, Zhang C, Toomu A, et al. SiglecF(HI) marks late-stage neutrophils of the infarcted heart: a single-cell transcriptomic analysis of neutrophil diversification. *J Am Heart Assoc.* 2021;10:e019019. [PubMed](#)

Porsche CE, Delproposto JB, Geletka L, O'Rourke R, Lumeng CN. Obesity results in adipose tissue T cell exhaustion. *JCI Insight.* 2021;6:e139793. [PubMed](#)

Takeuchi Y, Ohara D, Watanabe H, et al. Dispensable roles of Gsdmd and Ripk3 in sustaining IL-1 β production and chronic inflammation in Th17-mediated autoimmune arthritis. *Sci Rep.* 2021;11:18679. [PubMed](#)

Chen H-Y, Alamonte-Loya A, Lay F-Y, et al. Epigenetic remodeling by vitamin C potentiates the differentiation of mouse and human plasma cells. *bioRxiv.* [DOI](#)

Kim S, Vale N, Zacharakis N, et al. Adoptive cell therapy targeting common p53 neoantigens in human solid cancers. *Research Square.* [DOI](#)

Porsche CE, Delproposto JB, Patrick E, Zamarron BF, Lumeng CN. Adipose tissue dendritic cell signals are required to maintain T cell homeostasis and obesity-induced expansion. *Mol Cell.* 2020;505:110740. [PubMed](#)

Han M, Ishikawa T, Bermick JR, et al. IL-1 β prevents ILC2 expansion, type 2 cytokine secretion, and mucus metaplasia in response to early-life rhinovirus infection in mice. *Allergy.* 2020;75:2005-2019. [PubMed](#)

Fabian KP, Padget MR, Donahue RN, et al. PD-L1 targeting high-affinity NK (t-haNK) cells induce direct antitumor effects and target suppressive MDSC populations. *J Immunother Cancer.* 2020;8:e000450. [PubMed](#)

Sekine T, Perez-Potti A, Nguyen S, et al. TOX is expressed by exhausted and polyfunctional human effector memory CD8+ T cells. *Sci Immunol.* 2020;5:eaba7918. [PubMed](#)

Ma H, Song X, Zhang W, et al. Formaldehyde inhibits development of T lymphocytes in mice. *Toxicological & Environmental Chemistry.* 2020;102:473-489. DOI: 10.1080/02772248.2020.1815202 [DOI](#)

Fergusson JR, Wallace Z, Connolly MM, et al. Immune-mobilising monoclonal T cell receptors mediate specific and rapid elimination of hepatitis B-infected cells. *Hepatology.* 2020;72:1528-1540. DOI: 10.1002/hep.31503 [PubMed](#)

Gaballa A, Arruda LCM, Rådestad E, Uhlin M. CD8+ $\gamma\delta$ T cells are more frequent in CMV seropositive bone marrow grafts and display phenotype of an adaptive immune response. *Stem Cells Int.* 2019;6348060. [PubMed](#)

Rao A, Strauss O, Kokkinou E, et al. Cytokines regulate the antigen-presenting characteristics of human circulating and tissue-resident intestinal ILCs. *Nat Commun.* 2020;11:2049. [PubMed](#)

Yamauchi T, Hoki T, Oba T, et al. CX3CR1-CD8+ T cells are critical in antitumor efficacy but functionally suppressed in the tumor microenvironment. *JCI Insight.* 2020;5:e133920. [PubMed](#)

Quiniou V, Barennes P, Martina F, et al. Human thymopoiesis selects unconventional CD8+ α/β T cells that respond to multiple viruses. *bioRxiv.* DOI: 10.1101/2020.07.27.223354 [DOI](#)

Cell Biology

Kobayashi T, Castillo-Venzor A, Penfold CA, et al. Tracing the emergence of primordial germ cells from bilaminar disc rabbit embryos and pluripotent stem cells. *Cell Rep.* 2021;37:109812. [PubMed](#)

Schoutrop E, El-Serafi I, Poiret T, et al. Mesothelin-specific CAR T cells target ovarian cancer. *Cancer Res.* 2021;81:3022-3035. [PubMed](#)

Troutman TD, Bennett H, Sakai M, Seidman JS, Heinz S, Glass CK. Purification of mouse hepatic non-parenchymal cells or nuclei for use in ChIP-seq and other next-generation sequencing approaches. *STAR Protoc.* 2021;2:100363. [PubMed](#)

Ogata K, Liu Y, Ohara A, et al. Club cells are the primary target for permethrin-induced mouse lung tumor formation. *Toxicol Sci.* 2021;184:15-32. [PubMed](#)

Alvarez-Benayas J, Trasanidis N, Katsarou A, et al. Chromatin-based, in cis and in trans regulatory rewiring underpins distinct oncogenic transcriptomes in multiple myeloma. *Nat Commun.* 2021;12:5450. [PubMed](#)

Chan JM, Quintanal-Villalonga Á, Gao VR, et al. Signatures of plasticity, metastasis, and immunosuppression in an atlas of human small cell lung cancer. *Cancer Cell*. 2021;39:1479-1496. e18. [PubMed](#)

Holloway EM, Wu JH, Czerwinski M, et al. Differentiation of human intestinal organoids with endogenous vascular endothelial cells. *Dev Cell*. 2020;54:516-528.e7. [PubMed](#)

Gao H, Luo Z, Jin Z, Ji Y, Ying W. Adipose tissue macrophages orchestrate β cell adaptation in obesity through secreting miRNA-containing extracellular vesicles. *bioRxiv*. DOI:10.1101/2020.06.12.148809. [DOI](#)

Pine AR, Cirigliano SM, Nicholson JG, et al. Tumor microenvironment is critical for the maintenance of cellular states found in primary glioblastomas. *Cancer Discov*. 2020;10:964-979. [PubMed](#)

Worthen CA, Cui Y, Orringer JS, Johnson TM, Voorhees JJ, Fisher GJ. CD26 identifies a subpopulation of fibroblasts that produce the majority of collagen during wound healing in human skin. *J Invest Dermatol*. 2020;140:2515-2524.e3. [PubMed](#)

Calcagno DM, Ng Jr RP, Toomu A, et al. Type I interferon responses to ischemic injury begin in the bone marrow of mice and humans and depend on Tet2, Nrf2, and Irf3. *bioRxiv*. DOI: 10.1101/765404. [DOI](#)

Microbiology

Lal PB, Wells FM, Lyu Y, Ghosh IN, Landick R, Kiley PJ. A markerless method for genome engineering in *Zymomonas mobilis* ZM4. *Front Microbiol*. 2019;10:2216. [PubMed](#)

Infectious Disease

Scheid JF, Barnes CO, Eraslan B, et al. B cell genomics behind cross-neutralization of SARS-CoV-2 variants and SARS-CoV. *Cell*. 2021;184:3205-3221.e24. [PubMed](#)

Kruglova N, Siniavin A, Gushchin V, Mazurov D. Different neutralization sensitivity of SARS-CoV-2 cell-to-cell and cell-free modes of infection to convalescent sera. *Viruses*. 2021;13:1133. [PubMed](#)

Madan B, Reddem ER, Wang P, et al. Antibody screening at reduced pH enables preferential selection of potentially neutralizing antibodies targeting SARS-CoV-2. *AIChE J*. 2021;67:e17440. [DOI](#)

Sokal A, Chappert P, Barba-Spaeth G, et al. Maturation and persistence of the anti-SARS-CoV-2 memory B cell response. *Cell*. 2021;184:1201-1213.e14. [PubMed](#)

Witkowski M, Tizian C, Ferreira-Gomes M, et al. Untimely TGF β responses in COVID-19 limit antiviral functions of NK cells. *Nature*. [PubMed](#)

Ferreira-Gomes M, Kruglov A, Durek P, et al. SARS-CoV-2 in severe COVID-19 induces a TGF- β -dominated chronic immune response that does not target itself. *Nat Commun*. 2021;12:1961. [PubMed](#)

Zhang Y, Guo R, Kim SH, et al. SARS-CoV-2 hijacks folate and one-carbon metabolism for viral replication. *Nat Commun*. 2021;12:1676. [PubMed](#)

Learn more at: www.sonybiotechnology.com/ma900