



3PEAT™ TRIPLE ACTIVE COMBINATION ORAL DRENCH FOR SHEEP

ACTIVE CONSTITUENTS: 34 g/L Levamisole (equivalent to 40 g/L Levamisole hydrochloride), 25 g/L Albendazole, 1 g/L Abamectin, 0.21 g/L Cobalt (as Cobalt Sulphate Heptahydrate), 0.76 g/L Iodine (as EDDI stabilised), 0.5 g/L Selenium (as Sodium Selenate) and 0.6 g/L Zinc (as Zinc Sulphate Monohydrate)

Powerful Triple Active! The Sheep Drench that Works.

For the treatment and control of major gastrointestinal parasites in sheep, including strains with single or dual resistance to benzimidazole, levamisole/morantel and avermectin/milbemycin families, susceptible to the combination of Abamectin, Levamisole and Albendazole.

- ✿ Potent Triple actives formulation
- ✿ Low volume and stable formulation with no mixing
- ✿ Broad spectrum - Kills common susceptible parasites in sheep, effective against single resistant parasites and susceptibility to Abamectin, Levamisole or Benzimidazole
- ✿ Treats & controls:
 - Internal parasites from eggs through to mature adults
 - Lungworm, itch mite, larval stages of nasal bot
- ✿ Cost effective choice
- ✿ Supplemental Iodine, Cobalt, Selenium and Zinc to help keep your mob healthy, reproductive & producing quality wool
- ✿ Available in 1L Backpack, 5L Backpack and 20L Drum



Supplemental
Cobalt, Iodine,
Selenium, Zinc
for a healthy
mob!



	3 PEAT™ TRIPLE ACTIVE COMBINATION ORAL DRENCH FOR SHEEP	PYRIMIDE®** 3-WAY COMBINATION DRENCH FOR SHEEP	®TRIGUARD*** TRIPLE COMBINATION DRENCH FOR SHEEP
Active Constituents	40g/L Levamisole hydrochloride, 25g/L Albendazole, 1g/L Abamectin, with 0.5g/L selenium, 0.21g/L cobalt, 0.6g/L zinc and 0.76g/L Iodine	0.8g/L Abamectin, 25.5g/L levamisole (as levamisole hydrochloride), 20g/L albendazole, 0.4g/L selenium (as sodium selenate), 1.7g/L cobalt (as cobalt EDTA)	ABAMECTIN 1.0g/L, OXFENDAZOLE 22.7g/L, LEVAMISOLE 33.9g/L (as levamisole hydrochloride), SELENIUM 0.5g/L (as sodium selenate), COBALT 2.2g/L (as cobalt disodium EDTA)
Species	Sheep	Sheep	Sheep
Dose rate	1mL / 5kg body weight	1mL / 4kg body weight	1mL / 5kg body weight
Application	Oral Drench	Oral Drench	Oral Drench
Withholding Period			
Meat	14 days	14 days	14 Days
Milk	DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption	DO NOT USE in sheep which are producing or may in the future produce milk where the milk or milk products may be used for human consumption	DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption
ESI	28 days	42 days	28 days

3PEAT™ is a trademark of Abbey Laboratories Pty Ltd

**PYRIMIDE® is a registered trademark of Pharm Smart Australia Pty Ltd, Ashgrove, Australia

***TRIGUARD is a registered trademark of Merial Limited. ©2017 MERIAL Limited. All rights reserved. Merial is now part of Boehringer Ingelheim. TRIG-17100

3PEAT™ TRIPLE ACTIVE COMBINATION ORAL DRENCH FOR SHEEP



Powerful Triple Active! The Sheep Drench that Works. For the treatment and control of major gastrointestinal parasites in sheep

INDICATIONS

For the effective control of mature and immature stages of susceptible strains of the following parasites -

Gastrointestinal Parasite:

Barber's pole worm *Haemonchus contortus* (Including inhibited L4 stage)

Small brown stomach worm *Teladorsagia circumcincta* (Including inhibited L4 stage)

Stomach hair worm *Trichostrongylus axei*

Black scour worm *Trichostrongylus colubriformis*, *Trichostrongylus vitrinus*

Thin necked intestinal worm *Nematodirus spathiger*, *Nematodirus filicollis*

Small intestinal hair worm *Cooperia curticei*, *Cooperia oncophora*

Intestinal threadworm *Strongyloides papillosus*

Large mouthed bowel worm *Chabertia ovina*

Nodule worm *Oesophagostomum columbianum*

Whipworm *Trichuris ovis*

Nasal bot *Oestrus ovis* (parasitic larval stages only)

Large lungworm *Dictyocaulus filaria*

Itchmite *Psorergates ovis*

Resistance may develop to any chemical.

Ask your local Veterinary practitioner or Animal Health Advisor for recommended management practices. Check the effectiveness of anthelmintic at least annually and it is advised that a drench resistance test be conducted before any drench is used.

DIRECTIONS FOR USE

Restraints

MILK: DO NOT USE in ewes which are producing or may in the future produce milk that may be used or processed for human consumption.

Contraindications

This product is contraindicated for use in lambs under 6 weeks of age or less than 15 kg body weight.

This product should not be used when selenium intake from pasture is high or selenium is provided by other means (such as top dressing, vaccine, pellets or selenium drenches) if blood selenium levels at treatment are high. Users can determine selenium status by consulting their veterinarian.

This product is contraindicated for use unless cobalt deficiency is confirmed.

This product is contraindicated for use in sheep that are dehydrated or unwell.

This product is contraindicated for use unless mineral deficiency is confirmed.

Dogs, horses or any other pets should not be drenched with this product.

This product is not recommended for use in goats as safety and efficacy have not been evaluated.

Precautions

Recommended dose should not be exceeded. Minerals such as cobalt, selenium, iron, manganese and molybdenum compete with each other for metabolically important processes in the animal. Users should seek veterinary advice before treatment.

Fatal interactions may occur between levamisole and organophosphate dips.

Exercise care in handling weak, pregnant and young animals to avoid unnecessary stress. Avoid yarding animals off-feed overnight and ensure animals have access to water when yarded prior to drenching.

Do not treat sheep at intervals of less than 3 weeks, due to selenium content.

DOSAGE AND ADMINISTRATION

SHAKE WELL BEFORE USE.

For oral administration only.

Dose rate is 1ml per 5 kg bodyweight equivalent to 8 mg Levamisole hydrochloride, 5 mg Albendazole, 0.2 mg abamectin per kg bodyweight.

Body weight (kg)	Dose (mL)	1L Backpack	5L Backpack	20L Drum
15	3	333	1667	6667
16-20	4	250	1250	5000
21-25	5	200	1000	4000
26-30	6	167	833	3333
31-35	7	143	714	2857
36-40	8	125	625	2500
41-45	9	111	556	2222
46-50	10	100	500	2000
51-55	11	91	455	1818
56-60	12	83	417	1667
61-65	13	77	385	1538
66-70	14	71	357	1429
71-75	15	67	333	1333
Over 75kg add 1mL for every 5kg bodyweight				

A representative sample of animals should be weighed before treatment. Dose the mob to the heaviest animal by live weight in each group (ewes, wethers, rams, lambs); (bucks, does, kids). Where there is a large variation in size within the group, dose rate should be based on the label directions for each weight range. Do not underdose. Drafting into two or more lines may be appropriate, to avoid excessive overdosing.

DO NOT RE-TREAT ANIMALS FOR 28 DAYS AFTER LAST TREATMENT.

NOT TO BE USED FOR ANY PURPOSE, OR IN ANY

MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.

WITHHOLDING PERIODS

MEAT: DO NOT USE less than 14 days before slaughter for human consumption.

TRADE ADVICE-EXPORT SLAUGHTER INTERVAL (ESI)

DO NOT USE less than 28 days before slaughter for export. Before using this product confirm the current ESI from Abbey Animal Health Pty Ltd on 02 8088 0720 or the APVMA website www.apvma.gov.au/residues.

SAFETY DIRECTIONS

Harmful if swallowed. Will irritate the eyes and skin. Repeated exposure may cause allergic disorders. Avoid contact with eyes and skin. When opening the container and using the product wear cotton overalls buttoned to the neck and wrist (or equivalent clothing) and elbow-length chemical resistant gloves and face shield or goggles. If product in eyes, wash it out immediately with water. Wash hands after use. After each day's use, wash gloves, face shield or goggles and contaminated clothing.

FIRST AID

If poisoning occurs, contact a doctor or Poisons Information Centre. Phone Australia 131126. If in eyes, hold eyes open, flood with water for at least 15 minutes and see a doctor.

Additional information is listed in Safety Data Sheet.

ENVIRONMENT PROTECTION

Abamectin is extremely toxic to aquatic species. Do not contaminate dams, rivers, streams or other waterways with the chemical or used container.

DISPOSAL

Triple-rinse container into the medicated drench. Do not dispose of undiluted chemicals on-site. If recycling, replace cap and return clean container to recycler or designated collection point. If not recycling, break, crush, or puncture container and deliver to an approved waste management facility. If an approved waste management facility is not available, bury the broken, crushed or punctured containers 500 mm below the surface in a disposal pit specifically marked and set up for this purpose clear of waterways, vegetation and tree roots, in compliance with relevant local, state or territory government regulations. Do not burn empty containers or product.

STORAGE

Store below 25°C (air conditioning). Do Not Freeze. Protect from light.

APVMA Approval Number: 93487 / 139321