

FreeStyle
Libre

112  mg/dL



DIACAST
#02

Consequências da variabilidade glicêmica na pessoa com diabetes tipo 2 de alto risco cardiovascular

Com Dr. Fernando Valente e Dra. Nara Alves Buriti

FreeStyle Libre apresenta sua nova dose de conhecimento de CGM

Prevenir o desenvolvimento de complicações crônicas e agudas estão entre os objetivos do tratamento da pessoa com diabetes.

Por isso, convidamos o **Dr. Fernando Valente** e a **Dra. Nara Alves Buriti** para discutirem o papel do FreeStyle Libre na redução da hemoglobina glicada e complicações cardiovasculares em pessoas com diabetes tipo 2.


Ouçá agora!

Agora é fácil saber os padrões de glicose ocultos dos seus pacientes.

Agora Você Sabe



Escute agora o
DIACAST #02



Abbott

life. to the fullest.™

Zoungas S. et al. Severe Hypoglycemia and Risks of Vascular Events and Death. New England Journal of Medicine. 2010;363(15):1410-1418. Amiel S et al. Hypoglycaemia, cardiovascular disease, and mortality in diabetes: epidemiology, pathogenesis, and management. The Lancet Diabetes; Endocrinology. 2019;7(5):385-396. Zinman B. et al. Day-to-day fasting glycaemic variability in DEVOTE: associations with severe hypoglycaemia and cardiovascular outcomes (DEVOTE 2). Diabetologia. 2017;61(1):48-57. Miller E, Brandner L, Wright E. HbA1c reduction after initiation of the FreeStyle Libre system in type 2 diabetes patients on long-acting insulin or non-insulin therapy [84-LB]. Poster presented at: 80th Scientific Sessions of the American Diabetes Association: June 12-16, 2020; Virtual. Imagens meramente Ilustrativas. Divulgação Diacast Volume 2_ ADC-57456_V1. Leitor FreeStyle Libre - RMS ANVISA: 80146501903. ANATEL: 4072-14-9992. Sensor FreeStyle Libre - RMS ANVISA: 80146502021. FreeStyle LibreLink - RMS ANVISA: 80146502168. FreeStyle, Libre, e marcas relacionadas são marcas da Abbott. Abbott FreeStyle: 0800 703 0128.