



**COLUMBIA UNIVERSITY
CENTER FOR SHOULDER,
ELBOW AND SPORTS
MEDICINE**

Christopher S. Ahmad, MD
Office (212) 305-5561
Fax (212) 305-4040

Louis U. Bigliani, MD
Office (212) 305-5564
Fax (212) 305-0999

Edwin Cadet, MD
Office (212) 305-4626
Fax (212) 305-4040

William N. Levine, MD
Office (212) 305-0762
Fax (212) 305-4040

Charles A. Popkin, MD
Office (646) 317-4656
Fax (212) 305-8271

Appointment Scheduling
(212) 305-4565

Mailing Address:
622 West 168th Street, PH-11
New York, NY 10032

Office Locations:
51 West 51st Street, Ste 370
New York, NY 10019

161 Ft. Washington Ave.
New York, NY 10032

500 Grand Avenue
Englewood, NJ 07631

www.nyp.org/columbiaortho

**OATS PROCEDURE
PHYSICAL THERAPY PROTOCOL**

POST-OPERATIVE PHASE I: Weeks 0-6

PRIMARY GOALS: protect healing tissue, decrease pain and effusion, gradually improve knee flexion, restore passive knee extension, regain quad control

BRACE: locked at 0 degrees for WB activities, sleep in locked brace for 4 weeks

WEIGHT-BEARING: NWB (if graft site is MFC), WBAT (if graft site is patella or trochlea)

ROM: full passive knee extension immediately
initiate CPM POD 1 for a total of 6 hours per day x2-3 weeks
progress CPM ROM as tolerated 5-10 degrees per day, up to 6w
Knee flexion ROM goal: 90 degrees by 1-2w
Knee flexion ROM goal: 105 degrees by 3-4w, 120 degrees by 5-6w
patellar mobilization
stretch hamstrings and calf

STRENGTHENING: ankle pump using rubber tubing, quad sets, multi-angle isometrics, active knee extension (no resistance), SLR, stationary bike when ROM allows, biofeedback and electrical muscle stimulation, prn; isometric leg presses by week 4, may begin use of pool for gait training and exercises by week 4.

POST-OPERATIVE PHASE II: Weeks 6-12

PRIMARY GOALS: gradually increase ROM, improve quad strength/endurance, increase in functional activities

BRACE: D/C by week

WEIGHTBEARING: progress WBAT, FWB by 8-9 weeks, D/C crutches by 9 weeks (if graft site is MFC)

ROM: maintain full passive knee extension, progress to flexion to 125-135 degrees by week 8, continue stretching program as well as patellar mobilization

STRENGTHENING: initiate weight shifts week 6, mini-squats 0-45 degrees by week 8, closed kinetic chain (leg press), toe calf raises by week 8, stationary bicycle (low resistance), treadmill walking by weeks 10-12, balance and proprioception drills, initiate front and lateral step-ups and wall squats by weeks 8-10, continue use of biofeedback and electrical stimulation prn, use of pool for gait training and exercise

POST-OPERATIVE PHASE III: Weeks 12-26

ROM: 0-125 to 135 degrees

Incorporate home exercise program for strengthening and maintenance 3-4 times per week. Progress resistance as tolerated, agility and balance drills, sports programs depending on patient.

Gradual return to full unrestricted functional activities by 6 months and beyond.