



References for Herbal Mechanisms in Insulin Resistance

- Abdel-Barry JA, Abdel-Hassan IA, Jawad AM, al-Hakim MH. Hypoglycaemic effect of aqueous extract of the leaves of *Trigonella foenum-graecum* in healthy volunteers. *East Mediterr Health J* 2000 Jan;6(1):83-8
- Achrekar S, Kaklij GS, Pote MS, Kelkar SM. Hypoglycemic activity of *Eugenia jambolana* and *Ficus bengalensis*: mechanism of action. *In Vivo* 1991 Mar-Apr;5(2):143-7
- Agrawal P, Rai V, Singh RB. Randomized placebo-controlled, single blind trial of holy basil leaves in patients with noninsulin-dependent diabetes mellitus. *Int J Clin Pharmacol Ther* 1996 Sep;34(9):406-9
- Aguiyi JC, Obi CI, Gang SS, Igweh AC. Hypoglycaemic activity of *Ocimum gratissimum* in rats. *Fitoterapia* 2000 Aug;71(4):444-6
- Ahmad N, Hassan MR, Halder H, Bennoor KS. Effect of *Momordica charantia* (Karolla) extracts on fasting and postprandial serum glucose levels in NIDDM patients. *Bangladesh Med Res Counc Bull* 1999 Apr;25(1):11-3
- Bagdade JD, Bierman EL, Porter D. The significance of basal insulin levels in evaluation of the insulin response in diabetic and non-diabetic subjects. 1967. *J Clin Invest*(NY) 46(10), 1553.
- Baskaran K, Kizar Ahamath B, Radha Shanmugasundaram K, Shanmugasundaram ER. Antidiabetic effect of a leaf extract from *Gymnema sylvestre* in non-insulin-dependent diabetes mellitus patients. *J Ethnopharmacol* 1990 Oct;30(3):295-300
- Berrio LF, Polansky MM, Anderson RA. Insulin activity: stimulatory effects of cinnamon and brewer's yeast as influenced by albumin. *Horm Res* 1992;37(6):225-9
- Broadhurst CL, Polansky MM, Anderson RA. Insulin-like biological activity of culinary and medicinal plant aqueous extracts in vitro. *J Agric Food Chem* 2000 Mar;48(3):849-52



- Chattopadhyay RR. Hypoglycemic effect of Ocimum sanctum leaf extract in normal and streptozotocin diabetic rats. Indian J Exp Biol 1993 Nov;31(11):891-3
- Cheng JT, Liu IM, Chi TC, Su HC, Chang CG. Metformin-like effects of Quei Fu Di Huang Wan, a Chinese herbal mixture, on streptozotocin-induced diabetic rat. Horm Metab Res 2001 Dec;33(12):727-32
- Chung SH, Choi CG, Park SH. Comparisons between white ginseng radix and rootlet for antidiabetic activity and mechanism in KKAY mice. Arch Pharm Res 2001 Jun;24(3):214-8
- Cornfield J. The University Group Diabetes Program. A further statistical analysis of the mortality findings. JAMA. 1971 Sep 20 ;21 7(12):1676-87.
- Crellin JK and Philpott J. A Reference Guide to Medicinal Plants: Herbal Medicine Past and Present. Durham, North Carolina: Duke University Press, 1989
- El Gammel SY Antidiabetic herbs in history, in: Adly A. [Ed] The History of Medicinal Plants: Proceedings of the Second International Congress Organized by the Arab Society for the History of Pharmacy in co-operation with the World Union of Societies of the History of Pharmacy. Alexandria Egypt, December 15th to 19th 1980. Karachi, Hamdard Foundation Press, 1980
- Graham DJ, Drinkard CR, Shatin D. Incidence of idiopathic acute liver failure and hospitalized liver injury in patients treated with troglitazone. Am J Gastroenterol 2003 Jan;98(1):175-9
- Grover JK, Yadav S, Vats V. Medicinal plants of India with anti-diabetic potential. J Ethnopharmacol 2002 Jun;81(1):81-100
- Grover JK, Vats V, Rathi SS, Dawar R. Traditional Indian anti-diabetic plants attenuate progression of renal damage in streptozotocin induced diabetic mice. J Ethnopharmacol 2001 Aug;76(3):233-8
- Grover JK, Vats V, Rathi SS. Anti-hyperglycemic effect of Eugenia jambolana and Tinospora cordifolia in experimental diabetes and their effects on key metabolic enzymes involved in carbohydrate metabolism. J Ethnopharmacol 2000 Dec;73(3):461-70



- Gupta A, Gupta R, Lal B. Effect of *Trigonella foenum-graecum* (fenugreek) seeds on glycaemic control and insulin resistance in type 2 diabetes mellitus: a double blind placebo controlled study. *J Assoc Physicians India* 2001 Nov;49:1057-61 Said HM [Editor] *Hamdard Pharmacopoeia of Eastern Medicine*. Delhi; Sri Satguru Publications, 1970.
- Horio H, Ohtsuru M. Maitake (*Grifola frondosa*) improve glucose tolerance of experimental diabetic rats. *J Nutr Sci Vitaminol (Tokyo)* 2001 Feb;47(1):57-63
- Imparl-Radosevich J, Deas S, Polansky MM, Baedke DA, Ingebritsen TS, Anderson RA, Graves DJ. Regulation of PTP-1 and insulin receptor kinase by fractions from cinnamon: implications for cinnamon regulation of insulin signalling. *Horm Res* 1998 Sep;50(3):177-82
- Jarvill-Taylor KJ, Anderson RA, Graves DJ. A hydroxychalcone derived from cinnamon functions as a mimetic for insulin in 3T3-L1 adipocytes. *J Am Coll Nutr* 2001 Aug;20(4):327-36 Justice, J.W. Use of Devil's Club in Southeast Alaska. *Alaska Medicine* 8(2) 36-39
- Kar A, Choudhary BK, Bandyopadhyay NG. Comparative evaluation of hypoglycaemic activity of some Indian medicinal plants in alloxan diabetic rats. *J Ethnopharmacol* 2003 Jan;84(1):105-8
- Khan A, Bryden NA, Polansky MM, Anderson RA. Insulin potentiating factor and chromium content of selected foods and spices. *Biol Trace Elem Res* 1990 Mar;24(3):183-8
- Khan A, Safdar M, Ali Khan MM, Khattak KN, Anderson Cinnamon improves glucose and lipids of people with type 2 diabetes. *Diabetes Care*. 2003 Dec; 26(12): 3215-8.
- Kimura M, Waki I, Chujo T, Kikuchi T, Effects of hypoglycemic components in ginseng radix on blood insulin level in alloxan diabetic mice and on insulin release from perfused rat pancreas. *J Pharmacobiodyn* 1981 Jun;4(6):410-7
- Konno S, Tortorelis DG, Fullerton SA, Samadi AA, Hettiarachchi J, Tazaki H. A possible hypoglycaemic effect of maitake mushroom on Type 2 diabetic patients. *Diabet Med* 2001 Dec;18(12):1010
- Kubo K, Aoki H, Nanba H. Anti-diabetic activity present in the fruit body of *Grifola frondosa* (Maitake). *Biol Pharm Bull* 1994 Aug;17(8):1106-10



- Manohar V, Talpur NA, Echard BW, Lieberman S, Preuss HG. Effects of a water-soluble extract of maitake mushroom on circulating glucose/insulin concentrations in KK mice. *Diabetes Obes Metab* 2002 Jan;4(1):43-8
- Miura T, Itoh C, Iwamoto N, Kato M, Kawai M, Park SR, Suzuki I. Hypoglycemic activity of the fruit of the *Momordica charantia* in type 2 diabetic mice. *J Nutr Sci Vitaminol (Tokyo)* 2001 Oct;47(5):340-4
- Moore, M. *Medicinal Plants of the Desert and Canyon West*. Santa Fe: Museum of New Mexico Press, 1989
- Nadkarni KM *Indian Materia Medica*. Bombay: Popular Prakashan Perez-Gutierrez RM, Perez- Gonzalez C, Zavala-Sanchez MA, Perez-Gutierrez S. Hypoglycemic activity of *Bouvardia terniflora*, *Brickellia veronicaefolia*, and *Parmentiera edulis*. *Salud Publica Mex* 1998 Jul- Aug;40(4):354-8
- Pepato MT, Folgado VB, Kettelhut IC, Brunetti IL. Lack of antidiabetic effect of a *Eugenia jambolana* leaf decoction on rat streptozotocin diabetes. *Braz J Med Biol Res* 2001 Mar;34(3):389-95
- Persaud SJ, Al-Majed H, Raman A, Jones PM. *Gymnema sylvestre* stimulates insulin release in vitro by increased membrane permeability. *J Endocrinol* 1999 Nov;163(2):207-12
- Petlevski R, Hadzija M, Slijepcevic M, Juretic D. Effect of 'antidiabetis' herbal preparation on serum glucose and fructosamine in NOD mice. *J Ethnopharmacol* 2001 May;75(2-3):181-4
- Rai V, Iyer U, Mani UV. Effect of Tulasi (*Ocimum sanctum*) leaf powder supplementation on blood sugar levels, serum lipids and tissue lipids in diabetic rats. *Plant Foods Hum Nutr* 1997;50(1):9-16
- Rao BK, Rao CH. Hypoglycemic and antihyperglycemic activity of *Syzygium alternifolium* (Wt.) Walp. seed extracts in normal and diabetic rats. *Phytomedicine* 2001 Mar;8(2):88-93
- Rathi SS, Grover JK, Vikrant V, Biswas NR. Prevention of experimental diabetic cataract by Indian Ayurvedic plant extracts. *Phytother Res* 2002 Dec;16(8):774-7



- Savickiene N, Dagilyte A, Lukosius A, Zitkevicius V. Importance of biologically active components and plants in the prevention of complications of diabetes mellitus. *Medicina (Kaunas)* 2002;38(10):970-5
- Schor S. The University Group Diabetes Program. A statistician looks at the mortality results.
- JAMA. 1971 Sep 20;217(12):1671-5.
- Sowmya P, Rajyalakshmi P. Hypcholesterolemic effect of germinated fenugreek seeds in human subjects. *Plant Foods Hum Nutr* 1999;53(4):359-65
- Sharma RD, Raghuram TC, Rao NS. Effect of fenugreek seeds on blood glucose and serum lipids in type I diabetes. *Eur J Clin Nutr* 1990 Apr;44(4):301-6
- Sharma RD, Sarkar A, Hazra DK, Mishra B, et al. Use of fenugreek seed powder in the management of non-insulin dependent diabetes mellitus. *Nut. Res.* 16, 1331-1339
- Shanmugasundaram ER, Rajeswari G, Baskaran K, Rajesh Kumar BR, Radha Shanmugasundaram K, Kizar Ahmath B. Use of *Gymnema sylvestre* leaf extract in the control of blood glucose in insulin-dependent diabetes mellitus. *J Ethnopharmacol* 1990 Oct;30(3):281- 94
- Shimizu K, Ozeki M, Tanaka K, Itoh K, Nakajyo S, Urakawa N, Atsuchi M. Suppression of glucose absorption by extracts from the leaves of *Gymnema inodorum*. *J Vet Med Sci* 1997 Sep;59(9):753-7
- Sotaniemi EA, Haapakoski E, Rautio A. Ginseng therapy in non-insulin-dependent diabetic patients. *Diabetes Care* 1995 Oct;18(10):1373-5
- Talpur NA, Echard BW, Fan AY, Jaffari O, Bagchi D, Preuss HG. Antihypertensive and metabolic effects of whole Maitake mushroom powder and its fractions in two rat strains. *Mol Cell Biochem* 2002 Aug;237(1-2):129-36
- Thie, K. *Medicinal Plants of the Pacific Northwest: A Digest of Anthropological Writings about Native American Uses White Salmon*, Washington: Longevity Herb Press, 1999
- Thommassen HV, Wilson RA, McIlwain RG. Effect of devil's club on blood glucose levels in diabetes mellitus. *Canadian Family Physician* (1990) 36



- Teixeira CC, Rava CA, Mallman da Silva P, Melchior R, Argenta R, Anselmi F, Almeida CR, Fuchs FD. Absence of antihyperglycemic effect of jambolan in experimental and clinical models. *J Ethnopharmacol* 2000 Jul; 71(1-2): 343-7
- Teixeira CC, Pinto LP, Kessler FH, Knijnik L, Pinto CP, Gastaldo GJ, Fuchs FD. The effect of *Syzygium cumini* (L.) skeels on post-prandial blood glucose levels in non-diabetic rats and rats with streptozotocin-induced diabetes mellitus. *J Ethnopharmacol* 1997 May;56(3):209-13
- Tominaga M, Kimura M, Sugiyama K, Abe T, Igarashi K, Igarashi M, Eguchi H, Sekikawa A, Ogawa A, Manaka H, et al. Effects of seishin-renshi-in and *Gymnema sylvestre* on insulin resistance in streptozotocin-induced diabetic rats. *Diabetes Res Clin Pract* 1995 Jul;29(1):11-7
- University Group Diabetes Program Study pertaining to phenformin. *JAMA*. 1971 Aug 9;217(6):817.
- Vats V, Grover JK, Rathi SS. Evaluation of anti-hyperglycemic and hypoglycemic effect of *Trigonella foenum-graecum* Linn, *Ocimum sanctum* Linn and *Pterocarpus marsupium* Linn in normal and alloxanized diabetic rats. *J Ethnopharmacol* 2002 Jan;79(1):95-100
- Vikrant V, Grover JK, Tandon N, Rathi SS, Gupta N. Treatment with extracts of *Momordica charantia* and *Eugenia jambolana* prevents hyperglycemia and hyperinsulinemia in fructose fed rats. *J Ethnopharmacol* 2001 Jul;76(2):139-43
- Viseshakul D, Premvatana P, Chularojmontri V, Kewsiri D, Tinnarat P. Improved glucose tolerance induced by long term dietary supplementation with hairy basal seeds (*Ocimum canum sim*) in diabetics. *J Med Assoc Thai* 1985 Aug;68(8):408-11
- Vuksan V, Sievenpiper JL, Wong J, Xu Z, et al. American ginseng (*Panax quinquefolius* L.) attenuates postprandial glycemia in a time-dependent but not dose-dependent manner in healthy individuals. *Am J Clin Nutr* 2001 Apr;73(4):753-8
- Vuksan V, Sievenpiper JL, Koo VY, Francis T American ginseng (*Panax quinquefolius* L) reduces postprandial glycemia in nondiabetic subjects and subjects with type 2 diabetes mellitus. *Arch Intern Med* 2000(b) Apr 10;160(7):1009-13
- Vuksan V, Stavro MP, Sievenpiper JL, Beljan-Zdravkovic U, et al. Similar postprandial glycemic reductions with escalation of dose and administration time of American ginseng in type 2 diabetes. *Diabetes Care* 2000 (b) Sep;23(9):1221-6



- Waki I, Kyo H, Yasuda M, Kimura M. Effects of a hypoglycemic component of ginseng radix on insulin biosynthesis in normal and diabetic animals. J Pharmacobiodyn 1982 Aug;5(8):547-54
- Weiss, RF. Weiss's Herbal Medicine: Classic Edition. Stuttgart:Theime, 2001
- Yoshikawa M, Murakami T, Kadoya M, Li Y, Murakami N, Yamahara J, Matsuda H. Medicinal foodstuffs. IX. The inhibitors of glucose absorption from the leaves of *Gymnema sylvestre* R. BR. (Asclepiadaceae): structures of gymnemosides a and b. Chem Pharm Bull (Tokyo) 1997 Oct;45(10):1671-6