



**INE** | INSTITUTE OF  
NUTRITIONAL  
ENDOCRINOLOGY

## Micronutrients: Vitamin B12

Dr. Ritamarie Loscalzo

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

**Medical Disclaimer:** The information in this presentation is not intended to replace a one-on-one relationship with a qualified health care professional and is not intended as medical advice. It is intended as a sharing of knowledge and information from the research and experience of Dr. Ritamarie Loscalzo, drritamarie.com, and the experts who have contributed. We encourage you to make your own health care decisions based upon your research and in partnership with a qualified health care professional. This presentation is provided for informational purposes only and no guarantees, promises, representations or warranties of any kind regarding specific or general benefits, have been or will be made by Dr. Ritamarie Loscalzo, her affiliates or their officers, principals, representatives, agents or employees. Dr. Ritamarie Loscalzo is not responsible for, and shall have no liability for any success or failure, acts and/or omissions, the appropriateness of the participant's decisions, or the use of or reliance on this information.

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Vitamin B12 General Info

- ✓ Water-soluble vitamin
- ✓ The largest and most complex chemical structure of all the vitamins
- ✓ Contains a metal ion – cobalt – thus cobalamin
- ✓ Group of molecules called corrinoids
- ✓ Methylcobalamin and 5-deoxyadenosylcobalamin are the forms of vitamin B12 used in the human body
- ✓ Cofactor for two enzymes
  - Methionine synthase
  - L-methylmalonyl-coenzyme A mutase



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

### Vitamin B12 Structure

**Core is Corrin Ring**

- 4 reduced pyrrole rings
- Cobalt at center
- Nucleotide 5,6-dimethylbenzimidazole
- One of the following
  - CN – cyanide (cyanocobalamin)
  - OH – hydroxyl (hydroxycobalamin)
  - H<sub>2</sub>O – water (aquocobalamin)
  - NO<sub>2</sub> – nitric acid (nitritocobalamin)
  - 5'-deoxyadenosyl (adenosylcobalamin)
  - -CH<sub>3</sub> – methyl (methylcobalamin)

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Vitamin B12 Structure Up-close

One of the following:

- CN – cyanide (cyanocobalamin)
- OH – hydroxyl (hydroxycobalamin)
- H<sub>2</sub>O – water (aquocobalamin)
- NO<sub>2</sub> – nitric acid (Nitritocobalamin)
- 5'-deoxyadenosyl (adenosylcobalamin)
- -CH<sub>3</sub> – methyl (methylcobalamin)

- Corrin Ring
- Cobalt
- Pyrrole
- 5,6-dimethylbenzimidazole

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Vitamin B12 Structures Compared

Schwerner HA, 2012 \*

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Vitamin B12 Absorption and Storage

✓ **Stomach acid needed for absorption:**

- Bound to proteins and is released from the proteins by the action of a high concentration of hydrochloric acid
- Results in the free form of the vitamin, which is immediately bound to a mixture of glycoproteins secreted by the stomach and salivary glands (R-proteins, aka cobalophilins or haptocorrins)
- R-proteins protect vitamin B12 from chemical denaturation in stomach



✓ **Binds to Intrinsic Factor (IF) in duodenum (IF created by stomach):**

- Pancreatic proteases hydrolyze R-Proteins and free cobalamin is released
- IF binds B12 and carries to ileum for absorption via receptors (cubulins) – mostly distal 1/3
- 1 to 3% of intake can be absorbed by passive diffusion

✓ **Overall absorption from diet: 11 to 65% (average 50%)**

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Vitamin B12 Transport and Storage

- ✓ Appears in blood **3 to 4 hours after ingestion**
- ✓ Transported by **transcobalamin** (TCI, TCII, TCIII)
- ✓ **TCII is main protein that carries newly absorbed B12** to tissues
- ✓ **80% of B12 is carried by TCI**, considered a circulating storage form
- ✓ **All cells have TCII-cobalamin receptors**
- ✓ **2 to 4 mg stored in liver**; small amounts stored in muscle, bone, kidneys, heart, brain and spleen, mostly as **adenosylcobalamin**
- ✓ About **1.4 mcg per day of liver stores are secreted into bile** to provide B12 even in absence of dietary intake



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---


---

---

---

---

### Vitamin B12 Absorption Flowchart



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

---

---



---

---

---


---

---

---

---

---



---

---

---

---

---

---

---

---

### Metabolism of Vitamin B12

- ✓ **Hydroxycobalamin**
  - Cytosolic methylation to methylcobalamin
  - Reduction and reaction with ATP to produce adenosylcobalamin
- ✓ **Methylcobalamin**
  - Active cofactor in methylation pathways
- ✓ **Adenosylcobalamin**
  - Active cofactor for formation of succinyl CoA
- ✓ **Cyanocobalamin**
  - "Expensive" and "messy" conversion to methyl

www.DrRitamarie.com © Dr. Ritamarie Luszczko, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---


### Influences on Vitamin B12 Absorption

**Decreases Absorption:**

- ✓ **Age:** Many people over the age of 50 lose the ability to absorb B12 from foods
- ✓ **Gastrointestinal surgery:** i.e., Weight loss surgery
- ✓ **Digestive disorders:** Celiac disease or Crohn's
- ✓ **H. pylori infection and ulcers**
- ✓ **Pancreatic insufficiency**

**Increases Absorption:**

- ✓ **Adequate stomach acid**
- ✓ **Intrinsic factor**
- ✓ **Large supplemental doses**
- ✓ **Good pancreatic function**



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

### Functions of Vitamin B12

- ✓ **Energy** production
- ✓ Reduces **homocysteine**
- ✓ Helps **protect nerve cells**
- ✓ Supports **replenishment of the myelin sheath** and maintenance of the central nervous system
- ✓ Important for **protein metabolism**
- ✓ Supports the **production and repair** of DNA and RNA
- ✓ **Works with B9/folate:**
  - Maturation of red blood cells
  - Helps iron work more efficiently
  - Produce S-adenosylmethionine (SAME)



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Active Co-Enzyme Vitamin B12 Forms and Functions

- ✓ **Methylcobalamin** (60 - 80% of B12 in blood)
  - Used by the enzyme **methionine synthase** to turn homocysteine (HCV) into methionine
  - Methionine is further converted to the most important methyl donor, S-adenosylmethionine (SAM, aka SAME)
- ✓ **5'-deoxyadenosylcobalamin** (up to 20% of B12 in blood)
  - Used by the enzyme **L-methylmalonyl-CoA mutase** to convert methylmalonyl-CoA to succinyl-CoA (magnesium and biotin required to produce L-methylmalonyl-CoA)
  - Occurs in mitochondria
  - Used by the enzyme **leucine aminomutase** to convert B-leucine into L-leucine and vice-versa
  - B12 deficiency leads to a build-up of methylmalonic acid



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Vitamin B12 and Homocysteine

- ✓ Homocysteine is a **nerve and vessel toxin**
- ✓ Methionine is an essential amino acid
- ✓ Some **methionine is turned into homocysteine**
- ✓ The **body normally converts homocysteine to other molecules**, one of which is back into methionine
- ✓ If this pathway is blocked, homocysteine increases
- ✓ **Methylcobalamin (B12)** is needed by methionine synthase to convert homocysteine into methionine
- ✓ Thus, in B12 deficiency, homocysteine levels increase, and so does risk of cardiovascular and nerve damage



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Homocysteine and Heart Disease

- ✓ Homocysteine is thought to cause **cardiovascular disease** by oxidative stress and vessel wall damage
- ✓ With high homocysteine, **risk of developing coronary artery disease doubles**
- ✓ 2.5 times increased risk of **stroke**
- ✓ Risk factor for cardiovascular disease, blood clotting abnormalities, and atherosclerosis
- ✓ **Related to deficiencies of vitamin B6 and Folate (B9)** in addition to vitamin B12



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---


---

---

---

---

---



---

---

---

---

---

---


---

---


### Vitamin B12 and Anemia

- ✓ **Macrocytic anemia:**  
Mean Corpuscular Volume (MCV) increased
- ✓ **Megaloblastic anemia:**  
Abnormally large red blood cells are observed under a microscope

Normal blood cells



Megaloblastic anemia cells



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---


---

---


### Folate, Vitamin B12 and Megaloblastic Anemia

- ✓ Folate needed to turn **uracil into thymidine**, an essential building block of DNA
- ✓ **DNA needed for new red blood cell production and division**
- ✓ In creating methylcobalamin, a methyl group is donated from folate and folate converts from the **5-MTHF form to the THF form** needed to make DNA
- ✓ Without adequate B12, the methyl groups become "trapped" (known as the methyl-folate trap) and the **THF form of folate is not available for DNA synthesis and repair**
- ✓ Without THF form of folate, thymidine is not produced, and DNA is not available to facilitate the maturation of red blood cells, leading to **megaloblastic anemia**

Normal



Megaloblastic Changes



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---


---

---

---

---

---




---

---

---

---

---

---

---

---

---

---


---

---

### Role of B12 and Folate in Megaloblastic Anemia

**DNA needed for new red blood cell production and division**

- ✓ Without THF form of folate, DNA is not available to facilitate the maturation of red blood cells, leading to **megaloblastic anemia**
- ✓ Without adequate folate, **new red blood cells** (which start out as large cells called reticulocytes) **divide slowly**, as they are dependent on DNA for division
- ✓ Only RNA, not DNA, is **needed to produce the hemoglobin** found in the red blood cells
- ✓ Since **RNA does not require thymidine**, and hemoglobin is only dependent on RNA, it's produced at a normal rate which causes the production of **large red blood cells known as macrocytes**
- ✓ The result is **macrocytic anemia**



normal red cells

macrocytic red cells

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Masking of B12 Deficiency by Folate and Iron

- ✓ If the diet contains large amounts of folate (i.e., lots of uncooked greens), **adequate THF regeneration occurs to produce DNA, even in the absence of B12**
- ✓ **High intake of folate (>800 mcg) can "mask" a B12 deficiency** since megaloblastic anemia is avoided leading to possible nerve damage as deficiency goes unnoticed
- ✓ B12 deficiency can make itself worse because it can **prevent the production of the intestinal cells needed to absorb B12**, which are rapidly dying and need to be replaced using DNA
- ✓ **Iron deficiency leads to small red blood cells** so the MCV looks normal when megaloblastic anemia is present



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### B12 Deficiency Without Megaloblastic Anemia

- ✓ **Megaloblastic, aka macrocytic, anemia** is indicative of a B12 and/or folate deficiency
- ✓ **Neurological disorders due to B12 deficiency** often occur in the absence of a macrocytic anemia (sometimes due to "masking" by adequate folate)



These are faces of vitamin B12 deficiency  
It is commonly misdiagnosed and under treated.  
There are many symptoms and causes.

www.b12deficiency.info

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

### B12 Deficiency Without Megaloblastic Anemia: Cases

✓ Lindenbaum examined 141 cases of neurological problems due to B12 deficiency. 40 (28%) had no macrocytic anemia (iron deficiency may have contributed to a lack in 6 patients, and folate therapy could account for 2 others).


➤ **Labs:**

- Very high serum MMA
- High homocysteine
- Borderline (and sometimes normal) serum B12 levels

➤ **Symptoms:**

- Sensory loss
- Ataxia
- Dementia
- Psychiatric disorders

✓ 2011 study from Korea, 35 patients with vitamin B12 deficiency, most with neurological symptoms; none had anemia.



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---


### Vitamin B12 and Pernicious Anemia

✓ **Cause:** Stomach cells unable to make intrinsic factor

✓ **Symptoms:**

- Weakness
- Pale skin
- Diarrhea
- Weight loss
- Fever
- Numbness or tingling sensation in the hands and feet
- Loss of balance
- Confusion, memory loss, and moodiness

✓ **Solution:** B12 in high doses – ideally injections



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Symptoms of Vitamin B12 Deficiency

✓ Fatigue	✓ Disorientation
✓ Weakness	✓ Memory loss
✓ Shortness of breath	✓ Diarrhea
✓ Skin pallor	✓ Nervousness
✓ Loss of balance	✓ Numbness and tingling in fingers and toes
✓ Palpitations	✓ Anemia and pernicious anemia
✓ Insomnia	✓ Swelling of myelin fibers
✓ Abnormal gait	✓ Possible dementia
✓ Loss of concentration	



**Severe deficiency can cause irreversible nerve damage**

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

### Vitamin B12 Deficiency Risk Factors

- ✓ **Impaired stomach function**
  - H. pylori or ulcers: damage to stomach cells that make intrinsic factor
  - Atrophic gastritis
- ✓ **Intestinal malabsorption**
  - Crohn's disease and other inflammatory bowel diseases
  - Pancreatic disease
  - Bariatric surgery
  - Medications
- ✓ **Specific populations**
  - Vegan and vegetarians
  - The elderly
- ✓ **Specific conditions**
  - Eating disorders
  - HIV
  - Diabetes



www.DrRitamarie.com © Dr. Ritamarie Luscato, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### B12 and Fatigue

- ✓ Most common symptom of B12 deficiency
- ✓ Chronic fatigue syndrome (CFS) and fibromyalgia
  - B12 is nitric oxide scavenger, and high levels of nitric oxide and peroxynitrite linked to fibromyalgia and CFS
  - Benefit from B12 shots



Pall ML. Elevated, sustained peroxynitrite level as the cause of chronic fatigue syndrome. Medical Hypotheses 2000;54:115-125. Pall ML.

www.DrRitamarie.com © Dr. Ritamarie Luscato, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### B12 and Breast Cancer

- ✓ B12 and folate decrease risk of breast cancer
  - Deficiency leads to impaired DNA synthesis and repair
- ✓ **Postmenopausal women**
  - Shown to have lower B12 levels
  - Increased risk of breast cancer
- ✓ **Women with breast cancer**
  - Tend to have lower vitamin B12 levels than women without breast cancer



Wu K, Heitschouer KJ, Comstock GW, Hoffman SC, Nadeau MR, Selhub J. A prospective study on folate, B<sub>12</sub>, and pyridoxal 5'-phosphate (B6) and breast cancer. Cancer Epidemiol Biomarkers Prev. 1998;8(3):209-217.

www.DrRitamarie.com © Dr. Ritamarie Luscato, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### B12 and Male Fertility

- ✓ **B12 supplements**
  - Improve sperm count
  - Improve sperm motility
- ✓ **Infertility**
  - Deficiencies in B12 lead to genetic damage of the sperm cells
- ✓ **Erectile dysfunction**
  - Insufficient B12 can damage the nerves within the penis



"Sandier B, Faragher B. (1984). Treatment of oligospermia with vitamin B12. Infertility, Volume 7:133-8."

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Vitamin B12 and Macular Degeneration

- ✓ **B12 deficiency**
  - Renders nerves, including the optic nerve, more brittle
  - Can cause damage to the optic nerve, resulting in decreased central vision
- ✓ **Vitamin B12 supplementation**
  - 1,000 mcg of vitamin B12 along with 2500 mcg of folic acid and 500 mg of vitamin B6 daily reduced risk of developing loss of vision



Christen WG, Glynn RJ, Chew EY, Albert CM, Manson JE. Folic acid, pyridoxine, and cyanocobalamin combination treatment and age-related macular degeneration in women: The Women's Antioxidant and Folic Acid Cardiovascular Study. Arch Intern Med. 2009;169(4):335-41.

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Impact of Vitamin B12 Excess

- ✓ Well tolerated in general
- ✓ Rarely, excess B12 supplementation can cause some **numbness or tingling in the arm, hands and face**
- ✓ According to the NIH, people with an **hereditary eye disease called Leber's Disease** can have damage to the optic nerve from excess vitamin B12
- ✓ Some studies looked at **increased risk of prostate cancer**
  - They are retrospective
  - Look at blood levels
  - Are without specific ranges
  - Used cyanocobalamin and synthetic folic acid
- ✓ **High-dose supplementation or injections** can cause diarrhea, itching, blood clots, and allergic reactions in some people
- ✓ People who are **allergic to cobalt** should also avoid taking cobalamin



<http://www.drRitamarie.com/so/PMID15499634>

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---



### Food Sources of Vitamin B12

- ✓ Vitamin B12 is found only in animal foods
- ✓ **Microorganisms**, and especially bacteria and fungi, are the only organisms definitively known to produce vitamin B12
- ✓ **Germ-phobia and triple washing of produce** eliminates these organisms
- ✓ Nutritional yeast grown on a molasses medium is an example of a food-based, quasi-supplement that would provide a vegan source of vitamin B12
- ✓ Some reports of B12 in herbs and microalgae – likely B12 analogues not the “real” thing



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---


---

---

---

### Food Forms of Vitamin B12

- ✓ **Hydroxycobalamin:** Poultry, fish, shellfish, eggs, dairy
- ✓ **Adenosylcobalamin:** Poultry, fish, shellfish, eggs
- ✓ **Methylcobalamin:** Dairy
- ✓ **Cyanocobalamin:** Not naturally occurring in food; cyanide in tobacco can cause formation



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Food Sources of Vitamin B12

Food	Serving Size	Cals	Amount (mcg)	DR/DV (%)
Sardines	3.20 oz	188.7	8.11	338
Salmon	4 oz	157.6	5.67	236
Tuna	4 oz	147.4	2.66	111
Cod	4 oz	96.4	2.62	109
Lamb	4 oz	310.4	2.51	105
Scallops	4 oz	125.9	2.44	102
Shrimp	4 oz	134.9	1.88	78
Beef	4 oz	175.0	1.44	60
Yogurt	1 cup	149.4	0.91	38
Cow's milk	4 oz	74.4	0.55	23
Eggs	1 each	77.5	0.55	23
Turkey	4 oz	166.7	0.42	18
Chicken	4 oz	187.1	0.39	16
Cheese	1 oz	114.2	0.24	10
Mushrooms, Crimini	1 cup	15.8	0.07	3

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

---

---

### Herbs Reported to Be High in Vitamin B12



Alfalfa      Dandelion      Hawthorn berries

Hops      Bladderwrack      White oak bark

Reports are unreferenced and anecdotal at best.  
Likely, these are B12 analogues.  
<http://www.drRitamarie.com/go/HerbsEqualTonsOfVitaminsToo>

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---


---

---

---

---

### Assessing Status of Vitamin B12



- ✓ **Clinical Assessment**
  - Signs and symptoms
  - Food journals and diet history
- ✓ **Blood Testing**
  - **Serum B12 test:** not accurate, reflects only current, not active forms
  - **Mean corpuscular volume (MCV)** (functional): good indicator but many interfering factors
  - **Homocysteine:** good indicator of B12, folate, and/or vitamin B6
  - **Methylmalonic acid (MMA), serum:** good indicator but not as good as urine MMA
- ✓ **Urine**
  - **Methylmalonic acid (MMA), urine:** Norman Clinical Labs – gold standard ([www.B12.com](http://www.B12.com))
  - **Schilling test:** multiple stages with supplementation – invasive, requires injection of radioactive B12 and 24-hour collection

Good comparison of testing methods at: <http://www.drRitamarie.com/go/B12TestComparison>

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---


---

---

---

---

### B12 and Methylmalonic Acid



- ✓ **Mechanism:**
  - Adenosylcobalamin: cofactor for conversion of methylmalonyl-CoA to succinyl-CoA
  - When B12 is not available, methylmalonyl-CoA levels increase
  - Methylmalonyl-CoA is then converted to methylmalonic acid (MMA) which then accumulates in the blood and urine
  - Since B12 is the only coenzyme required in this pathway, MMA levels are the best indicators of a B12 deficiency
- ✓ **High MMA levels can also (but rarely) be caused by:**
  - Genetic defects, kidney failure, low blood volume, gut bacteria changes, pregnancy, and thyroid disease
- ✓ **Normal serum MMA: .07 to .27 µmol/l**
  - Patients with serum MMA levels in the range of .76 to 187 µmol/l had neurological problems
- ✓ **Normal urine MMA: 0.0 to 3.5**
  - Test developed by Eric J. Norman, Ph.D., in collaboration with the late M. Drue Denton, M.D., and co-workers at the Hematology Division of the University of Cincinnati College of Medicine, USA

Yamauchi H, Onime M, Tsukamoto N, et al. Urinary methylmalonic acid excretion and clinical features in megaloblastic anemia due to vitamin B12 deficiency. *Jpn J Clin Hematol* 1989; 30:835-839

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### Vitamin B12 Supplement Forms

- ✓ **Cyanocobalamin:**  
Least effective, not easily converted to methyl or adenosyl
- ✓ **Methylcobalamin:**  
Metabolically active
- ✓ **Adenosylcobalamin:**  
Metabolically active
- ✓ **Hydroxycobalamin:**  
Easily converted to methyl and adenosyl



Caution with high dose methylcobalamin and certain genetic SNPs, like COM-T – can lead to methyl trapping

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---

### Vitamin B12 Supplementation: Methods Administration

- ✓ Intramuscular
- ✓ Intranasal
- ✓ Topical patch
- ✓ Topical cream
- ✓ Liposomal
- ✓ Sublingual drops
- ✓ Sublingual lozenges
- ✓ Capsules
- ✓ Tablets
- ✓ Soft gels
- ✓ Lozenges
- ✓ Gum



Take with water after eating. Doses range from 150 mcg up to 15 gm

www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---


---

---

---

### References

- ✓ **Advanced Nutrition and Human Metabolism** – Gropper, Smith and Groff.
- ✓ Scalabrino G. The neurotrophic action of cobalamin (vitamin B12) revisited. <http://www.drritamarie.com/go/PMID11272999>
- ✓ Seetharam B, Li N. Transcobalamin II and its cell surface receptor. <http://www.drritamarie.com/go/PMID10714245>
- ✓ Hackam DG, Peterson JC, Spence JD. **What level of plasma homocysteine should be treated?** Effects of vitamin therapy on progression of carotid atherosclerosis in patients with homocysteine levels above and below 14 micromol/L. <http://www.drritamarie.com/go/PMID10678280>
- ✓ Guyton AC, Hall JE. *Textbook of Medical Physiology*, 9th ed.
- ✓ Murray RK, Granner DK, Mayes PA, Rodwell VW. *Harper's Biochemistry*, 24th ed.
- ✓ **Linus Pauling Institute** - <http://www.drritamarie.com/go/LPIVitaminB12>



www.DrRitamarie.com © Dr. Ritamarie Loscalzo, MS, DC, CCN, DACBN, Institute of Nutritional Endocrinology (INE)

---

---

---

---

---

---

---

---