



INE | INSTITUTE OF
NUTRITIONAL
ENDOCRINOLOGY

Micronutrients: Vitamin B Complex

Dr. Ritamarie Loscalzo



Medical Disclaimer: The information in this presentation is not intended to replace a one-on-one relationship with a qualified health care professional and is not intended as medical advice. It is intended as a sharing of knowledge and information from the research and experience of Dr. Ritamarie Loscalzo, drritamarie.com, and the experts who have contributed. We encourage you to make your own health care decisions based upon your research and in partnership with a qualified health care professional.



B Vitamins At a Glance

Letter	Names	Notes/Actions
B1	Thiamin, Benfotiamine	Energy, heart, muscle, and nerve function
B2	Riboflavin, R 5'-Phosphate	Energy, red blood cells, vision
B3	Niacin, Nicotinic Acid, Niacinamide	Energy, nerve function, circulation and heart
B4	Choline, Adenine, Carnitine	Cell membranes, memory, neuromuscular
B5	Pantothenic Acid	Coenzyme A, adrenals, skin
B6	Pyridoxine, Pyridoxal 5'-Phosphate	Brain and nerve, hormones, protein synthesis
B7	Biotin	Hair, metabolism
B8	Inositol	Loosely considered a B vitamin
B9	Folate, Methylfolate, Folinic Acid	Red blood cell production, DNA repair, brain
B10	Pteroylmonoglutamic Acid	Really a form of folate, skin protector
B11	Salicylic Acid	Not technically a vitamin, loosely categorized
B12	Cobalamin	Red blood cells, DNA repair, nervous system



Vitamin	Name	Structure	Molecular Function
Vitamin B ₁	Thiamine		Thiamine plays a central role in the generation of energy from carbohydrates. It is involved in RNA and DNA production, as well as nerve function. Its active form is a coenzyme called thiamine pyrophosphate (TPP), which takes part in the conversion of pyruvate to acetyl coenzyme A (CoA) in metabolism.
Vitamin B ₂	Riboflavin		Riboflavin is involved in the energy production for the electron transport chain, the citric acid cycle, as well as the catabolism of fatty acids (beta oxidation)
Vitamin B ₃	Niacin		Niacin is composed of two structures: nicotinic acid and nicotinamide. There are two coenzyme forms of niacin: nicotinamide adenine dinucleotide (NAD) and nicotinamide adenine dinucleotide phosphate (NADP). Both play an important role in energy transfer reactions in the metabolism of glucose, fat and alcohol. NAD carries hydrogens and their electrons during metabolic reactions, including the pathway from the citric acid cycle to the electron transport chain. NADP is a coenzyme in lipid and nucleic acid synthesis.
Vitamin B ₅	Pantothenic acid		Pantothenic acid is involved in the oxidation of fatty acids and carbohydrates. Coenzyme A, which can be synthesized from pantothenic acid, is involved in the synthesis of amino acids, fatty acids, ketones, cholesterol, phospholipids, steroid hormones, neurotransmitters (such as acetylcholine), and antibodies.

Chart derived from: <http://www.drritamarie.com/go/WikiBVitamins>







Vitamin	Name	Structure	Molecular Function
Vitamin B ₆	pyridoxine, pyridoxal, pyridoxamine		The active form pyridoxal 5'-phosphate (PLP) (depicted) serves as a cofactor in many enzyme reactions mainly in amino acid metabolism including biosynthesis of neurotransmitters.
Vitamin B ₇	Biotin		Biotin plays a key role in the metabolism of lipids, proteins and carbohydrates. It is a critical co-enzyme of four carboxylases: acetyl CoA carboxylase, which is involved in the synthesis of fatty acids from acetate; propionyl CoA carboxylase, involved in gluconeogenesis; β -methylcrotonyl CoA carboxylase, involved in the metabolism of leucine; and pyruvate CoA carboxylase, which is involved in the metabolism of energy, amino acids and cholesterol.
Vitamin B ₉	Folate		Folate acts as a co-enzyme in the form of tetrahydrofolate (THF), which is involved in the transfer of single-carbon units in the metabolism of nucleic acids and amino acids. THF is involved in pyrimidine nucleotide synthesis, so is needed for normal cell division, especially during pregnancy and infancy, which are times of rapid growth. Folate also aids in erythropoiesis, the production of red blood cells.
Vitamin B ₁₂	Cobalamin		Vitamin B12 is involved in the cellular metabolism of carbohydrates, proteins and lipids. It is essential in the production of blood cells in bone marrow, and for nerve sheaths and proteins. Vitamin B12 functions as a co-enzyme in intermediary metabolism for the methionine synthase reaction with methylcobalamin, and the methylmalonyl CoA mutase reaction with adenosylcobalamin.

Chart derived from: <http://www.drritamarie.com/go/WikiBVitamins>





Vitamin	Name	Deficiency effects – Classic Syndromes
Vitamin B ₁	Thiamine	Deficiency causes beriberi. Symptoms of this disease of the nervous system include weight loss, emotional disturbances, Wernicke's encephalopathy (impaired sensory perception), weakness and pain in the limbs, periods of irregular heartbeat, and edema (swelling of bodily tissues). Heart failure and death may occur in advanced cases. Chronic thiamin deficiency can also cause <u>Korsakoff's</u> syndrome, an irreversible dementia characterized by amnesia and compensatory confabulation.
Vitamin B ₂	Riboflavin	Deficiency causes <u>riboflavinosis</u> . Symptoms may include <u>cheilosis</u> (cracks in the lips), high sensitivity to sunlight, angular <u>cheilitis</u> , glossitis (inflammation of the tongue), seborrheic dermatitis or pseudo-syphilis (particularly affecting <u>the scrotum</u> or labia majora and the mouth), pharyngitis (sore throat), hyperemia, and edema of the pharyngeal and oral mucosa.
Vitamin B ₃	Niacin	Deficiency, along with a deficiency of tryptophan causes pellagra. Symptoms include aggression, dermatitis, insomnia, weakness, mental confusion, and diarrhea. In advanced cases, pellagra may lead to dementia and death (the 3(+1) Ds: dermatitis, diarrhea, dementia, and death).
Vitamin B ₅	Pantothenic acid	Deficiency can result in acne and paresthesia, although it is uncommon.
Vitamin B ₆	pyridoxine, pyridoxal, <u>pyridoxamine</u>	The classic clinical syndrome for vitamin B6 deficiency is a <u>seborrhoeic</u> dermatitis-like eruption, atrophic glossitis with ulceration, angular <u>cheilitis</u> , conjunctivitis, <u>intertrigo</u> , and neurologic symptoms of somnolence, confusion, and neuropathy[16] (due to impaired <u>sphingosin</u> synthesis) and <u>sideroblastic</u> anemia (due to impaired <u>heme</u> synthesis).
Vitamin B ₇	Biotin	Deficiency does not typically cause symptoms in adults but may lead to impaired growth and neurological disorders in infants. Multiple carboxylase deficiency, an inborn error of metabolism, can lead to biotin deficiency even when dietary biotin intake is normal.
Vitamin B ₉	Folate	Deficiency results in a macrocytic anemia, and elevated levels of homocysteine. Deficiency in pregnant women can lead to birth defects. Supplementation is often recommended during pregnancy. Researchers have shown that folic acid might also slow the insidious effects of age on the brain.
Vitamin B ₁₂	Cobalamin	Deficiency results in a macrocytic anemia, elevated homocysteine, peripheral neuropathy, memory loss and other cognitive deficits. It is most likely to occur among elderly people, as absorption through the gut declines with age; the autoimmune disease pernicious anemia is another common cause. It can also cause symptoms of mania and psychosis. In rare extreme cases, paralysis can result.

Vitamin	Name	Tolerable Upper Intake Level	Harmful effects
Vitamin B ₁	Thiamine	None	No known toxicity from oral intake. There are some reports of anaphylaxis caused by high-dose thiamin injections into the vein or muscle. However, the doses were greater than the quantity humans can physically absorb from oral intake.
Vitamin B ₂	Riboflavin	None.	No evidence of toxicity based on limited human and animal studies. The only evidence of adverse effects associated with riboflavin comes from in vitro studies showing the production of reactive oxygen species (free radicals) when riboflavin was exposed to intense visible and UV light.
Vitamin B ₃	Niacin	35 mg/day from supplements, drugs or fortified food	Intake of 3000 mg/day of nicotinamide and 1500 mg/day of nicotinic acid are associated with nausea, vomiting, and signs and symptoms of liver toxicity. Other effects may include glucose intolerance, and (reversible) ocular effects. Additionally, the nicotinic acid form may cause vasodilatory effects, also known as flushing, including redness of the skin, often accompanied by an itching, tingling, or mild burning sensation, which is also often accompanied by pruritus, headaches, and increased intracranial blood flow, and occasionally accompanied by pain. Medical practitioners prescribe recommended doses up to 2000 mg per day of niacin, usually in time release format, to combat arterial plaque development in cases of high lipid levels.
Vitamin B ₅	Pantothenic acid	None	No known toxicity
Vitamin B6	pyridoxine, pyridoxal, pyridoxamine	200mg/day	Primary symptoms are pain and numbness of the extremities. In severe cases, motor neuropathy may occur causing difficulty in walking. Sensory neuropathy typically develops at doses of pyridoxine in excess of 1,000 mg per day, but adverse effects can occur with much less, so doses over 200 mg are not considered safe.[33] Symptoms among women taking lower doses have been reported
Vitamin B ₇	Biotin	None	No known toxicity
Vitamin B ₉	Folate	1 mg/day	Masks B12 deficiency, which can lead to permanent neurological damage.
Vitamin B ₁₂	Cobalamin	None established.	Skin and spinal lesions. Acne-like rash [causality is not conclusively established].



Chart derived from: <http://www.drritamarie.com/go/WikiBVitamins>

Resources

- ✓ *Advanced Nutrition and Human Metabolism – Gropper, Smith and Groff*
- ✓ *Linus Pauling Institute: An Evidence-based Approach to Vitamins and Minerals: Health Benefits and Intake Recommendations.*
- ✓ <http://www.drritamarie.com/go/WikiB Vitamins>

