

Diagnos-Techs, Inc.

Clinical & Research Laboratory
 19110 66th Ave S. Bldg. G, Kent WA 98023
 Tel: (425) 251-0596
 CLIA License # 50D0630141

Reported :

DR RITA MARIE LLC
 Accession #

Age/DOB:56 /

Received : 03
 Completed:

Sex:Female

RITA MARIE LOSCALZ

Results For:

Specimen Collected:

Dx Code: Not Provided

Test	Description	Result	Ref Values
------	-------------	--------	------------

Free Cortisol Rhythm - Saliva

ASI Adrenal Stress Index (Original) - Saliva

TAP

06:00 - 08:00 AM 24 Normal 13-24 nM

11:00 - 1:00 PM 5 Normal 5-10 nM


04:00 - 05:00 PM 4 Normal 3-8 nM

10:00 - Midnight 3 Normal 1-4 nM.
 The cortisol load reflects the area under the cortisol curve. This is an indicator of overall cortisol exposure, where high values favor a catabolic state, and low values are sign of adrenal deterioration. **22 - 46 nM**

Figure 2.


The Cortisol release inducers fall into 4 broad categories shown in the adjacent flowchart. Long term adrenal axis maintenance and restoration, require optimization of all the cortisol inducers.

Figure 1. Circadian Cortisol Profile



The Inducers of Cortisol Release

Inducers below must be individually examined for successful restoration of adrenals.




Test	Accession: Description	Result	Ref Values	C
DHEA	Dehydroepiandrosterone Free Pooled Value	[DHEA + DHEA-S] 2	Depressed Adults (M/F): 3-10 ng/ml	

Figure 3 shows your cortisol-DHEA correlation was in:

↳ Zone 4 - Maladapted phase II

This zone represents a marginal cortisol output with reduced DHEA levels reflecting a limited adrenal response. The utilization of the precursor pregnenolone is usually limited and the adrenal cortex may show hypertrophic changes. Under stress most patients in maladaptation phase II will have a suboptimal response to stress. This suboptimal response is any response not consistent with a normal diurnal cortisol production pattern. This condition is usually the outcome of chronic and protracted stress exposure.



CORTISOL-DHEA CORRELATION SPECTRUM

1. Adapted to stress.
2. Adapted with DHEA slump.
3. Maladapted Phase I.
- 4. Maladapted Phase II.**
5. Non-adapted, Low Reserves
6. High DHEA.
7. Adrenal Fatigue.

ISN	Insulin - Saliva			
	Fasting	<3		Normal: 3-12 uIU/mL
	Non-Fasting	<3	Depressed	Optimal: 5-20 uIU/mL

Depressed Non Fasting insulin within four hours after meal. This may be caused by a small carbohydrate load in the preceding challenge meal or a reduction in pancreatic insulin release or synthesis. Consider a closer examination of challenge meal composition to rule out pre-diabetic tendencies.

Why Test for Insulin?

Insulin activity is affected by the stress and cortisol responses. Chronic stress with cortisol elevation antagonizes insulin, and may cause functional insulin resistance. Furthermore, chronic hypercortisol causes hyperinsulin responses to carbohydrate intake. Chronic insulin resistance and overproduction lead to pancreatic exhaustion.

Basic facts about insulin values.

Fasting: This insulin value is elevated in cases of insulin resistance.

Non Fasting: This insulin value varies with type of meal and time of sample collection. See figure 4b. Adapted, Br. J. Nutr. 2003, 90:853. For an after meal insulin, instruct patient to eat 50g of carbohydrate or what is equivalent to 200 calories about 45-90 minutes before noon sample collection. Examples: 2 slices of white bread and 1 cup of orange juice OR 1 cup of cooked oatmeal and 1 cup of orange juice OR 2 ounces of corn flakes snack.

Figure 4a. Insulin Levels





Figure 4b. Serum Insulin - Time Curve




Shaded area is optimal period of post-prandial collection.

Accession:

Test	Description	Result	Ref Values
P17-OH 17-OH Progesterone	18	Depressed	Adults Optimal: 22-100 pg/ml Borderline: 101-130 pg/ml Elevated: >130 pg/ml

Figure 5. Adrenal Steroid Synthesis Pathway



MB2S	Total Salivary SIgA	<5	Depressed
------	---------------------	----	-----------

A depressed mucosal SIgA may be attributed to one or more of the following reasons:

- 1- Excessive chronic cortisol output causes reduction in SIgA production due to low counts of SIgA immunocytes. Appropriate restorative treatments have been shown to produce incremental improvements in SIgA.
- 2- A short imbalance in sympathetic to parasympathetic activity rapidly inhibits SIgA release from the mucosal immunocytes for several hours.
- 3- Chronic deficits in cortisol and/or DHEA levels.
- 4- Possible systemic deficit in capacity to produce IgA - an inherited problem. Rule out possibility with a serum IgA test. A normal finding rules out this possibility.


Normal: 25-60 mg/dl

Borderline: 20-25 mg/dl

Basic Facts About SIgA

1. Secretory IgA (SIgA) is secreted by the various mucosal surfaces. It is mostly a dimeric molecule. Less than 2% of Saliva is of serum origin. The secretory component of SIgA stabilizes it against enzymatic and bacterial degradation.
2. The main functions of SIgA include Immune Exclusion, Viral and Toxin Neutralization, Plasmid Elimination, and Inhibition of Bacterial Colonization. SIgA immune complexes are not inflammatory to the mucosal surfaces.

Figure 6. Effect of Emotion on SIgA Release
Anger versus Care



Diagnos-Techs, Inc.

Accession:

Test	Description	Result	Ref Values
FI4	Gliadin Ab, SIgA (Saliva) 6	Negative	Borderline: 13-15 U/ml Positive: >15 U/ml Notes on Gliadin Ab Test Gliadins are polypeptides found in wheat, rye, oat, barley, and other grain glutes, and are toxic to the intestinal mucosa in susceptible individuals. Healthy adults and children may have a positive antigliadin test because of subclinical gliadin intolerance. Some of their symptoms include mild enteritis, occasional loose stools, fat intolerance, marginal vitamin and mineral status, fatigue, or accelerated osteoporosis. Scan. J. Gastroenterol. 29:248(1994).

COURTESY INTERPRETATION of test and technical support are available upon request, to Physicians Only