



In association with  
**Mayo Clinic**

## Baseline Test Result

Patient Name: \_\_\_\_\_

Baseline (seconds): \_\_\_\_\_ Baseline Test Date: \_\_\_\_\_ Valid Until: \_\_\_\_\_

### **Next Steps:**

#### **Activate your King-Devick Test Sideline Concussion Screening Account**

1. Go to [www.kingdevicktest.com/for-concussions/purchase](http://www.kingdevicktest.com/for-concussions/purchase)
2. Purchase a King-Devick Test Subscription  
Compatible with internet connected computer and/or standard-sized iPad
3. Login in to your King-Devick Test account and enter baseline information above

## CONCUSSION ACTION PLAN

### 1. **Recognize**

#### a. Failed King-Devick Test

-OR-

#### b. Signs or symptoms of suspected concussion

##### Signs - Observed

- Appears dazed or stunned
- Is confused about assignment or position
- Forgets an instruction
- Is unsure of game, score, or opponent
- Moves clumsily
- Answers questions slowly
- Loses consciousness (even briefly)
- Mood, behavior, or personality changes
- Can't recall events prior to or after hit or fall

##### Symptoms - Reported

- Headache or "pressure" in head
- Nausea or vomiting
- Balance problems or dizziness
- Double or blurry vision
- Sensitivity to light or noise
- Feeling sluggish, hazy, foggy, or groggy
- Concentration or memory problems
- Confusion
- Just not "feeling right" or is "feeling down"

### 2. **Remove From Play**

- a. If you suspect that an athlete has a concussion, remove the athlete from play
- b. Ensure that the athlete is evaluated by a health care professional

## RED FLAGS

If any of the following symptoms are present, call 911 or take the athlete to the nearest emergency room:

- ✓ Worsening headaches
- ✓ Seizures
- ✓ Repeated vomiting
- ✓ Slurred speech
- ✓ Increasing confusion or irritability
- ✓ Can't recognize people or places
- ✓ Neck Pain
- ✓ Decreasing alertness