



**CHOOSE EPCLUSA FOR  
PEOPLE WITH CHRONIC HCV  
WHO INJECT DRUGS**

Actor portrayals.

**YOU ARE INVITED TO ATTEND  
A SPEAKER PROGRAM**

**LYNN FRAZIER, NP  
APRN/Director  
Liver Wellness Center**

**Tuesday, May 9, 2023  
6:30 PM**

**Waterfront Grill  
5201 DeSaird Street  
Monroe, Louisiana  
(318) 345-0064**

**PLEASE RSVP TO  
Suzanne Hebert at (337) 278-0334 or via e-mail at [suzanne.hebert@gilead.com](mailto:suzanne.hebert@gilead.com)  
by 5/6/2023**

SPONSORED BY:



**INDICATION**

EPCLUSA is indicated for the treatment of adults with chronic hepatitis C virus (HCV) genotype 1-6 infection without cirrhosis or with compensated cirrhosis.

**IMPORTANT SAFETY INFORMATION**

**WARNING: RISK OF HEPATITIS B VIRUS REACTIVATION IN PATIENTS COINFECTED WITH HCV AND HBV: Hepatitis B virus (HBV) reactivation has been reported, in some cases resulting in fulminant hepatitis, hepatic failure, and death.**

Please see additional Important Safety Information on the next page, and click the link to view the full Prescribing Information for [EPCLUSA](#), including **BOXED WARNING ON HEPATITIS B REACTIVATION**.

Please be aware that if you are a licensed US healthcare professional, any meal or other transfers of value associated with this program may be reportable under the federal Open Payments/Sunshine Act and/or state law. Gilead does not provide or pay for alcohol at Speaker Programs. Gilead policy and the PhRMA Code on Interactions with Health Care Professionals limit attendance at this promotional educational program to healthcare professionals. Accordingly, attendance by guests who are non-healthcare professionals is not appropriate and cannot be accommodated.

## IMPORTANT SAFETY INFORMATION (CONT'D)

### BOXED WARNING: RISK OF HEPATITIS B VIRUS REACTIVATION IN HCV/HBV COINFECTED PATIENTS

Test all patients for evidence of current or prior hepatitis B virus (HBV) infection before initiating treatment with EPCLUSA. HBV reactivation has been reported in HCV/HBV coinfecting patients who were undergoing or had completed treatment with HCV direct-acting antivirals (DAAs) and were not receiving HBV antiviral therapy. Some cases have resulted in fulminant hepatitis, hepatic failure, and death. Cases have been reported in patients who are HBsAg positive, in patients with serologic evidence of resolved HBV, and also in patients receiving certain immunosuppressant or chemotherapeutic agents; the risk of HBV reactivation associated with treatment with HCV DAAs may be increased in patients taking these other agents. Monitor HCV/HBV coinfecting patients for hepatitis flare or HBV reactivation during HCV treatment and post-treatment follow-up. Initiate appropriate patient management for HBV infection as clinically indicated.

### WARNINGS AND PRECAUTIONS

- **Serious Symptomatic Bradycardia When Coadministered with Amiodarone:** Amiodarone is not recommended for use with EPCLUSA due to the risk of symptomatic bradycardia, particularly in patients also taking beta blockers or with underlying cardiac comorbidities and/or with advanced liver disease. A fatal cardiac arrest was reported in a patient taking amiodarone who was coadministered a sofosbuvir-containing regimen. In patients without alternative viable treatment options, cardiac monitoring is recommended. Patients should seek immediate medical evaluation if they develop signs or symptoms of bradycardia.
- **Risk of Reduced Therapeutic Effect Due to Concomitant Use of EPCLUSA with P-gp Inducers and/or Moderate to Strong Inducers of CYP2B6, CYP2C8 or CYP3A4:** Rifampin, St. John's wort, and carbamazepine are not recommended for use with EPCLUSA as they may significantly decrease sofosbuvir and/or velpatasvir plasma concentrations.

### ADVERSE REACTIONS

- The most common adverse reactions ( $\geq 10\%$ , all grades) with EPCLUSA were headache and fatigue.

### DRUG INTERACTIONS

- Coadministration of EPCLUSA is not recommended with topotecan due to increased concentrations of topotecan.
- Coadministration of EPCLUSA is not recommended with proton-pump inhibitors, phenobarbital, phenytoin, rifabutin, rifapentine, efavirenz, and tipranavir/ritonavir due to decreased concentrations of sofosbuvir and/or velpatasvir.

Consult the full Prescribing Information for EPCLUSA for more information on potentially significant drug interactions, including clinical comments.

Please click the link to view the full Prescribing Information for [EPCLUSA](#), including **BOXED WARNING**.

EPCLUSA, the EPCLUSA logo, GILEAD, and the GILEAD logo are trademarks of Gilead Sciences, Inc., or its related companies.  
©2023 Gilead Sciences, Inc. All rights reserved. US-EPCL-0296 03/23

