

BOEHRINGER INGELHEIM PHARMACEUTICALS, INC.
& LILLY USA, LLC CORDIALLY INVITE YOU TO A PRESENTATION



IN ADULTS WITH HEART FAILURE OR ADULTS WITH TYPE 2
DIABETES MELLITUS AND ESTABLISHED CARDIOVASCULAR DISEASE

THE POWER OF SGLT2is: A GUIDELINE- RECOMMENDED THERAPY

FEATURED GUEST SPEAKER	DATE
	TIME
LOCATION	

RSVP	RSVP BY
------	---------

This program is open to healthcare practitioners (HCPs) for whom the information presented is relevant to their practice. Spouses or guests cannot be accommodated.

By registering for this event, I agree to allow Boehringer Ingelheim Pharmaceuticals, Inc. (BIPI), Lilly USA, LLC, and third parties associated with the execution of this program to contact me by phone, fax, email, or in person.

BIPI and Lilly USA, LLC have adopted the PhRMA (Pharmaceutical Research and Manufacturers of America) Code on Interactions with Healthcare Professionals that went into effect on July 1, 2002. Pursuant to the Code, inclusion of healthcare professionals' spouses or guests is not permitted. We appreciate your understanding and support of our commitment to following the highest ethical standards as related to interactions with healthcare professionals. To comply with federal regulations, attendance at the entire session is required and early departures are not permitted.

In accordance with Boehringer Ingelheim, Inc. and Lilly USA, LLC Policy and the updated PhRMA Code on Interactions with HCPs, Boehringer Ingelheim, Inc. and Lilly USA, LLC will not provide or pay for alcohol at this educational event.

INDICATIONS AND LIMITATIONS OF USE

JARDIANCE is indicated:

- as an adjunct to diet and exercise to improve glycemic control in adults and pediatric patients aged 10 years and older with type 2 diabetes mellitus
- to reduce the risk of cardiovascular death in adults with type 2 diabetes mellitus and established cardiovascular disease
- to reduce the risk of cardiovascular death and hospitalization for heart failure in adults with heart failure
- to reduce the risk of sustained decline in eGFR, end-stage kidney disease, cardiovascular death, and hospitalization in adults with chronic kidney disease at risk of progression

JARDIANCE is not recommended for use to improve glycemic control in patients with type 1 diabetes mellitus. It may increase their risk of diabetic ketoacidosis.

JARDIANCE is not recommended for use to improve glycemic control in patients with type 2 diabetes mellitus with an eGFR <30 mL/min/1.73 m². JARDIANCE is likely to be ineffective in this setting based upon its mechanism of action.

JARDIANCE is not recommended for the treatment of chronic kidney disease in patients with polycystic kidney disease or patients requiring or with a recent history of intravenous immunosuppressive therapy or greater than 45 mg of prednisone or equivalent for kidney disease. JARDIANCE is not expected to be effective in these populations.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATIONS: Hypersensitivity to empagliflozin or any of the excipients in JARDIANCE, reactions such as angioedema have occurred.

Please see additional Important Safety Information on the next page and accompanying Prescribing Information, including Medication Guide for JARDIANCE.



Copyright © 2025 Boehringer Ingelheim Pharmaceuticals, Inc.
All rights reserved. (01/25) PC-US-141810-A

IMPORTANT SAFETY INFORMATION (cont'd)

WARNINGS AND PRECAUTIONS

Diabetic Ketoacidosis in Patients with Type 1 Diabetes Mellitus and Other Ketoacidosis: JARDIANCE increases the risk of life-threatening ketoacidosis in patients with type 1 diabetes and fatal ketoacidosis has occurred with JARDIANCE. Type 2 diabetes and pancreatic disorders are also risk factors for ketoacidosis and fatal events of ketoacidosis have been reported in patients with type 2 diabetes using JARDIANCE. Precipitating conditions for diabetic ketoacidosis or other ketoacidosis include under-insulinization due to insulin dose reduction or missed insulin doses, acute febrile illness, reduced caloric intake, ketogenic diet, surgery, volume depletion, and alcohol abuse. Signs and symptoms of diabetic ketoacidosis are consistent with dehydration and severe metabolic acidosis and include nausea, vomiting, abdominal pain, generalized malaise, and shortness of breath. Assess patients who present with signs and symptoms of metabolic ketoacidosis, regardless of blood glucose levels. If suspected, discontinue JARDIANCE, treat promptly and monitor for resolution before restarting. Consider ketone monitoring in patients with type 1 diabetes mellitus as well as in others at risk for ketoacidosis. Withhold JARDIANCE in clinical situations known to predispose to ketoacidosis and resume when clinically stable. Educate all patients on the signs and symptoms of ketoacidosis and instruct patients to discontinue JARDIANCE and seek medical attention immediately if signs and symptoms occur.

Volume Depletion: JARDIANCE can cause intravascular volume depletion which may manifest as symptomatic hypotension or acute transient changes in creatinine. Acute kidney injury requiring hospitalization and dialysis has been reported in patients with type 2 diabetes receiving SGLT2 inhibitors, including JARDIANCE. Before initiating, assess volume status and renal function in patients with impaired renal function (eGFR <60 mL/min/1.73 m²), elderly patients or patients on loop diuretics. In patients with volume depletion, correct this condition before initiating JARDIANCE. After initiating, monitor for signs and symptoms of volume depletion and renal function.

Urosepsis and Pyelonephritis: Serious urinary tract infections including urosepsis and pyelonephritis requiring hospitalization have been identified in patients receiving JARDIANCE. Treatment with JARDIANCE increases the risk for urinary tract infections. Evaluate for signs and symptoms of urinary tract infections and treat promptly.

Hypoglycemia: In adult patients, the use of JARDIANCE in combination with insulin or insulin secretagogues can increase the risk of hypoglycemia. In pediatric patients aged 10 years and older, the risk of hypoglycemia was higher with JARDIANCE regardless of insulin use. The risk of hypoglycemia may be lowered by a reduction in the dose of sulfonylurea (or other concomitantly administered insulin secretagogues) or insulin.

Necrotizing Fasciitis of the Perineum (Fournier's Gangrene): Serious, life-threatening cases requiring urgent surgical intervention have occurred in both females and males receiving JARDIANCE. Serious outcomes have included hospitalization, multiple surgeries and death. Assess patients presenting with pain or tenderness, erythema, or swelling in the genital or perineal area, along with fever or malaise. If suspected, institute prompt treatment and discontinue JARDIANCE.

Genital Mycotic Infections: JARDIANCE increases the risk for genital mycotic infections, especially in patients with prior infections. Monitor and treat as appropriate.

Lower Limb Amputation: Lower limb amputations have been observed in patients with chronic kidney disease taking JARDIANCE. Peripheral artery disease, and diabetic foot infection (including osteomyelitis), were the most common precipitating medical events leading to the need for an amputation. The risk of amputation was highest in patients with a baseline history of diabetic foot, peripheral artery disease (including previous amputation) or diabetes. Counsel patients receiving JARDIANCE about the importance of routine preventative foot care and monitor for signs and symptoms of diabetic foot infection (including osteomyelitis), new pain or tenderness, sores or ulcers involving the lower limbs, and institute appropriate treatment.

Hypersensitivity Reactions: Serious hypersensitivity reactions have occurred with JARDIANCE (angioedema). If hypersensitivity reactions occur, discontinue JARDIANCE, treat promptly, and monitor until signs and symptoms resolve.

MOST COMMON ADVERSE REACTIONS (≥5%): Urinary tract infections and female genital mycotic infections.

DRUG INTERACTIONS

Diuretics: Coadministration with diuretics may enhance the potential for volume depletion. Monitor for signs and symptoms.

Lithium: Concomitant use with lithium may decrease serum lithium concentrations. Monitor more frequently during JARDIANCE initiation and dosage changes.

USE IN SPECIAL POPULATIONS

Pregnancy: JARDIANCE is not recommended during the second and third trimesters.

Lactation: JARDIANCE is not recommended while breastfeeding.

Geriatric Use: JARDIANCE is expected to have diminished glycemic efficacy in elderly patients with renal impairment. Assess renal function more frequently in elderly patients. The incidence of volume depletion-related adverse reactions and urinary tract infections increased in T2D patients ≥75 years treated with empagliflozin.

CL-JAR-100162 09.21.2023

Please see accompanying Prescribing Information, including Medication Guide, for JARDIANCE.