

PLEASE JOIN US FOR A LIVE PRESENTATION ON A Treatment Option for Appropriate Patients With Certain Skin Diseases Driven in Part by Type 2 Inflammation

INDICATIONS

Atopic Dermatitis: DUPIXENT is indicated for the treatment of adult and pediatric patients aged 6 months and older with moderate-to-severe atopic dermatitis whose disease is not adequately controlled with topical prescription therapies or when those therapies are not advisable. DUPIXENT can be used with or without topical corticosteroids.

Prurigo Nodularis: DUPIXENT is indicated for the treatment of adult patients with prurigo nodularis (PN).

Chronic Spontaneous Urticaria: DUPIXENT is indicated for the treatment of adult and pediatric patients aged 12 years and older with chronic spontaneous urticaria (CSU) who remain symptomatic despite H1 antihistamine treatment.

Limitations of Use: DUPIXENT is not indicated for treatment of other forms of urticaria.

Bullous Pemphigoid: DUPIXENT is indicated for the treatment of adult patients with bullous pemphigoid (BP).

IMPORTANT SAFETY INFORMATION

CONTRAINDICATION: DUPIXENT is contraindicated in patients with known hypersensitivity to dupilumab or any of its excipients.

Please see additional Important Safety Information throughout and accompanying full Prescribing Information.

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, this Program is limited to U.S. Healthcare Professionals and persons with bona fide professional interest in the information presented. Attendance at this Program by guests or spouses is not permitted unless they would qualify as an appropriate attendee on their own.

If a meal is provided, actively licensed Minnesota and Vermont prescribers may attend, but not partake in the meal. Full-time Federal Employees may attend and partake in the meal if the Program is considered widely attended (50 or more attendees). If not, they may attend, but not partake in the meal. Part-time Federal Employees acting in their civilian capacity may attend the Program and partake in the meal. Alcohol is not permitted in connection with Speaker Programs and will not be provided. The value of any meal provided in connection with the Program may be reported in accordance with federal and state laws and regulations.

sanofi | REGENERON®

© 2025 Sanofi and Regeneron Pharmaceuticals, Inc. All Rights Reserved.
DUPIXENT is a registered trademark of Sanofi or an affiliate.

US.DUP.25.07.0208

DUPIXENT®
(dupilumab) Injection
200mg · 300mg



PRESENTED BY

Geeta Patel, DO
River Oaks Dermatology
Houston, TX

HOSTED BY

Lea Smolik
lea.smolik@regeneron.com
281-690-2228
RGN0042312

You may RSVP to your program host.

PROGRAM AGENDA

May 19, 2026
6:00 PM Central Presentation
Grotto
4715 Westheimer Rd
Houston, Texas 77027

Please arrive 30 minutes prior to registration.

TO REGISTER FOR THE EVENT

To register for this event, please call 281-690-2228
and reference RGN0042312, or visit
<https://tpprod-regeneron.physiciansworld.com/3xS8D6>

IMPORTANT SAFETY INFORMATION (CONT.)

WARNINGS AND PRECAUTIONS

Hypersensitivity: Hypersensitivity reactions, including anaphylaxis, acute generalized exanthematous pustulosis (AGEP), serum sickness or serum sickness-like reactions, angioedema, generalized urticaria, rash, erythema nodosum, and erythema multiforme have been reported. A case of AGEP was reported in an adult subject who participated in the bullous pemphigoid development program. If a clinically significant hypersensitivity reaction occurs, institute appropriate therapy and discontinue DUPIXENT.

Conjunctivitis and Keratitis: Conjunctivitis and keratitis occurred more frequently in AD and BP subjects who received DUPIXENT versus placebo, with conjunctivitis being the most frequently reported eye disorder in AD. Conjunctivitis also occurred more frequently in PN subjects who received DUPIXENT compared to those who received placebo. Conjunctivitis and keratitis have been reported with DUPIXENT in postmarketing settings, predominantly in AD patients. Some patients reported visual disturbances (e.g., blurred vision) associated with conjunctivitis or keratitis. Advise patients or their caregivers to report new-onset or worsening eye symptoms. Consider ophthalmological examination for patients who develop conjunctivitis that does not resolve following standard treatment or signs and symptoms suggestive of keratitis, as appropriate.

Risk Associated with Abrupt Reduction of Corticosteroid Dosage: Do not discontinue systemic, topical, or inhaled corticosteroids abruptly upon initiation of DUPIXENT. Reductions in corticosteroid dose, if appropriate, should be gradual and performed under the direct supervision of a healthcare provider. Reduction in corticosteroid dose may be associated with systemic withdrawal symptoms and/or unmask conditions previously suppressed by systemic corticosteroid therapy.

Patients with Co-morbid Asthma: Advise patients not to adjust or stop their asthma treatments without consultation with their physicians.

Psoriasis: Cases of new-onset psoriasis have been reported with the use of DUPIXENT for the treatment of atopic dermatitis, including in patients without a family history of psoriasis. In postmarketing reports, these cases resulted in partial or complete resolution of psoriasis with discontinuation of dupilumab, with or without use of supplemental treatment for psoriasis (topical or systemic). Those who continued dupilumab received supplemental treatment for psoriasis to improve associated symptoms. Advise patients to report new-onset psoriasis symptoms. If symptoms persist or worsen, consider dermatologic evaluation and/or discontinuation of DUPIXENT.

Arthralgia and Psoriatic Arthritis: Arthralgia has been reported with the use of DUPIXENT with some patients reporting gait disturbances or decreased mobility associated with joint symptoms; some cases resulted in hospitalization. Cases of new-onset psoriatic arthritis requiring systemic treatment have been reported with the use of DUPIXENT. Advise patients to report new-onset or worsening joint symptoms. If symptoms persist or worsen, consider rheumatological evaluation and/or discontinuation of DUPIXENT.

Parasitic (Helminth) Infections: It is unknown if DUPIXENT will influence the immune response against helminth infections. Treat patients with pre-existing helminth infections before initiating therapy with DUPIXENT. If patients become infected while receiving treatment with DUPIXENT and do not respond to anti-helminth treatment, discontinue treatment with DUPIXENT until the infection resolves.

Vaccinations: Consider completing all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating DUPIXENT. Avoid use of live vaccines during treatment with DUPIXENT.

ADVERSE REACTIONS:

Most common adverse reactions are:

- **Atopic Dermatitis** (incidence $\geq 1\%$): injection site reactions, conjunctivitis, blepharitis, oral herpes, keratitis, eye pruritus, other herpes simplex virus infection, dry eye, and eosinophilia. The safety profile in pediatric patients through Week 16 was similar to that of adults with AD. In an open-label extension study, the long-term safety profile of DUPIXENT \pm TCS in pediatric patients observed through Week 52 was consistent with that seen in adults with AD, with hand-foot-and-mouth disease and skin papilloma (incidence $\geq 2\%$) reported in patients 6 months to 5 years of age. These cases did not lead to study drug discontinuation.
- **Prurigo Nodularis** (incidence $\geq 2\%$): nasopharyngitis, conjunctivitis, herpes infection, dizziness, myalgia, and diarrhea.
- **Chronic Spontaneous Urticaria** (incidence $\geq 2\%$): injection site reactions.
- **Bullous Pemphigoid** (incidence $\geq 2\%$): arthralgia, conjunctivitis, vision blurred, herpes viral infections, keratitis.

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** A pregnancy exposure registry monitors pregnancy outcomes in women exposed to DUPIXENT during pregnancy. To enroll or obtain information call 1-877-311-8972 or go to <https://mothertobaby.org/ongoing-study/dupixent/>. Available data from case reports and case series with DUPIXENT use in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. Human IgG antibodies are known to cross the placental barrier; therefore, DUPIXENT may be transmitted from the mother to the developing fetus.
- **Lactation:** There are no data on the presence of DUPIXENT in human milk, the effects on the breastfed infant, or the effects on milk production. Maternal IgG is known to be present in human milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for DUPIXENT and any potential adverse effects on the breastfed child from DUPIXENT or from the underlying maternal condition.

Please see accompanying full [Prescribing Information](#)