



# CABENUVA

cabotegravir 200 mg/mL; rilpivirine 300 mg/mL  
extended-release injectable suspensions

## ViiV Healthcare Presents:

### PRESENTED BY:



EVENT DATE:

EVENT LOCATION:



Scan QR code to  
register for this program.

Please RSVP by contacting your ViiV Representative:

Speaker(s) compensated or employed  
by ViiV Healthcare.

### INDICATION

CABENUVA is indicated as a complete regimen for the treatment of HIV-1 infection in adults and adolescents 12 years of age and older and weighing at least 35 kg to replace the current antiretroviral regimen in those who are virologically suppressed (HIV-1 RNA <50 copies/mL) on a stable antiretroviral regimen with no history of treatment failure and with no known or suspected resistance to either cabotegravir or rilpivirine.

### IMPORTANT SAFETY INFORMATION

#### CONTRAINDICATIONS

- Do not use CABENUVA in patients with previous hypersensitivity reaction to cabotegravir or rilpivirine
- Do not use CABENUVA in patients receiving carbamazepine, oxcarbazepine, phenobarbital, phenytoin, rifabutin, rifampin, rifapentine, systemic dexamethasone (>1 dose), and St John's wort

Please see additional Important Safety Information for CABENUVA on page 2.

Please see accompanying full [Prescribing Information](#) for CABENUVA.

Please RSVP by



## IMPORTANT SAFETY INFORMATION (cont'd)

### WARNINGS AND PRECAUTIONS

#### Hypersensitivity Reactions:

- Serious or severe hypersensitivity reactions, including Stevens-Johnson syndrome (SJS)/toxic epidermal necrolysis (TEN) and drug reaction with eosinophilia and systemic symptoms (DRESS), have been reported with CABENUVA or its components. While some skin reactions were accompanied by constitutional symptoms such as fever, other skin reactions were associated with organ dysfunctions, including elevations in hepatic serum biochemistries
- Discontinue CABENUVA immediately if signs or symptoms of hypersensitivity reactions develop. Clinical status, including liver transaminases, should be monitored and appropriate therapy initiated. Cabotegravir and rilpivirine oral lead-in may be used to help identify patients who may be at risk of a hypersensitivity reaction

#### Post-Injection Reactions:

- Serious post-injection reactions (reported in less than 1% of subjects) were reported within minutes after the injection of rilpivirine, including dyspnea, bronchospasm, agitation, abdominal cramping, rash/urticaria, dizziness, flushing, sweating, oral numbness, changes in blood pressure, and pain (e.g., back and chest). These events may have been associated with accidental intravenous administration and began to resolve within a few minutes after the injection
- Carefully follow the Instructions for Use when preparing and administering CABENUVA. The suspensions should be injected slowly via intramuscular injection and avoid accidental intravenous administration. Observe patients briefly (approximately 10 minutes) after the injection. If a post-injection reaction occurs, monitor and treat as clinically indicated

#### Hepatotoxicity:

- Hepatotoxicity has been reported in patients receiving cabotegravir or rilpivirine with or without known pre-existing hepatic disease or identifiable risk factors
- Patients with underlying liver disease or marked elevations in transaminases prior to treatment may be at increased risk for worsening or development of transaminase elevations
- Monitoring of liver chemistries is recommended and treatment with CABENUVA should be discontinued if hepatotoxicity is suspected

#### Depressive Disorders:

- Depressive disorders (including depressed mood, depression, major depression, mood altered, mood swings, dysphoria, negative thoughts, suicidal ideation, suicide attempt) have been reported with CABENUVA or the individual products
- Promptly evaluate patients with depressive symptoms

#### Risk of Adverse Reactions or Loss of Virologic Response Due to Drug Interactions:

- The concomitant use of CABENUVA and other drugs may result in known or potentially significant drug interactions (see Contraindications and Drug Interactions)
- Rilpivirine doses 3 and 12 times higher than the recommended oral dosage can prolong the QTc interval
- CABENUVA should be used with caution in combination with drugs with a known risk of Torsade de Pointes

#### Long-Acting Properties and Potential Associated Risks with CABENUVA:

- Residual concentrations of cabotegravir and rilpivirine may remain in the systemic circulation of patients for prolonged periods (up to 12 months or longer). Select appropriate patients who agree to the required monthly or every-2-month injection dosing schedule because non-adherence could lead to loss of virologic response and development of resistance
- To minimize the potential risk of developing viral resistance, it is essential to initiate an alternative, fully suppressive antiretroviral regimen no later than 1 month after the final injection doses of CABENUVA when dosed monthly and no later than 2 months after the final injections of CABENUVA when dosed every 2 months. If virologic failure is suspected, switch the patient to an alternative regimen as soon as possible

### ADVERSE REACTIONS

- The most common adverse reactions (incidence  $\geq 2\%$ , Grades 1 to 4) with CABENUVA were injection site reactions, pyrexia, fatigue, headache, musculoskeletal pain, nausea, sleep disorders, dizziness, and rash

### DRUG INTERACTIONS

- Refer to the applicable full Prescribing Information for important drug interactions with CABENUVA, VOCABRIA (cabotegravir), or EDURANT (rilpivirine)
- Because CABENUVA is a complete regimen, coadministration with other antiretroviral medications for the treatment of HIV-1 infection is not recommended
- Drugs that are strong inducers of UGT1A1 or UGT1A9 are expected to decrease the plasma concentrations of cabotegravir. Drugs that induce or inhibit CYP3A may affect the plasma concentrations of rilpivirine
- CABENUVA should be used with caution in combination with drugs with a known risk of Torsade de Pointes

### USE IN SPECIFIC POPULATIONS

- **Pregnancy:** There are insufficient human data on the use of CABENUVA during pregnancy to adequately assess a drug-associated risk for birth defects and miscarriage. Discuss the benefit-risk of using CABENUVA during pregnancy and conception and consider that cabotegravir and rilpivirine are detected in systemic circulation for up to 12 months or longer after discontinuing injections of CABENUVA. An Antiretroviral Pregnancy Registry has been established
- **Lactation:** Potential risks of breastfeeding include HIV-1 transmission, developing viral resistance in HIV-positive infants, and adverse reactions in a breastfed infant

Please see additional Important Safety Information on page 1.

Please see accompanying full [Prescribing Information](#) for CABENUVA.



Trademarks are property of their respective owners.  
©2025 ViiV Healthcare or licensor.  
PMUS-CBRLBND240042 June 2025  
Produced in USA.



**CABENUVA**  
cabotegravir 200 mg/mL; rilpivirine 300 mg/mL  
extended-release injectable suspensions