

Madrigal cordially invites you to attend an educational program on

Rezdiffra® resmetirom tablets

60mg • 80mg • 100mg

Rezdiffra: Unlocking the potential of a liver-directed treatment

The first FDA-approved treatment, in conjunction with diet and exercise, for adults with noncirrhotic MASH with moderate to advanced fibrosis

This indication is approved under accelerated approval based on improvement of MASH and fibrosis. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials. Limitation of Use: Avoid use in patients with decompensated cirrhosis.

Presented by



Andres Henriquez, MD

Endocrinologist
Private Practice
Pembroke Pines, FL



Thursday, November 12, 2026



Arrival: 6:15 PM Start: 6:30 PM



**Vic's on the River
26 E. Bay Street
Savannah, GA 31401**

Registration:

We look forward to your participation at this educational program.

If you have any questions, please contact your Madrigal representative, Chris Gaffney at 843-696-8666 or via email at cgaffney@madrigalpharma.com.

Scan Here



INDICATION AND IMPORTANT SAFETY INFORMATION

INDICATION

Rezdiffra is indicated in conjunction with diet and exercise for the treatment of adults with noncirrhotic metabolic dysfunction-associated steatohepatitis (MASH) with moderate to advanced liver fibrosis (consistent with stages F2 to F3 fibrosis).

This indication is approved under accelerated approval based on improvement of MASH and fibrosis. Continued approval for this indication may be contingent upon verification and description of clinical benefit in confirmatory trials.

Limitation of Use: Avoid use in patients with decompensated cirrhosis.

WARNINGS AND PRECAUTIONS

Hepatotoxicity

Hepatotoxicity has been observed with the use of Rezdiffra. One patient developed substantial elevations of liver biochemistries that resolved when treatment was interrupted. *Please see full Prescribing Information for more details on this specific case of Hepatotoxicity [see Warnings and Precautions (5.1)].*

Monitor for elevations in liver tests, liver-related adverse reactions, and symptoms/signs of hepatotoxicity (eg, fatigue, nausea, vomiting, right upper quadrant pain or tenderness, jaundice, fever, rash, and/or eosinophilia [$>5\%$]). If hepatotoxicity is suspected, discontinue Rezdiffra and monitor. If laboratory values return to baseline, weigh the potential risks against the benefits of restarting Rezdiffra. If laboratory values do not return to baseline, consider drug-induced autoimmune-like hepatitis (DI-ALH) or autoimmune liver disease in the evaluation of elevations in liver tests.

Gallbladder-Related Adverse Reactions

Cholelithiasis, acute cholecystitis, and obstructive pancreatitis (gallstone) were observed more often in Rezdiffra-treated patients than in placebo-treated patients. The exposure-adjusted incidence rates (EAIRs) for these events were less than 1 per 100 person-years (PY) for all treatment arms. If cholelithiasis is suspected, gallbladder diagnostic studies and appropriate clinical follow-up are indicated. If an acute gallbladder event is suspected, interrupt treatment until the event is resolved.

Please see additional Important Safety Information on back and accompanying full Prescribing Information for Rezdiffra or visit <https://www.madrigalpharma.com/wp-content/uploads/2026/01/Rezdiffra-USPI-approved-Dec-2025.pdf>

Visit RezdiffraHCP.com to learn more and get patients started.



IMPORTANT SAFETY INFORMATION (cont.)

WARNINGS AND PRECAUTIONS (cont.)

Drug Interaction with Certain Statins

An increase in exposure of atorvastatin, pravastatin, rosuvastatin and simvastatin was observed when concomitantly administered with Rezdiffra, which may increase the risk of adverse reactions related to these drugs.

Dosage adjustment for certain statins is recommended. Monitor for statin-related adverse reactions including, but not limited to, elevation of liver tests, myopathy, and rhabdomyolysis.

ADVERSE REACTIONS

The most common adverse reactions with Rezdiffra (reported in $\geq 5\%$ of patients and higher compared to placebo) are diarrhea, nausea, pruritus, vomiting, constipation, abdominal pain, and dizziness. Diarrhea and nausea were the most common causes of treatment discontinuation.

DRUG INTERACTIONS

Clinically Significant Interaction Effects of Strong or Moderate CYP2C8 Inhibitors on Rezdiffra

- Concomitant use with strong CYP2C8 inhibitors (eg, gemfibrozil) is not recommended. Reduce Rezdiffra dosage if used concomitantly with a moderate CYP2C8 inhibitor (eg, clopidogrel).

Clinically Significant Interactions Affecting Other Drugs

- Statins:** Limit daily rosuvastatin and simvastatin dosage to 20 mg. Limit pravastatin and atorvastatin dosage to 40 mg.
- CYP2C8 Substrates:** Monitor patients more frequently for substrate-related adverse reactions if Rezdiffra is co-administered with CYP2C8 substrates where minimal concentration changes may lead to serious adverse reactions.

USE IN SPECIFIC POPULATIONS

Pregnancy

There are no available data on Rezdiffra use in pregnant women to evaluate for a drug-associated risk of major birth defects, miscarriage, or other adverse maternal or fetal outcomes. Report pregnancies to Madrigal Pharmaceuticals, Inc. Adverse Event Reporting line at 1-800-905-0324 and visit <https://pregnancyregistry.madrigalpharma.com> for information about a pregnancy safety study.

Lactation

There is no information regarding the presence of Rezdiffra in human or animal milk, the effects on the breast-fed infant, or the effects on milk production. The developmental and health benefits of breastfeeding should be considered along with the mother's

clinical need for Rezdiffra and any potential adverse effects on the breastfed infant from Rezdiffra or from the underlying maternal condition.

Geriatric Use

Numerically higher incidence of adverse reactions have been observed in patients ≥ 65 years of age compared to younger adult patients.

Renal Impairment

The recommended dosage of Rezdiffra in patients with mild, moderate, or severe renal impairment is the same as in patients with normal kidney function.

Hepatic Impairment

Avoid use in patients with decompensated cirrhosis (consistent with moderate to severe hepatic impairment). Moderate or severe hepatic impairment (Child-Pugh Class B or C) may increase the risk of adverse reactions.

The safety and effectiveness of Rezdiffra have not been established in patients with cirrhosis.

DISCLOSURE

This promotional educational activity is not accredited. The program content is developed by Madrigal Pharmaceuticals, Inc. Speakers are being compensated for their time by Madrigal Pharmaceuticals, Inc., present on behalf of the company, and are required to present information in compliance with FDA requirements for communications about its medicine.

In adherence with PhRMA guidelines, spouses or other guests are not permitted to attend company-sponsored programs, unless spouse or other guest(s) independently qualify as an attendee.

For all attendees, please be advised that information, such as your name, and the value and purpose of any educational item, meal, or other items of value you receive, may be publicly disclosed. If you are licensed in any state or other jurisdiction, or are an employee or contractor of any organization or governmental entity that limits or prohibits meals from pharmaceutical companies, please identify yourself so that you (and we) are able to comply with such requirements.

Please note that the company prohibits the offering of gifts, gratuities, or meals to federal government employees/officials. Thank you for your cooperation.

Please see accompanying full Prescribing Information for Rezdiffra at

<https://www.madrigalpharma.com/wp-content/uploads/2026/01/Rezdiffra-USPI-approved-Dec-2025.pdf>

FDA=Food and Drug Administration; MASH=metabolic dysfunction-associated steatohepatitis, formerly known as NASH or nonalcoholic steatohepatitis.



© 2026 Madrigal Pharmaceuticals, Inc. All rights reserved. US-PP-RES-00560-v3 2/26
Rezdiffra® and its related logo are trademarks of Madrigal Pharmaceuticals, Inc.

Rezdiffra®
resmetirom tablets
60mg - 80mg - 100mg