



You are cordially invited to attend a **Milestone Speaker Program:**

# **CARDAMYST: The First and Only FDA-Approved Nasal Spray for Adults with Acute, Symptomatic PSVT**

**Speaker:** Neil Sanghvi, MD  
UF Health St. Johns  
St. Augustine, FL

**Location, Date, and Time:** Thursday, November 5, 2026 | 6:00 PM ET  
Seasons 52  
5096 Big Island Drive  
Jacksonville, FL 32246

**Rep Contact:** Valerie Morvant  
Territory Manager  
914-420-5430

**RSVP:**



Registration required. Please RSVP by:  
10/29/2026

Please note this program is intended for US healthcare professionals only. This speaker program is sponsored by Milestone Pharmaceuticals and is not eligible for CME credits. The featured speaker is a paid consultant of Milestone Pharmaceuticals.

## **IMPORTANT SAFETY INFORMATION FOR CARDAMYST (etripamil)**

### **INDICATIONS AND USAGE**

CARDAMYST is indicated for the conversion of acute symptomatic episodes of paroxysmal supraventricular tachycardia (PSVT) to sinus rhythm in adults.

### **CONTRAINDICATIONS**

CARDAMYST is contraindicated in patients with:

- Hypersensitivity to CARDAMYST or any of its components.

**Please see additional Important Safety Information on back. Please see accompanying full Prescribing Information, including Patient Information.**



Product is not actual size.

## IMPORTANT SAFETY INFORMATION FOR CARDAMYST (etripamil) (cont)

### CONTRAINDICATIONS (cont)

- Heart failure - New York Heart Association (NYHA) Class II to IV.
- Wolff-Parkinson-White (WPW), Lown-Ganong-Levine (LGL) syndromes, or manifest pre-excitation (delta wave) on a 12-lead electrocardiogram (ECG).
- Sick sinus syndrome without a permanent pacemaker.
- Second degree atrioventricular (AV) Mobitz 2 block or higher degree of AV block.

### WARNINGS AND PRECAUTIONS

#### Syncope Related to Hemodynamic Effects

- Because of effects on blood pressure, heart rate, and cardiac conduction, CARDAMYST may cause dizziness and/or syncope, especially in patients with a history of syncope and high-grade AV block or sinus node dysfunction, or those with a history of syncope during an episode of PSVT.
- In clinical trials, a small percentage of patients (0.4%) experienced clinically significant hypotension during test dosing prior to randomization, which precluded further participation in the study. Patients with a history of hypotensive episodes or those at increased risk for hemodynamic instability should be monitored appropriately when initiating CARDAMYST.
- If syncope occurs, patients should be placed in the recumbent position and treated supportively.
- Patients should be cautioned about these possible adverse effects and advised to administer CARDAMYST in a sitting position, and in a location where the risk of fall is minimal.

**ADVERSE REACTIONS:** The most common adverse reactions (incidence  $\geq 5\%$ ) observed in randomized controlled studies are nasal discomfort, nasal congestion, rhinorrhea, throat irritation, and epistaxis.

**Lactation:** A lactating woman should pump and discard breastmilk for 12 hours after CARDAMYST administration in order to minimize exposure to a breastfed infant.

**Please see accompanying full Prescribing Information, including Patient Information.**