



# Diabetes Management in Primary Care



Kristin Pratt, RN, MSN, FNP-C  
Huron Regional Medical Center

# Disclosures

- PRN Employee with Horizon Health
- Current Board Member of the SDDC

# Learning Objectives

1. Understand the role of primary care providers in diabetes management.
2. Identify evidence-based approaches to diabetes treatment and monitoring.
3. Address common challenges in diabetes care and patient engagement.

# Prevalence of Diabetes in SD


In South Dakota, 10% of adults were living with a diagnosis of diabetes.

This ranged from 7% to 22% of adults across counties in the state.

Every year, an estimated 4,100 adults in South Dakota are diagnosed with diabetes.

Please note: This information is from the **2024 Annual Data Release** and uses data from **2021** for this measure.

Reported DM DX 10  
22%



SD Population  
WITHOUT DM Dx  
78%

# So, what is the role of the Primary Care Clinician?

- Likely to be the first and most frequent point of contact for individuals with Type 2 Diabetes.
- Play a pivotal role in early diagnosis, establishment of effective treatment strategies, and coordination of care.
- Aim to prevent but also screen and identify any secondary complications.



# Key Components of Diabetes Care in the Primary Care Setting

1. Complete comprehensive diabetes medical evaluation at initial, follow-up, and annual visits
2. Screen, prevent, and delay complications
3. Facilitate Positive Health Behaviors and Well-Being to Improve Health Outcomes (education and lifestyle change program, nutrition, any other pertinent referrals)
4. Establish glycemic goals, time-in-range goals, and plans for hypo/hyperglycemia
5. Consider what diabetes technology will be used and prescribed

| Guidelines                                                                                                                                                | Details                                                                                                                                                                                | Action Plan                                                                                                  |       |       |       |        |       |                                                                                                                                                                                                                                                |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------|-------|-------|-------|--------|-------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Diabetes Medicines</b><br>                                          | <b>My insulin dosing schedule</b><br>Type of insulin: _____ #units _____ Time to inject: _____<br><br><b>My diabetes medications</b><br>Rx Name: _____ Dose: _____ Time to take: _____ | <b>Target blood sugars</b><br><br>Before meals: _____<br><br>2 hours after meal: _____<br><br>Bedtime: _____ |       |       |       |        |       |                                                                                                                                                                                                                                                |
| <b>Eating Schedule</b><br>                                             | <b>Meals and snacks:</b>                                                                                                                                                               | <b>Notes:</b><br><div style="border: 1px solid black; height: 40px; width: 100%;"></div>                     |       |       |       |        |       |                                                                                                                                                                                                                                                |
| <b>Carbohydrate Budget</b><br>Starches, fruits, milk, and sweets<br> | <table border="1"> <tr><td>Breakfast</td></tr> <tr><td>Snack</td></tr> <tr><td>Lunch</td></tr> <tr><td>Snack</td></tr> <tr><td>Dinner</td></tr> <tr><td>Snack</td></tr> </table>       | Breakfast                                                                                                    | Snack | Lunch | Snack | Dinner | Snack | <b>15g of Carbohydrate:</b><br>1 cup milk<br>½ cup cooked rice, pasta,<br>½ cup dry cereal<br>½ cup cooked oatmeal<br>1 oz slice bread<br>1 medium fruit<br>3 cups popped popcorn<br>½ cup fruit juice<br>½ cup tomato sauce<br>1 small potato |
| Breakfast                                                                                                                                                 |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |
| Lunch                                                                                                                                                     |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |
| Dinner                                                                                                                                                    |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                                              |       |       |       |        |       |                                                                                                                                                                                                                                                |

# Key Components of Diabetes Care in the Primary Care Setting (cont.)

6. Consider specific goals and plans for patients who are overweight or obese/overall weight management discussions
7. Pharmacologic approach and plan
8. Cardiovascular disease and risk management
9. Chronic kidney disease and risk management
10. Retinopathy, neuropathy, and foot care
11. Considerations for older adults
12. Recognize and address any barriers/challenges to care

| Guidelines                                                                                                                                                | Details                                                                                                                                                                                | Action Plan                                                                               |       |       |       |        |       |                                                                                                                                                                                                                                         |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------|-------|-------|-------|--------|-------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Diabetes Medicines</b><br>                                          | <b>My insulin dosing schedule</b><br>Type of insulin: _____ #units _____ Time to inject: _____<br><br><b>My diabetes medications</b><br>Rx Name: _____ Dose: _____ Time to take: _____ | Target blood sugars<br>Before meals: _____<br>2 hours after meal: _____<br>Bedtime: _____ |       |       |       |        |       |                                                                                                                                                                                                                                         |
| <b>Eating Schedule</b><br>                                             | Meals and snacks:                                                                                                                                                                      | <b>Notes:</b><br>_____<br>_____<br>_____                                                  |       |       |       |        |       |                                                                                                                                                                                                                                         |
| <b>Carbohydrate Budget</b><br>Starches, fruits, milk, and sweets<br> | <table border="1"> <tr><td>Breakfast</td></tr> <tr><td>Snack</td></tr> <tr><td>Lunch</td></tr> <tr><td>Snack</td></tr> <tr><td>Dinner</td></tr> <tr><td>Snack</td></tr> </table>       | Breakfast                                                                                 | Snack | Lunch | Snack | Dinner | Snack | 15g of Carbohydrate:<br>1 cup milk<br>½ cup cooked rice, pasta,<br>½ cup dry cereal<br>½ cup cooked oatmeal<br>1 oz slice bread<br>1 medium fruit<br>3 cups popped popcorn<br>½ cup fruit juice<br>½ cup tomato sauce<br>1 small potato |
| Breakfast                                                                                                                                                 |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |
| Lunch                                                                                                                                                     |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |
| Dinner                                                                                                                                                    |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |
| Snack                                                                                                                                                     |                                                                                                                                                                                        |                                                                                           |       |       |       |        |       |                                                                                                                                                                                                                                         |

# Comprehensive Diabetes Medical Evaluation


# Comprehensive Diabetes Medical Evaluation



- Initial visit: complete H&P, ask about DM hx, co-morbid conditions, inquire about behavioral factors, medications, immunizations, technology use, social life, physical exam, lab evaluation, make follow-up appointments
- Every follow-up: interval changes, any updates to hx, last dental/vision exams, changes in medications, social/behavioral status, review glucose logs or CGM data, physical exam, follow-up labs

# Screen, Prevent, and Delay Complications

Consider having  
patients take the 60-  
Second Risk Rest for  
Type 2 Diabetes |  
UndotheRisk |  
[www.undotheRisk.com](http://www.undotheRisk.com)



The screenshot shows the top of the UndotheRisk website. At the top is the South Dakota Department of Health logo, which consists of an orange 'H' icon and the text 'SOUTH DAKOTA DEPARTMENT OF HEALTH'. To the right of the logo is a navigation menu with links: 'Topics', 'Programs', 'Licensing & Records', and 'Healthcare Professionals', each followed by a downward arrow. Below the navigation bar is a large blue banner featuring a close-up, profile view of a young girl and a woman smiling at each other. Overlaid on the banner is the text 'Undo The Risk' in white, with a small orange horizontal line underneath. Above this text is a breadcrumb trail: '> Topics > Diseases > Chronic Disease'. Below the banner is a section with a heart icon containing a red checkmark, followed by the text 'Check. Know. Act'. Underneath this is a heading 'A simple test can preserve everything you have.' and a paragraph of text: 'Chronic health conditions like diabetes, heart disease, and stroke can be avoided or even undone, but only if you take control. Start today by taking this quick and easy risk assessment to see where you stand.' At the bottom of this section is a dark blue button with the text 'Take the Assessment' in white.

SOUTH DAKOTA  
DEPARTMENT OF HEALTH

Topics ▾ Programs ▾ Licensing & Records ▾ Healthcare Professionals ▾

> Topics > Diseases > Chronic Disease

**Undo The Risk**

 **Check. Know. Act**

**A simple test can preserve everything you have.**

Chronic health conditions like diabetes, heart disease, and stroke can be avoided or even undone, but only if you take control. Start today by taking this quick and easy risk assessment to see where you stand.

[Take the Assessment](#)

# ADA Adult Screening Recommendations

Criteria for screening for diabetes or prediabetes in asymptomatic adults:

1. Testing should be considered in adults with overweight or obesity ( $\text{BMI} \geq 25 \text{ kg/m}^2$  or  $\geq 23 \text{ kg/m}^2$  in individuals of Asian ancestry) who have one or more of the following risk factors:
  - First-degree relative with diabetes
  - High-risk race, ethnicity, and ancestry (e.g., African American, Latino, Native American, Asian American)
  - History of cardiovascular disease
  - Hypertension ( $\geq 130/80 \text{ mmHg}$  or on therapy for hypertension)
  - HDL cholesterol level  $< 35 \text{ mg/dL}$  ( $< 0.9 \text{ mmol/L}$ ) and/or triglyceride level  $> 250 \text{ mg/dL}$  ( $> 2.8 \text{ mmol/L}$ )
  - Individuals with polycystic ovary syndrome
  - Physical inactivity
  - Other clinical conditions associated with insulin resistance (e.g., severe obesity, acanthosis nigricans, metabolic dysfunction – associated steatotic liver disease)

# ADA Adult Screening Recommendations (cont.)

2. People with prediabetes (A1C  $\geq$  5.7% [ $\geq$ 39 mmol/mol], IGT or IFG)\* should be tested yearly.
3. People who were diagnosed with GDM\* should have testing at least every 1-3 years.
4. For all other people, testing should begin at age 35 years.
5. If results are normal, testing should be repeated at a minimum of 3-year intervals, with consideration of more frequent testing depending on initial results and risk status.
6. Individuals in other high-risk groups (e.g., people with HIV, exposure to high-risk medicines, evidence of periodontal disease, history of pancreatitis) should also be closely monitored.

\*Impaired glucose tolerance (IGT), impaired fasting glucose (IFG), gestational diabetes mellitus (GDM)



Adults with overweight or obesity (BMI  $\geq 25$  kg/m<sup>2</sup> or  $\geq 23$  kg/m<sup>2</sup> in individuals of Asian ancestry) who have one or more of the following risk factors:

- ☐ First-degree relative with diabetes
- ☐ High-risk race, ethnicity, and ancestry (e.g., African American, Latino, Native American, and Asian American)
- ☐ History of cardiovascular disease
- ☐ Hypertension ( $\geq 130/80$  mmHg or on therapy for hypertension)
- ☐ HDL cholesterol  $< 35$  mg/dL ( $< 0.9$  mmol/L) and/or triglyceride level  $> 250$  mg/dL ( $> 2.8$  mmol/L)
- ☐ Polycystic ovary syndrome
- ☐ Physical inactivity
- ☐ Other clinical conditions associated with insulin resistance (e.g., severe obesity, acanthosis nigricans, and metabolic dysfunction-associated steatotic liver disease)



#### Clinical Notes

- If results are normal, repeat screening at least every 3 years (annually for those with prediabetes), or sooner with symptoms or changes in risk factors for diabetes.

# ADA Children/Adolescent Screening Recommendations

- Risk-based screening for Type 2 Diabetes or prediabetes in asymptomatic children and adolescents in a clinical setting.
- Screening should be considered in youth\* who have overweight ( $\geq 85$ th percentile) or obesity ( $\geq 95$ th percentile) **and** who have one or more additional risk factors:
  - Maternal history of diabetes or GDM during the child's gestation
  - Family history of Type 2 Diabetes in first- or second-degree relative
  - High-risk race, ethnicity, and ancestry (see Table 2.5)
  - Signs of insulin resistance or conditions associated with insulin resistance (acanthosis nigricans, hypertension, dyslipidemia, polycystic ovary syndrome, large- or small-for-gestational-age birth weight)

\* After the onset of puberty or after 10 years of age, whichever occurs earlier. If tests are normal, repeat testing at a minimum of 3-year intervals (or more frequently if BMI is increasing or risk factor profile is deteriorating) is recommended. Reports of Type 2 Diabetes before age 10 years exist, and this can be considered with numerous risk factors.

# Key Strategies for Diabetes Prevention



01

## Healthy Eating Habits

Focus on balanced, nutritious meals daily. Encourage more plant-based foods, healthy fats, and skip fad diets and encourage lifestyle changes.



02

## Regular Physical Activity

Engage in exercise for at least 30 minutes of aerobic activity most days for a total of 150 minutes a week. Encourage strength training, balance training for older adults, and overall less time sitting.

03

## Losing Weight if You're Overweight or Obese

The American Diabetes Association recommends that people with prediabetes lose at least 5% to 7% of their body weight to prevent diabetes. More weight loss can turn into even greater benefits.

People in one large study lowered their risk of developing Type 2 Diabetes by almost 60% over three years. In this study, they lost about 7% of their body weight with changes in physical activity and diets.

|                                                                                                                                                             |                                                                                                                                                                                                                                                                  |                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>1. Why Screen?</b> <ul style="list-style-type: none"> <li>• Lab testing is safe and cost-effective.</li> <li>• Screening presents an opportunity to address cardiovascular risk factors (e.g., hypertension and dyslipidemia).</li> </ul> | <b>2. How to Screen</b> <ul style="list-style-type: none"> <li>• Conduct an assessment of risk factors.</li> </ul> <p>—or—</p> <ul style="list-style-type: none"> <li>• Use an assessment tool such as the <a href="#">American Diabetes Association risk test</a>.</li> <li>• Consider diagnostic testing based on assessment results.</li> </ul> | <b>3. When to Screen</b> <ul style="list-style-type: none"> <li>• Monitor people with prediabetes at least annually.</li> <li>• In those without prediabetes who have normal results, repeat screening at least every 3 years.</li> <li>• Screen after the onset of puberty or after the age of 10 years in children and adolescents with overweight (BMI ≥85th percentile) or obesity (BMI ≥95th percentile) who have at least one risk factor.</li> <li>• Screen whenever there are symptoms suggestive of diabetes or changes in risk.</li> </ul> |

#### Nutrition and Physical Activity Recommendations for Adults at Risk for Type 2 Diabetes

| Follow a Healthy Eating Pattern                                                                                                                                                                                                                                                                                                                                                                                                                                                                                           | Get Regular Physical Activity                                                                                                                                                                                                                                                                                             |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• Emphasize whole grains, legumes, nuts, fruits, and vegetables and minimize refined and processed foods</li> <li>• A variety of healthy eating patterns include:               <ul style="list-style-type: none"> <li>» Mediterranean-style</li> <li>» Low-carbohydrate</li> <li>» Vegetarian or plant-based</li> <li>» DASH (Dietary Approaches to Stop Hypertension)</li> </ul> </li> </ul>  | <ul style="list-style-type: none"> <li>• ≥150 min/week of moderate-intensity physical activity, such as brisk walking</li> <li>• May include resistance or strength training</li> <li>• Break up prolonged sedentary time</li> </ul>  |

#### Weight Management for Type 2 Diabetes Prevention


| Where to Refer                                                                                                                                                                                                                                                                                                                                                                      | What is the Diabetes Prevention Program?                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>• Refer adults with overweight or obesity who are at high risk for type 2 diabetes to a recognized diabetes prevention lifestyle change program (<a href="http://www.cdc.gov/diabetes-prevention/lifestyle-change-program/find-a-program.html">www.cdc.gov/diabetes-prevention/lifestyle-change-program/find-a-program.html</a>).</li> </ul> | <ul style="list-style-type: none"> <li>• The Diabetes Prevention Program (DPP) study demonstrated that intensive lifestyle intervention could reduce the risk of type 2 diabetes by 58% over 3 years. The two major goals of the DPP intensive lifestyle intervention were to achieve and maintain ≥7% weight loss and ≥150 min/week of moderate-intensity physical activity, such as brisk walking.</li> <li>• Technology-assisted programs using smartphones, web apps, and telehealth platforms can effectively deliver the DPP lifestyle change program, overcoming barriers, especially for people with low income and/or in rural locations.</li> <li>• DPP lifestyle change programs are covered by Medicare, Medicaid in certain states, and some commercial payors.</li> </ul>  |

# Prediabetes

- More than 1/3 American adults have prediabetes
- Identify patients with prediabetes
  - Make appropriate referrals, weight loss goals, and decide if medication tx should be offered, make plan for monitoring
    - Consider offering Metformin to:
      - Adults aged 25–59 years with a BMI  $\geq 35$  kg/m<sup>2</sup>
      - Those with higher A1C (e.g.,  $\geq 6.0\%$  [ $\geq 42$  mmol/mol])
      - Individuals with higher FPG (e.g.,  $\geq 110$  mg/dL [ $\geq 6$  mmol/L])
      - Individuals with a history of GDM

# ADA Recommendations for Suspected Type 1 Adults

- In primary care, our role is generally to recognize Type 2 Diabetes and refer patients to endocrinology when appropriate.
- Be a point of contact/coordination depending on location, but ideally not managing these patients (provider comfort level).
- For this presentation, Type 2 is the focus.




# ADA Recommend Screening Modalities for Type 2 Diabetes Management

Criteria for the diagnosis of diabetes in nonpregnant individuals

- **A1C  $\geq 6.5\%$  ( $\geq 48$  mmol/mol).** The test should be performed in a laboratory using a method that is NGSP certified and standardized to the DCCT assay.\*
- **FPG  $\geq 126$  mg/dL ( $\geq 7.0$  mmol/L).** Fasting is defined as no caloric intake for at least 8 h.\*
- **2-h PG  $\geq 200$  mg/dL ( $\geq 11.1$  mmol/L) during OGTT.** The test should be performed as described by the WHO, using a glucose load containing the equivalent of 75 g anhydrous glucose dissolved in water.
- In an individual with **classic symptoms of hyperglycemia or hyperglycemic crisis, a random plasma glucose  $\geq 200$  mg/dL ( $\geq 11.1$  mmol/L).** Random is any time of the day without regard to time since previous meal.

DCCT, Diabetes Control and Complications Trial; FPG, fasting plasma glucose; OGTT, oral glucose tolerance test; NGSP, National Glycohemoglobin Standardization Program; WHO, World Health Organization; 2-h PG, 2-h plasma glucose. \*In the absence of unequivocal hyperglycemia, diagnosis requires two abnormal results from different tests which may be obtained at the same time (e.g., A1C and FPG), or the same test at two different time points.

# What is the Diagnosis?



"Patient's understanding of their diabetes empowers them to take control of their health and well-being."



A1C of  
6.5%....  
Now what?



**DIAGNOSIS**

Name: \_\_\_\_\_ Age: \_\_\_\_\_


Doctor's name: \_\_\_\_\_

**Diabetes Mellitus**

☐ Type 1

☒ Type 2

# Decision Cycle for Person-Centered Glycemic Management in Type 2 Diabetes



The background is a solid blue color with a subtle, darker blue wavy pattern. The text is white and centered.

Facilitate Positive Health  
Behaviors and Well-Being to  
Improve Health Outcomes

## Facilitate Positive Health Behaviors and Well-Being to Improve Health Outcomes

- Building positive health behaviors and maintaining psychological well-being are foundational for achieving diabetes management goals and maximizing quality of life.
- Essential tasks to help people with diabetes achieve their health goals:
  - Refer for diabetes self-management education and support (DSMES)
  - Counsel on and support cessation of tobacco products and vaping
  - Refer for medical nutrition therapy (MNT)
  - Counsel on health behaviors and sleep quality
  - Counsel on routine physical activity
  - Support and refer to behavioral health professionals for psychosocial care



## SITTING/BREAKING UP PROLONGED SITTING

- Limit sitting. Breaking up prolonged sitting (at least every 30 min) with short regular bouts of slow walking or simple resistance exercises can improve glucose metabolism.

## STEPPING

- An increase of only 500 steps/day is associated with 2-9% decreased risk of cardiovascular morbidity and all-cause mortality.
- A 5-to 6-min brisk-intensity walk per day equates to ~4 years' greater life expectancy.

## SLEEP

Aim for consistent, uninterrupted sleep, even on weekends.

**Quantity** - Long (>8 h) and short (<6 h) sleep durations negatively impact A1C.

**Quality** - Irregular sleep results in poorer glycemic levels, likely influenced by the increased prevalence of insomnia, obstructive sleep apnea, and restless leg syndrome in people with type 2 diabetes.

**Chronotype** - Evening chronotypes (i.e., night owl: go to bed late and get up late) may be more susceptible to inactivity and poorer glycemic levels than morning chronotypes (i.e., early bird: go to bed early and get up early).

## SWEATING (MODERATE-TO-VIGOROUS ACTIVITY)

- Encourage ≥150 min/week of moderate-intensity physical activity (i.e., uses large muscle groups, rhythmic in nature) OR ≥75 min/week vigorous-intensity activity spread over ≥3 days/week, with no more than 2 consecutive days of inactivity. Supplement with two to three resistance, flexibility, and/or balance sessions.
- As little as 30 min/week of moderate-intensity physical activity improves metabolic profiles.

## Physical function/ frailty/sarcopenia

- The frailty phenotype in type 2 diabetes is unique, often encompassing obesity alongside physical frailty, at an earlier age. The ability of people with type 2 diabetes to undertake simple functional exercises in middle-age is similar to that in those over a decade older.

## STRENGTHENING

Resistance exercise (i.e., any activity that uses the person's own body weight or works against a resistance) also improves insulin sensitivity and glucose levels; activities like tai chi and yoga also encompass elements of flexibility and balance.


|                                          | Glucose/insulin | Blood pressure | A1C | Lipids | Physical function | Depression | Quality of life |
|------------------------------------------|-----------------|----------------|-----|--------|-------------------|------------|-----------------|
| SITTING/BREAKING UP PROLONGED SITTING    | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| STEPPING                                 | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| SWEATING (MODERATE-TO-VIGOROUS ACTIVITY) | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| STRENGTHENING                            | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| ADEQUATE SLEEP DURATION                  | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| GOOD SLEEP QUALITY                       | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |
| CHRONOTYPE/CONSISTENT TIMING             | ⬆               | ⬆              | ⬆   | ⬆      | ⬆                 | ⬆          | ⬆               |

## IMPACT OF PHYSICAL BEHAVIORS ON CARDIOMETABOLIC HEALTH IN PEOPLE WITH TYPE 2 DIABETES

- ⬆ Higher levels of improvement (physical function, quality of life) ⬆ Lower levels of improvement (glucose/insulin, blood pressure, A1C, lipids, depression)
- ⬆ No data available
- ⬆ Green arrows = strong evidence ⬆ Yellow arrows = medium-strength evidence ⬆ Red arrows = limited evidence

# Assessment of Glycemic Status

- A1C typically every 3, 4, or 6 months
- AM fasting goal <80-130
- 2-hour post prandial goal of <180
- CGM goal-typically TIR around 70% or greater



## Clinical Notes



### Limitations of A1C testing:

- ▶ Accuracy can be affected by conditions that affect red blood cell turnover (e.g., hemolytic and other anemias, glucose-6-phosphate dehydrogenase deficiency, recent blood transfusion, use of drugs that stimulate erythropoiesis, end-stage kidney disease, and pregnancy).
- ▶ Some hemoglobin variants can interfere with some A1C assays, although this problem has been minimized with newer assays.
- ▶ A1C cannot be used for people with sickle cell disease or other homozygous hemoglobin variants.
- ▶ A1C does not assess glycemic variability or hypoglycemia.
- ▶ Consider alternative measures, such as fructosamine and glycated albumin, when necessary.
- ▶ Assess glycemic variability using blood glucose monitoring/CGM and A1C values.

# Assessment of Glycemic Status

- Glucose Assessment via CGM: The Ambulatory Glucose Profile (AGP) Report


1. Ensure sufficient data for analysis; CGM active at least 70% of the time over 14 days is recommended.

2. Discuss individuals' daily self-management routine.


3. Ask individuals to identify and explain what they observe on the report.

4. Check overall CGM metrics:


- ✓ TBR: percentage of readings and time with glucose <70 mg/dL (<3.9 mmol/L)
- ✓ TIR: percentage of readings and time with glucose 70–180 mg/dL (3.9–10.0 mmol/L)
- ✓ TAR: percentage of reading and time with glucose >180 mg/dL (>10.0 mmol/L)
- ✓ Average glucose
- ✓ Glucose management indicator: an estimate of A1C
- ✓ Glycemic variability: expressed as percent coefficient of variation; target ≤36%



5. Identify any hypoglycemia first, and then look for hyperglycemia patterns. Review the time spent in these patterns on the overall profile and daily graphs.



6. Discuss identified patterns and ask individuals to reflect on potential causes and possible solutions.
7. If you use CGM metrics to assess glycemia, goals for nonpregnant adults or those with frailty will vary; set glycemic goals based on the general guidelines on the next page.
8. Use shared decision-making to develop an action plan.



# Taking Control of Your Diabetes (TCOYD)

- Non-profit for diabetes education by 2 endocrinologists who have had Type 1 since their teen years
- Resources for providers and patients
- Podcast
- Free CME on rapid fire CGM interpretation
- <https://tcoyd.org/cme-enduring-cgm-data/>


# Correlation Between A1C and Estimated Average Glucose (eAG)

| A1C (%) | mg/dL         | mmol/L           |
|---------|---------------|------------------|
| 5       | 97 (76–120)   | 5.4 (4.2–6.7)    |
| 6       | 126 (100–152) | 7.0 (5.5–8.5)    |
| 7       | 154 (123–185) | 8.6 (6.8–10.3)   |
| 8       | 183 (147–217) | 10.2 (8.1–12.1)  |
| 9       | 212 (170–249) | 11.8 (9.4–13.9)  |
| 10      | 240 (193–282) | 13.4 (10.7–15.7) |
| 11      | 269 (217–314) | 14.9 (12.0–17.5) |
| 12      | 298 (240–347) | 16.5 (13.3–19.3) |

A calculator for converting A1C results into eAG, in either mg/dL or mmol/L, is available at [professional.diabetes.org/eAG](http://professional.diabetes.org/eAG).

Data in parentheses are a 95% CI. Adapted from Nathan DM, Kuenen J, Borg R, Zheng H, Schoenfeld D; A1c-Derived Average Glucose Study Group. Translating the A1C assay into estimated average glucose values. *Diabetes Care* 2008;31:1473–1478.

# Establishing Personalized A1C/TIR Goals



# Hypoglycemia and Hyperglycemia


- Educate patients on symptoms of each
- Include an individualized plan as to what to do in each situation
- Review impact of medication/awareness





Consider What Diabetes  
Technology Will Be Used  
and Prescribed

## Consider What Diabetes Technology Will Be Used and Prescribed

- Mainly use traditional finger sticks and CGMs in primary care
- Consider cost and insurance coverage

|                                                                                                                                                           |                                                                                                                                                                            |                                                                                                                               |                                                                                                                                                            |                                                                                                                                                                                                                     |
|-----------------------------------------------------------------------------------------------------------------------------------------------------------|----------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
|                                                                          |                                                                                           |                                             |                                                                         |                                                                                                                                  |
| <p><b>Insulin pumps</b> (also called continuous subcutaneous insulin infusion [CSII] systems) are insulin delivery devices that are worn on the body.</p> | <p><b>Connected insulin pens and pen caps</b> are insulin delivery pens or related devices that can record and/or send insulin dose data and may also calculate doses.</p> | <p><b>Continuous glucose monitoring (CGM) systems</b> and <b>glucose meters</b> are devices to monitoring glucose levels.</p> | <p><b>Automated insulin delivery (AID) systems</b> connect a CGM system and an insulin pump with a control algorithm to deliver insulin automatically.</p> | <p><b>Diabetes self-management support software</b> includes apps or online platforms that are intended to treat a medical or psychological condition or assist with data management or lifestyle modification.</p> |





Consider Specific Goals and Plans for  
Patients Who are Overweight or Obese

Overall Weight Management Discussions

# Consider Specific Goals and Plans for Patients who are Overweight or Obese


## Overall Weight Management Discussions

Nutrition, Physical Activity, and Behavioral Therapy for People With Overweight and Obesity

### Recommendations:

- Nutrition, physical activity, and behavioral therapy are recommended for people with Type 2 Diabetes and overweight or obesity to achieve both weight and health outcome goals.
- Frequent counseling ( $\geq 16$  sessions in 6 months) focusing on nutrition, exercise, and behavioral strategies to achieve a 500–750 kcal/day energy deficit, is beneficial for weight loss and recommended if available.
- Long-term ( $\geq 1$  year) weight maintenance programs are advised for those meeting weight loss goals, offering monthly support, at least weekly body weight monitoring, self-monitoring strategies, and regular physical activity (200–300 min/week).
- Short-term, structured very-low-calorie diets (800–1,000 kcal/day) should be reserved for carefully selected individuals, be prescribed by trained practitioners in medical settings with close monitoring, and include counseling for long-term weight maintenance.

When developing a care plan, consider systemic, structural, cultural, and socioeconomic factors that may affect nutrition patterns and food choices, such as food insecurity and hunger, access to healthful food options, and other social determinants of health.



| Medication                                                                                                     | Common side effects                                                                                             | Possible safety concerns/considerations                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                            |
|----------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| Short-term treatment (12 weeks)                                                                                |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Sympathomimetic amine anorectic                                                                                |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Phentermine<br>               | Dry mouth, insomnia, dizziness, irritability, increased blood pressure, and elevated heart rate                 | <ul style="list-style-type: none"><li>• Contraindicated for use in combination with monoamine oxidase inhibitors</li><li>• Caution with cardiovascular disease</li><li>• Do not use if at high risk for glaucoma due to risk of acute angle-closure glaucoma</li></ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                             |
| Long-term treatment (52–72 weeks)*                                                                             |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Lipase inhibitor                                                                                               |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Orlistat<br>                  | Abdominal pain, flatulence, and fecal urgency                                                                   | <ul style="list-style-type: none"><li>• Potential malabsorption of fat-soluble vitamins (A, D, E, and K) and of certain medications (e.g., cyclosporine, thyroid hormone, and anticonvulsants)</li><li>• Rare cases of severe liver injury reported</li><li>• Cholelithiasis</li><li>• Nephrolithiasis</li></ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                   |
| Sympathomimetic amine anorectic/antiepileptic combination                                                      |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Phentermine/topiramate ER<br> | Constipation, paresthesia, insomnia, nasopharyngitis, xerostomia, increased blood pressure, and nephrolithiasis | <ul style="list-style-type: none"><li>• Contraindicated for use in combination with monoamine oxidase inhibitors</li><li>• Birth defects</li><li>• Cognitive impairment</li><li>• Caution with cardiovascular disease</li><li>• Do not use if at high risk for glaucoma due to risk of acute angle-closure glaucoma</li></ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                      |
| Opioid antagonist/antidepressant combination                                                                   |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Naltrexone/bupropion ER<br>   | Constipation, nausea, headache, xerostomia, insomnia, elevated heart rate and blood pressure                    | <ul style="list-style-type: none"><li>• Contraindicated in people with unmanaged hypertension and/or seizure disorders</li><li>• Contraindicated for use with chronic opioid therapy</li><li>• Acute angle-closure glaucoma</li><li>• <b>BLACK BOX WARNING:</b> Risk of suicidal behavior/ideation in people &lt;24 years of age who have depression</li></ul>                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                     |
| Glucagon-like peptide 1 (GLP-1) receptor agonists (RAs)                                                        |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Liraglutide<br>              | Gastrointestinal side effects (nausea, vomiting, diarrhea, and esophageal reflux)                               | <ul style="list-style-type: none"><li>• Hypoglycemia (with concomitant use of insulin or sulfonylurea)</li><li>• Pancreatitis has been reported in clinical trials, but causality has not been established. Discontinue if pancreatitis is suspected</li><li>• Use caution in people with kidney disease when initiating dose due to increased risk of gastrointestinal side effects and potential risk of acute kidney injury from dehydration</li><li>• May cause cholelithiasis and gallstone-related complications</li><li>• Gastrointestinal disorders (severe constipation and small-bowel obstruction/ileus progression)</li><li>• Monitor for potential consequences of delayed absorption of oral medications</li><li>• May cause injection site reactions</li><li>• May cause elevated heart rate</li><li>• <b>BLACK BOX WARNING:</b> Risk of thyroid C-cell tumors in rodents; human relevance not determined; do not use in individuals with a personal or family history of medullary thyroid cancer or multiple endocrine neoplasia type 2</li></ul> |
| Semaglutide<br>             |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Dual glucose-dependent insulinotropic polypeptide and GLP-1 RA                                                 |                                                                                                                 |                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                                    |
| Tirzepatide<br>             | Gastrointestinal side effects (nausea, vomiting, diarrhea, and esophageal reflux)                               | <ul style="list-style-type: none"><li>• Same as for GLP-1 RAs</li><li>• Monitor effects of oral medications with narrow therapeutic index (warfarin) or whose efficacy is dependent on threshold concentration</li><li>• Advise females using oral contraceptives to switch to a nonoral contraceptive or add a barrier method of contraception for 4 weeks after initiation and for 4 weeks after each dose escalation</li><li>• <b>BLACK BOX WARNING:</b> Risk of thyroid C-cell tumors in rodents; human relevance not determined; do not use in individuals with a personal or family history of medullary thyroid cancer or multiple endocrine neoplasia type 2</li></ul>                                                                                                                                                                                                                                                                                                                                                                                     |

**Table 8.2—Median monthly (30-day) AWP and NADAC of maximum or maintenance dose of obesity pharmacotherapies**

| Medication name                                                   | Typical adult maintenance dose           | AWP (median and range for 30-day supply) | NADAC (median and range for 30-day supply) |
|-------------------------------------------------------------------|------------------------------------------|------------------------------------------|--------------------------------------------|
| Sympathomimetic amine anorectic: approved for short-term use only |                                          |                                          |                                            |
| Phentermine                                                       | 8–37.5 mg daily                          | \$43 (\$3–\$58)*                         | \$2 (\$2–\$3)*                             |
| Lipase inhibitor                                                  |                                          |                                          |                                            |
| Orlistat                                                          | 60 mg t.i.d. (OTC)<br>120 mg t.i.d. (Rx) | \$58 (\$41–\$90)<br>\$675 (\$520–\$781)  | NA<br>\$514 (\$416–\$611)                  |
| Sympathomimetic amine anorectic/antiepileptic combination         |                                          |                                          |                                            |
| Phentermine/topiramate ER                                         | 7.5 mg/46 mg daily                       | \$238 (\$238–\$251)                      | NA                                         |
| Opioid antagonist/antidepressant combination                      |                                          |                                          |                                            |
| Naltrexone/bupropion ER                                           | 16 mg/180 mg b.i.d.                      | \$750                                    | NA                                         |
| GLP-1 receptor agonist                                            |                                          |                                          |                                            |
| Liraglutide                                                       | 3 mg daily                               | \$1,619                                  | \$1,303                                    |
| Semaglutide                                                       | 2.4 mg once weekly                       | \$1,619                                  | \$1,302                                    |
| Dual GIP and GLP-1 receptor agonist                               |                                          |                                          |                                            |
| Tirzepatide                                                       | 5, 10, or 15 mg once weekly              | \$1,304**                                | \$1,022                                    |

The costs listed in this table are representative of costs at a national level. These costs may not be representative of an individual's cost and do not account for medication coverage or available discounts. AWP, average wholesale price; b.i.d., twice daily; ER, extended release; GIP, glucose-dependent insulintropic polypeptide; GLP-1, glucagon-like peptide 1; NA, data not available; NADAC, National Average Drug Acquisition Cost; OTC, over the counter; Rx, prescription; t.i.d., three times daily. AWP and NADAC prices are for a 30-day supply of maximum or maintenance dose as of 15 July 2025 ([190,191](#)). \*Data are for 37.5 mg q.d. dose. \*\*Pricing listed for tirzepatide is for the pens only, not the cost of vials available from the manufacturer.

# Pharmacologic Approach and Plan

# Pharmacologic Approach and Plan

- Make sure your patient knows you are their champion and have bought into their treatments as best as possible.
- Identify and address any barriers while in office.
- Look at comorbidities and overall A1C goals to decide.

| Intervention                                                                                      | Expected decrease in A1C with monotherapy (%) | Advantages                                                                                                                                                                   | Disadvantages                                                                                                                                        |
|---------------------------------------------------------------------------------------------------|-----------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------|
| <b>Initial therapy</b>                                                                            |                                               |                                                                                                                                                                              |                                                                                                                                                      |
| Lifestyle change to decrease weight and increase activity                                         | 1 to 2                                        | Broad benefits                                                                                                                                                               | May be insufficient for most within first year due to limited adherence, inadequate weight loss, and/or weight regain                                |
| Metformin                                                                                         | 1 to 2                                        | Weight loss to weight neutral, low cost, low risk of hypoglycemia, generally well tolerated, well-studied in combination with other agents                                   | GI side effects, contraindicated with impaired kidney function (eGFR <30 mL/min/1.73 m <sup>2</sup> )*                                               |
| <b>Additional therapy<sup>‡</sup></b>                                                             |                                               |                                                                                                                                                                              |                                                                                                                                                      |
| Insulin (usually with a single daily injection of intermediate- or long-acting insulin initially) | 1.5 to 3.5                                    | No dose limit, rapidly effective, improved lipid profile                                                                                                                     | Hypoglycemia, may require more than 1 daily injection, requires home glucose monitoring, weight gain, analogs are expensive                          |
| Dual GLP-1 and GIP receptor agonist (once-weekly injections)                                      | 2 to 2.5                                      | Weight loss                                                                                                                                                                  | Requires injection, frequent GI side effects, very expensive                                                                                         |
| GLP-1 receptor agonist (oral or daily to weekly injections)                                       | 0.5 to 2                                      | Weight loss, reduction in major adverse cardiovascular events (liraglutide, subcutaneous semaglutide, dulaglutide) in patients with established CVD and at high risk for CVD | Most agents require injection, frequent GI side effects, very expensive                                                                              |
| SGLT2 inhibitor                                                                                   | 0.5 to 0.7                                    | Weight loss, reduction in systolic blood pressure, reduced heart failure and cardiovascular mortality, improved kidney outcomes in patients with nephropathy                 | Mycotic genital infection, DKA; SGLT2 inhibitors have also been associated with urinary tract infections, bone fractures, and lower limb amputations |
| Sulfonylurea (shorter-acting agents preferred)                                                    | 1 to 2                                        | Rapidly effective                                                                                                                                                            | Hypoglycemia (especially with glyburide/glibenclamide or chlorpropamide), weight gain                                                                |
| Glinide                                                                                           | 0.5 to 1.5 <sup>Δ</sup>                       | Rapidly effective                                                                                                                                                            | Hypoglycemia, weight gain, may require 3 times daily dosing                                                                                          |
| Pioglitazone                                                                                      | 0.5 to 1.4                                    | Improved lipid profile, potential decrease in MI and stroke                                                                                                                  | Fluid retention, HF, weight gain, bone fractures, and bladder cancer; side effects minimized at doses of 15 to 30 mg                                 |
| DPP-4 inhibitor                                                                                   | 0.5 to 0.8                                    | Weight neutral                                                                                                                                                               | Possible increased risk of HF with saxagliptin, expensive                                                                                            |
| Alpha-glucosidase inhibitor                                                                       | 0.5 to 0.8                                    | Weight neutral                                                                                                                                                               | Frequent GI side effects limit use, 3 times daily dosing                                                                                             |

A1C: glycated hemoglobin; CVD: cardiovascular disease; DKA: diabetic ketoacidosis; DPP-4: dipeptidyl peptidase 4; eGFR: estimated glomerular filtration rate; GI: gastrointestinal; GIP: glucose-dependent insulinotropic polypeptide; GLP-1: glucagon-like peptide 1; HF: heart failure; MI: myocardial infarction; SGLT2: sodium-glucose cotransporter 2.

\* Initiation is contraindicated with eGFR <30 mL/min/1.73 m<sup>2</sup> and not recommended with eGFR 30 to 45 mL/min/1.73 m<sup>2</sup>.

‡ The order of listing of additional therapies does not indicate a preferred order of selection. The choice of additional therapy should be based on criteria discussed in the UpToDate topics on the management of hyperglycemia in diabetes mellitus.


Δ Repaglinide is more effective in lowering A1C than nateglinide.

Adapted from:

1. Nathan DM, Buse JB, Davidson MB, et al. Medical Management of Hyperglycemia in Type 2 Diabetes: A Consensus Algorithm for the Initiation and Adjustment of Therapy: A consensus statement of the American Diabetes Association and the European Association for the Study of Diabetes. Diabetes Care 2009; 32:193.


With additional data from:

| <p>❓ Which therapy should I start first?</p>                                                                                                                                                                                                                                                                                                                                    | <p>❓ When should I start insulin first?</p>                                                                                                                                                                                                                                                                                                                           | <p>❓ Can I use combination insulin and noninsulin injectable therapy?</p>                                                                                                                                                                                                                                                                                                          | <p>❓ When would I use combination insulin and noninsulin injectable therapy?</p>                                                                                                           | <p>❓ When should I modify a patient's injectable therapy?</p>                                                                                                                                                           |
|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-----------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <ul style="list-style-type: none"> <li>✔ Treatment with a glucagon-like peptide 1 (GLP-1) receptor agonist (RA) or a dual glucose-dependent insulintropic polypeptide (GIP)/GLP-1 RA is preferred before insulin therapy because of its ability to achieve both glycemic and weight management goals.</li> <li>✔ Some GLP-1 RAs also provide cardiovascular benefit.</li> </ul> | <ul style="list-style-type: none"> <li>✔ If there are symptoms of hyperglycemia (polyuria or polydipsia) or there is evidence of catabolism (unexpected weight loss).</li> <li>✔ When A1C or blood glucose levels are very high (A1C &gt;10% [<math>&gt;86</math> mmol/mol] or blood glucose <math>\geq 300</math> mg/dL [<math>\geq 16.7</math> mmol/L]).</li> </ul> | <ul style="list-style-type: none"> <li>✔ Yes; combination therapy with insulin and a noninsulin injectable is recommended for greater glycemic effectiveness and beneficial effects on weight and hypoglycemia risk.</li> <li>✔ If insulin is already being used, insulin dosing should be reassessed upon addition or dose escalation of a GLP-1 or dual GIP/GLP-1 RA.</li> </ul> | <ul style="list-style-type: none"> <li>✔ Consider combination insulin and GLP-1 or dual GIP/GLP-1 RA therapy when individualized goals are not met using either one separately.</li> </ul> | <ul style="list-style-type: none"> <li>✔ Intensify or deintensify therapy when an individual is not meeting treatment goals, including management of hyperglycemia and weight and avoidance of hypoglycemia.</li> </ul> |



# HEALTHY LIFESTYLE BEHAVIORS; DIABETES SELF-MANAGEMENT EDUCATION AND SUPPORT; SOCIAL DETERMINANTS OF HEALTH

To avoid  
therapeutic  
inertia, reassess  
and modify  
treatment  
regularly  
(3-6 months)



\* In people with HF, CKD, established CVD, or multiple risk factors for CVD, the decision to use a GLP-1 RA or SGLT2i with proven benefit should be made irrespective of background use of metformin or A1C.

† ASCVD: Defined differently across CVOTs but all included individuals with established CVD (e.g., MI, stroke, and arterial revascularization procedure) and variably included conditions such as transient ischemic attack, unstable angina, amputation, and symptomatic or asymptomatic coronary artery disease. Indicators of high risk: While definitions vary, most comprise ≥55 years of age with two or more additional risk factors (including obesity, hypertension, smoking, dyslipidemia, or albuminuria).

‡ A strong recommendation is warranted for people with CVD and a weaker recommendation for those with indicators of high-risk CVD. Moreover, a higher absolute risk reduction and thus lower numbers needed to treat are seen at higher levels of baseline risk and should be factored into the shared decision-making process. See text for details.

§ For GLP-1 RAs, CVOTs demonstrate their efficacy in reducing composite MACE, CV death, all-cause mortality, MI, stroke, and kidney end points in individuals with T2D with established or high risk of CVD. One kidney outcome trial demonstrated benefit in reducing persistent eGFR reduction and CV death for a GLP-1 RA in individuals with CKD and T2D.

¶ For SGLT2is, CV and kidney outcomes trials demonstrate their efficacy in reducing the risks of composite MACE, CV death, all-cause mortality, MI, HF, and kidney outcomes in individuals with T2D and established or high risk of CVD.

⌘ Low-dose pioglitazone may be better tolerated and similarly effective as higher doses.

# Name That Price... \$

**Table 9.3—Median monthly (30-day) AWP and NADAC of maximum approved daily dose of noninsulin glucose-lowering agents in the U.S.**

| Class                          | Compound        | Dosage strength/<br>product (if applicable) | Maximum approved<br>daily dose† | Median AWP<br>(min, max)* | Median NADAC<br>(min, max)* |
|--------------------------------|-----------------|---------------------------------------------|---------------------------------|---------------------------|-----------------------------|
| Biguanides                     | • Metformin     | 500 mg (IR)                                 | 2,000 mg                        | \$85 (\$3, \$216)         | \$2                         |
|                                |                 | 500 mg (ER)                                 | 2,000 mg                        | \$89 (\$5, \$6,719)       | \$3                         |
|                                |                 | 850 mg (IR)                                 | 2,550 mg                        | \$108 (\$3, \$189)        | \$2                         |
|                                |                 | 1,000 mg (IR)                               | 2,000 mg                        | \$87 (\$2, \$144)         | \$1                         |
|                                |                 | 1,000 mg (ER)                               | 2,000 mg                        | \$1,884 (\$242, \$7,214)  | \$26 (\$24, \$28)           |
|                                |                 | 500 mg (Sol)                                | 2,000 mg                        | \$810 (\$810, \$1,478)    | \$417                       |
| Sulfonylureas (2nd generation) | • Glimepiride   | 4 mg                                        | 8 mg                            | \$73 (\$71, \$198)        | \$2                         |
|                                |                 | 10 mg (IR)                                  | 40 mg                           | \$72 (\$67, \$91)         | \$6                         |
|                                |                 | 10 mg (XL/ER)                               | 20 mg                           | \$48                      | \$9                         |
|                                | • Glyburide     | 6 mg (micronized)                           | 12 mg                           | \$54 (\$48, \$71)         | \$13                        |
|                                |                 | 5 mg                                        | 20 mg                           | \$88 (\$63, \$432)        | \$7                         |
|                                |                 |                                             |                                 |                           |                             |
| Thiazolidinedione              | • Pioglitazone  | 45 mg                                       | 45 mg                           | \$348 (\$7, \$349)        | \$4                         |
| α-Glucosidase inhibitors       | • Acarbose      | 100 mg                                      | 300 mg                          | \$106 (\$104, \$378)      | \$22                        |
|                                | • Miglitol      | 100 mg                                      | 300 mg                          | \$294 (\$241, \$346)      | \$183                       |
| Meglitinides                   | • Nateglinide   | 120 mg                                      | 360 mg                          | \$104                     | \$16                        |
|                                | • Repaglinide   | 2 mg                                        | 16 mg                           | \$878 (\$799, \$1,728)    | \$28                        |
| DPP-4 inhibitors               | • Alogliptin    | 25 mg                                       | 25 mg                           | \$234                     | \$143                       |
|                                | • Linagliptin   | 5 mg                                        | 5 mg                            | \$630                     | \$504                       |
|                                | • Saxagliptin   | 5 mg                                        | 5 mg                            | \$524 (\$523, \$524)      | \$179                       |
|                                | • Sitagliptin   | 100 mg                                      | 100 mg                          | \$341                     | \$303                       |
|                                |                 | 25 mg/mL                                    | 100 mg                          | \$354                     | NA                          |
| SGLT2 inhibitors               | • Bexagliflozin | 20 mg                                       | 20 mg                           | \$47                      | NA                          |
|                                | • Canagliflozin | 300 mg                                      | 300 mg                          | \$718                     | \$575                       |
|                                | • Dapagliflozin | 10 mg                                       | 10 mg                           | \$664                     | \$345                       |
|                                | • Empagliflozin | 25 mg                                       | 25 mg                           | \$629                     | \$604                       |
|                                | • Ertugliflozin | 15 mg                                       | 15 mg                           | \$428                     | \$343                       |
| GLP-1 RAs                      | • Dulaglutide   | 4.5 mg pen                                  | 4.5 mg†                         | \$1,185                   | \$953                       |
|                                | • Liraglutide   | 18 mg/3 mL pen                              | 1.8 mg                          | \$929 (\$845, \$929)      | \$577                       |
|                                | • Semaglutide   | 2 mg pen                                    | 2 mg†                           | \$1,197                   | \$966                       |
|                                |                 | 14 mg (tablet)                              | 14 mg                           | \$1,197                   | \$965                       |
| Dual GIP and GLP-1 RA          | • Tirzepatide   | 15 mg pen                                   | 15 mg†                          | \$1,296                   | \$1,041                     |
| Bile acid sequestrant          | • Colesevelam   | 625 mg tabs                                 | 3.75 g                          | \$692 (\$674, \$712)      | \$56                        |
|                                |                 | 3.75 g suspension                           | 3.75 g                          | \$674 (\$673, \$675)      | \$93                        |
| Dopamine-2 agonist             | • Bromocriptine | 0.8 mg                                      | 4.8 mg                          | \$1,188                   | \$957                       |

AWP, average wholesale price; DPP-4, dipeptidyl peptidase 4; ER and XL, extended release; GIP, glucose-dependent insulinotropic polypeptide; GLP-1 RA, glucagon-like peptide 1 receptor agonist; IR, immediate release; max, maximum; min, minimum; NA, data not available; NADAC, National Average Drug Acquisition Cost; SGLT2, sodium-glucose cotransporter 2; Sol, solution. AWP and NADAC prices as of 15 July 2025.

\*Calculated for 30-day supply (AWP [222] or NADAC [223] unit price × number of doses required to provide maximum approved daily dose × 30 days); median AWP or NADAC listed alone when only one product and/or price. †Used to calculate median AWP and NADAC (min, max); generic prices used, if available commercially. ‡Administered once weekly.

# Name That Price...(cont.) \$

**Table 9.4—Median cost of insulin products in the U.S. calculated as AWP and NADAC per 1,000 units of specified dosage form/product**

| Insulins                           | Compounds                                | Dosage form/product      | Median AWP (min, max)* | Median NADAC (min, max)* |
|------------------------------------|------------------------------------------|--------------------------|------------------------|--------------------------|
| Rapid-acting                       | • Aspart                                 | U-100 vial               | \$87†                  | \$70†                    |
|                                    |                                          | U-100 cartridge          | \$107†                 | \$86†                    |
|                                    |                                          | U-100 prefilled pen      | \$112†                 | \$90†                    |
|                                    | • Aspart biosimilars#                    | U-100 vial               | \$83                   | NA                       |
|                                    |                                          | U-100 prefilled pen      | \$107                  | NA                       |
|                                    | • Aspart ("faster acting product")       | U-100 vial               | \$347                  | \$278                    |
|                                    |                                          | U-100 cartridge          | \$430                  | \$344                    |
|                                    |                                          | U-100 prefilled pen      | \$447                  | \$358                    |
|                                    | • Glulisine                              | U-100 vial               | \$102                  | \$82                     |
|                                    |                                          | U-100 prefilled pen      | \$132                  | \$105                    |
|                                    | • Inhaled insulin                        | Inhalation cartridges    | \$1,578                | \$1,265                  |
|                                    | • Lispro                                 | U-100 vial               | \$30†                  | \$24†                    |
|                                    |                                          | U-100 cartridge          | \$123                  | \$98                     |
|                                    |                                          | U-100 prefilled pen      | \$127†                 | \$102†                   |
|                                    | • Lispro-aabc                            | U-200 prefilled pen      | \$424                  | \$339                    |
|                                    |                                          | U-100 vial               | \$330                  | \$263                    |
|                                    |                                          | U-100 prefilled pen      | \$424                  | \$339                    |
|                                    | • Lispro follow-on product               | U-200 prefilled pen      | \$424                  | \$339                    |
|                                    |                                          | U-100 vial               | \$118                  | \$110                    |
|                                    |                                          | U-100 prefilled pen      | \$151                  | \$121                    |
| Short-acting                       | • Human regular                          | U-100 vial               | \$56 (\$54, \$58)†     | \$44 (\$43, \$46)†       |
|                                    |                                          | U-100 prefilled pen      | \$73                   | \$58                     |
| Intermediate-acting                | • Human NPH                              | U-100 vial               | \$58 (\$54, \$58)†     | \$45 (\$43, \$46)†       |
|                                    |                                          | U-100 prefilled pen      | \$93 (\$73, \$113)     | \$74 (\$58, \$91)        |
| Concentrated human regular insulin | • U-500 human regular insulin            | U-500 vial               | \$178                  | \$143                    |
|                                    |                                          | U-500 prefilled pen      | \$230                  | \$183                    |
| Long-acting                        | • Degludec                               | U-100 vial               | \$142†                 | \$114†                   |
|                                    |                                          | U-100 prefilled pen      | \$142†                 | \$114†                   |
|                                    |                                          | U-200 prefilled pen      | \$142†                 | \$114†                   |
|                                    | • Glargine                               | U-100 vial               | \$77                   | \$62                     |
|                                    |                                          | U-100 prefilled pen      | \$77                   | \$62                     |
|                                    |                                          | U-300 prefilled pen      | \$152                  | \$122†                   |
|                                    | • Glargine biosimilar/follow-on products | U-100 vial               | \$76†                  | \$61†                    |
|                                    |                                          | U-100 prefilled pen      | \$74 (\$74,† \$261)    | \$59 (\$59,† \$209)      |
| Premixed insulin products          | • Aspart 70/30                           | U-100 vial               | \$87†‡                 | \$69†‡                   |
|                                    |                                          | U-100 prefilled pen      | \$112†‡                | \$90†‡                   |
|                                    | • Lispro 50/50                           | U-100 vial               | \$102                  | NA                       |
|                                    |                                          | U-100 prefilled pen      | \$127                  | \$102                    |
|                                    | • Lispro 75/25                           | U-100 vial               | \$102                  | \$82                     |
|                                    |                                          | U-100 prefilled pen      | \$127†                 | \$102†                   |
|                                    | • NPH/regular 70/30                      | U-100 vial               | \$56 (\$54, \$58)      | \$45 (\$43, \$46)        |
|                                    |                                          | U-100 prefilled pen      | \$93 (\$73, \$113)‡    | \$74 (\$58, \$90)‡       |
| Premixed insulin/GLP-1 RA products | • Degludec/liraglutide                   | 100/3.6 mg prefilled pen | \$1,073                | \$859                    |
|                                    | • Glargine/lixisenatide                  | 100/33 mg prefilled pen  | \$713                  | \$571                    |

AWP, average wholesale price; GLP-1 RA, glucagon-like peptide 1 receptor agonist; NA, data not available; NADAC, national average drug acquisition cost. AWP (222) and NADAC (223) prices as of 15 July 2025. \*AWP or NADAC calculated as in Table 9.3. †Unbranded product prices used when available. ‡AWP and NADAC data presented do not include human insulins (approximately \$25/vial or \$43/box of 5 pens) or select analog insulins (approximately \$73/vial or \$86/box of 5 pens) available at Walmart; median listed alone when only one product and/or price. #Pricing for aspart-xjhz not available on 15 July 2025.

# Insulin Considerations

- Provide or refer people with diabetes and caregivers for education about injection technique and timing and problem-solving for issues related to insulin therapy (e.g., hypoglycemia, missed or incorrect doses, and dose adjustments).
- Ensure that individuals have all supplies necessary for injections (e.g., pen needles for insulin pens or appropriate syringes for insulin dose size or concentration when using insulin in vials) and glucose monitoring.
- Evaluate individuals with Type 2 Diabetes to determine whether they are candidates for GLP-1 or dual GIP/GLP-1 RA therapy.
- Evaluate all people on insulin therapy to determine whether they could benefit from continuous glucose monitoring.
- Ensure that people on insulin therapy have the education and supplies needed to prevent and treat hypoglycemia, including glucagon, glucose monitoring supplies, and appropriate sources of carbohydrates to treat low glucose levels.
- Schedule timely and routine follow-up visits to reassess therapy and adjust care plans to avoid therapeutic inertia.

# Cardiovascular Disease and Risk Management






Figure 10.1—Multifactorial approach to reduction in risk of diabetes complications.

## Recommendations for the Treatment of Confirmed Hypertension in Nonpregnant People With Diabetes





## Who to Screen?



### Coronary Artery Disease

Routine screening is not recommended for asymptomatic individuals.

If signs or symptoms of cardiac or associated vascular disease, or electrocardiogram abnormalities are present, screen using routine methods.

### Heart Failure

Screen all adults with diabetes, which increases risks for asymptomatic (stage B) and symptomatic (stage C) HF.

Measure BNP or N-terminal pro-BNP. Echocardiography is recommended for those with abnormal BNP levels.

### Peripheral Artery Disease

Screen individuals with diabetes and age  $\geq 65$  years, any microvascular disease, foot complications, or end-stage organ damage from diabetes, if a PAD diagnosis would change management. Consider screening anyone with a diabetes duration  $\geq 10$  years.

Screen with ankle-brachial index testing.

Recommendations to reduce the risk of symptomatic heart failure in people with diabetes

ACEi or ARB and  $\beta$ -blocker for hypertension, or recent history of myocardial infarction/acute coronary syndrome, or reduced LVEF  $\leq 40\%$


If high risk for or established CVD

SGLT2 inhibitor

If CKD

SGLT2 or SGLT1/2 inhibitor

Finerenone



# Lipids

- How often should you check lipids?
- For people with prediabetes or diabetes not on lipid-lowering therapy, check at diagnosis and at least annually thereafter.
- Check lipids at initiation of lipid-lowering therapy, 4–12 weeks after initiation or dose changes, and annually thereafter.



## ASCVD Primary Prevention

Lipid management for primary prevention of atherosclerotic cardiovascular disease events in people with diabetes in addition to healthy behavior modification




**Figure 10.3**—Recommendations for primary prevention of atherosclerotic cardiovascular disease (ASCVD) in people with diabetes using cholesterol-lowering therapy. \*ASCVD risk factors include older age, hypertension, dyslipidemia, smoking, chronic kidney disease, or obesity. Adapted from "Standards of Care in Diabetes—2024 Abridged for Primary Care Professionals" (315).

# Lipids (cont.)

## ASCVD Secondary Prevention

Use lifestyle and high-intensity statin therapy to reduce LDL cholesterol by  $\geq 50\%$  from baseline to a goal of  $< 55$  mg/dL ( $< 1.4$  mmol/L).



Add ezetimibe or a PCSK9-directed therapy with demonstrated benefit if LDL cholesterol goals are not met on maximum tolerated statin therapy.



Use an alternative lipid-lowering treatment for those who are statin intolerant:

- PCSK9 inhibitor with monoclonal antibody treatment
- Bempedoic acid
- PCSK9 inhibitor with siRNA inclisiran

# HF and DM




**Figure 10.5**—Overview of recommendations for the prevention and treatment of symptomatic heart failure in people with diabetes. ACEi, ACE inhibitor; ARB, angiotensin receptor blocker; ARNI, angiotensin receptor neprilysin inhibitor; BNP, B-type natriuretic peptide; NT-proBNP, N-terminal pro-B-type natriuretic peptide; CKD, chronic kidney disease; CVD, cardiovascular disease; EF, ejection fraction; eGFR, estimated glomerular filtration rate; GDMT, guideline-directed medical therapy; GIP/GLP-1 RA, glucose-dependent insulintropic polypeptide and glucagon-like peptide-1 receptor agonist; GLP-1 RA, glucagon-like peptide-1 receptor agonist; HF, heart failure; HFmrEF, heart failure with mildly reduced ejection fraction; HFpEF, heart failure with preserved ejection fraction; HFrEF, heart failure with reduced ejection fraction; HTN, hypertension; MI, myocardial infarction; MRA, mineralocorticoid receptor antagonist; nsMRA, nonsteroidal mineralocorticoid receptor antagonist; SGLT1/2i, sodium-glucose cotransporter 1 and 2 inhibitor; SGLT2i, sodium-glucose cotransporter 2 inhibitor; TTE, transthoracic echocardiography; UACR, urine albumin-to-creatinine ratio.

# Chronic Kidney Disease and Diabetes Management

# Chronic Kidney Disease (CKD) and Diabetes Management (DM)



## Who?

- ✓ Everyone with type 2 diabetes
- ✓ Everyone with type 1 diabetes for  $\geq 5$  years



## How?

- ✓ Urinary albumin-to-creatinine ratio (UACR)
- ✓ Estimated glomerular filtration rate (eGFR)



## How often?

Annually

### Monitoring Established CKD

**How?** UACR and eGFR. Use the CKD Epidemiology Collaboration's CKD-EPI Refit equation, which eliminates race as a variable, as the eGFR formula for all individuals.

**How often?** One to four times per year, depending on the stage of the disease.



- Reduces cardiovascular risks
- Improves patient outcomes

### Classification of CKD

CKD is classified based on:

- GFR (G)
- Albuminuria (A)

|                                                                       |     |                                  |           | Albuminuria categories     |                             |                                      |
|-----------------------------------------------------------------------|-----|----------------------------------|-----------|----------------------------|-----------------------------|--------------------------------------|
|                                                                       |     |                                  |           | Description and range      |                             |                                      |
|                                                                       |     |                                  |           | A1                         | A2                          | A3                                   |
|                                                                       |     |                                  |           | Normal to mildly increased | Moderately increased        | Severely increased                   |
|                                                                       |     |                                  |           | <30 mg/g<br><3 mg/mmol     | 30-299 mg/g<br>3-29 mg/mmol | $\geq 300$ mg/g<br>$\geq 30$ mg/mmol |
| GFR categories (mL/min/1.73 m <sup>2</sup> )<br>Description and range | G1  | Normal or high                   | $\geq 90$ | Screen 1                   | Treat 1                     | Treat and refer 2                    |
|                                                                       | G2  | Mildly decreased                 | 60-89     | Screen 1                   | Treat 1                     | Treat and refer 2                    |
|                                                                       | G3a | Mildly to moderately decreased   | 45-59     | Treat 1                    | Treat 2                     | Treat and refer 3                    |
|                                                                       | G3b | Moderately to severely decreased | 30-44     | Treat 2                    | Treat and refer 3           | Treat and refer 3                    |
|                                                                       | G4  | Severely decreased               | 15-29     | Treat and refer 3          | Treat and refer 3           | Treat and refer 4+                   |
|                                                                       | G5  | Kidney failure                   | <15       | Treat and refer 4+         | Treat and refer 4+          | Treat and refer 4+                   |

■ Low risk (if no other markers of kidney disease, no CKD) ■ Moderately increased risk ■ High risk ■ Very high risk

# CKD and DM




**Figure 11.2**—Holistic approach for improving outcomes in people with diabetes and CKD. Icons presented indicate the following benefits: BP cuff, BP lowering; glucose meter, glucose lowering; heart, cardioprotection; kidney, kidney protection; scale, weight management. eGFR is presented in units of mL/min/1.73 m<sup>2</sup>. †ACEi or ARB (at maximal tolerated doses) should be first-line therapy for hypertension when albuminuria is present. Otherwise, dihydropyridine calcium channel blocker or diuretic can also be considered; all three classes are often needed to attain BP targets. ‡Finerenone is currently the only nsMRA with proven clinical kidney and cardiovascular benefits. ACEi, angiotensin-converting enzyme inhibitor; ACR, albumin-to-creatinine ratio; ARB, angiotensin receptor blocker; ASCVD, atherosclerotic cardiovascular disease; BP, blood pressure; CCB, calcium channel blocker; CVD, cardiovascular disease; eGFR, estimated glomerular filtration rate; GLP-1 RA, glucagon-like peptide 1 receptor agonist; HTN, hypertension; MRA, mineralocorticoid receptor antagonist; nsMRA, nonsteroidal mineralocorticoid receptor antagonist; PCSK9i, proprotein convertase subtilisin/kexin type 9 inhibitor; RAS, renin-angiotensin system; SGLT2i, sodium-glucose cotransporter 2 inhibitor; T1D, type 1 diabetes; T2D, type 2 diabetes.


## CKD and DM (cont.)

- Monitor for  $\uparrow$  serum creatinine and for  $\uparrow$  serum  $K^+$  levels when ACEi, ARBs, and MRAs are used, or for  $\downarrow K^+$  levels when diuretics are used at routine visits and 7–14 days after initiation or after a dose change.
- Continue ACE inhibitor or ARB therapy for  $\leq 30\%$  increases in serum creatinine in the absence of volume depletion.
- Aim for a urinary albumin reduction  $\geq 30\%$  in people with CKD and urinary albumin  $\geq 300$  mg/g to slow CKD progression.
- Optimize blood pressure management (aim for  $<130/80$  mmHg).

# Retinopathy, Neuropathy, and Foot Care

# Retinopathy, Neuropathy, and Foot Care

If PCPs can optimize the management of glycemia, blood pressure, and lipids -this can reduce or slow the progression of microvascular complications of diabetes.


| How?                                                                                                                                                                                                                                 | When?                                                                                                                                                                                                                                                                                                                                       | Follow-Up Eye Exam Schedule                                                                                                                                                                                                                                                                                                                                                                           |
|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|---------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|-------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|
| <br>Dilated and comprehensive eye exam<br><br>Retinal photography* | <br>For people with type 1 diabetes: within 5 years after the onset of diabetes<br><br><b>NOW</b><br>For people with type 2 diabetes: at the time of diabetes diagnosis | <ul style="list-style-type: none"><li>• At least annually for people with any level of retinopathy</li><li>• Every 1–2 years for those with no retinopathy for one or more annual exams and well-managed glycemia</li><li>• More frequently for those with progressing or sight-threatening retinopathy</li></ul>  |

\*Retinal photography with remote reading or use of an authorized artificial intelligence tool can expand access to screening where qualified eye care professionals are not available. When abnormalities are detected, in-person exams will be needed.

## Treatment


Promptly refer to an ophthalmologist who is knowledgeable and experienced in managing DR any individuals with:

- Any level of diabetic macular edema
- Moderate or worse nonproliferative DR (a precursor of proliferative DR)
- Any proliferative DR



# Retinopathy, Neuropathy, and Foot Care

All patients should be evaluated for DPN at dx and at least annually.



Consider if any referrals are needed - podiatry, vascular



Check feet at least annually for risk factors for ulcers and amputations; more frequent examinations are needed for those with loss of protective sensation, peripheral artery disease, or other serious complications.

# How Often Should Foot Screenings Be Performed?

**Table 12.1—International Working Group on Diabetic Foot risk stratification system and corresponding foot screening frequency**


| Category | Ulcer risk | Characteristics                                                                                                                                                                      | Examination frequency* |
|----------|------------|--------------------------------------------------------------------------------------------------------------------------------------------------------------------------------------|------------------------|
| 0        | Very low   | No LOPS and no PAD                                                                                                                                                                   | Annually               |
| 1        | Low        | LOPS or PAD                                                                                                                                                                          | Every 6–12 months      |
| 2        | Moderate   | LOPS + PAD, or<br>LOPS + foot deformity, or<br>PAD + foot deformity                                                                                                                  | Every 3–6 months       |
| 3        | High       | LOPS or PAD and one or more of the following: <ul style="list-style-type: none"><li>• History of foot ulcer</li><li>• Amputation (minor or major)</li><li>• Kidney failure</li></ul> | Every 1–3 months       |

Adapted with permission from Schaper et al. (123). LOPS, loss of protective sensation; PAD, peripheral artery disease. \*Examination frequency suggestions are based on expert opinion and person-centered requirements.

# Considerations for Older Adults

# Considerations for Older Adults

- Where is the patient at functionally?
- Is this goal reasonable?
- Do medication risks outweigh benefits/patient desired wishes?
- How much assistance do they need to give this medication or adhere to this plan?
- Can the plan be simplified?
- Simplify complex insulin regimens.
- Consider recent hospitalizations.
- Respect your patient and be rigorous in reevaluating health status.



Recognize and Address Any  
Barriers/Challenges to Care

# Recognize and Address Any Barriers/Challenges to Care

Most common in my practice area include:

- Cost/Insurance coverage
- Lack of understanding
- Missed follow-ups
- Language or cultural concerns
- Anxieties around healthcare in general



# Local Resources for Patients/Providers

# KNOW YOUR CARE

Learn more about how diabetes impacts your body and the importance of receiving routine diabetes care. The following information provides recommendations for routine diabetes care, but it is important to follow your provider's orders regarding the suggested frequency of exams, tests, and other care needs. For more information about insurance coverage regarding routine diabetes care, turn to the back page and visit [doh.sd.gov/topics/diseases/chronic-disease/diabetes/diabetes-resources/](http://doh.sd.gov/topics/diseases/chronic-disease/diabetes/diabetes-resources/).

High blood sugar can cause people with diabetes to experience blurry vision and eye disease.

**It is recommended to have an eye exam, including a retinal check, once a year.**

Individuals with diabetes may have additional complications with certain illnesses and should get immunizations, as appropriate.

**Talk with your provider about the frequency of important immunizations.**

High blood sugar can harm the vessels in the kidneys that filter blood and may lead to the need for dialysis.


**Make sure to have lab work done once a year to check kidney function.**

High blood sugar can lead to diabetes complications throughout the body.

**You should have your A1C checked\* every 3-6 months, and check your blood sugar at regular times as recommended by your provider.**

High blood sugar can lead to nerve damage and poor blood flow, which can lead to infections and amputations.

**Inspect your feet routinely at home. Get a comprehensive foot exam once a year and a foot check at each visit.**



High blood sugar can damage blood vessels and cause an increased risk for stroke.

**Be sure to have your blood pressure checked at each visit.**

The most common oral health complication of high blood sugar is gum disease. Diabetes can also cause a dry mouth and infections.

**Schedule a dental exam and cleaning every 3 to 6 months (depending on a dentist's recommendation and dental plan coverage).**

High blood sugar can increase cholesterol making the heart work harder to pump blood, increasing the risk for heart disease and heart attacks.

**Be sure to have your cholesterol checked each year.**

Understanding diabetes care and management is important to prevent or reduce the risk for complications. Individuals with diabetes should attend diabetes education:

- At time of diagnosis
- If complications appear
- When transitions in care occur

**Follow up with your health care provider for questions or concerns regarding your diabetes care and management.**



All recommendations above are sources from the American Diabetes Association Standards of Care in Diabetes - 2025. \* Frequency may vary based on provider's care plan for diabetes management.

# KNOW YOUR PLAN


It is important to understand your insurance coverage for your diabetes care – including office visits, screenings, lab tests, and pharmacy coverage.

## BEFORE YOU CALL:

- Consider checking your insurance company's website or app — many answers are available there.

- If you need additional assistance, call the phone number listed on your insurance card.

- Check with your insurance company to understand if there are different costs for in-network and out-of-network visits.



**Your medical and/or other insurance providers may provide the following benefits:**

- Diabetes Self-Management Education Services (DSMES)
- Dietitian services/ Medical Nutrition Therapy (MNT)
- Continuous Glucose Monitor (CGM)
- Blood sugar testing supplies and test strips
- Immunizations
- Specialty care
- Pharmacy benefits
- Mental health screenings and coverage
- Adaptive shoes
- Foot exams/foot care
- Lab work
- Dilated eye exams
- Dental exams/cleanings

**Many insurance companies offer additional free services, including, but not limited to:**

- Annual wellness visits
- Mental health screenings
- Nutrition counseling
- Fitness & wellness resources
- Fitness/gym membership reimbursement
- Health coaching/case management/community health worker services for diabetes management or prevention (participation may include free testing supplies and resources)

Visit [doh.sd.gov/topics/diseases/chronic-disease/diabetes/diabetes-resources/](http://doh.sd.gov/topics/diseases/chronic-disease/diabetes/diabetes-resources/) to learn more and to download a sample script to assist you in calling your insurance company.



This resource is meant to guide users through contacting their insurance company and is not a guarantee of coverage or benefits.



## Additional Know Your... Resources

In addition to Know Your Care and Know Your Plan, the SD DOH and SDDC are developing the following resources:

- Know Your Medications
- Know Your Team
- Know Your Numbers
- Know Your Emergency Plan
- Know Your Sick Day
- Know Your Travel
- Know Your Technology
- And complimenting fillable forms!

<https://doh.sd.gov/topics/diseases/chronic-disease/diabetes/diabetes-management/>



Topics ▾ Programs ▾ Licensing & Records ▾ Healthcare Professionals ▾

### Talking with Your Doctor

Being prepared with your questions and goals in mind will help you and your care team or doctor make treatment decisions and a diabetes care plan that works for you.

#### Find Resources to Help Prepare for a Doctor's Visit:

[NIDDK | Getting Ready for Your Diabetes Care Visit](#) [↗](#) – You are the person in charge of your diabetes. Being prepared for your healthcare visit can help you get the information you need to manage your disease.

[Your Health Care Team](#) [↗](#) – Information on the members of your diabetes care team and what they do, from the American Diabetes Association.

[Know Your Care](#) [📄](#) – Learn more about how diabetes impacts your body and the importance of receiving routine diabetes care.

# South Dakota Diabetes Coalition



ABOUT PATIENTS EDUCATORS PROVIDERS CALENDAR OF EVENTS STAY CONNECTED CONTACT US



The mission of the South Dakota Diabetes Coalition (SDDC) is to collaboratively improve the quality of life for all South Dakotans at risk for, or affected by, diabetes.

The SDDC is a volunteer-based, non-profit 501c3 organization in South Dakota.

ABOUT THE SDDC

**For Patients and Family Members**

Are you or a loved one affected by diabetes? Whether prediabetes, Type 1 Diabetes, Type 2 Diabetes, or Gestational

**For Educators**

Diabetes Educators play an important role in caring for and educating those with diabetes and prediabetes in South Dakota.

**For Providers**

Providers (from family medicine, to endocrinology, to podiatrists, dentists, and eye doctors) play an important role in



ABOUT PATIENTS EDUCATORS PROVIDERS CALENDAR OF EVENTS STAY CONNECTED CONTACT US

## Empower SD Educational Sessions

### Upcoming Educational Sessions

Beginning in October 2026, Educational Sessions will resume for individuals with Type 1 and Type 2 Diabetes and for their family members and caregivers who support them.

Sign up to receive updates for dates and times for the Empower SD Sessions.

**TYPE 1 DIABETES**  
Educational Sessions for Type 1 Diabetes will be on the first **Tuesday of the month** at 7 PM CT/ 6 PM MT.

SIGN UP FOR TYPE 1 EMPOWER SD SESSION UPDATES

**TYPE 2 DIABETES**  
Educational Sessions for Type 2 Diabetes will be on the **second Tuesday of the month** at 7 PM CT/ 6 PM MT.

SIGN UP FOR TYPE 2 EMPOWER SD SESSION UPDATES

### Recorded Educational Sessions

[Overview](#)

## Case Study – Joe

- Joe, a 56-year-old, Hispanic male, presents to your clinic as a new patient with c/o fatigue, blurry vision, a non-healing wound to his LLE, and overall, just not feeling well.
- He has no known medical history as he admits he hasn't seen a provider since he had a laceration in the ER a few years ago. He is a smoker, denies alcohol or drug use, and works in a factory setting. He is wondering what he can do to have more energy and heal his leg.
- His vitals are: BP 150/90; HR 88; Resp 21; BMI 33.0

### WHERE DO YOU START?

- Finish getting your history, perform PE, and start lab work-up.
- Be mindful you likely aren't going to fix all of Joe's issues in one visit and he will need education on adherence to treatment plan and importance of follow-up visits.

Your nurse walks in as you are assessing his LLE wound and updates you his A1C is back and is **9% – WHAT NEXT?**

- Time to start in with education, choosing medications, and don't forget you still have to address his wound and BP. Keep in mind your next visit showed up and it's 2 siblings here for well child checks. Welcome to Primary Care :)



Thank You!



# Resources

- <https://diabetesjournals.org/clinical/article/doi/10.2337/cd25-aint/157719/Standards-of-Care-in-Diabetes-2025-Abridged-for?searchresult=1>
- <https://diabetesjournals.org/collection/2720/2025-Abridged-Standards-of-Care>
- [https://www.countyhealthrankings.org/health-data/south-dakota?measure=Diabetes+Prevalence\\*&year=2024](https://www.countyhealthrankings.org/health-data/south-dakota?measure=Diabetes+Prevalence*&year=2024)
- <https://diabetesjournals.org/clinical/article/40/4/401/144956/Role-of-Primary-Care-Clinicians-in-the-Management>
- [https://diabetesjournals.org/care/article/48/Supplement\\_1/S27/157566/2-Diagnosis-and-Classification-of-Diabetes](https://diabetesjournals.org/care/article/48/Supplement_1/S27/157566/2-Diagnosis-and-Classification-of-Diabetes)
- <https://www.cdc.gov/diabetes/risk-factors/index.html> <https://diabetes.org/about-diabetes/diabetes-prevention>
- <https://www.mayoclinic.org/diseases-conditions/type-2-diabetes/in-depth/diabetes-prevention/art-20047639>
- <https://www.cdc.gov/diabetes-prevention/lifestyle-change-program/lifestyle-change-program-details.html>
- [https://www.uptodate.com/contents/initial-management-of-hyperglycemia-in-adults-with-type-2-diabetes-mellitus?search=type%20%20dm%20medication&source=search\\_result&selectedTitle=2%7E150&usage\\_type=default&display\\_rank=2](https://www.uptodate.com/contents/initial-management-of-hyperglycemia-in-adults-with-type-2-diabetes-mellitus?search=type%20%20dm%20medication&source=search_result&selectedTitle=2%7E150&usage_type=default&display_rank=2)
- <https://pmc.ncbi.nlm.nih.gov/articles/PMC4346284/>