RYBELSUS®
semaglutide tablets 7mg | 14mg

you're invited.

Join us for an upcoming event.

Discover When RYBELSUS® Should Be Considered for Adults With T2D

Presenter: TANYA MUNGER, NP

Thursday, March 31, 2022 06:15 PM - 07:15 PM CDT

GreenFire

Host:
ELISE REMSEN

6795 East Riverside Boulevard

Rockford
Illinois
61114

To RSVP or if you have questions about this program, please call **ELISE REMSEN** at **+1 779-238-2966** or email **ezre@novonordisk.com**

Indications and Usage

RYBELSUS® (semaglutide) tablets 7 mg or 14 mg is indicated as an adjunct to diet and exercise to improve glycemic control in adults with type 2 diabetes.

Limitations of Use

- RYBELSUS® is not recommended as a first-line therapy for patients who have inadequate glycemic control on diet and exercise because of the uncertain relevance of rodent C-cell tumor findings to humans
- RYBELSUS® has not been studied in patients with a history of pancreatitis. Consider other antidiabetic therapies in patients with a history of pancreatitis
- RYBELSUS® is not indicated for use in patients with type 1 diabetes

Important Safety Information

WARNING: RISK OF THYROID C-CELL TUMORS

- **In rodents, semaglutide causes dose-dependent and treatment-duration-dependent thyroid C-cell tumors at clinically relevant exposures. It is unknown whether RYBELSUS® causes thyroid C-cell tumors, including medullary thyroid carcinoma (MTC), in humans as human relevance of semaglutide-induced rodent thyroid C-cell tumors has not been determined**
- **RYBELSUS® is contraindicated in patients with a personal or family history of MTC and in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2). Counsel patients regarding the potential risk for MTC with the use of RYBELSUS® and inform them of symptoms of thyroid tumors (e.g. a mass in the neck, dysphagia, dyspnea, persistent hoarseness). Routine monitoring of serum calcitonin or using thyroid ultrasound is of uncertain value for early detection of MTC in patients treated with RYBELSUS®**

Contraindications

- RYBELSUS® is contraindicated in patients with a personal or family history of medullary thyroid carcinoma (MTC) or in patients with Multiple Endocrine Neoplasia syndrome type 2 (MEN 2), and in patients with a prior serious hypersensitivity reaction to semaglutide or to any of the excipients in RYBELSUS®. Serious hypersensitivity reactions including anaphylaxis and angioedema have been reported with RYBELSUS®

Please see additional Important Safety Information on following page
Please see accompanying Prescribing Information, including Boxed Warning.

Important Safety Information

Warnings and Precautions

- **Risk of Thyroid C-Cell Tumors:** Patients should be further evaluated if serum calcitonin is measured and found to be elevated or thyroid nodules are noted on physical examination or neck imaging
- **Pancreatitis:** Has been reported in clinical trials. Observe patients carefully for signs and symptoms of pancreatitis (including persistent severe abdominal pain, sometimes radiating to the back and which may or may not be accompanied by vomiting). If pancreatitis is suspected, discontinue RYBELSUS® and initiate appropriate management; if confirmed, do not restart RYBELSUS®
- **Diabetic Retinopathy Complications:** In a pooled analysis of glycemic control trials with RYBELSUS®, patients reported diabetic retinopathy related adverse reactions during the trial (4.2% with RYBELSUS® and 3.8% with comparator). In a 2-year trial with semaglutide injection involving patients with type 2 diabetes and high cardiovascular risk, more events of diabetic retinopathy complications occurred in patients treated with semaglutide injection (3.0%) compared to placebo (1.8%). The absolute risk increase for diabetic retinopathy complications was larger among patients with a history of diabetic retinopathy at baseline than among patients without a known history of diabetic retinopathy.
  Rapid improvement in glucose control has been associated with a temporary worsening of diabetic retinopathy. Patients with a history of diabetic retinopathy should be monitored for progression of diabetic retinopathy
- **Hypoglycemia:** Patients receiving RYBELSUS® in combination with an insulin secretagogue (e.g., sulfonylurea) or insulin may have an increased risk of hypoglycemia, including severe hypoglycemia. The risk of hypoglycemia may be lowered by a reduction in the dose of sulfonylurea (or other concomitantly administered insulin secretagogue) or insulin. Inform patients using these concomitant medications of the risk of hypoglycemia and educate them on the signs and symptoms of hypoglycemia
- **Acute Kidney Injury:** There have been postmarketing reports of acute kidney injury and worsening of chronic renal failure, which may sometimes require hemodialysis, in patients treated with GLP-1 receptor agonists, including semaglutide. Some of these events have been reported in patients without known underlying renal disease. A majority of the reported events occurred in patients who had experienced nausea, vomiting, diarrhea, or dehydration. Monitor renal function when initiating or escalating doses of RYBELSUS® in patients reporting severe adverse gastrointestinal reactions
- **Hypersensitivity:** Serious hypersensitivity reactions (e.g., anaphylaxis, angioedema) have been reported in patients treated with RYBELSUS®. If hypersensitivity reactions occur, discontinue use of RYBELSUS®, treat promptly per standard of care, and monitor until signs and symptoms resolve. Use caution in a patient with a history of angioedema or anaphylaxis with another GLP-1 receptor agonist

Adverse Reactions

- The most common adverse reactions, reported in ≥5% of patients treated with RYBELSUS® are nausea, abdominal pain, diarrhea, decreased appetite, vomiting and constipation

Drug Interactions

- When initiating RYBELSUS®, consider reducing the dose of concomitantly administered insulin secretagogue (such as sulfonylureas) or insulin to reduce the risk of hypoglycemia
- RYBELSUS® delays gastric emptying and has the potential to impact the absorption of other oral medications. Closely follow RYBELSUS® administration instructions when coadministering with other oral medications and consider increased monitoring for medications with a narrow therapeutic index, such as levothyroxine

Use in Specific Populations

- **Pregnancy:** Available data with RYBELSUS® are not sufficient to determine a drug-associated risk for major birth defects, miscarriage, or other adverse maternal or fetal outcomes. Based on animal reproduction studies, there may be risks to the fetus from exposure to RYBELSUS®. Use only if the potential benefit justifies the potential risk to the fetus
- **Lactation:** There are no data on the presence of semaglutide in human milk, the effects on the breastfed infant, or the effects on milk production. Because of the unknown potential for serious adverse reactions in the breastfed infant due to the possible accumulation of salcaprozate sodium (SNAC), an absorption enhancer in RYBELSUS®, from breastfeeding and because there are alternative formulations of semaglutide that can be used during lactation, advise patients that breastfeeding is not recommended during treatment with RYBELSUS®
- Discontinue RYBELSUS® in women at least 2 months before a planned pregnancy due to the long washout period for semaglutide
- **Pediatric Use:** Safety and efficacy of RYBELSUS® have not been established in pediatric patients (younger than 18 years)

Please see additional Important Safety Information on previous page.
Please see accompanying Prescribing Information, including Boxed Warning.

For additional promotional information or patient resources, visit rybelsuspro.com.

Novo Nordisk is subject to federal and state laws that require the disclosure of items of value, such as meals, provided to certain health care professionals. By participating in this activity, the value of the meal will be reported as required, and that information may be publicly available. If you prefer to attend the program and not accept a meal, there is a box to check at the event sign-in, indicating that you did not accept a meal.

Novo Nordisk will report all transfers of value, including meals, as required by law. Please refrain from consuming any meals available at this event if you are a health care provider licensed in a state that prohibits, or applies limits to, such interactions.

In Accordance with the PhRMA Code on Interactions with Healthcare Professionals, Novo Nordisk will not provide alcohol at its programs and attendance at this program is limited to healthcare professionals with an educational need to receive the information. Additionally, attendance by guests or spouses is not appropriate and cannot be accommodated.

We appreciate your understanding and apologize for any inconvenience this may cause. Please contact NNI Ethics & Compliance with any questions at NNIComplianceAssist@novonordisk.com.

novo nordisk®

RYBELSUS® is a registered trademark of Novo Nordisk A/S.
Novo Nordisk is a registered trademark of Novo Nordisk A/S.
© 2021 Novo Nordisk. Printed in the U.S.A.
US21RYB00216 June 2021