

Please Join Us for a Discussion

The DUPIXENT Difference: Directly Inhibiting IL-4 and IL-13 Signaling in Appropriate Asthma Patients

The mechanism of dupilumab action has not been definitively established



INDICATIONS

Asthma: DUPIXENT is indicated as an add-on maintenance treatment of adult and pediatric patients aged 6 years and older with moderate-to-severe asthma characterized by an eosinophilic phenotype or with oral corticosteroid dependent asthma.

Limitations of Use: DUPIXENT is not indicated for the relief of acute bronchospasm or status asthmaticus.

Chronic Obstructive Pulmonary Disease: DUPIXENT is indicated as an add-on maintenance treatment of adult patients with inadequately controlled chronic obstructive pulmonary disease (COPD) and an eosinophilic phenotype.

Limitations of Use: DUPIXENT is not indicated for the relief of acute bronchospasm.

IMPORTANT SAFETY INFORMATION

CONTRAINDICATION: DUPIXENT is contraindicated in patients with known hypersensitivity to dupilumab or any of its excipients.

Please see additional Important Safety Information below.

PROGRAM AGENDA

8/27/2026

6:00 PM Central

French market bistro

Please arrive 30 minutes before the program.

16645 highland road

Baton Rouge, Louisiana 70810

PRESENTED BY

David Kaufman, MD

LSU Health Shreveport

HOSTED BY

Kendra Thompson

kendra.thompson@regeneron.com

337-349-6358

RGN0044018

You may RSVP to your program host.

CLICK TO REGISTER!

<https://tpprod-regeneron.physiciansworld.com/1IG7ED>

IMPORTANT SAFETY INFORMATION (CONT'D)

WARNINGS AND PRECAUTIONS

Hypersensitivity: Hypersensitivity reactions, including anaphylaxis, serum sickness or serum sickness-like reactions, angioedema, generalized urticaria, rash, erythema nodosum, and erythema multiforme have been reported. If a clinically significant hypersensitivity reaction occurs, institute appropriate therapy and discontinue DUPIXENT.

Please see additional Important Safety Information on next page.

DUPIXENT[®]
(dupilumab) Injection
200mg • 300mg

IMPORTANT SAFETY INFORMATION (CONT'D)

WARNINGS AND PRECAUTIONS

Conjunctivitis, Keratitis, and Blepharitis: Conjunctivitis and keratitis occurred more frequently in COPD subjects who received DUPIXENT versus placebo. Conjunctivitis, keratitis, and blepharitis have been reported with DUPIXENT in postmarketing settings. Some patients reported varying degrees of transient or ongoing visual impairment including blindness associated with conjunctivitis, keratitis, or blepharitis leading to discontinuation of DUPIXENT and/or surgical intervention. Advise patients or their caregivers to promptly report new onset or worsening eye symptoms to their healthcare provider. Consider discontinuation of DUPIXENT and prompt ophthalmological examination for patients who develop signs and symptoms suggestive of keratitis, or when conjunctivitis or blepharitis do not resolve following standard treatment, as appropriate. Use with caution in patients with significant dry eye disease, history of significant lid abnormalities/surgeries, or history of nasolacrimal surgery.

Eosinophilic Conditions: Patients being treated for asthma may present with clinical features of eosinophilic pneumonia or eosinophilic granulomatosis with polyangiitis (EGPA). These events may be associated with the reduction of oral corticosteroid therapy. Healthcare providers should be alert to vasculitic rash, worsening pulmonary symptoms, cardiac complications, kidney injury, and/or neuropathy presenting in their patients with eosinophilia. Cases of eosinophilic pneumonia were reported in adults who participated in the asthma development program and cases of EGPA have been reported with DUPIXENT in adults who participated in the asthma development program as well as in adult subjects with co-morbid asthma in the CRSwNP development program. Advise patients to report signs of eosinophilic pneumonia and EGPA. Consider withholding DUPIXENT if eosinophilic pneumonia or EGPA are suspected.

Acute Symptoms of Asthma or Chronic Obstructive Pulmonary Disease or Acute Deteriorating Disease: Do not use DUPIXENT to treat acute symptoms or acute exacerbations of asthma or COPD, acute bronchospasm, or status asthmaticus. Patients should seek medical advice if their asthma or COPD remains uncontrolled or worsens after initiation of DUPIXENT.

Risk Associated with Abrupt Reduction of Corticosteroid Dosage: Do not discontinue systemic, topical, or inhaled corticosteroids abruptly upon initiation of DUPIXENT. Reductions in corticosteroid dose, if appropriate, should be gradual and performed under the direct supervision of a healthcare provider. Reduction in corticosteroid dose may be associated with systemic withdrawal symptoms and/or unmask conditions previously suppressed by systemic corticosteroid therapy.

Patients with Co-morbid Asthma: Advise patients with co-morbid asthma not to adjust or stop their asthma treatments without consultation with their physicians.

Psoriasis: Cases of new-onset psoriasis have been reported with the use of DUPIXENT for the treatment of asthma, including in patients without a family history of psoriasis. In postmarketing reports, these cases resulted in partial or complete resolution of psoriasis with discontinuation of dupilumab, with or without use of supplemental treatment for psoriasis (topical or systemic). Those who continued dupilumab received supplemental treatment for psoriasis to improve associated symptoms. Advise patients to report new-onset psoriasis symptoms. If symptoms persist or worsen, consider dermatologic evaluation and/or discontinuation of DUPIXENT.

Arthralgia and Psoriatic Arthritis: Arthralgia has been reported with use of DUPIXENT with some patients reporting gait disturbances or decreased mobility associated with joint symptoms; some cases resulted in hospitalization. Cases of new-onset psoriatic arthritis requiring systemic treatment have been reported with the use of DUPIXENT. Advise patients to report new onset or worsening joint symptoms. If symptoms persist or worsen, consider rheumatological evaluation and/or discontinuation of DUPIXENT.

Parasitic (Helminth) Infections: It is unknown if DUPIXENT will influence the immune response against helminth infections. Treat patients with pre-existing helminth infections before initiating therapy with DUPIXENT. If patients become infected while receiving treatment with DUPIXENT and do not respond to anti-helminth treatment, discontinue treatment with DUPIXENT until the infection resolves. Helminth infections (5 cases of enterobiasis and 1 case of ascariasis) were reported in pediatric patients 6 to 11 years old in the pediatric asthma development program.

Vaccinations: Consider completing all age-appropriate vaccinations as recommended by current immunization guidelines prior to initiating DUPIXENT. Avoid use of live vaccines during treatment with DUPIXENT.

ADVERSE REACTIONS:

Most common adverse reactions:

Asthma (incidence $\geq 1\%$): injection site reactions, oropharyngeal pain, and eosinophilia.

Chronic Obstructive Pulmonary Disease (incidence $\geq 2\%$): viral infection, headache, nasopharyngitis, back pain, diarrhea, arthralgia, urinary tract infection, local administration reactions, rhinitis, eosinophilia, toothache, and gastritis.

USE IN SPECIFIC POPULATIONS

- **Pregnancy:** A pregnancy exposure registry monitors pregnancy outcomes in women exposed to DUPIXENT during pregnancy. Available data from case reports and case series with DUPIXENT use in pregnant women have not identified a drug-associated risk of major birth defects, miscarriage or adverse maternal or fetal outcomes. Human IgG antibodies are known to cross the placental barrier; therefore, DUPIXENT may be transmitted from the mother to the developing fetus.
- **Lactation:** There are no data on the presence of DUPIXENT in human milk, the effects on the breastfed infant, or the effects on milk production. Maternal IgG is known to be present in human milk. The developmental and health benefits of breastfeeding should be considered along with the mother's clinical need for DUPIXENT and any potential adverse effects on the breastfed child from DUPIXENT or from the underlying maternal condition.

Please see accompanying full Prescribing Information.

In accordance with the PhRMA Code on Interactions with Healthcare Professionals, this Program is limited to U.S. Healthcare Professionals and persons with bona fide professional interest in the information presented. Attendance at this Program by guests or spouses is not permitted unless they would qualify as an appropriate attendee on their own. If a meal is provided, actively licensed Minnesota and Vermont prescribers may attend, but not partake in the meal. Full-time Federal Employees may attend and partake in the meal if the Program is considered widely attended (50 or more attendees). If not, they may attend, but not partake in the meal. Part-time Federal Employees acting in their civilian capacity may attend the Program and partake in the meal. Alcohol is not permitted in connection with Speaker Programs and will not be provided. The value of any meal provided in connection with the Program may be reported in accordance with federal and state laws and regulations.