



LKT Laboratories, Inc.

Vitamin D2


Phone: 888-558-5227
651-644-8424
Fax: 888-558-7329
Email: getinfo@lktlabs.com
Web: lktlabs.com

Product Information

Product ID V3476
CAS No. 50-14-6
Chemical Name (3 β ,5Z,7E,22E)-9,10-Secoergosta-5,7-10(19),22-tetra- en-3-ol

Synonym Calciferol, Ergocalciferol, Viosterol, Condol, Deltalin, Diactol, Ergorone, Infron, Mulsiferol, Ostelin, Radsterin

Formula C₂₈H₄₄O
Formula Wt. 396.65
Melting Point 115-118°C
Purity ≥91%
Solubility Soluble in organic solvents. Insoluble in water.



Pricing and Availability

Bulk quantities available upon request

Product ID	Size	List Price
V3476	1 g	\$50.90
V3476	5 g	\$183.60

Store Temp -20°C

Ship Temp Ambient

Description Calciferol is a Vitamin D prodrug found in fungi and alfalfa; it is commercially used as a vitamin D supplement for bone strength but also exhibits antioxidative and anticancer chemotherapeutic activities. In leukemia cells, calciferol induces apoptosis, decreases the mitochondrial membrane potential, and increases release of cytochrome C, cleavage of PARP, and activation of caspases. In animal models of breast cancer, calciferol decreases tumor growth.

References Bikle DD. Vitamin D metabolism, mechanism of action, and clinical applications. Chem Biol. 2014 Mar 20;21(3):319-29. PMID: 24529992.

Chen WJ, Huang YT, Wu ML, et al. Induction of apoptosis by vitamin D2, ergocalciferol, via reactive oxygen species generation, glutathione depletion, and caspase activation in human leukemia Cells. J Agric Food Chem. 2008 May 14;56(9):2996-3005. PMID: 18386902.

Zinser GM, Tribble E, Valrance M, et al. 1,24(S)-dihydroxyvitamin D2, an endogenous vitamin D2 metabolite, inhibits growth of breast cancer cells and tumors. Anticancer Res. 2005 Jan-Feb;25(1A):235-41. PMID: 15816543.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.