

Phone: 888-558-5227

651-644-8424

888-558-7329 Fax: Email: getinfo@lktlabs.com

Web: lktlabs.com

## **Product Information**

Product ID R2112

CAS No.

**Chemical Name** 

Synonym

Formula C<sub>22</sub>H<sub>33</sub>N<sub>7</sub>O<sub>8</sub>

Formula Wt. 523.5

**Melting Point** 

Purity ≥98%

Solubility Soluble in water (1 mg/mL).

Arg-Phe-Asp-Ser

## **Pricing and Availability**

Bulk quanitites available upon request

Product ID Size **List Price** R2112 \$106.10 5 mg

Store Temp -20°C Ship Temp Ambient

**Description** RFDS is often used as a negative control for RGD peptide activity. RFDS exhibits immunomodulatory activity; it

contains a peptide sequence homologous with the B1-domain region of a MHC class II antigen from CD4.

References Iwao Y, Fujimura T. Activation of Xenopus eggs by RGD-containing peptides accompanied by intracellular Ca2+ release. Dev Biol. 1996 Aug 1;177(2):558-67. PMID: 8806831.

> Mazerolles F, Durandy A, Piatier-Tonneau D, et al. Immunosuppressive properties of synthetic peptides derived from CD4 and HLA-DR antigens. Cell. 1988 Nov 4;55(3):497-504. PMID: 3263212.

Auffray C, Novotny J. Speculations on sequence homologies between the fibronectin cell-attachment site, major histocompatibility antigens, and a putative AIDS virus polypeptide. Hum Immunol. 1986 Apr;15(4):381-90. PMID: 3009364.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.