Phone: 888-558-5227

651-644-8424

Fax: 888-558-7329 Email: getinfo@lktlabs.com

Web: lktlabs.com

Product Information

Product ID N1755 CAS No. 1405-10-3

Chemical Name 5-amino-2-(aminomethyl)-6-[4,6-diamino-2-[4-[3-amino-6-

(aminomethyl)-4,

5-dihydroxyoxan-2-yl]oxy-3-hydroxy-5-(hydroxymethyl)oxolan-2-yl]oxy

Synonym

Formula $C_{23}H_{46}N_6O_{13} \cdot H_2SO_4$

Formula Wt. 908.9

Melting Point

Purity ≥98%

Solubility Soluble in water (1 g/mL).

HO,NH ₂		
HO - ○ • H ₂ SO ₄		
H ₂ N O, NH ₂		
OH OH OH OH		
HO HO NH₂		
HO NH ₂		
Neomycin B: $R^1 = H$, $R^2 = CH_2NH_2$ Neomycin C: $R^2 = CH_2NH_2$, $R^1 = H$		

Pricing and Availability

Bulk quanitites available upon request

Product ID	Size	List Price
N1755	1 g	\$27.60
N1755	5 g	\$35.80
N1755	10 g	\$41.10
N1755	25 g	\$82.70
N1755	100 g	\$242.80

Store Temp Ambient Ship Temp Ambient

Description Neomycin is a polycationic aminoglycoside antibiotic that exhibits antibacterial and neuromodulatory activities; it was originally found in species of Streptomyces. Neomycin binds to 23S rRNA, constraining tRNA in the P-site, inhibiting translocation and ribosomal recycling; together, this results in inhibition of protein synthesis. Neomycin B binds and inhibits bacterial RNase P and bacterial T box antiterminator RNA. Neomycin B inhibits mammalian RNA splicing as well. Neomycin also noncompetitively inhibits cation channels, including P2X receptors and transient receptor potential vanilloid receptor 1 (TRPV1) channels. Additionally, this compound inhibits Ca2+ influx, preventing presynaptic release of acetylcholine and norepinephrine in feline superior cervical ganglia.

References Wang L, Pulk A, Wasserman MR, et al. Allosteric control of the ribosome by small-molecule antibiotics. Nat Struct Mol Biol. 2012 Sep;19(9):957-63. PMID: 22902368.

> Bongartz EV, Rettinger J, Hausmann R. Aminoglycoside block of P2X2 receptors heterologously expressed in Xenopus laevis oocytes. Purinergic Signal. 2010 Dec;6(4):393-403. PMID: 21437010.

Aukema KG, Chohan KK, Plourde GL, et al. Small molecule inhibitors of yeast pre-mRNA splicing. ACS Chem Biol. 2009 Sep 18;4(9):759-68. PMID: 19634919.

Anupam R, Denapoli L, Muchenditsi A, et al. Identification of neomycin B-binding site in T box antiterminator model RNA. Bioorg Med Chem. 2008 Apr 15;16(8):4466-70. PMID: 18329274.

Kawamoto SA, Sudhahar CG, Hatfield CL, et al. Studies on the mechanism of inhibition of bacterial ribonuclease P by aminoglycoside derivatives. Nucleic Acids Res. 2008 Feb;36(2):697-704. PMID: 18084035.

Raisinghani M, Premkumar LS. Block of native and cloned vanilloid receptor 1 (TRPV1) by aminoglycoside antibiotics. Pain. 2005 Jan;113(1-2):123-33. PMID: 15621372.

Wright JM, Collier B. The effects of neomycin upon transmitter release and action. J Pharmacol Exp Ther. 1977 Mar; 200(3):576 -87. PMID: 191590.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.