

Phone: 888-558-5227

651-644-8424

Fax: 888-558-7329 Email: getinfo@lktlabs.com

Web: lktlabs.com

Product Information

Product ID B1653 CAS No. 622-78-6

Chemical Name Benzene, (isothiocyanatomethyl)-

Synonym Isothiocyanic acid benzyl ester, 2-Bromo-4-trifluoro- methyl-6-nitrophenyl-


isothiocyanate

Formula C₈H₇NS Formula Wt. 149.22 **Melting Point**

Purity ≥97%

Solubility Soluble in DMSO, ethanol,

chloroform.



Pricing and Availability

Bulk quanitites available upon request

Product ID	Size	List Price
B1653	5 g	\$45.90
B1653	10 g	\$81.50

Store Temp 4°C Ship Temp Ambient

Description Benzyl isothiocyanate (BITC) is an isothiocyanate originally found in cruciferous vegetables that exhibits immunomodulatory, anti-parasitic, antibiotic, antioxidative, anti-atherosclerotic, anti-angiogenic, anti-metastatic, anticancer chemotherapeutic, and chemopreventive activities. BITC inhibits IL-13 expression in basophils. BITC induces phase II enzymes, increasing levels of heme oxygenase 1 (HO-1), glutathione, and glutamate cysteine ligase; it also decreases levels of ROS, activation of NF-kB, adhesion of monocytes, and expression of ICAM-1, VCAM-1, and E-selectin. This compound inhibits growth of Trypanosoma and displays antibacterial efficacy against Campylobacter by disrupting metabolic processes. In vitro and in vivo, BITC decreases high fat diet-induced tumor growth and multiplicity, also inhibiting macrophage migration and lipid droplet accumulation. This compound inhibits squamous cell carcinoma cell invasion and migration and induces apoptosis and autophagy in prostate cancer cells. In various animal models, BITC inhibits development of mammary gland tumors and decreases activation of STAT3 and secretion of VEGF, MMP-2, and HIF-1 α .

References Tang Y, Abe N, Yoshimoto M, et al. Benzyl isothiocyanate inhibits IL-13 expression in human basophilic KU812 cells. Biosci Biotechnol Biochem. 2014 Sep 25:1-5. PMID: 25253661.

> Kim M, Cho HJ, Kwon GT, et al. Benzyl isothiocyanate suppresses high-fat diet-stimulated mammary tumor progression via the alteration of tumor microenvironments in obesity-resistant BALB/c mice. Mol Carcinog. 2014 Apr 11. [Epub ahead of print]. PMID: 24729546.

> Steverding D, Michaels S, Read KD. In vitro and in vivo studies of trypanocidal activity of dietary isothiocyanates. Planta Med. 2014 Feb;80(2-3):183-6. PMID: 24452460.

> Dufour V, Stahl M, Rosenfeld E, et al. Insights into the mode of action of benzyl isothiocyanate on Campylobacter jejuni. Appl Environ Microbiol. 2013 Nov;79(22):6958-68. PMID: 24014524.

> Huang CS, Lin AH, Liu CT, et al. Isothiocyanates protect against oxidized LDL-induced endothelial dysfunction by upregulating Nrf2-dependent antioxidation and suppressing NFkB activation. Mol Nutr Food Res. 2013 Nov;57(11):1918-30. PMID: 23836589.

Lin JF, Tsai TF, Liao PC, et al. Benzyl isothiocyanate induces protective autophagy in human prostate cancer cells via inhibition of mTOR signaling. Carcinogenesis. 2013 Feb;34(2):406-14. PMID: 23172666.

Boreddy SR, Sahu RP, Srivastava SK. Benzyl isothiocyanate suppresses pancreatic tumor angiogenesis and invasion by inhibiting

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.