Phone: 888-558-5227

651-644-8424

888-558-7329 Fax: Email: getinfo@lktlabs.com

Web: lktlabs.com

Product Information

Product ID C0150 CAS No. 7689-03-4

Chemical Name (S)-4-Ethyl-4-hydroxy-1H-pyrano[3',4':6,7]indolizino- [1,2-b]quinoline

-3,14(4H,12H)-dione

Synonym

Formula C₂₀H₁₆N₂O₄ Formula Wt. 348.35

Melting Point 264-267°C(dec.)

Purity ≥98%

Solubility Soluble in DMSO (10

mg/mL), methanol (40 mg/mL), 0.1 N NaOH (50 mg/mL). Insoluble in water.

Store Temp -20°C

Ship Temp Ambient

 $\mathsf{OH} \ \mathsf{O}$

Pricing and Availability

Bulk quanitites available upon request

Size	List Price
25 mg	\$30.00
100 mg	\$60.00
250 mg	\$147.20
500 mg	\$268.40
	25 mg 100 mg 250 mg

anticancer chemotherapeutic activity, inhibiting topoisomerase I and inducing double-stranded DNA breaks. Camptothecin is extremely cytotoxic and only its derivatives are clinically used as chemotherapeutics.

Description Camptothecin is a quinolone alkaloid precursor of irinotecan; it is originally produced by Camptotheca. Camptothecin exhibits

References Rodríguez-Berna G, Mangas-Sanjuán V, Gonzalez-Alvarez M, et al. A promising camptothecin derivative: Semisynthesis, antitumor activity and intestinal permeability. Eur J Med Chem. 2014 Jun 25;83C:366-373. PMID: 24980118.

> Berniak K, Rybak P, Bernas T, et al. Relationship between DNA damage response, initiated by camptothecin or oxidative stress, and DNA replication, analyzed by quantitative 3D image analysis. Cytometry A. 2013 Jul 11. [Epub ahead of print]. PMID: 23846844.

> Redinbo MR, Stewart L, Kuhn P, et al. Crystal structures of human topoisomerase I in covalent and noncovalent complexes with DNA. Science. 1998 Mar 6;279(5356):1504-13. PMID: 9488644.

Caution: This product is intended for laboratory and research use only. It is not for human or drug use.