



FIGHT-BAC™ RTU



Disinfectant • Cleaner • Tuberculocidal • *Virucide • Fungicide (against pathogenic fungi)

This product is a one-step disinfectant cleaner and deodorant designed for general cleaning and disinfecting on hard non-porous inanimate surfaces. Quickly removes dirt, grime, mold, mildew, food residue, blood and other organic matter commonly found in health care facilities. When used as directed, this product is highly effective against a broad spectrum of pathogenic microorganisms.



Uses

- Appliances
- Cabinets
- Floors
- Plastic surfaces
- Bathrooms
- Garbage cans
- Kitchen counters
- Metal surfaces
- Sinks
- Walls

Benefits

- Eliminates odors, leaving restrooms smelling clean and fresh.
- Meets requirements of OSHA Bloodborne Pathogen Standard for decontamination of working surfaces. Kills TB as well as HBV and HIV.
- Broad Spectrum Disinfectant – Kills TB, Hepatitis A, B and C Viruses, MRSA, VRE, Norovirus and Rotavirus.
- Virucidal against HIV-1 (AIDS virus), Canine Parvovirus, HBV and HCV on pre-cleaned environmental surfaces.
- Fungicidal against pathogenic fungi.
- Nonflammable.

Properties

Color	Clear
Film residue	Free rinsing
Fragrance	Pleasant
pH	11.7
Shelf life	1 year
Specific gravity	1.02
Viscosity	Water-thin
ACTIVE INGREDIENTS:	
n-Alkyl (60% C ₁₄ , 30% C ₁₆ , 5% C ₁₂ , 5% C ₁₈)	0.105%
dimethyl benzyl ammonium chlorides	0.105%
n-Alkyl (68% C ₁₂ , 32% C ₁₄) dimethyl ethylbenzyl ammonium chlorides	0.105%

Safety

CAUTION: KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment. **IF IN EYES:** Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice. **IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything by mouth to an unconscious person. **IF INHALED:** Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice. **NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

Available In

Case/12 Quarts (946 mL); Item #31112-00

Acceptance

Registered with the U.S. Environmental Protection Agency, EPA Reg. No. 1839-83-4170.

Directions For Use

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. This product is not to be used as a terminal sterilant/high-level disinfectant on any surface or instrument that (1) is introduced directly into the human body, either into or in contact with the bloodstream or normally sterile areas of the body, or (2) contacts intact mucous membranes but which does not ordinarily penetrate the blood barrier or otherwise enter normally sterile areas of the body. **DISINFECTION AND CLEANING:** Remove gross filth or heavy soil prior to application of the product. Hold container six to eight inches from surface to be treated. Spray area until it is covered with the solution. Allow product to penetrate and the surface to remain wet for **3 minutes** (for Bloodborne Pathogens: allow surface to remain wet for HIV-1, **1 minute** and for HBV and HCV, **5 minutes**). No scrubbing is necessary. Wipe off with a clean cloth, mop or sponge. This product will not leave grit or soap scum. **Disinfection of animal quarters and kennels:** For disinfection of pre-cleaned animal quarters and kennels, apply this product. Remove all animals and feed from premises. Remove all litter and droppings from floors, walls and surfaces of facilities occupied or traversed by animals. Empty all troughs, racks and other feeding and watering appliances. Thoroughly clean all surfaces with soap or detergent and rinse with water. Saturate the surfaces with the disinfecting solution for a period of **10 minutes**. Ventilate building and other closed spaces. Do not house animals or employ equipment until treatment has been absorbed, set or dried. All treated equipment that will contact feed or drinking water must be rinsed with potable water before reuse. Using approved AOAC test methods, in the presence of 5% blood serum and a 3 minute contact time, unless otherwise noted, this product kills the following organisms on hard non-porous inanimate surfaces: **BACTERICIDAL ACTIVITY (BACTERIA):** Pseudomonas aeruginosa, Listeria monocytogenes, Staphylococcus aureus, Salmonella choleraesuis, Enterococcus faecium, Escherichia coli, Escherichia coli O157:H7, Streptococcus pyogenes (Necrotizing Fasciitis-Group A). **ANTIBIOTIC-RESISTANT BACTERICIDAL ACTIVITY:** Methicillin resistant Staphylococcus aureus (MRSA), Vancomycin resistant Enterococcus faecalis (VRE), Vancomycin intermediate resistant Staphylococcus aureus (VISA). **TUBERCULOCIDAL ACTIVITY** - This product exhibits disinfectant efficacy against Mycobacterium tuberculosis BCG at 20 degrees Centigrade when the treated surface is allowed to remain wet for **5 minutes**. ***VIRUCIDAL ACTIVITY** - This product kills on hard, non-porous inanimate surfaces when allowed to remain wet for a **10 minute** contact time against: Hepatitis A Virus (HAV), Poliovirus Type 1, Canine Parvovirus. This product, when used on environmental, inanimate hard surfaces exhibits virucidal activity against: **Hepatitis B Virus (HBV), Hepatitis C Virus (HCV) and Bovine Viral Diarrhea Virus (BVDV)** when the treated surface is allowed to remain wet for **5 minutes**; **Human Coronavirus, SARS associated Coronavirus and Avian Influenza A for 2 minutes; Rhinovirus type 39, and Rotavirus (Norwalk virus); HIV-1 (associated with AIDS) for 1 minute (60 seconds); and Norovirus (Norwalk virus), Feline Calicivirus and Rabies Virus for 30 seconds. KILLS HIV-1, HBV and HCV ON PRE-CLEANED ENVIRONMENTAL SURFACES/OBJECTS PREVIOUSLY SOILED WITH BLOOD/BODY FLUIDS** in health care settings (Hospitals, Nursing Homes) or other settings in which there is an expected likelihood of soiling of inanimate surfaces/objects with blood or body fluids, and in which the surfaces/objects likely to be soiled with blood or body fluids can be associated with the potential for transmission of Human Immunodeficiency Virus Type 1 (HIV-1) (associated with AIDS), Hepatitis B Virus (HBV) and Hepatitis C Virus (HCV). **SPECIAL INSTRUCTIONS FOR CLEANING AND DECONTAMINATION AGAINST HIV-1, HBV, and HCV ON SURFACES/OBJECTS SOILED WITH BLOOD/BODY FLUIDS: PERSONAL PROTECTION:** When handling items soiled with blood or body fluids use disposable latex gloves, gowns, masks, or eye coverings. **CLEANING PROCEDURES:** Blood and other body fluids must be thoroughly cleaned from surfaces and objects before application of this product. **CONTACT TIME:** Allow surface to remain wet for HBV and HCV 5 minutes, for HIV-1 1 minute. These contact times will not control other common types of viruses and bacteria. **DISPOSAL OF INFECTIOUS MATERIALS:** Blood and other body fluids should be autoclaved and disposed of according to local regulations for infectious waste disposal. **FUNGICIDAL ACTIVITY** - This product is fungicidal against the pathogenic fungus, Trichophyton mentagrophytes (Athlete's Foot Fungus) (a cause of ring worm of the foot), when used with a 10 minute contact time on clean, non-porous, hard surfaces as found in bathrooms, shower stalls, locker rooms, or other areas commonly contacted by bare feet. **EFFICACY TESTS HAVE DEMONSTRATED THAT THIS IS AN EFFECTIVE BACTERICIDE, AND VIRUCIDE IN THE PRESENCE OF ORGANIC SOIL (5% BLOOD SERUM).**

Literature Item #31192-92

