

SAFETY INFORMATION

Welcome to the Occipital Bridge. Bridge Explorer Edition gives you, as a developer, the unique opportunity to integrate true 6-DoF positional tracking into the mixed reality experiences you create.

As a developer, you are responsible for ensuring that you develop your applications in a way consistent with industry best practices and for keeping informed regarding relevant industry regulations and scientific literature regarding safety. In addition, you must ensure that all content that you develop complies with all relevant health and safety standards. This includes ensuring that any content you develop includes appropriate consumer notifications.

The unique 6-DoF and 3D-mapping capabilities of Bridge create an opportunity to integrate mixed reality and real-world elements into the virtual reality environment for such applications as collision avoidance and interaction with real-world elements. However, this mixed reality can be confusing and disorienting to users and can create physical hazards. Your development should carefully account for these risks and your consumer warnings should incorporate these potential hazards. You should specifically enumerate the correct user operating environment to ensure physical safety of your users.

In addition to notifying your consumers of all applicable safety risks, you should observe these for yourself and anyone testing your content. Of course, you should also always use good judgment in the development process and avoid prolonged use without breaks. The following is a summary of the relevant health and safety warnings. Our health and safety warnings are continuously being updated. You should regularly check occipital.com/bridge/safety for the latest version.

READ THE FOLLOWING WARNINGS AND INFORMATION BEFORE USE OF THE PRODUCT AND FOLLOW ALL PRODUCT SAFETY AND OPERATING INSTRUCTIONS. FAILURE TO DO SO MAY CAUSE BODILY INJURY, ELECTRIC SHOCK, FIRE, AND OTHER DAMAGE.

Use by children.

Bridge was not designed to be used by children under the age of 13. Children are still in a stage of visual and neurological development, so they should not be permitted to use Bridge. Do not leave the product within the reach of children or allow them to use or play with it.

If children above the age of 13 are permitted to use Bridge, they should do so only under the direct supervision of adults. Adults should monitor older children for any of the

symptoms or risks described here, should limit the time these children are permitted to use Bridge and should ensure that proper breaks are taken from use. Prolonged use by children can result in significant negative impacts to their development.

Health conditions and concerns.

- Do not use Bridge if you are sick, tired or under the influence of drugs or alcohol.
- Consult a physician prior to use if you are pregnant, elderly or have pre-existing medical conditions (such as a heart condition or psychiatric condition).
- If you have had negative physical or psychological reactions (such as PTSD, elevated heart rate, panic attacks, or fainting) to real-life situations do not use Bridge to reproduce similar virtual situations as you may have the same reactions.
- Bridge may trigger seizures, fainting, or severe dizziness even in people who have no history of these conditions. If you have a previous history of epilepsy or seizures, loss of awareness, or other symptoms linked to an epileptic condition, consult your doctor before using the product.

- Overuse of the product may cause repetitive stress injuries. It may also negatively impact hand-eye coordination and balance. Using the product frequently and for prolonged periods of time can make your muscles, joints or skin hurt. If any part of your body becomes tired or sore while using the headset or its components, or if you feel symptoms such as tingling, numbness, burning or stiffness, stop and rest for several hours before using it again.
- If you have any of the following symptoms or other discomfort during or after use, stop use and see a doctor. Following any of these symptoms, do not drive, operate machinery, or take part in activities that may have potentially serious consequences until you have consulted with a doctor.
 1. Seizures, loss of awareness, convulsions, involuntary movements, dizziness, disorientation, nausea, lightheadedness, drowsiness, or fatigue;
 2. Eye pain or discomfort, eyestrain, eye twitching, or vision abnormalities (such as altered, blurred, or double vision);
 3. Excessive sweating, increased salivation, impaired sense of balance, impaired hand-eye coordination, or other symptoms similar to motion sickness.

Bridge can interfere with implanted medical devices, such as pacemakers. If

you have an implanted medical device do not use Bridge until you have consulted with your doctor.

Physical safety.

Content used on Bridge may or may not incorporate some or all of your physical surroundings into the user experience. While using Bridge, you can only see those elements of your physical surroundings that have been incorporated into the content by the developer; as a result, you are susceptible to tripping over or colliding with real-world objects that have not been incorporated into the content you are viewing. You should only use Bridge in a safe environment that has been adjusted to accommodate the content that you are using. You should take special care that you are not near other people, stairs, balconies, windows or other areas that could cause you to fall. You should ensure that you only move in the ways specified by the content that you are using and that any physical objects not incorporated into the content you are using have been removed from the space. Never use the headset while moving in space not designed for your specific content or while operating or sitting in any kind of vehicle.