

## **PA Criteria**

<b>Prior Authorization Group</b>	ACITRETIN
<b>Drug Names</b>	ACITRETIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have severe disease. Must have trial of methotrexate or cyclosporine with inadequate response or significant side effect/toxicity or have a contraindication to these therapies. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Dermatologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ACNE PRODUCTS
<b>Drug Names</b>	AVITA, TAZAROTENE, TAZORAC, TRETINOIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Diagnoses not covered: solar elastosis, sun damage, wrinkles, actinic damage, melasma, lentigines / freckles (hyperpigmented macules, liver spots), heliodermatitis, dermatoheliosis
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ACTIMMUNE
<b>Drug Names</b>	ACTIMMUNE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Diagnoses not covered: basal cell carcinoma of the skin, breast cancer, burn infection, chronic myeloid leukemia, condyloma acuminatum, graft vs. host disease, idiopathic pulmonary fibrosis, Kaposi's sarcoma, malignant mesothelioma, mycobacteriosis, ovarian cancer, rheumatoid arthritis, scleroderma, chronic hepatitis B, Whipple's disease
<b>Required Medical Information</b>	Diagnosis. For severe malignant osteopetrosis: must have diagnosis confirmed by radiological evidence.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	For chronic granulomatous disease: by or in consultation with immunologist, hematologist, rheumatologist, or infectious disease physician. For severe malignant osteopetrosis: by or in consultation with orthopedic surgeon, hematologist, endocrinologist or oncologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ACUTE HAE
<b>Drug Names</b>	ICATIBANT ACETATE, RUCONEST, SAJAZIR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of HAE confirmed by following laboratory values on 2 separate instances (copy of laboratory reports required, must include reference ranges): low C4 complement level in mg/dL, normal C1q complement component level in mg/dL (C1q complement component level not required for patients under age of 18 or patients whose symptoms began before age 18), and either low C1 esterase inhibitor antigenic level in mg/dL or low C1 esterase inhibitor functional level expressed as a percent. Must have chart documentation indicating member has received at least one dose of requested product as treatment for HAE attack in past, responded to medication, and was able to tolerate medication. For reauth, must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or under the direction of a HAE specialist (defined as an allergist/immunologist who attests to clinical experience in HAE).
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	If clinical documentation confirms the required criteria, the requested medication will be approved after consultation with a Medical Director.

**Prior Authorization Group****Drug Names****PA Indication Indicator****Off-label Uses****Exclusion Criteria****Required Medical Information**

ADEFOVIR

ADEFOVIR DIPIVOXIL

All FDA-approved Indications

-

Hepatitis B Virus Drug Resistance panel showing resistance to prior tx w/ adefovir  
Diagnosis. Must have documentation of results of Hep B Virus Drug Resistance panel if previously received antiviral tx regimen for Hep B. Must have documentation of baseline eval and results for following tests: Hep B virus (HBV) DNA viral load, hepatitis B e antigen (HBeAg), antibody to hepatitis B e antigen (anti-HBe), hepatitis B surface antigen (HBsAg), antibody to hepatitis surface antigen (anti-HBs), liver biopsy (if available), alanine aminotransferase (ALT) level and assay reference range. Must have a trial of entecavir with inadequate response, significant side effect/toxicity, contraindication, or documented viral resistance to entecavir or have clinical rationale to support use of adefovir over entecavir. For reauth: must have doc from prescriber indicating continued benefit from tx, doc of recent HBV DNA level, chart doc of HBV Drug Resistance panel if mbr has evidence of virologic breakthrough (greater than 10-fold increase in serum HBV DNA from nadir during tx in mbr who had initial virologic response), and doc of HBeAg/Anti-HBe/HBsAg/Anti-HBs (for mbrs with HBeAg positive and for mbrs with HBeAg negative not falling under any other indications).

**Age Restrictions**

No Age Restrictions

**Prescriber Restrictions**

Infectious disease physician, gastroenterologist, hepatologist, or transplant physician

**Coverage Duration**

365 days or until disease progression or clearance

**Other Criteria**

Regimens/requirements based upon AASLD Practice Guidelines for Chronic Hepatitis B. For HBeAg+ chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN OR evidence of moderate/severe inflammation or signif. fibrosis on biopsy) and have HBV DNA level greater than 20,000 IU/mL (not required for pediatric patients if ALT greater than or equal to 2xULN for longer than 6 months). For HBeAg- chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN, ALT greater than 1xULN w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, ALT less than or equal to ULN w/ ALT increased over time) and 1 HBV DNA criterion (HBV DNA greater than 20,000 IU/mL, HBV DNA greater than 2,000 IU/mL w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, HBV DNA less than or equal to 2,000 IU/mL w/ HBV DNA increased over time). For cirrhosis w/ HBV: must have HBV DNA greater than 2,000 IU/mL OR detectable HBV DNA level w/ elevated ALT. For HBV mbr who had liver txfr for HBV or who received solid organ txfr from HBV+ donor: approve regardless of HBV DNA and ALT levels. For HBV carrier who needs immunosuppressive or cytotoxic tx: must be HBsAg+, have planned course of cancer chemo tx or immunosuppressive tx. Reauth for HBeAg+: approve x1 year until all of following are met (loss of HBeAg, undetectable serum HBV DNA, completed 6-12 months of additional tx after appearance of anti-HBe. Reauth for HBeAg-: approve x1 yr until loss of HBsAg. Reauth for cirrhosis, for liver txfr for HBV, or for solid organ txfr from HBV+ donor: long-term tx approvable. Reauth for HBV carriers receiving immunosuppressive or cytotoxic tx: mbr w/ baseline HBV DNA less than 2,000 IU/mL

should continue x6 months after completion of chemo tx or immunosuppressive tx, mbr w/ baseline HBV DNA greater than 2,000 IU/mL should continue until reach therapeutic endpoints for immunocompetent HBV as listed above.

**Prior Authorization Group**

**Drug Names**

**PA Indication Indicator**

**Off-label Uses**

**Exclusion Criteria**

**Required Medical Information**

ADEMPAS

ADEMPAS

All FDA-approved Indications

-

Current use of nitrate product or phosphodiesterase inhibitor (i.e. sildenafil or tadalafil).  
Diagnosis. Must have baseline negative pregnancy test prior to initiation of riociguat (if a female of childbearing potential). For PAH (WHO Group I), must have diagnosis confirmed by right heart catheterization, must have inadequate response or intolerance to sildenafil (Revatio), AND must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. For reauth: must have chart documentation from prescriber indicating improvement in condition. For CTEPH (WHO Group 4), must be refractory to surgical treatment (i.e. pulmonary endarterectomy) or have inoperable CTEPH, must have chart documentation showing CTEPH confirmed through ventilation-perfusion scanning or pulmonary angiography AND a right heart catheterization that indicates the following hemodynamic values at least 90 days after start of full anticoagulation or 180 days after pulmonary endarterectomy unless there is clinical evidence of right heart failure and pulmonary hypertension on clinical exam and echocardiogram: mean pulmonary arterial pressure greater than 25mmHg and pulmonary vascular resistance greater than 3 Wood units. For reauth: must have documentation from prescriber indicating improvement in condition.

**Age Restrictions**

**Prescriber Restrictions**

**Coverage Duration**

**Other Criteria**

No Age Restrictions

Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.

Initial: 90 days. Reauth: 365 days.

Not Applicable

<b>Prior Authorization Group</b>	AIMOVIG
<b>Drug Names</b>	AIMOVIG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had one of the following: an inadequate response or intolerance to an antiepileptic drug, beta blocker, or antidepressant unless contraindicated, OR at least 3 months previous treatment with the requested drug with a reduction in migraine days per month from baseline. For reauth: must have documentation of decrease in frequency and/or severity of headaches as a result of therapy.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Neurologist, headache specialist, or pain specialist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 1 year.
<b>Other Criteria</b>	Not applicable
<b>Prior Authorization Group</b>	ALECENSA
<b>Drug Names</b>	ALECENSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming ALK mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ALOSETRON
<b>Drug Names</b>	ALOSETRON HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Male gender. Constipation. Anatomical or biochemical abnormalities of the gastrointestinal tract. Concomitant use of fluvoxamine. History of chronic or severe constipation or sequelae from constipation, intestinal obstruction, stricture, toxic megacolon, gastrointestinal perforation, and/or adhesions ischemic colitis, impaired intestinal circulation, thrombophlebitis, or hypercoagulable state, Crohn's disease or ulcerative colitis, diverticulitis, severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis. Must have chronic IBS symptoms. Must have chart documentation of how diagnosis was confirmed. Must have trial of loperamide AND an antispasmodic (e.g. dicyclomine) with inadequate response or significant side effect/toxicity or have a contraindication. For reauth: must have documentation from prescriber indicating improvement in condition and no evidence of constipation or ischemic colitis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Gastroenterologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ALPHA1-PROTEINASE INHIBITORS
<b>Drug Names</b>	ARALAST NP, PROLASTIN-C, ZEMAIRA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Confirmed diagnosis of congenital alpha1-antitrypsin deficiency with clinically evident emphysema or airflow obstruction. Alpha1-antitrypsin phenotype of PI*ZZ, PI*ZNull or PI*NullNull. Baseline (pretreatment) serum alpha1-antitrypsin concentration of less than 11 micromol/L as documented by either of the following: less than 50mg/dL as determined by nephelometry OR less than 80mg/dL as determined by radial immunodiffusion. Must not have selective IgA deficiencies with known antibodies against IgA (anti-IgA antibodies). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a pulmonologist
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ALUNBRIG
<b>Drug Names</b>	ALUNBRIG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming ALK mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	AMBRISANTAN
<b>Drug Names</b>	AMBRISANTAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have baseline negative pregnancy prior to initiation of therapy if a female of child-bearing potential. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ANDROGENS
<b>Drug Names</b>	ANDRODERM, METHYLTESTOSTERONE, TESTOSTERONE, TESTOSTERONE CYPIONATE, TESTOSTERONE ENANTHATE, TESTOSTERONE PUMP
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	APOKYN
<b>Drug Names</b>	APOKYN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Concomitant therapy with 5-HT3 antagonist (e.g. ondansetron)
<b>Required Medical Information</b>	Diagnosis. Must be on concomitant therapy with carbidopa/levodopa AND one of the following numbered options: (1)a dopamine agonist (e.g. ropinirole or pramipexole), (2)a monoamine oxidase-B inhibitor (e.g. rasagiline or selegiline), or (3)a catechol O-methyltransferase inhibitor (e.g. entacapone). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	APTOM
<b>Drug Names</b>	APTOM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 generic antiepileptic drugs (e.g., oxcarbazepine, carbamazepine, lamotrigine, valproic acid, levetiracetam, zonisamide). If using eslicarbazepine as adjunctive therapy to other antiepileptic drugs, cannot be used with oxcarbazepine. Must have documentation of baseline transaminase and bilirubin levels.
<b>Age Restrictions</b>	Age 4 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	ARANESP
<b>Drug Names</b>	ARANESP ALBUMIN FREE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Uncontrolled hypertension, known hypersensitivity to active substance or any excipients of product.
<b>Required Medical Information</b>	Diagnosis. Must have Hgb less than 10g/dL. For anemia due to chemotx for nonmyeloid malignancy: must have documentation of a minimum of 2 more months of chemotx planned. All dx: Must have iron status evaluated before and during treatment with EPO. Reauth for CKD on dialysis: must have Hgb less than 11g/dL. Reauth for CKD not on dialysis: must have Hgb less than 10g/dL. Reauth for anemia due to chemotx for nonmyeloid malignancy: must have Hgb less than 12g/dL and documentation of a minimum 2 more months of chemotx planned. Reauth for other dx: must have Hgb less than 12g/dL.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a nephrologist, hematologist/oncologist, or transplant physician
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 90 days for chemotx, 180 days for other dx.
<b>Other Criteria</b>	Part B versus Part D determination will made at time of prior authorization review per CMS guidance to establish if the drug prescribed is to be used for an ESRD-related condition. If the drug is determined not to be ESRD-related, criteria apply.

<b>Prior Authorization Group</b>	ARCALYST
<b>Drug Names</b>	ARCALYST
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of TNF-blocking or other biologic agent in combination with rilonacept.
<b>Required Medical Information</b>	Diagnosis. Negative tuberculosis skin test, baseline lipid panel assessment. For Muckle-Wells: must have chart doc of diagnosis confirmed by genetic test (must have documentation of lab result confirming mutation in NLRP3 gene) or a clinical diagnosis (must have 3 of following: autosomal dominant pattern of disease inheritance, presence of severe fatigue, presence of musculoskeletal symptoms, presence of ocular symptoms, presence of erythematous rash, duration of most febrile episodes lasting greater than 24 hours, presence of amyloidosis, presence of hearing loss). For Familial Cold Autoinflammatory Syndrome: must have chart doc of diagnosis confirmed by genetic test (must have documentation of lab result confirming mutation in NLRP3 gene) or a clinical diagnosis (must have 4 of following: recurrent intermittent episodes of fever and rash that primarily follow natural/experimental/both types of generalized cold exposures, autosomal dominant pattern of disease inheritance, age of onset less than 6 months of age, duration of most attacks less than 24 hours, presence of conjunctivitis associated with attacks, absence of deafness/periorbital edema/lymphadenopathy/serositis). For recurrent pericarditis: must have a trial and failure of an NSAID, colchicine, or corticosteroids unless intolerant or contraindicated. For reauth: must have documentation from prescriber indicating improvement in condition and assessment of lipid panel within 3 months (1st reauth) and regularly thereafter.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Rheumatologist, dermatologist, immunologist, cardiologist, or genetic specialist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ARIKAYCE
<b>Drug Names</b>	ARIKAYCE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must be used as part of a combination antibacterial drug regimen. Must have failed at least 6 months of treatment with a multidrug regimen. Failure defined as not achieving negative sputum cultures. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	180 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ARIPIPRAZOLE
<b>Drug Names</b>	ARIPIPRAZOLE, ARIPIPRAZOLE ODT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For Major Depressive Disorder without psychosis, must have adequate trial and failure or inadequate response, duration of at least 4 weeks, or intolerance to monotherapy with 2 different antidepressant therapies (e.g. SSRIs or SNRIs) and must be on concomitant therapy with an SSRI or SNRI as adjunctive treatment (which can include medication from monotherapy trial above)
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ARMODAFINIL
<b>Drug Names</b>	ARMODAFINIL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of sleep study confirming diagnosis for narcolepsy and OSA. Must have trial of modafinil with an inadequate response OR have had a significant side effect/toxicity to modafinil. For narcolepsy: must have trial and failure of CNS stimulant (e.g. amphetamine salts, dextroamphetamine, methylphenidate). For shift-work sleep disorder (SWSD), must meet International Classification of Sleep Disorders criteria for SWSD (either primary complaint of excessive sleepiness or insomnia temporarily associated w/ work period that occurs during habitual sleep phase OR polysomnography and Multiple Sleep Latency Test demonstrate loss of normal sleep-wake pattern, no other medical or mental disorders account for symptoms, and symptoms do not meet criteria for any other sleep disorder producing insomnia or excessive sleepiness such as time zone change syndrome) and must provide chart documentation of shift work schedule showing 5 or more night shifts per month (defined as at least 4 hours of shift occurring between 10pm and 8am). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	SWSD: 180 days. Narcolepsy, OSA: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ASENAPINE
<b>Drug Names</b>	ASENAPINE MALEATE SL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have tried and failed 2 atypical antipsychotics.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	AURYXIA
<b>Drug Names</b>	AURYXIA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Diagnoses not covered: iron deficiency anemia in chronic kidney disease not on dialysis.
<b>Required Medical Information</b>	Diagnosis. For hyperphosphatemia in CKD: must be on dialysis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	AUVELITY
<b>Drug Names</b>	AUVELITY
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure or an inadequate response or intolerance to 2 generic formulary antidepressants (e.g., sertraline, citalopram, fluoxetine) unless contraindicated.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	AYVAKIT
<b>Drug Names</b>	AYVAKIT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For unresectable or metastatic gastrointestinal stromal tumor (GIST): must have documentation of platelet-derived growth factor receptor alpha (PDGFRA) exon 18 mutation, including PDGFRA D842V mutations.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist, hematologist, allergist, or immunologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not applicable
<b>Prior Authorization Group</b>	B VS. D
<b>Drug Names</b>	ABELCET, ACETYLCYSTEINE, ACYCLOVIR SODIUM, ALBUTEROL SULFATE, AMBISOME, AMINOSYN M, AMINOSYN-PF 7%, AMPHOTERICIN B, APREPITANT, ARZERRA, AZATHIOPRINE, BUDESONIDE, CASPOFUNGIN ACETATE, CINACALCET HYDROCHLORIDE, CLINIMIX 4.25%/DEXTROSE 1, CLINIMIX 4.25%/DEXTROSE 5, CLINIMIX 5%/DEXTROSE 15%, CLINIMIX 5%/DEXTROSE 20%, CLINIMIX E 2.75%/DEXTROSE, CLINIMIX E 4.25%/DEXTROSE, CLINIMIX E 5%/DEXTROSE 15, CLINIMIX E 5%/DEXTROSE 20, CLINISOL SF 15%, CLONIDINE HCL, CROMOLYN SODIUM, CYCLOPHOSPHAMIDE, CYCLOSPORINE, CYCLOSPORINE MODIFIED, CYTARABINE, EMEND, ENGERIX-B, EPOPROSTENOL SODIUM, FOSCARNET SODIUM, GABLOFEN, GENGRAF, HEPATAMINE, INTRALIPID, IPRATROPIUM BROMIDE, IPRATROPIUM BROMIDE/ALBUT, KABIVEN, LEVALBUTEROL, LEVALBUTEROL HCL, LEVALBUTEROL HYDROCHLORID, METHOTREXATE, METHOTREXATE SODIUM, MYCOPHENOLATE MOFETIL, MYCOPHENOLIC ACID DR, ONDANSETRON HCL, ONDANSETRON HYDROCHLORIDE, ONDANSETRON ODT, PENTAMIDINE ISETHIONATE, PERIKABIVEN, PLENAMINE, PORTRAZZA, PREHEVBRIO, PREMASOL, PROGRAF, PROSOL, RECOMBIVAX HB, SIMULECT, TACROLIMUS, TOBRAMYCIN, TRAVASOL, TROPHAMINE, VECTIBIX, VELETRI, XATMEP
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	-
<b>Required Medical Information</b>	-
<b>Age Restrictions</b>	-
<b>Prescriber Restrictions</b>	-
<b>Coverage Duration</b>	NA
<b>Other Criteria</b>	-

<b>Prior Authorization Group</b>	BALVERSA
<b>Drug Names</b>	BALVERSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For locally advanced or metastatic urothelial carcinoma: must have 1) susceptible FGFR3 or FGFR2 genetic alterations AND 2) progressed during or following at least one line of prior platinum-containing chemotherapy including within 12 months of neoadjuvant or adjuvant platinum-containing chemotherapy.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	BARBITURATES
<b>Drug Names</b>	PHENOBARBITAL, PHENOBARBITAL SODIUM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	BENLYSTA
<b>Drug Names</b>	BENLYSTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Severe active lupus nephritis or severe active central nervous system lupus. Evidence of infection. On concomitant therapy with biologic therapies, including B-cell targeted therapies, or IV cyclophosphamide.
<b>Required Medical Information</b>	Diagnosis. Must be auto-antibody positive, as evidenced through documentation of having one of the following laboratory markers: positive antinuclear antibodies titer greater than or equal to 1:80 or anti-double stranded DNA greater than or equal to 30 IU/mL. Must have trial of hydroxychloroquine, azathioprine, methotrexate, or mycophenolate with inadequate response or significant side effect/toxicity or have a contraindication to these therapies. Must be on concomitant therapy with any of the following (alone or in combination): corticosteroids, antimalarials, NSAIDs, and/or immunosuppressants. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Rheumatologist or nephrologist
<b>Coverage Duration</b>	Initial 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	BESREMI
<b>Drug Names</b>	BESREMI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection
<b>Required Medical Information</b>	Must have a baseline platelet count of at least 50,000 cells/mm <sup>3</sup> prior to initiation. Must currently require phlebotomy and must have trial of hydroxyurea with an inadequate response or significant side effect/toxicity unless contraindicated.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	BOSULIF
<b>Drug Names</b>	BOSULIF
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	BRAFTOVI AND MEKTOVI
<b>Drug Names</b>	BRAFTOVI, MEKTOVI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming BRAFV600E or BRAFV600K mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	BRIVIACT
<b>Drug Names</b>	BRIVIACT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to two other antiepileptic drugs. Must have an evaluation by a psychiatrist and be followed concurrently by a psychiatrist if the member has a history of psychiatric symptoms including anger, aggression, hostility, irritability, suicidal ideation, and homicidal ideation OR if the member is currently undergoing psychiatric treatment.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	BRUKINSA
<b>Drug Names</b>	BRUKINSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For mantle cell lymphoma (MCL): must have received at least one prior therapy. For relapsed or refractory marginal zone lymphoma (MZL): must have received at least one anti-CD20-based regimen.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not applicable
<b>Prior Authorization Group</b>	CABOMETYX
<b>Drug Names</b>	CABOMETYX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For hepatocellular carcinoma: must have been previously treated with sorafenib (Nexavar).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with an oncologist or hematologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CALQUENCE
<b>Drug Names</b>	CALQUENCE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No exclusion criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of trial with at least 1 prior therapy for mantle cell lymphoma. For other diagnoses, no prior therapies required.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	CAPLYTA
<b>Drug Names</b>	CAPLYTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure or an inadequate response or intolerance to 2 oral generic atypical antipsychotics.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CAPRELSA
<b>Drug Names</b>	CAPRELSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CARBAGLU
<b>Drug Names</b>	CARGLUMIC ACID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Confirmed diagnosis of one of the following: N-acetylglutamate synthase (NAGS) deficiency, N-acetylglutamate (NAG) deficiency, carbamoyl phosphate synthetase 1 (CPS 1) deficiency, propionic acidemia (PA), or methylmalonic acidemia (MMA). Must have chart documentation describing how diagnosis was confirmed (e.g. genetic testing results, enzyme assays, ammonia levels, progress notes, etc.). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a physician who specializes in the treatment of inherited metabolic disorders.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	CERDELGA
<b>Drug Names</b>	CERDELGA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	On concomitant therapy with a CYP2D6 inhibitor (e.g. paroxetine) and a strong or moderate CYP3A inhibitor (e.g. ketoconazole) if a CYP2D6 extensive or intermediate metabolizer. On concomitant therapy with a strong CYP3A inhibitor (e.g. ketoconazole) if a CYP2D6 intermediate or poor metabolizer. CYP2D6 ultra-rapid metabolizer.
<b>Required Medical Information</b>	Diagnosis of mild to moderate Type I Gaucher disease with any of the following: hepatomegaly (defined as liver size greater than or equal to 1.25 times normal), splenomegaly (defined as spleen size greater than 0.2% of body weight), bone disease (defined as having one of the following: avascular necrosis, Erlenmeyer flask deformity, lytic disease, marrow infiltrations, osteopenia, osteosclerosis, pathological fracture, or radiological evidence of joint deterioration), or bone marrow disease (defined as having anemia or thrombocytopenia). Must have chart documentation of FDA-cleared test confirming CYP2D6 extensive metabolizer (EM), intermediate metabolizer (IM), or poor metabolizer (PM). For reauth: must have documentation from prescriber indicating improvement in condition and that member is being monitored for neurological side effects of eliglustat (Cerdelga).
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CHEMET
<b>Drug Names</b>	CHEMET
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have blood lead level greater than 45 micrograms per deciliter. Must have chart documentation of identification and removal of the cause of lead exposure. For reauth: must meet initial authorization criteria and have clinical rationale from the prescriber for continuation of treatment.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Toxicologist or other clinician who has experience with chelating agents
<b>Coverage Duration</b>	30 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	CLOBAZAM
<b>Drug Names</b>	CLOBAZAM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Lennox-Gastaut syndrome. Must have had an inadequate response or intolerance to 2 generic antiepileptic drugs (e.g. lamotrigine, topiramate, felbamate) and be using clobazam as adjunctive therapy to other anti-epileptic drugs.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CLONIDINE ER
<b>Drug Names</b>	CLONIDINE HYDROCHLORIDE E
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Attention Deficit Hyperactivity Disorder. Must have trial and failure of clonidine (non-extended release) with inadequate response or significant side effects/toxicity unless contraindicated. Must have trial of a CNS stimulant (e.g. methylphenidate, amphetamine salts) with inadequate response or significant side effects/toxicity unless contraindicated. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	COMETRIQ
<b>Drug Names</b>	COMETRIQ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	COPIKTRA
<b>Drug Names</b>	COPIKTRA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For treatment of relapsed or refractory chronic lymphocytic leukemia (CLL) or small lymphocytic leukemia (SLL): must have tried and failed at least two prior therapies.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CORLANOR
<b>Drug Names</b>	CORLANOR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Blood pressure less than 90/50mmHg. Current acute decompensated heart failure. Sick sinus syndrome, sinoatrial block, or 3rd degree AV block, unless a functioning demand pacemaker is present. Severe hepatic impairment. Dependence on a pacemaker, where heart rate is maintained exclusively by the pacemaker, such as ventricular or atrioventricular pacing more than 40% of the day or demand pacemakers set to a rate greater than 60 beats per minute.
<b>Required Medical Information</b>	Diagnosis. Must currently be taking a beta-blocker (e.g., metoprolol succinate sustained-release, carvedilol, bisoprolol) at maximum tolerated dose for heart failure unless a prior trial with beta-blocker therapy resulted in significant side effect/toxicity or there is a contraindication to use of beta-blocker therapy (e.g., bronchospastic disease such as chronic obstructive pulmonary disease and asthma, severe hypotension or bradycardia). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	COTELLIC
<b>Drug Names</b>	COTELLIC
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For diagnosis of metastatic melanoma with a BRAF V600E or V600K mutation: must have chart documentation of lab result confirming BRAFV600E or BRAFV600K mutation. For diagnosis of histiocytic neoplasms: no other information required.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CRESEMBA
<b>Drug Names</b>	CRESEMBA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	CYSTAGON
<b>Drug Names</b>	CYSTAGON
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must confirm that clinical work-up has been performed to rule out other diagnoses. Diagnosis must be confirmed by having all of the following: elevated white blood cell cystine levels greater than 2nmol per 1/2 cystine per mg of protein, laboratory result confirming CTNS gene mutation, and clinical symptoms of nephropathic cystinosis including electrolyte imbalances and polyuria. For reauth: must have documentation from prescriber indicating improvement in condition and a reduction in WBC cystine levels since starting treatment with oral cysteamine.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a nephrologist or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	CYSTARAN
<b>Drug Names</b>	CYSTARAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis (dx). Must confirm that clinical work-up has been performed to rule out other dx. Must have chart documentation of elevated baseline white blood cell (WBC) cystine level greater than 1 nmol per 1/2 cystine per mg of protein, laboratory result confirming CTNS gene mutation, clinical symptoms consistent with dx (i.e. photophobia, corneal erosions, keratopathies), AND ophthalmologic exam confirming dx. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with an ophthalmologist or a physician who specializes in the treatment of inherited metabolic disorders.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	DALFAMPRIDINE
<b>Drug Names</b>	DALFAMPRIDINE ER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Moderate to severe renal impairment (CrCl less than or equal to 50mL/min), history of seizure, on concomitant therapy with other forms of 4-aminopyridine.
<b>Required Medical Information</b>	Diagnosis. Chart documentation of baseline motor disability or dysfunction. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Neurologist or Physical Medicine and Rehabilitation physician in consultation with the member's treating Neurologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	DALIRESP
<b>Drug Names</b>	DALIRESP, ROFLUMILAST
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Moderate to severe liver impairment.
<b>Required Medical Information</b>	Diagnosis of GOLD Stage III or IV COPD associated with chronic bronchitis. Documentation of COPD exacerbation within the past year. Must have trial and failure of inhaled long-acting beta-agonist or inhaled long-acting anticholinergic or a contraindication to these agents. Must have trial and failure of inhaled glucocorticosteroid or a contraindication to these agents. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	DAURISMO
<b>Drug Names</b>	DAURISMO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	DEFERASIROX
<b>Drug Names</b>	DEFERASIROX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Concomitant advanced malignancy or high-risk myelodysplastic syndrome. Serum creatinine greater than 2 times the age-appropriate upper limit of normal or creatinine clearance less than 40mL/min.
<b>Required Medical Information</b>	Diagnosis. Must have platelet count greater than or equal to 50,000. For treatment of chronic iron overload due to non-transfusion dependent thalassemia syndromes: must have liver iron concentration of at least 5mg of iron per gram dry weight and serum ferritin greater than 300mcg/L. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Due to transfusions: age 2 years or older. Not due to transfusions: age 10 years or older.
<b>Prescriber Restrictions</b>	Hematologist, oncologist, or hepatologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	DEPEN
<b>Drug Names</b>	PENICILLAMINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have baseline (within 6 months) urinalysis, complete blood cell count, platelet count, and hemoglobin. For Wilson's disease, must have chart documentation of how diagnosis was confirmed including at least one of the following: hepatic parenchymal copper content greater than or equal to 250 micrograms per gram dry weight, presence of Kayser-Fleischer Ring in cornea, serum ceruloplasmin level less than 50mg/L, basal 24-hour urinary excretion of copper greater than 100 micrograms (1.6 millimoles), or genetic testing indicating mutation in ATP7B gene. For Cystinuria: must have chart documentation of how diagnosis was confirmed. For Rheumatoid Arthritis: must have severely active disease, must have a trial of methotrexate with inadequate response or significant side effects or toxicity or have a contraindication, and must have a trial of leflunomide, hydroxychloroquine, minocycline, or sulfasalazine with inadequate response or significant side effects or toxicity or have a contraindication. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Wilson's disease: by or in consultation with gastroenterologist or physician who specializes in the treatment of inherited metabolic disorders. Cystinuria: by or in consultation with nephrologist or physician who specializes in the treatment of inherited metabolic disorders. RA: rheumatologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	DIACOMIT
<b>Drug Names</b>	DIACOMIT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 other antiepileptic drugs (such as valproate and topiramate). Must be used as adjunctive therapy with Onfi (clobazam).
<b>Age Restrictions</b>	Age 6 months or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	DRONABINOL
<b>Drug Names</b>	DRONABINOL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For chemotherapy induced nausea and vomiting: 1) must be receiving chemotherapy, 2) must have a trial and failure of a 5HT-3 receptor antagonist (e.g., ondansetron) unless intolerant or contraindicated. For AIDS anorexia, patient must have diagnosis of AIDS with anorexia and weight loss. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, gastroenterologist, or infectious disease physician
<b>Coverage Duration</b>	180 days
<b>Other Criteria</b>	B vs. D determination will be made prior to clinical criteria being applied.
<b>Prior Authorization Group</b>	DUAVEE
<b>Drug Names</b>	DUAVEE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Undiagnosed abnormal uterine bleeding. Known, suspected, or past history of breast cancer. Known or suspected estrogen-dependent neoplasia. Active or past history of venous thromboembolism and/or arterial thromboembolism. Known hepatic impairment or disease. Known protein C, protein S, or antithrombin deficiency or other known thrombophilic disorders. Pregnancy, women who may become pregnant, and nursing mothers.
<b>Required Medical Information</b>	Diagnosis. For moderate to severe vasomotor symptoms associated with menopause: must have documentation of clinical rationale for continued use of Duavee (including an explanation of the member's specific benefit of the drug and how that benefit outweighs the potential risk) AND documentation of previous trial of Femring with an inadequate response or significant side effect/toxicity. For osteoporosis prophylaxis: must have trials with a bisphosphonate (e.g. alendronate) and raloxifene with inadequate responses or significant side effects/toxicities unless contraindicated. For vasomotor symptom reauth: must have documentation of clinical rationale for continued use of Duavee (including an explanation of the member's specific benefit of the drug and how that benefit outweighs the potential risk). For osteoporosis reauth: must have documentation indicating continued benefit with use of Duavee.
<b>Age Restrictions</b>	Age 65 years or older: criteria apply. Age less than 65 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ENBREL
<b>Drug Names</b>	ENBREL, ENBREL MINI, ENBREL SURECLICK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of TNF-blocking or other biologic agent in combination with etanercept.
<b>Required Medical Information</b>	Diagnosis. Negative tuberculosis skin test. For RA and JIA: must have diagnosis of moderately to severely active disease, must have trial of methotrexate with inadequate response (if significant side effects/toxicity or contraindication to methotrexate must have trial of hydroxychloroquine, leflunomide, or sulfasalazine for RA and of leflunomide or sulfasalazine for JIA). For psoriatic arthritis (peripheral disease): must have moderately to severely active psoriatic arthritis AND must have trial of 1 NSAID at target anti-inflammatory dose and of 1 conventional systemic therapy (e.g. methotrexate, cyclosporine, leflunomide, sulfasalazine) with inadequate responses or significant side effects/toxicities or have contraindication to these therapies. For psoriatic arthritis (axial, skin, nail, enthesitis, or dactylitis dominant): must have a trial of 2 NSAIDs at target anti-inflammatory dose with inadequate response or sig. side effects/toxicities unless contraindicated. For ankylosing spondylitis: must have active disease and must have trial of 2 NSAID at target anti-inflammatory dose with inadequate response or significant side effects/toxicity or have a contraindication. For plaque psoriasis: must have chronic moderate to severe plaque psoriasis, must have minimum BSA involvement of at least 5% (not required if plaque psoriasis on palms, soles, head/neck, or genitalia), must have trial of 1 topical treatment or phototherapy or photochemotherapy with inadequate response or significant side effects/toxicity or have a contraindication, and must have trial of 1 conventional systemic therapy (e.g. methotrexate, acitretin, cyclosporine) with inadequate response or significant side effects/toxicity or have a contraindication. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	RA, JIA, ankylosing spondylitis: rheumatologist. Psoriatic arthritis: rheumatologist or dermatologist. Plaque psoriasis: dermatologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ENTECAVIR
<b>Drug Names</b>	BARACLUDE, ENTECAVIR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of results of Hep B Virus Drug Resistance panel if previously received antiviral tx regimen for Hep B. Must have documentation of baseline eval and results for following tests: Hep B virus (HBV) DNA viral load, hepatitis B e antigen (HBeAg), antibody to hepatitis B e antigen (anti-HBe), hepatitis B surface antigen (HBsAg), antibody to hepatitis surface antigen (anti-HBs), liver biopsy (if available), alanine aminotransferase (ALT) level and assay reference range. For reauth: must have doc from prescriber indicating continued benefit from tx, doc of recent HBV DNA level, chart doc of HBV Drug Resistance panel if mbr has evidence or virologic breakthrough (greater than 10-fold increase in serum HBV DNA from nadir during tx in mbr who had initial virologic response), and doc of HBeAg/Anti-HBe/HBsAg/Anti-HBs (for mbrs with HBeAg positive and for mbrs with HBeAg negative not falling under any other indications).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Infectious disease physician, gastroenterologist, hepatologist, or transplant physician
<b>Coverage Duration</b>	365 days or until disease progression or clearance
<b>Other Criteria</b>	Regimens/requirements based upon AASLD Practice Guidelines for Chronic Hepatitis B. For HBeAg+ chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN OR evidence of moderate/severe inflammation or signif. fibrosis on biopsy) and have HBV DNA level greater than 20,000 IU/mL (not required for pediatric patients if ALT greater than or equal to 2xULN for longer than 6 months). For HBeAg- chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN, ALT greater than 1xULN w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, ALT less than or equal to ULN w/ ALT increased over time) and 1 HBV DNA criterion (HBV DNA greater than 20,000 IU/mL, HBV DNA greater than 2,000 IU/mL w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, HBV DNA less than or equal to 2,000 IU/mL w/ HBV DNA increased over time). For cirrhosis w/ HBV: must have HBV DNA greater than 2,000 IU/mL OR detectable HBV DNA level w/ elevated ALT. For HBV mbr who had liver txfr for HBV or who received solid organ txfr from HBV+ donor: approve regardless of HBV DNA and ALT levels. For HBV carrier who needs immunosuppressive or cytotoxic tx: must be HBsAg+, have planned course of cancer chemotx or immunosuppressive tx. Reauth for HBeAg+: approve x1 year until all of following are met (loss of HBeAg, undetectable serum HBV DNA, completed 6-12 months of additional tx after appearance of anti-HBe. Reauth for HBeAg-: approve x1 yr until loss of HBsAg. Reauth for cirrhosis, for liver txfr for HBV, or for solid organ txfr from HBV+ donor: long-term tx approvable. Reauth for HBV carriers receiving immunosuppressive or cytotoxic tx: mbr w/ baseline HBV DNA less than 2,000 IU/mL should continue x6 months after completion of chemotx or immunosuppressive tx, mbr w/ baseline HBV DNA greater than 2,000 IU/mL should continue until reach therapeutic



endpoints for immunocompetent HBV as listed above.

<b>Prior Authorization Group</b>	EPCLUSA
<b>Drug Names</b>	EPCLUSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of chronic Hep C. Doc of prior treatment (tx) for Hep C. Chart doc of lab genotype (GT) result, detectable baseline HCV RNA level (incl. assay date, ref. range), test indicating presence or absence of cirrhosis (e.g. F4 score on liver biopsy from within past 3 years, MRI, ultrasound, CT scan).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Infectious disease physician, gastroenterologist, hepatologist, HIV specialist, or transplant physician
<b>Coverage Duration</b>	12 or 24 weeks based on GT, prior tx, presence of cirrhosis
<b>Other Criteria</b>	Regimens/requirements based on AASLD/IDSA Hep C Tx Guidelines found at <a href="https://www.hcvguidelines.org/">https://www.hcvguidelines.org/</a> .
<b>Prior Authorization Group</b>	EPIDIOLEX
<b>Drug Names</b>	EPIDIOLEX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 generic antiepileptic drugs (e.g. lamotrigine, topiramate, felbamate).
<b>Age Restrictions</b>	Age 1 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	EPOETIN ALFA
<b>Drug Names</b>	PROCRT, RETACRIT
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Uncontrolled hypertension, known hypersensitivity to active substance or any excipients of product.
<b>Required Medical Information</b>	Diagnosis. For Epogen or Procrit requests only: must have had trial of Retacrit with inadequate response or significant side effect/toxicity or contraindication to use of Retacrit. Initial for ribavirin-induced anemia: must have Hgb less than 10g/dL or a 3g/dL decrease from baseline with anemia symptoms and documentation that dose reduction of ribavirin did not resolve anemia. Initial to reduce risk of allogenic blood transfusions: must have Hgb 10-13g/dL and be at high risk for perioperative transfusion due to significant anticipated blood loss and be scheduled to undergo elective, non-cardiac, or nonvascular surgery. Initial for other dx: must have Hgb less than 10g/dL. Initial for anemia due to chemotx for nonmyeloid malignancy: must have documentation of a minimum 2 more months of chemotx planned. Must have iron status evaluated before and during treatment with EPO. Reauth for CKD on dialysis: must have Hgb less than 11g/dL. Reauth for CKD not on dialysis: must have Hgb less than 10g/dL. Reauth for pediatric CKD: must have Hgb less than 12 g/dL. Reauth for anemia due to chemotx for nonmyeloid malignancy: must have Hgb less than 10g/dL and documentation of a minimum 2 more months of chemotx planned. Reauth for other dx: must have Hgb less than 12g/dL.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a nephrologist, hematologist/oncologist, gastroenterologist, hepatologist, transplant physician, surgeon, or an infectious disease physician
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 90 days for d/t chemotx, 180 days for other dx.
<b>Other Criteria</b>	Part B versus Part D determination will made at time of prior authorization review per CMS guidance to establish if the drug prescribed is to be used for an ESRD-related condition. If the drug is determined not to be ESRD-related, criteria apply.
<b>Prior Authorization Group</b>	ERIVEDGE
<b>Drug Names</b>	ERIVEDGE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ERLEADA
<b>Drug Names</b>	ERLEADA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ERLOTINIB
<b>Drug Names</b>	ERLOTINIB HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For 1st-line treatment of patients w/ metastatic NSCLC whose tumors have EGFR exon 19 deletions or exon 21 substitution mutations: must have chart documentation of laboratory result confirming EGFR mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ESBRIET AND OFEV
<b>Drug Names</b>	ESBRIET, OFEV, PIRFENIDONE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For diagnosis of idiopathic pulmonary fibrosis: must have definitive diagnosis of idiopathic pulmonary fibrosis confirmed by either high-resolution computed tomography (HRCT) or surgical lung biopsy, must have all other diagnoses ruled out (e.g., domestic and occupational environmental exposures, connective tissue disease, and drug toxicity), and must submit documentation of baseline liver function testing, including alanine aminotransferase (ALT), aspartate aminotransferase (AST), and bilirubin. For reauth: must have documentation from prescriber indicating that member still is a candidate for treatment and showing that liver function tests (including ALT, AST, and bilirubin) are being monitored regularly.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Pulmonologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 180 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	EUCRISA
<b>Drug Names</b>	EUCRISA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure of moderate to high potency topical corticosteroid or have a contraindication to this therapy (such as dermatitis on face, genitalia) AND must have trial and failure of topical tacrolimus with inadequate response or significant side effect/toxicity or have a contraindication to this therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	EVEROLIMUS ONCOLOGY
<b>Drug Names</b>	EVEROLIMUS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, or neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	EVEROLIMUS TRANSPLANT
<b>Drug Names</b>	EVEROLIMUS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Must have undergone solid organ transplant. Must have trial and failure (defined as intolerance to regimen or inability of regimen to prevent rejection at appropriate therapeutic dosing) of anti-rejection regimen containing at least 2 drugs (including cyclosporine, tacrolimus, azathioprine, mycophenolate mofetil, mycophenolate sodium) unless contraindicated.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a transplant specialist
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	B vs. D determination will be made prior to clinical criteria being applied.

<b>Prior Authorization Group</b>	EXKIVITY
<b>Drug Names</b>	EXKIVITY
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	FANAPT
<b>Drug Names</b>	FANAPT, FANAPT TITRATION PACK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure of risperidone and 1 other atypical antipsychotic (including, but not limited to: aripiprazole, olanzapine, quetiapine, ziprasidone) with inadequate responses or intolerance.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	FARESTON
<b>Drug Names</b>	TOREMIFENE CITRATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have previous inadequate response or intolerance to tamoxifen.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	FENTANYL CITRATE
<b>Drug Names</b>	FENTANYL CITRATE ORAL TRA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Treatment of acute or postoperative pain
<b>Required Medical Information</b>	Diagnosis. Must be opioid tolerant, defined as requiring medication for a week or longer containing at least 60mg/day of morphine. Must currently be using a long-acting opioid. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or pain specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	FETZIMA
<b>Drug Names</b>	FETZIMA, FETZIMA TITRATION PACK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Concomitant therapy with monoamine oxidase inhibitor, linezolid, or intravenous methylene blue.
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure of one generic serotonin norepinephrine reuptake inhibitor (such as venlafaxine ER) indicated for the treatment of major depressive disorder AND one generic selective serotonin reuptake inhibitor (such as citalopram or fluoxetine). If transitioning from a monoamine oxidase inhibitor to levomilnacipran, must have at least a 14-day washout period in between.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

**Prior Authorization Group****Drug Names****PA Indication Indicator****Off-label Uses****Exclusion Criteria****Required Medical Information**

FILGRASTIM

GRANIX, NIVESTYM, ZARXIO

All Medically-accepted Indications

-

No Exclusion Criteria

Diagnosis. Primary prophylaxis of FN: must be receiving myelosuppressive chemo with greater than 20% FN risk OR non-myelosuppressive chemo (less than or equal to 20% FN risk) and considered high risk for chemo-induced FN or infection with at least 1 risk factor (see other criteria) OR dose-dense chemo for tx of node + breast ca, small-cell lung ca, or diffuse aggressive Non-Hodgkin's Lymphoma. Secondary prophylaxis of FN: must have experienced neutropenic complication from prior chemo cycle for which primary prophylaxis not received and in which reduced dose may compromise disease-free or overall survival or tx outcome. Tx of febrile pts w/ neutropenia: must have fever and neutropenia and be at high risk for infection-related complications or have prognostic factors predictive of poor clinical outcomes, have at least one risk factor (see other criteria), AND not have received prophylactic pegfilgrastim during current chemo cycle. Bone marrow txfr: must be used after autologous peripheral blood progenitor cell transplant OR mobilization of progenitor cells into peripheral blood (often in conjunction with chemo) for collection by leukapheresis. AML: must be receiving induction or consolidation tx. ALL: must be using after completion of initial 1st few days of chemo of initial induction or 1st post-remission course. Myelodysplastic syndrome: must have severe neutropenia and recurrent infection. Pts receiving radiation: must be receiving radiation tx w/o concomitant chemo w/ expected prolonged delays due to neutropenia. Older lymphoma pts: must have dx of acute aggressive lymphoma tx w/ curative chemo (CHOP or more aggressive regimen). Congenital, cyclic, or idiopathic neutropenia: must have symptomatic neutropenia. Drug-induced agranulocytosis: must have severe neutropenia w/ fever or serious infection as result of myelosuppressive regimen. For reauth: must have doc from prescriber indicating improvement in condition.

**Age Restrictions****Prescriber Restrictions****Coverage Duration****Other Criteria**

No Age Restrictions

No Prescriber Restrictions

90 days

% risk of FN based on ASCO or NCCN guidelines. Risk factors for primary prophylaxis of FN: age 65 years or older, poor performance status, previous episode of FN, extensive prior treatment including large radiation ports, previous chemotherapy or radiation therapy, pre-existing neutropenia, cytopenia due to bone marrow involvement by tumor, poor nutritional status, presence of open wounds or active infection, recent surgery, advanced cancer, liver dysfunction such as elevated bilirubin, or other serious comorbidities. Risk factors for tx of febrile pts w/ neutropenia: sepsis syndrome, expected prolonged neutropenia for greater than 10 days, severe neutropenia with ANC less than 100/microliter, age 65 years or older, uncontrolled primary disease, pneumonia, hypotension and multi-organ dysfunction (sepsis syndrome), invasive fungal infection, other clinically documented infections, hospitalization at time of fever,

prior episode of febrile neutropenia. For treatment of febrile patients w/ neutropenia: must not have received pegfilgrastim during current chemotx cycle.

<b>Prior Authorization Group</b>	FINTEPLA
<b>Drug Names</b>	FINTEPLA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 other antiepileptic drugs (such as valproate and topiramate).
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	FIRDAPSE
<b>Drug Names</b>	FIRDAPSE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	History of seizures
<b>Required Medical Information</b>	Diagnosis. Reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	FIRST GENERATION ANTIHISTAMINES
<b>Drug Names</b>	CYPROHEPTADINE HCL, CYPROHEPTADINE HYDROCHLOR, HYDROXYZINE HCL, HYDROXYZINE HYDROCHLORIDE, HYDROXYZINE PAMOATE, PROMETHAZINE HCL, PROMETHAZINE HCL PLAIN, PROMETHAZINE HYDROCHLORID, PROMETHEGAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Approve cyproheptadine, hydroxyzine, and promethazine if prior trial and failure, intolerance, toxicity or contraindication of levocetirizine for allergic rhinitis, allergic conditions, or urticaria. Approve promethazine if prior trial and failure, intolerance, toxicity or contraindication of ondansetron for nausea and vomiting. Approve hydroxyzine if prior trial and failure, intolerance, toxicity or contraindication of two therapies (such as SSRIs and SNRIs) for anxiety. For all other FDA-approved indications, no prior drug trials are required.
<b>Age Restrictions</b>	Age 65 years or older: criteria apply. Age less than 65 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	FOTIVDA
<b>Drug Names</b>	FOTIVDA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of a previous trial and failure of at least two systemic therapies for renal cell carcinoma.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	FYCOMPA
<b>Drug Names</b>	FYCOMPA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 other antiepileptic drugs (such as carbamazepine, oxcarbazepine, or phenytoin). Must have documentation indicating the member will be monitored for the psychiatric side effects of perampanel. For reauth: must have documentation from prescriber indicating improvement in condition and monitoring for psychiatric side effects.
<b>Age Restrictions</b>	Age 4 years or older
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	GALAFOLD
<b>Drug Names</b>	GALAFOLD
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For males: must have diagnosis of Fabry disease based upon clinical symptoms or by genetic testing. For females: must have presumed symptoms of Fabry disease (heterozygous carriers) based on family history and/or genetic testing. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	GARDASIL
<b>Drug Names</b>	GARDASIL 9
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	None
<b>Age Restrictions</b>	Between the ages of 9 and 45 years
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	3 doses per 365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	GATTEX
<b>Drug Names</b>	GATTEX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Active intestinal obstruction or active malignancy.
<b>Required Medical Information</b>	Diagnosis of short bowel syndrome. Must provide baseline parenteral or intravenous nutrition (PN/IV) support schedule including frequency and volume, colonoscopy within 6 months before starting teduglutide (if appropriate), and baseline (within 6 months) lab monitoring of bilirubin, alkaline phosphatase, lipase, and amylase. Must be receiving parenteral or intravenous nutrition support at least 3 times per week. For reauth: must have documentation from prescriber indicating improvement in condition, that member has weaned off or decreased PN/IV requirements, that the member had a colonoscopy (if appropriate) after 1 year of teduglutide treatment and at least every 5 years after the 1st year, and that member is undergoing laboratory testing of bilirubin, alkaline phosphatase, lipase, and amylase every 6 months.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a gastroenterologist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	GAVRETO
<b>Drug Names</b>	GAVRETO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Chart documentation of diagnosis with an FDA approved test
<b>Age Restrictions</b>	Age 12 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	GILOTRIF
<b>Drug Names</b>	GILOTRIF
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For first line treatment of metastatic non-small cell lung cancer, must have chart documentation of lab result confirming non-resistant epidermal growth factor receptor (EGFR) mutations. Documentation of mutation(s) not required for treatment of metastatic, squamous NSCLC progressing after platinum-based chemotherapy.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	GLEOSTINE
<b>Drug Names</b>	GLEOSTINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	GRALISE
<b>Drug Names</b>	GRALISE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Postherpetic Neuralgia. Must have trial and failure of tricyclic antidepressant unless contraindicated. Must have trial and failure of gabapentin defined as either failure due to insufficient efficacy at dose of at least 1800mg/day OR chart documented failure due to intolerance despite slow dose titration.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

**Prior Authorization Group****Drug Names****PA Indication Indicator****Off-label Uses****Exclusion Criteria****GROWTH HORMONE**

GENOTROPIN, GENOTROPIN MINIQUICK

All Medically-accepted Indications

-

Active malignancy in the past year. Active proliferative or severe non-proliferative diabetic retinopathy. For Prader-Willi: severely obese (BMI greater than or equal to 97th percentile for age/gender or BMI greater than or equal to 35), history of upper airway obstruction or sleep apnea, or severe respiratory impairment.

**Required Medical Information**

Diagnosis. All dx: chart doc of present height, %, height SD score, pre-tx growth velocity (initial auth), growth velocity on tx (reauth), recent skeletal bone age. Classic growth hormone deficiency (GHD): names/dates of specific GH stim tests, history of irradiation or multiple pituitary hormone deficiency. For chronic renal insufficiency (CRI): estimated date of renal transplant (txfr). Prader-Willi: BMI. For child born small for gestational age (SGA): GA, birth weight and length, height or weight % or SD at birth. Child w/ SHOX deficiency: chart doc of lab of SHOX mutation. Child w/ idiopathic short stature (ISS): doc of growth rates unlikely to permit attainment of adult height w/i target range based on parental heights. Adults: doc of GHD during childhood and cause of GHD (if applicable), serum IGF-I level while not on GH (if applicable), names and dates of specific GH stim tests (if applicable), whether there is pituitary adenoma (and if so, if tumor size has remained stable x1 yr), doc of possible cause of GH deficiency (severe GH deficiency as child d/t genetic cause, severe GH and receipt of high-dose cranial radiation tx, structural hypothalamic-pituitary disease, CNS tumor, deficiencies in pituitary hormones such as ACTH/TSH/prolactin/gonadotropins/arginine vasopressin). GH stim tests accepted for adults: insulin tolerance test (ITT) required unless contraind (pts w/ known or at high risk for CAD, hx of seizures, severe panhypopituitarism/hypoadrenalism) w/ neg response of peak GH less than or equal to 5ug/L, if ITT contraind glucagon test req unless contraind (malnourished or have not eaten in 48 hrs, pheochromocytoma, insulinoma, severe hypocortisolemia) w/ neg response of peak GH less than or equal to 3ug/L, if ITT and glucagon contraind arginine test req w/ neg response of peak GH less than or equal to 0.4ug/L.

**Age Restrictions**

No Age Restrictions

**Prescriber Restrictions**

Endocrinologist, nephrologist, pediatric endocrinologist, or pediatric nephrologist dependent upon diagnosis.

**Coverage Duration**

Idiopathic short stature: 180 days. Other dx: 365 days.

**Other Criteria**

Child w/ Classic GHD: must have doc of failure to respond to 2 GH provocative tests (1 test if h/o irradiation or multiple pituitary hormone deficiency) w/ serum peak GH level less than 10ng/mL on stim tests (insulin, levodopa, arginine, clonidine, glucagon), must have at least 2 of following (present height less than 3rd % or greater than 2 SD below mean for gender/age, pre-tx growth velocity less than 7cm/yr for child less than 3 yrs OR less than 4cm for child 3 yrs and older OR less than 10th % for gender/age based on at least 6 months of growth data for child of any age, comparison of skeletal/bone age by x-ray of left hand and wrist greater than 2 SD below chronological age). Growth retardation d/t CRI, Prader-Willi Syndrome, SHOX Deficiency: documented dx of CRI

up to time of renal txfr (CRI only), must have at least 1 of following (present height less than 3rd % or greater than 2 SD below mean for gender/age, pre-tx growth velocity less than 7cm/yr for child less than 3 yrs OR less than 4cm for child 3 yrs and older OR less than 10th % for gender/age based on at least 6 months of growth data for child of any age). Turner's Syndrome (females), Noonan Syndrome (males, females), must have 1 of following: present height less than 5th % or greater than 2 SD below mean for gender/age, pre-tx growth velocity less than 7cm/yr for child less than 3 yrs OR less than 4cm for child 3 yrs and older OR less than 10th % for gender/age based on at least 6 months of growth data for child of any age. ISS: must have height SD score of less than -2.25cm/yr. SGA: must have low birth weight (either birth weight less than 2500g at GA of more than 37 wks OR birth weight and length less than 3rd % or less than -2 SD for GA), must have failed to achieve catch-up growth by ages 2-4 with baseline pre-tx height SD score less than -2.5 SD for age/gender. Adult w/ GHD, childhood onset: must stop GH tx x1 mon after completion of linear growth and have GH levels reassessed (not req if high likelihood of GHD defined as IGF-1 less than 84ug/L while off GH tx AND at least 1 of following: severe GHD as child d/t genetic cause, structural hypothalamic-pituitary disease, CNS tumors, severe GHD and receipt of high-dose cranial radiation tx, deficiencies in at least 3 pituitary hormones), must have GHD reassessed w/ 1 GH stim test if IGF-1 less than 84ug/L while not on GH tx and w/ 2 GH stim tests if IGF-1 normal while not on GH tx. Adult w/ GHD, adult onset: if panhypopituitarism GH stim test not required if pt has deficiencies in at least 3 pituitary hormones and IGF-1 less than 84ug/L while not on GH tx, if no panhypopituitarism w/ low IGF-1 must have GH deficiency confirmed by 2 GH stim tests. For child reauth: d/c if growth velocity on GH tx less than 2.5cm/yr, reached adult height, growth plates fused, need for renal txfr (for CRI), bone age of 14 in females and 16 in males. For any age: must have doc from prescriber indicating improvement in condition.

**Prior Authorization Group**

**Drug Names**

**PA Indication Indicator**

**Off-label Uses**

**Exclusion Criteria**

**Required Medical Information**

HARVONI

HARVONI

All FDA-approved Indications

-

No Exclusion Criteria

Diagnosis of chronic Hep C. Doc of prior treatment (tx) for Hep C. Chart doc of lab genotype (GT) result, detectable baseline HCV RNA level (incl. assay date, ref. range), test indicating presence or absence of cirrhosis (e.g. F4 score on liver biopsy from within past 3 years, MRI, ultrasound, CT scan).

**Age Restrictions**

**Prescriber Restrictions**

Age 3 years or older

Infectious disease physician, gastroenterologist, hepatologist, HIV specialist, or transplant physician

**Coverage Duration**

**Other Criteria**

12 or 24 weeks based on GT, prior tx, presence of cirrhosis

Regimens/requirements based on AASLD/IDSA Hep C Tx Guidelines found at <https://www.hcvguidelines.org/>.

<b>Prior Authorization Group</b>	HETLIOZ
<b>Drug Names</b>	HETLIOZ, HETLIOZ LQ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must submit chart documentation describing how diagnosis was confirmed (e.g. sleep-wake logs, melatonin secretion abnormalities, progress notes, genetic testing for SMS, etc.). For diagnosis of non-24-hour sleep-wake disorder: patient must be totally blind with no perception of light. For diagnosis of Smith-Magenis Syndrome: no other clinical information required. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By a neurologist or physician who specializes in sleep medicine
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	HIGH RISK ANTIDEPRESSANTS
<b>Drug Names</b>	AMITRIPTYLINE HCL, AMITRIPTYLINE HYDROCHLORI, AMOXAPINE, CLOMIPRAMINE HYDROCHLORID, DESIPRAMINE HYDROCHLORIDE, DOXEPIN HCL, DOXEPIN HYDROCHLORIDE, IMIPRAMINE HCL, IMIPRAMINE HYDROCHLORIDE, IMIPRAMINE PAMOATE, NORTRIPTYLINE HCL, NORTRIPTYLINE HYDROCHLORI, PAROXETINE HCL, PAROXETINE HYDROCHLORIDE
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Approve amitriptyline, amoxapine, clomipramine, desipramine, doxepin (doses higher than 6mg/day), imipramine, imipramine pamoate, nortriptyline, or paroxetine if prior trial and failure of 2 of following for depression: SSRIs, venlafaxine, venlafaxine ER capsules, trazodone, mirtazapine, bupropion. Approve clomipramine if prior trial and failure of 2 of following for obsessive-compulsive disorder: citalopram, escitalopram, fluoxetine, fluvoxamine, sertraline, venlafaxine, venlafaxine ER capsules. Approve doxepin for urticaria/pruritus if prior trial and failure of levocetirizine. For all other FDA-approved indications, no prior drug trials are required.
<b>Age Restrictions</b>	Age 65 years or older: criteria apply. Age less than 65 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	HORIZANT
<b>Drug Names</b>	HORIZANT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For RLS: must have trial and failure of pramipexole or ropinirole (defined as insufficient efficacy of pramipexole 0.5mg per day or ropinirole 4mg per day or intolerance to these meds) AND must have trial and failure of gabapentin (defined as insufficient efficacy of gabapentin 1800mg per day or intolerance to med despite slow dose titration or contraindication). For PHN: must have trial and failure of gabapentin (defined as insufficient efficacy of gabapentin 1800mg per day or intolerance to med despite slow dose titration or contraindication) AND must have trial of tricyclic antidepressant unless intolerant or contraindicated. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	HUMIRA
<b>Drug Names</b>	HUMIRA, HUMIRA PEDIATRIC CROHNS D, HUMIRA PEN, HUMIRA PEN-CD/UC/HS START, HUMIRA PEN-PEDIATRIC UC S, HUMIRA PEN-PS/UV STARTER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of TNF-blocking or other biologic agent in combination with adalimumab.
<b>Required Medical Information</b>	Negative TB skin test. For plaque psoriasis (PS): must have chronic mod to severe dx. For ankylosing spondylitis (AS): must have active dx. For all other dx: must have mod to severely active dx. For RA, JIA: trial of methotrexate (MTX) with inadeq response (if sig. side effects/toxicity or contraindication (CI) to MTX must have trial of hydroxychloroquine for RA or leflunomide or sulfasalazine for RA/JIA). For psoriatic arthritis (PsA) (peripheral disease): must have trial of 1 NSAID at target anti-inflamm dose and of 1 conventional systemic therapy (e.g. methotrexate, cyclosporine, leflunomide, sulfasalazine) with inadeq responses or significant side effects/toxicities or have contraindication to these therapies. For PsA (axial, skin, nail, enthesitis, or dactylitis dominant): must have trial of 2 NSAIDs at target anti-inflammatory dose with inadeq response or sig. side effects/toxicities unless contraindicated. For AS: trial of 2 NSAIDs at target anti-inflam dose with inadeq response or sig. side effects/toxicity or have a CI. For PS: min BSA of at least 5% (not req if on palms, soles, head/neck, genitalia), trial of 1 topical treatment or phototx or photochemotx with inadeq response or sig. side effects/toxicity unless contraindicated, and trial of 1 conventional systemic therapy (e.g. MTX, acitretin, cyclosporine) with inadeq response or sig. side effects/toxicity unless CI. For Crohns, UC: trial of 1 conventional therapy incl corticosteroid, 5-ASA agent (UC only), or immunosuppressant with inadeq response or sig. side effects/toxicity unless CI. For hidradenitis suppurativa (HS): mod or severe dx (w/ 3 active abscesses, inflammatory nodules, or lesions). For uveitis: trial of 1 immunosuppressant (e.g. MTX, mycophenolate, tacrolimus, cyclosporine) with inadeq response or sig. side effects/toxicity unless CI. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	JIA, uveitis: age 2 years or older. Crohns: age 6 years or older. HS: age 12 years or older. Ulcerative colitis: age 5 years or older. Other dx: age 18 years or older.
<b>Prescriber Restrictions</b>	RA, JIA, ankylosing spondylitis: rheumatologist. Psoriatic arthritis: rheumatologist, dermatologist. Plaque psoriasis, HS: dermatologist. Crohn's, UC: gastroenterologist. Uveitis: ophthalmologist, rheumatologist.
<b>Coverage Duration</b>	HS initial: 90 days. HS reauth: 365 days. All other dx: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	HYPNOTICS
<b>Drug Names</b>	ZOLPIDEM TARTRATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a previous trial and failure of therapy with trazodone, Rozerem, or Silenor unless intolerant or contraindicated.
<b>Age Restrictions</b>	Age 65 years or older: criteria apply. Age less than 65 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	IBRANCE
<b>Drug Names</b>	IBRANCE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ICLUSIG
<b>Drug Names</b>	ICLUSIG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or Hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	IDHIFA
<b>Drug Names</b>	IDHIFA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of laboratory result confirming an isocitrate dehydrogenase-2 (IDH2) mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	IMATINIB
<b>Drug Names</b>	IMATINIB MESYLATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, allergist, or immunologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	IMBRUVICA
<b>Drug Names</b>	IMBRUVICA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For mantle cell lymphoma (MCL): must have received at least one prior therapy. For marginal zone lymphoma (MZL): must have received at least one prior anti-CD20-based regimen. For chronic graft vs host disease (cGVHD): must have tried and failed at least one line of systemic therapy (examples may include corticosteroids, cyclosporine, tacrolimus).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, or transplant specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	IMMUNE GLOBULINS
<b>Drug Names</b>	FLEBOGAMMA DIF, GAMMAGARD LIQUID, GAMMAGARD S/D IGA LESS TH, GAMMAKED, GAMUNEX-C, HIZENTRA, PRIVIGEN
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	<p>Diagnosis. Primary immunodeficiency: IgG less than 500mg/dL (clinical rationale for use required if IgG is 500mg/dL or greater) and history of at least 1 bacterial infection directly attributable to deficiency for initial auth and recent IgG level for reauth. Children w/ ITP: platelet count less than: 20,000 and significant mucous membrane bleeding, 10,000 and minor purpura, or 20,000 and inaccessibility or noncompliance is concern, OR need for any surgery, dental extraction, or other procedure likely to cause blood loss. Adults w/ ITP: plt count less than 30,000 and previous documented inadequate response or intolerance to corticosteroids OR need for surgery likely to cause blood loss (platelet count less than or equal to: 10,000 for dentistry, 30,000 for tooth extraction or regional dental block, 50,000 for minor surgery, 80,000 for major surgery). For pregnant women w/ ITP: plt count less than 100,000, history of splenectomy, or previous delivery of infant(s) w/ autoimmune thrombocytopenia. B-Cell CLL: IgG less than 500mg/dL and previous history of serious bacterial infection requiring antibiotics. CIDP: doc of electrodiagnostic testing confirming dx. Multifocal Motor Neuropathy: must have conduction block and progressive symptomatic disease diagnosed on basis of electrophysiologic findings to r/o other possible conditions. For reauth (all dx): must have documentation from prescriber indicating improvement in condition.</p>
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	<p>Primary immunodeficiency: by or in consultation w/ immunologist, hematologist. ITP: hematologist, oncologist. B-cell CLL: hematologist, oncologist, ID specialist. CIDP, Multifocal Motor Neuropathy: neurologist.</p>
<b>Coverage Duration</b>	ITP: 30 days. CIDP, multifocal motor neuropathy: 90 days. Other dx: 365 days.
<b>Other Criteria</b>	BvsD determination will be made prior to clinical criteria being applied.

<b>Prior Authorization Group</b>	INCRELEX
<b>Drug Names</b>	INCRELEX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Closed epiphyses, presence of active or suspected neoplasia, allergy to mecasermin, current treatment with growth hormone replacement therapy, secondary forms of IGF-1 deficiency (e.g. growth hormone deficiency, malnutrition not corrected prior to start of mecasermin, hypothyroidism not corrected prior to start of mecasermin, chronic treatment with pharmacological dose of anti-inflammatory steroids)
<b>Required Medical Information</b>	Diagnosis. For growth hormone deletion: must have growth hormone (GH) gene deletion in gene GH1 and developed neutralizing antibodies to GH therapy. For growth failure due to severe IGF-1 deficiency: must have dx of severe IGF-1 deficiency (defined as having all of the following: height standard deviation (SD) score less than or equal to -3.0 for age and sex, basal IGF-1 SD of less than or equal to -3.0 based on lab reference range, normal or elevated GH defined as stimulated serum GH level of greater than 10ng/mL or basal serum GH level greater than 5ng/mL). For reauth, must have documentation of recent progress note from prescriber indicating growth and maturation as a result of treatment and that epiphyses have not closed.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	Endocrinologist with appropriate endocrinologist follow-up.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	INLYTA
<b>Drug Names</b>	INLYTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or Hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	INQOVI
<b>Drug Names</b>	INQOVI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	INREBIC
<b>Drug Names</b>	INREBIC
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a baseline platelet count of at least 50,000 cells/mm <sup>3</sup> prior to initiation of fedratinib.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	IRESSA
<b>Drug Names</b>	IRESSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming epidermal growth factor receptor (EGFR) exon 19 deletion or exon 21 (L858R) substitution mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ITRACONAZOLE
<b>Drug Names</b>	ITRACONAZOLE
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For dermatological mycoses: must be too large to treat with topical antifungals, or have not responded, had an intolerance or contraindication to at least 1 topical antifungal agent (e.g. ciclopirox, clotrimazole, ketoconazole, or nystatin). For onychomycosis: must have trial and failure, intolerance, or contraindication to 1 course (3 months) of oral terbinafine. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	JAKAFI
<b>Drug Names</b>	JAKAFI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection
<b>Required Medical Information</b>	Diagnosis of intermediate or high-risk myelofibrosis (includes primary myelofibrosis, post-polycythemia vera myelofibrosis, and post-essential thrombocythemia myelofibrosis). Must have a baseline platelet count of at least 50,000 cells/mm <sup>3</sup> prior to initiation of ruxolitinib. For diagnosis of polycythemia vera: must currently require phlebotomy and must have trial of hydroxyurea with an inadequate response or significant side effect/toxicity unless contraindicated. For diagnosis of acute graft-versus-host disease: no other information required.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, or transplant specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	JUXTAPID
<b>Drug Names</b>	JUXTAPID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Members who are pregnant, who have moderate or severe hepatic impairment (Child-Pugh category B or C) or active liver disease, or who are on concomitant moderate or strong CYP3A4 inhibitors.
<b>Required Medical Information</b>	Diagnosis of homozygous familial hypercholesterolemia confirmed by genetic testing with functional mutation(s) in both LDL receptor alleles or alleles known to affect LDL receptor functionality or have clinical diagnosis defined as one of the following (1) untreated LDL greater than 500mg/dL AND untreated total cholesterol (TC) greater than 500mg/dL and triglycerides (TG) less than 300mg/dL with both parents with untreated TC greater than 250mg/dL, (2) skin fibroblast LDL receptor activity less than 20% of normal AND untreated TC greater than 500mg/dL and TG less than 300mg/dL with both parents with untreated TC greater than 250mg/dL, (3) presence of cutaneous and tendon xanthomas and corneal arcus in first decade of life AND untreated TC greater than 500mg/dL and TG less than 300mg/dL with both parents with untreated TC greater than 250mg/dL, (4) untreated LDL greater than 500mg/dL AND skin fibroblast LDL receptor activity less than 20% of normal, (5) untreated LDL greater than 500mg/dL AND presence of cutaneous and tendon xanthomas and corneal arcus in first decade of life. Must have chart documentation of clinical work-up to rule out other diagnoses. For initial auth: baseline negative pregnancy test with date of test and be on effective contraception for females of reproductive potential, and baseline laboratory monitoring of LDL, total cholesterol, triglycerides, transaminases, alkaline phosphatase, and bilirubin with date of test. Must be on statin (e.g. atorvastatin, simvastatin) unless intolerant or contraindicated and another LDL-lowering medication from a different class (e.g. ezetimibe, colestipol) prior to starting lomitapide. For reauth: documentation from prescriber indicating improvement in condition, documentation of follow-up LDL levels showing reduction in LDL level since starting treatment, and documentation of monitoring of transaminase, alkaline phosphatase, and bilirubin levels.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Lipidologist, cardiologist, endocrinologist, or geneticist
<b>Coverage Duration</b>	Initial: 120 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	KALYDECO
<b>Drug Names</b>	KALYDECO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Documentation of lab result confirming at least one copy of a mutation in the CFTR gene that is responsive to ivacaftor based on clinical and/or in vitro assay data. Baseline percent of predicted FEV1. For reauth: must have documentation from prescriber showing member benefit from treatment, clinical rationale to support continuation of therapy, and current percent predicted FEV1.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Pulmonologist or cystic fibrosis specialist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	KERENDIA
<b>Drug Names</b>	KERENDIA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must be using concurrently with maximally tolerated dose of an ACE inhibitor (e.g., lisinopril) or ARB (e.g., losartan) unless intolerant or contraindicated.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable

**Prior Authorization Group****Drug Names****PA Indication Indicator****Off-label Uses****Exclusion Criteria**

KEVEYIS

KEVEYIS

All FDA-approved Indications

-

Concomitant use with high dose aspirin. Severe pulmonary disease, limiting compensation to metabolic acidosis that may be caused by dichlorphenamide. Hepatic encephalopathy.

**Required Medical Information**

For primary hypokalemic periodic paralysis: must have documentation confirming diagnosis, defined as one of the following scenarios: two or more attacks of muscle weakness with documented serum K less than 3.5mEq/L, or one attack of muscle weakness in the member with documented serum potassium less 3.5mEq/L and one attack of weakness in a relative with a hx of the condition, or three of the following clinical/laboratory features: onset of symptoms in the first or second decade of life, duration of attack (muscle weakness involving one or more limbs) longer than two hours, presence of triggers (previous carbohydrate rich meal, symptom onset during rest after exercise or during stressful situations) for attacks, improvement in symptoms with potassium intake, family hx of the condition or genetically confirmed skeletal calcium or sodium channel mutation, positive long exercise test. For hyperkalemic periodic paralysis: must have documentation confirming diagnosis based on genetics or clinical presentation (see other coverage criteria).

**Age Restrictions**

No Age Restrictions

**Prescriber Restrictions**

No Prescriber Restrictions

**Coverage Duration**

Initial 90 days. Reauth: 365 days.

**Other Criteria**

For primary hypokalemic periodic paralysis: must have documentation excluding other causes of hypokalemia (renal, adrenal, thyroid dysfunction, renal tubular acidosis, diuretic and laxative abuse) and must currently be using a potassium supplement. For hyperkalemic periodic paralysis confirmed based on genetics testing: must have both of the following: family hx of the condition or genetically confirmed skeletal sodium channel mutation associated with hyperkalemic periodic paralysis and a hx of at least two attacks of flaccid limb weakness (which may also include weakness of the muscles of the eyes, throat, and trunk) or 1 attack with a family hx of attacks of hyperkalemic periodic paralysis. For hyperkalemic periodic paralysis confirmed based on clinical presentation must have all of the following: a hx of at least two attacks of flaccid limb weakness (which may also include weakness of the muscles of the eyes, throat, and trunk) or 1 attack with a family hx of attacks of hyperkalemic periodic paralysis, serum potassium greater than 5mEq/L or an increase of serum potassium concentration of at least 1.5 mEq/L during an attack of weakness and/or onset/worsening of an attack as a result of oral potassium intake, and presence of myotonia or any 3 of the following clinical features: typical attack duration less than 2 hours, onset before 30 years, positive long exercise test (greater than 40% decrement in CMAP), or typical external triggers (rest after exercise, potassium load, fasting). For hyperkalemic periodic paralysis: must have documentation of normal serum potassium concentration and muscle strength between attacks and electroencephalogram (ECG) recording for the

exclusion of long QTc interval and ventricular arrhythmias. For hyperkalemic periodic paralysis: must not have secondary hyperkalemic periodic paralysis due to ingestion of potassium or of a potassium sparing diuretic or paramyotonia (i.e. muscle stiffness that is worsening after exercise or cold-induced). For hyperkalemic periodic paralysis must have documentation of exclusion of other hereditary forms of hyperkalemia (i.e., Andersen-Tawil syndrome) and acquired forms of hyperkalemia (drug abuse, renal and adrenal dysfunction). For reauth: must have documentation from prescriber indicating improvement in condition.

***Prior Authorization Group***

***Drug Names***

***PA Indication Indicator***

***Off-label Uses***

***Exclusion Criteria***

***Required Medical Information***

***Age Restrictions***

***Prescriber Restrictions***

***Coverage Duration***

***Other Criteria***

KISQALI

KISQALI, KISQALI FEMARA 200 DOSE, KISQALI FEMARA 400 DOSE, KISQALI FEMARA 600 DOSE

All FDA-approved Indications

-

No Exclusion Criteria

Diagnosis.

No Age Restrictions

Oncologist or hematologist

365 days

Not Applicable

<b>Prior Authorization Group</b>	KORLYM
<b>Drug Names</b>	KORLYM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Pregnant. History of unexplained vaginal bleeding, endometrial hyperplasia with atypia, or endometrial carcinoma. Concomitant therapy with simvastatin, lovastatin, or CYP3A4 substrates with narrow therapeutic range (i.e. cyclosporine, tacrolimus). Concurrent long-term corticosteroid treatment.
<b>Required Medical Information</b>	Diagnosis. Must have failed surgery or not be a candidate for surgery (trans-sphenoidal surgery for pituitary dependent Cushing's or surgical removal of an adrenocortical tumor or a source of ectopic ACTH in malignant Cushing's). Female members of reproductive potential: must have baseline (within previous month, must include date of test) negative pregnancy test prior to starting mifepristone and must be using non-hormonal medically acceptable method of contraception (unless surgically sterilized) during treatment and for 1 month after mifepristone therapy. Must have baseline hemoglobin A1C level. For reauth: must have documentation from prescriber indicating improvement in condition, must have documentation of recent (within previous month) negative pregnancy test including date of test if female of reproductive potential, and must have documentation of improvement in hyperglycemia control as evidenced by a reduction in blood glucose levels, HbA1c, or anti-hyperglycemic medication doses or number of medications.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	KOSELUGO
<b>Drug Names</b>	KOSELUGO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	By or in consultation with an oncologist, ophthalmologist, hereditary disorder specialist, or neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	KRAZATI
<b>Drug Names</b>	KRAZATI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Hematologist or oncologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	KYNMOBI
<b>Drug Names</b>	KYNMOBI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Concomitant therapy with 5-HT3 antagonist (e.g. ondansetron)
<b>Required Medical Information</b>	Diagnosis. Must be on concomitant therapy with carbidopa/levodopa AND one of the following numbered options: (1)a dopamine agonist (e.g. ropinirole or pramipexole), (2)a monoamine oxidase-B inhibitor (e.g. rasagiline or selegiline), or (3)a catechol O-methyltransferase inhibitor (e.g. entacapone). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LATUDA
<b>Drug Names</b>	LATUDA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure or an inadequate response or intolerance to 2 generic antipsychotics (e.g., haloperidol, fluphenazine, chlorpromazine, perphenazine, aripiprazole, olanzapine, quetiapine, risperidone, ziprasidone).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	LENVIMA
<b>Drug Names</b>	LENVIMA 10 MG DAILY DOSE, LENVIMA 12MG DAILY DOSE, LENVIMA 14 MG DAILY DOSE, LENVIMA 18 MG DAILY DOSE, LENVIMA 20 MG DAILY DOSE, LENVIMA 24 MG DAILY DOSE, LENVIMA 4 MG DAILY DOSE, LENVIMA 8 MG DAILY DOSE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LEUKINE
<b>Drug Names</b>	LEUKINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For AML: must be receiving induction chemotherapy. For bone marrow transplant, must have one of the following: must require administration after autologous (not allogeneic) bone marrow transplant for NHL/ALL/Hodgkin's disease, must require mobilization of progenitor cells into peripheral blood (often in conjunction with chemotherapy) for collection by leukapheresis, must have undergone allogeneic bone marrow transplant from HLA-matched related donor, OR must have undergone allogeneic or autologous bone marrow transplantation where engraftment is delayed or has failed. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	AML: age 55 years or older. BMT: age 2 years or older.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	90 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	LEUPROLIDE AND DERIVATIVES
<b>Drug Names</b>	ELIGARD, FIRMAGON, LEUPROLIDE ACETATE, LUPANETA PACK, LUPRON DEPOT (1-MONTH), LUPRON DEPOT (3-MONTH), LUPRON DEPOT (4-MONTH), LUPRON DEPOT (6-MONTH), SYNAREL, TRELSTAR MIXJECT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For endometriosis: must have diagnosis confirmed by laparoscopy OR chart documentation of clinical work-up and clinical rationale for diagnosis, and must have trial and failure of oral contraceptives and/or progestins for mild disease. For fibroids: must be used preoperatively to maximize hemoglobin in patients with documented anemia (Hgb less than 11g/dL), or preoperatively to decrease size of fibroid uterus so less invasive route of hysterectomy can be attempted, or must provide clinical rationale if using outside context of preoperative adjuvant therapy in the surgical management of fibroids. For central precocious puberty: must have onset of secondary sexual characteristics earlier than age 8 years in females and 9 years in males. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Central Precocious Puberty: only be approved up to age 11 years in females and age 12 years in males
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Cancer, CPP: 365 dys. Endometriosis: 180 dys. Fibroid: 90 dys.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LIDOCAINE PATCH
<b>Drug Names</b>	LIDOCAINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	LOKELMA
<b>Drug Names</b>	LOKELMA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of elevated serum potassium. Must have trial and failure of therapy with sodium polystyrene sulfonate with inadequate response or significant side effects/toxicity unless contraindicated. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 1 year.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LONG ACTING OPIOIDS
<b>Drug Names</b>	FENTANYL, MORPHINE SULFATE ER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial of a short acting opioid with inadequate response or significant side effect/toxicity or have a contraindication to these therapies.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LONSURF
<b>Drug Names</b>	LONSURF
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. ECOG Performance Status.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	LORBRENA
<b>Drug Names</b>	LORBRENA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LUMAKRAS
<b>Drug Names</b>	LUMAKRAS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Chart documentation of lab results confirming KRAS G12C-mutation.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not applicable
<b>Prior Authorization Group</b>	LYBALVI
<b>Drug Names</b>	LYBALVI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have tried and failed 2 atypical antipsychotics.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	LYNPARZA
<b>Drug Names</b>	LYNPARZA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	MAVYRET
<b>Drug Names</b>	MAVYRET
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh class B or C)
<b>Required Medical Information</b>	Diagnosis of chronic Hep C. Doc of prior treatment (tx) for Hep C. Chart doc of lab genotype (GT) result, detectable baseline HCV RNA level (incl. assay date, ref. range), test indicating presence or absence of cirrhosis (e.g. F4 score on liver biopsy from within past 3 years, MRI, ultrasound, CT scan).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Infectious disease physician, gastroenterologist, hepatologist, HIV specialist, or transplant physician
<b>Coverage Duration</b>	8-16 weeks
<b>Other Criteria</b>	Regimens/requirements based on AASLD/IDSA Hep C Tx Guidelines found at <a href="https://www.hcvguidelines.org/">https://www.hcvguidelines.org/</a> .
<b>Prior Authorization Group</b>	MEGESTROL
<b>Drug Names</b>	MEGESTROL ACETATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, or HIV specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	MEKINIST
<b>Drug Names</b>	MEKINIST
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Treatment with prior BRAF-inhibitor therapy if using trametinib as monotherapy
<b>Required Medical Information</b>	Diagnosis. For unresectable, metastatic, or completely resected melanoma: must have chart documentation of lab result confirming BRAFV600E or BRAFV600K mutation. For metastatic non-small cell lung cancer or locally advanced or metastatic anaplastic thyroid cancer: must have chart documentation of lab result confirming BRAFV600E mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	MEMANTINE
<b>Drug Names</b>	MEMANTINE HCL TITRATION P, MEMANTINE HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of moderate to severe dementia of Alzheimer's type. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Must be age 18 or older. Age 18 to 40 years: criteria apply. Age greater than 40 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	METHAMPHETAMINE
<b>Drug Names</b>	METHAMPHETAMINE HCL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 6 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	METHOXSALEN
<b>Drug Names</b>	METHOXSALEN
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For psoriasis: must have severe, recalcitrant, disabling psoriasis confirmed by biopsy, must use in conjunction with UVA light therapy, and must have a trial of 2 topical treatments (e.g. calcipotriene, fluocinonide, betamethasone, hydrocortisone, clobetasol propionate) with an inadequate response or significant side effects /toxicity or have a contraindication. For vitiligo: must use in conjunction with UVA light therapy and must have a trial of calcipotriene with an inadequate response or significant side effect/toxicity or have a contraindication. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Psoriasis, vitiligo: dermatologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	MODAFINIL
<b>Drug Names</b>	MODAFINIL
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Chart documentation of sleep study confirming diagnosis for narcolepsy and OSA. For narcolepsy: must have trial and failure of CNS stimulant (e.g. amphetamine salts, dextroamphetamine, methylphenidate). For shift-work sleep disorder (SWSD): must meet International Classification of Sleep Disorders criteria for SWSD (either primary complaint of excessive sleepiness or insomnia temporarily associated w/ work period that occurs during habitual sleep phase OR polysomnography and Multiple Sleep Latency Test demonstrate loss of normal sleep-wake pattern, no other medical or mental disorders account for symptoms, and symptoms do not meet criteria for any other sleep disorder producing insomnia or excessive sleepiness such as time zone change syndrome) and must provide chart documentation of shift work schedule showing 5 or more night shifts per month (defined as at least 4 hours of shift occurring between 10pm and 8am). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	SWSD: 180 days. Narcolepsy, OSA: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	MOLINDONE
<b>Drug Names</b>	MOLINDONE HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure or an inadequate response or intolerance to 2 generic antipsychotics (e.g., haloperidol, fluphenazine, chlorpromazine, perphenazine, aripiprazole, olanzapine, quetiapine, risperidone, ziprasidone).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	MULTIPLE SCLEROSIS
<b>Drug Names</b>	BETASERON, COPAXONE, FINGOLIMOD, GILENYA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth, documentation from provider showing disease has improved or stabilized while on therapy.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Neurologist or gastroenterologist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	MYALEPT
<b>Drug Names</b>	MYALEPT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	HIV-related lipodystrophy
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of a clinical work-up to rule out other diagnoses and clinical rationale for the diagnosis and exclusion of other diagnoses. Must have severe insulin resistance resulting in diabetes mellitus (a hemoglobin A1c of at least 7% or fasting plasma glucose of at least 126mg/dL) with chart documentation showing a trial of diabetic pharmacotherapy (such as with an insulin product) that did not allow the member to achieve adequate glucose control AND/OR must have severe hypertriglyceridemia (triglyceride level of at least 500mg/dL) with chart documentation of a trial of lipid-lowering pharmacotherapy (such as a fibrate, omega-3 fatty acid, or statin) that did not allow the member to achieve adequate triglyceride control. For reauth: must have documentation from prescriber indicating benefit with metreleptin treatment (as evidenced by decrease in hemoglobin A1c, fasting plasma glucose, and/or triglyceride levels from baseline).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Endocrinologist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	MYTESI
<b>Drug Names</b>	MYTESI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must be actively utilizing antiretroviral agents to treat HIV/AIDS. Must have documentation of persistent loose stools despite regular use of at least one anti-diarrheal medication (e.g. loperamide, diphenoxylate/atropine). For reauth: must have documentation from prescriber indicating improvement in condition, decrease in number of watery bowel movements, and continuous use of antiretroviral agents for HIV/AIDS.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	HIV or infectious disease specialist
<b>Coverage Duration</b>	Initial: 30 days. Reauth: 180 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NATPARA
<b>Drug Names</b>	NATPARA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Pagets disease of the bone or unexplained elevations of alkaline phosphatase, hereditary disorders predisposing to osteosarcoma or prior external beam or implant radiation therapy involving the skeleton. Hypoparathyroidism caused by calcium-sensing receptor mutations. Acute (less than 6 months) post-surgical hypoparathyroidism.
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of a laboratory report (including reference range) of a recent parathyroid hormone level below the lower limit of normal. Must have uncontrolled hypocalcemia confirmed by chart documentation of a laboratory report (including reference range) of a recent calcium level below the lower limit of normal. Must have a baseline serum calcium concentration greater than 7.5mg/dL prior to initiating parathyroid hormone (Natpara) therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Endocrinologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	NERLYNX
<b>Drug Names</b>	NERLYNX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NEUPRO
<b>Drug Names</b>	NEUPRO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For Parkinson's disease: must have trial and failure of pramipexole or ropinirole. For Restless Legs Syndrome: must have moderate to severe dx, must have trial and failure of pramipexole or ropinirole (defined as insufficient efficacy of pramipexole 0.5mg per day or ropinirole 4mg per day, intolerance to a lower dose of one of these medications, or contraindication) AND must have trial and failure of gabapentin. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NEXAVAR
<b>Drug Names</b>	SORAFENIB TOSYLATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	NINLARO
<b>Drug Names</b>	NINLARO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NITISINONE
<b>Drug Names</b>	NITISINONE, ORFADIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Hereditary Tyrosinemia Type 1. Laboratory test of baseline succinylacetone (SA) level, liver evaluation, and ophthalmologic testing. For reauth: must have documentation from prescriber indicating improvement in condition, monitoring for hematologic and hepatic side effects, and laboratory test demonstrating progressive SA suppression.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a gastroenterologist, hematologist, nephrologist, or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	NORTHERA
<b>Drug Names</b>	DROXIDOPA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of symptomatic neurogenic hypotension caused by 1 of following: primary autonomic failure (e.g. Parkinson's disease, multiple system atrophy, pure autonomic failure), dopamine beta-hydroxylase deficiency, or non-diabetic autonomic neuropathy. Must have chart doc showing how diagnosis made, incl BP readings showing systolic blood pressure decrease of at least 20mmHg or diastolic blood pressure decrease of at least 10mmHg within 3 minutes of standing. Must have doc that member is symptomatic as result of low BP readings, including doc to support that member is experiencing at least 1 of the following symptoms: dizziness, lightheadedness, feeling faint, feeling like might black out. Must have chart doc indicating d/c or dose decrease of drugs which can cause orthostatic hypotension such as anti-hypertensives, nitrates, alpha-1 blockers (i.e. terazosin, prazosin), antiparkinsonian agents (i.e. levodopa, bromocriptine, ropinirole, pramipexole), diuretics, monoamine oxidase inhibitors, narcotics/tranquilizers/sedatives, drugs for erectile dysfunction, tricyclic antidepressants. Must have a trial of midodrine. For reauth: must have doc from prescriber indicating improvement in condition as evidenced by improvement in the symptoms member was experiencing (i.e. dizziness, lightheadedness, feeling faint, or feeling like might black out) and showing member is being monitored for adverse effects (i.e. supine hypertension) and additional drugs have not been added to the drug regimen that would cause supine hypertension.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Cardiologist or neurologist
<b>Coverage Duration</b>	Initial: 1 month. Reauth: 6 months.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NUBEQA
<b>Drug Names</b>	NUBEQA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	NUCALA
<b>Drug Names</b>	NUCALA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For severe persistent asthma: Must have blood eosinophil count of greater than 150 cells/uL within the past six weeks (while on corticosteroid) or greater than or equal to 300 cells/uL within past year, including date test performed. Must have trial of combination therapy with an ICS/LABA (inhaled corticosteroid/long-acting beta-agonist, such as Advair, Breo Ellipta, or Symbicort) AND either a LAMA (long-acting muscarinic antagonist, such as Spiriva) or a leukotriene receptor antagonist (such as montelukast) with inadequate response or significant side effects/toxicities or have a contraindication to these therapies. Must have asthma symptoms that continue to be uncontrolled on optimized medication therapy regimen (uncontrolled defined as hospitalization for asthma within past year, requirement for oral or parenteral corticosteroids to control exacerbations of asthma on 2 occurrences in the past year, or need for daily corticosteroid with inability to taper off). For eosinophilic granulomatosis with polyangiitis: Must have trial of an oral corticosteroid and an immunosuppressant (such as methotrexate). For chronic rhinosinusitis with nasal polyps (CRSwNP): Must have tried and failed therapy with an intranasal corticosteroid (e.g., fluticasone, flunisolide) unless intolerant or contraindicated, and must be used in combination with another agent for CRSwNP (can be an intranasal corticosteroid). For reauth: must have documentation from prescriber indicating improvement in condition (such as reduced exacerbations, hospitalizations, emergency department visits, requirement for oral corticosteroid therapy, or reduction in nasal polyps score, as appropriate for diagnosis).
<b>Age Restrictions</b>	Severe persistent asthma: Age 6 years or older, Eosinophilic granulomatosis with polyangiitis: Age 18 years or older
<b>Prescriber Restrictions</b>	Severe persistent asthma: By or in consultation with an allergist, an immunologist, or a pulmonologist, Eosinophilic granulomatosis with polyangiitis: By or in consultation with an allergist, immunologist, pulmonologist, rheumatologist, or hematologist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	NUEDEXTA
<b>Drug Names</b>	NUEDEXTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Members on quinidine, quinine, mefloquine, MAOIs in the last 14 days, drugs that prolong the QT interval and are metabolized by CYP2D6. History of hypersensitivity to quinidine, quinine, mefloquine, or dextromethorphan. Diagnosis of prolonged QT interval, congenital long QT syndrome or a history suggestive of torsades de pointes, heart failure, complete AV (atrioventricular) block without an implanted pacemaker, or high risk of complete AV block
<b>Required Medical Information</b>	Diagnosis of pseudobulbar affect (PBA) supported by chart documentation of the following: involuntary outbursts of laughing and/or crying that are incongruous or disproportionate to the patient's emotional state AND documentation of a clinical work-up, including clinical rationale for the PBA diagnosis and exclusion of other possible conditions that could result in emotional lability (e.g. depression, bipolar disorder, schizophrenia, epilepsy). Must have underlying neurological disorder such as amyotrophic lateral sclerosis, multiple sclerosis, Alzheimer's and related diseases, Stroke, Traumatic Brain Injury, or Parkinsonian Syndrome. For reauth: must have documentation from prescriber indicating decrease in number of laughing and/or crying episodes as a result of therapy.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	NUPLAZID
<b>Drug Names</b>	NUPLAZID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must provide chart documentation of clinical work-up to rule out other diagnoses. Clinical rationale for diagnosis and exclusion of other diagnoses must be provided. Must have tried to discontinue or reduce the dose of any medication(s) that may cause or contribute to hallucinations and delusions (i.e. dopamine agonists, amantadine, monoamine oxidase B inhibitors, anticholinergics) or provide clinical rationale indicating why dose reduction or discontinuation of applicable medications would not be appropriate.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a psychiatrist or neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	NURTEC
<b>Drug Names</b>	NURTEC
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For preventive treatment of migraine, must have had an inadequate response or intolerance to an antiepileptic drug, beta blocker, or antidepressant unless contraindicated or intolerant. For acute treatment of migraine: must have had an inadequate response or intolerance to TWO generic triptans (e.g., sumatriptan, rizatriptan, naratriptan, zolmitriptan) unless contraindicated or intolerant. For reauth: must have documentation of decrease in frequency and/or severity of headaches as a result of therapy.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Neurologist, headache specialist, or pain specialist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 1 year.
<b>Other Criteria</b>	Not applicable
<b>Prior Authorization Group</b>	OCALIVA
<b>Drug Names</b>	OCALIVA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Complete biliary obstruction
<b>Required Medical Information</b>	Diagnosis. Must have a diagnosis of primary biliary cholangitis (PBC) defined by meeting at least two of the following criteria: 1) chart doc of lab result showing elevated alkaline phosphatase (ALP) above the upper limit of normal (ULN) for at least 6 months based on the reference range provided by lab, 2) positive anti-mitochondrial antibody (AMA) titer, 3) liver biopsy consistent with PBC. Must have an adequate trial of at least 12 months with ursodiol at a dose of 13-15 mg/kg/day with an inadequate response (defined as ALP 1.5-times the ULN) or intolerance at any dose or must have a contraindication to ursodiol. Must be used in combination with ursodiol unless clinically contraindicated or intolerant to ursodiol. For reauth: Must have documentation from provider showing disease has improved while on therapy and monitoring of liver function tests occurring annually.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a gastroenterologist, hepatologist, or liver transplant specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ODOMZO
<b>Drug Names</b>	ODOMZO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ONUREG
<b>Drug Names</b>	ONUREG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	OPSUMIT
<b>Drug Names</b>	OPSUMIT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have baseline hemoglobin and liver function tests (AST, ALT) prior to initiation of therapy. Must have baseline negative pregnancy prior to initiation of therapy if a female of child-bearing potential. Must have previous inadequate response or intolerance to ambrisentan (Letairis). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ORAL BUDESONIDE
<b>Drug Names</b>	BUDESONIDE, BUDESONIDE ER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ORALAIR
<b>Drug Names</b>	ORALAIR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	History of any severe systemic allergic reaction. History of eosinophilic esophagitis. Severe, unstable or uncontrolled asthma. On concomitant immunotherapy.
<b>Required Medical Information</b>	Diagnosis. Must have grass pollen-induced allergic rhinitis with or without conjunctivitis. Must have diagnosis confirmed by positive skin test or in vitro testing for pollen-specific IgE antibodies for Timothy grass or cross-reactive grass pollens (i.e. Sweet Vernal, Orchard, Perennial Rye, Kentucky Blue Grass) and chart documentation demonstrating seasonal symptoms to grass-pollen from the previous pollen season. Must have chart documentation demonstrating daily concomitant use of an inhaled nasal corticosteroid (i.e. fluticasone) and an oral antihistamine (i.e. levocetirizine) during the previous pollen season with inadequate responses or significant side effects/toxicities or have contraindication to these therapies. Must have plan for first dose to be administered in physician office due to potential for life-threatening allergic reactions, including anaphylaxis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 5 years through 65 years
<b>Prescriber Restrictions</b>	Allergist or immunologist
<b>Coverage Duration</b>	180 days
<b>Other Criteria</b>	Therapy must be initiated 4 months prior to the onset of grass pollen season.

<b>Prior Authorization Group</b>	ORFADIN SUSPENSION
<b>Drug Names</b>	ORFADIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Hereditary Tyrosinemia Type 1. Laboratory test of baseline succinylacetone (SA) level, liver evaluation, and ophthalmologic testing. Must have chart documentation of the clinical rationale for why nitisinone capsule cannot be used. For reauth: must have documentation from prescriber indicating improvement in condition, monitoring for hematologic and hepatic side effects, and laboratory test demonstrating progressive SA suppression.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a gastroenterologist, hematologist, nephrologist, or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ORGOVYX
<b>Drug Names</b>	ORGOVYX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Must be prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ORKAMBI
<b>Drug Names</b>	ORKAMBI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Documentation of lab result confirming the following mutation in CFTR gene: F508del. Baseline percent of predicted FEV1. For reauth: must have chart documentation from prescriber showing member benefit from treatment, clinical rationale to support continuation of therapy, and current percent predicted FEV1.
<b>Age Restrictions</b>	Age 1 year or older
<b>Prescriber Restrictions</b>	Pulmonologist or cystic fibrosis specialist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	OTREXUP AND RASUVO
<b>Drug Names</b>	OTREXUP, RASUVO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial of oral methotrexate or generically-available subcutaneous methotrexate with an inadequate response OR have had a significant side effect/toxicity. For Otrexup requests must have a trial of Rasuvo with an inadequate response OR have had a significant side effect/toxicity. For reauth: must have documentation from the prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	RA, polyarticular JIA: rheumatologist. Psoriasis: dermatologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	OXANDROLONE
<b>Drug Names</b>	OXANDROLONE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Carcinoma of breast or prostate in male patients. Carcinoma of breast in female patients with hypercalcemia. Pregnancy. Nephrosis (i.e. nephrotic phase of nephritis). Hypercalcemia. Severe hepatic dysfunction.
<b>Required Medical Information</b>	Diagnosis. Must be used as adjunctive therapy to medically-accepted treatment for the diagnosis. For reauth: must have documentation from prescriber indicating improvement or stabilization in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Osteoporosis bone pain: endocrinologist or orthopedist. Chronic infection: immunologist or infectious disease specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PALIPERIDONE
<b>Drug Names</b>	PALIPERIDONE ER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of schizophrenia or schizoaffective disorder. Must have trial and failure of 2 oral atypical antipsychotics.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	PANRETIN
<b>Drug Names</b>	PANRETIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating stabilization or improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a dermatologist, oncologist, or HIV specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PEGFILGRASTIM
<b>Drug Names</b>	FULPHILA, NYVEPRIA, UDENYCA, ZIEXTENZO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must be receiving chemo regimen with dosing frequency of once every 2 wks or greater. For primary prophylaxis of febrile neutropenia (FN): must be receiving either myelosuppressive chemo regimen with greater than 20% risk of FN (per ASCO or NCCN guidelines) or non-myelosuppressive chemo regimen (less than or equal to 20% risk of FN) and considered to be at high risk for chemo-induced FN or infection with at least one risk factor (age 65 years or older, poor performance status, previous episode of FN, extensive prior treatment including large radiation ports, previous chemotherapy or radiation therapy, pre-existing neutropenia, cytopenia due to bone marrow involvement by tumor, poor nutritional status, presence of open wounds or active infection, recent surgery, advanced cancer, liver dysfunction such as elevated bilirubin, or other serious comorbidities). For secondary prophylaxis of FN: must have experienced a neutropenic complication from prior chemo cycle for which primary prophylaxis was not received and in which reduced dose may compromise disease-free or overall survival or treatment outcome. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	90 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	PEMAZYRE
<b>Drug Names</b>	PEMAZYRE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PHEOCHROMOCYTOMA
<b>Drug Names</b>	METYROSINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have surgical resection planned, have a contraindication to surgery, or have malignant pheochromocytoma. For reauth: must have documentation from prescriber indicating stabilization or improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a nephrologist, oncologist, endocrinologist, or endocrine surgeon
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PICATO
<b>Drug Names</b>	PICATO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Initial auth: actinic keratosis must be present on face, scalp, trunk, or extremities and pt must have adequate trial of topical fluorouracil or imiquimod 5% with inadequate response unless intolerant or contraindicated. For reauth: must meet initial auth criteria and must have either clinical rationale from the prescriber for continuation of treatment at the same site or documentation that therapy is required at an alternative site.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a dermatologist or oncologist
<b>Coverage Duration</b>	3 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	PIQRAY
<b>Drug Names</b>	PIQRAY 200MG DAILY DOSE, PIQRAY 250MG DAILY DOSE, PIQRAY 300MG DAILY DOSE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	POMALYST
<b>Drug Names</b>	POMALYST
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For diagnosis of multiple myeloma: must have received at least two prior therapies including lenalidomide and a proteasome inhibitor (example: Velcade).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	POSACONAZOLE
<b>Drug Names</b>	NOXAFIL, POSACONAZOLE DR
<b>PA Indication Indicator</b>	All Medically-accepted Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Members on the following medications: terfenadine, astemizole, cisapride, pimizide, halofantrine, quinidine, sirolimus.
<b>Required Medical Information</b>	Diagnosis. For prophylaxis of Aspergillus and Candida infections (tablet, injection, or suspension): must be severely immunocompromised. For treatment of oropharyngeal candidiasis (suspension): must have trial and failure of fluconazole or itraconazole for at least 10 days. For treatment of invasive aspergillosis: no other documentation required. For reauth: must have documentation from prescriber indicating clinical rationale for retreatment.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Prophy of Aspergill/Candida: 120 dys. Tx of aspergillosis: 90 dys. Oropharyngeal Candida: 30 dys.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	PRALUENT
<b>Drug Names</b>	PRALUENT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have confirmed diagnosis of primary hyperlipidemia (including heterozygous familial hypercholesterolemia), homozygous familial hypercholesterolemia, or clinical atherosclerotic cardiovascular disease (ASCVD). For ASCVD: must have chart documentation confirming history of at least one of the following: myocardial infarction or other acute coronary syndromes (including ST-elevation myocardial infarction, non-ST-elevation myocardial infarction, and unstable angina), coronary or other revascularization procedure, ischemic stroke or transient ischemic attack, atherosclerotic peripheral arterial disease (includes ankle/brachial index of less than 0.90), coronary artery calcium greater than or equal to 300 Agatston units or greater than or equal to 75th percentile for age/sex/ethnicity, carotid plaque greater than or equal to 50%, coronary atherosclerosis as demonstrated by angiography (cardiac CT angiography or conventional cardiac catheterization). Must have baseline and target LDL-cholesterol levels. Must have LDL-C level above target despite trial of 2 high intensity statins (atorvastatin 40-80mg daily and rosuvastatin 20-40mg daily), unless intolerant to statin treatment (defined as confirmed, intolerable statin-related adverse effects or biomarker abnormalities that improve or resolve with statin dose decrease or discontinuation) or statin treatment is contraindicated (defined as documented active liver disease, which may include unexplained persistent elevations in hepatic transaminase levels). If able to tolerate statin, must continue treatment with statin at maximally tolerated dose. For reauth: must have documentation of: (1) recent assessment of LDL-C level with decrease and (2) continued treatment with maximally tolerated dose of a statin (if applicable).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a cardiologist or an endocrinologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PREVMIS
<b>Drug Names</b>	PREVMIS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	100 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	PROLIA
<b>Drug Names</b>	PROLIA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure of bisphosphonate therapy unless intolerant or contraindicated. For postmenopausal osteoporosis in females or to increase bone mass in males with osteoporosis at high risk of fracture: must have bone mineral density T-score of less than or equal to -2.5 at conventional skeletal sites including the total hip, femoral neck, lumbar spine (post-anterior, not lateral) or radius OR must have history of fragility fracture as an adult. For females with breast cancer: must be receiving aromatase inhibitor therapy. For males with non-metastatic prostate cancer: must be receiving androgen deprivation therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	PROMACTA
<b>Drug Names</b>	PROMACTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For ITP: must have platelet count less than 30,000. For thrombocytopenia associated with chronic Hepatitis C: must have platelet count of less than 75,000 and currently be on treatment with or anticipating hepatitis C treatment with interferon product. For aplastic anemia: must have severe disease, must have platelet count less than 30,000, and must either be used in combination with standard immunosuppressive therapy or have previous inadequate response or intolerance to antithymocyte globulin-based immunosuppressive therapy (Atgam, Thymoglobulin). For reauth: must have documentation from prescriber indicating improvement in condition (all dx), improvement in platelet count from baseline (all dx), and hematologic response (aplastic anemia dx: increase in platelet count, increase in Hgb, increase in ANC, reduction in frequency of platelet or RBC transfusions).
<b>Age Restrictions</b>	ITP: age 1 year or older. Severe aplastic anemia: age 2 years or older. Other diagnoses: age 18 years or older.
<b>Prescriber Restrictions</b>	ITP, aplastic anemia: hematologist or oncologist. Hep C: gastroenterologist, hematologist, hepatologist, or infectious disease specialist.
<b>Coverage Duration</b>	90 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	PULMOZYME
<b>Drug Names</b>	PULMOZYME
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of cystic fibrosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Pulmonologist
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	BvsD determination will be made prior to clinical criteria being applied.
<b>Prior Authorization Group</b>	PURIXAN
<b>Drug Names</b>	PURIXAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure of mercaptopurine tablets or have chart documentation of the clinical rationale for why the tablet version cannot be used.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a hematologist or an oncologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	QINLOCK
<b>Drug Names</b>	QINLOCK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For diagnosis of advanced gastrointestinal stromal tumor (GIST): must have received prior treatment with 3 or more kinase inhibitors, including imatinib.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	QUETIAPINE
<b>Drug Names</b>	QUETIAPINE FUMARATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	QUETIAPINE ER
<b>Drug Names</b>	QUETIAPINE FUMARATE ER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For schizophrenia, bipolar disorder: must have previous trial and failure of immediate-release quetiapine. For major depressive disorder: must have adequate trial and failure (duration at least 4 weeks) with 2 different antidepressant therapies (e.g. SSRIs, SNRIs) with inadequate responses or intolerance.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	QUININE
<b>Drug Names</b>	QUININE SULFATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of malaria. For reauth: must have documentation from prescriber indicating continued benefit from therapy.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	RAVICTI
<b>Drug Names</b>	RAVICTI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation describing how diagnosis was confirmed (e.g. genetic testing results, enzyme assays, ammonia levels, progress notes, etc.). Must have chart documentation of a trial of sodium phenylbutyrate with either inadequate response despite dose titration or significant side effect/toxicity or have a contraindication to this therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation a with physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	REGRANEX
<b>Drug Names</b>	REGRANEX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Neoplasm at site of application
<b>Required Medical Information</b>	Diagnosis. For treatment of diabetic neuropathic ulcers, patient must have a lower extremity diabetic neuropathic ulcer. Treatment will be given in combination with ulcer wound care (i.e. debridement, infection control, and/or pressure relief). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	20 weeks
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	RELISTOR
<b>Drug Names</b>	RELISTOR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For opioid-induced constipation and advanced life-limiting illness: must have documentation of previous trial of lactulose. For opioid-induced constipation with chronic non-cancer pain: must have documentation of current and ongoing opioid therapy and must have trials of 2 of the following with inadequate responses or significant side effects/toxicity or have a contraindication to these therapies: naloxegol (Movantik), lubiprostone (Amitiza), or lactulose. For reauth: must have documentation from prescriber indicating improvement in condition (both diagnoses) and must continue to be on opioid therapy (non-cancer pain).
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 90 days (non-cancer pain), 120 days (life-limiting illness). Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	RETEVMO
<b>Drug Names</b>	RETEVMO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Current weight.
<b>Age Restrictions</b>	Thyroid cancer: 12 years of age or older. NSCLC: 18 years of age or older.
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	REVLIMID
<b>Drug Names</b>	LENALIDOMIDE, REVLIMID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	REXULTI
<b>Drug Names</b>	REXULTI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For Major Depressive Disorder: must have trial and failure or inadequate response or intolerance to aripiprazole and must be on concomitant therapy with an SSRI or SNRI as adjunctive treatment. For Schizophrenia: must have a trial and failure or inadequate response or intolerance to 2 generic oral atypical antipsychotics (e.g. aripiprazole, olanzapine, quetiapine, risperidone, ziprasidone).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	REZUROCK
<b>Drug Names</b>	REZUROCK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Trial and failure, or intolerance of at least two anti-rejection systemic therapies.
<b>Age Restrictions</b>	Age 12 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a transplant specialist or oncologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not applicable

<b>Prior Authorization Group</b>	RINVOQ
<b>Drug Names</b>	RINVOQ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of another JAK inhibitor or other biologic agent in combination with upadacitinib.
<b>Required Medical Information</b>	Negative TB skin test. For psoriatic arthritis (PsA) or ankylosing spondylitis (AS): must have active disease. For all other dx: must have mod to severely active dx. For RA: trial of methotrexate (MTX) and one TNF blocker with inadeq response (if sig. side effects/toxicity or contraindication (CI) to MTX must have trial of hydroxychloroquine, leflunomide, or sulfasalazine). For psoriatic arthritis (peripheral disease): must have moderately to severely active psoriatic arthritis AND must have trial of 1 NSAID at target anti-inflammatory dose and of 1 conventional systemic therapy (e.g. methotrexate, cyclosporine, leflunomide, sulfasalazine) with inadequate responses or significant side effects/toxicities or have contraindication to these therapies. For psoriatic arthritis (axial, skin, nail, enthesitis, or dactylitis dominant): must have a trial of 2 NSAIDs at target anti-inflammatory dose with inadeq response or sig. side effects/toxicities unless contraindicated. For atopic dermatitis: trial of 1 systemic drug product (e.g., methotrexate, azathioprine, or cyclosporine) unless CI. For UC: trial of 1 conventional therapy incl corticosteroid, 5-ASA agent (UC only), or immunosuppressant AND 1 TNF blocker with inadeq response or sig. side effects/toxicity unless CI. For active non-radiographic axial spondyloarthritis (nr-axSpA): no drug trials required. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Atopic dermatitis: age 12 years or older. Other dx: age 18 years or older.
<b>Prescriber Restrictions</b>	RA, nr-axSpA: rheumatologist. Psoriatic arthritis, atopic dermatitis: rheumatologist, dermatologist. UC: gastroenterologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ROZLYTREK
<b>Drug Names</b>	ROZLYTREK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	RUBRACA
<b>Drug Names</b>	RUBRACA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	RUFINAMIDE
<b>Drug Names</b>	RUFINAMIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Lennox-Gastaut Syndrome. Must have had an inadequate response or intolerance to 2 generic antiepileptic drugs (e.g. lamotrigine, topiramate, felbamate) and be using rufinamide as adjunctive therapy to other antiepileptic drugs (which can include medication from trial above).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	RYDAPT
<b>Drug Names</b>	RYDAPT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For acute myeloid leukemia: must have chart documentation of lab result confirming FLT3 mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SAPROPTERIN
<b>Drug Names</b>	JAVYGTOR, SAPROPTERIN DIHYDROCHLORI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Baseline serum phenylalanine level. For reauth: must have documentation from prescriber indicating response to therapy, follow-up serum phenylalanine level.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 60 days. Reauth: 365 days.
<b>Other Criteria</b>	Continuation/Discontinuation criteria: lab reassessment will be conducted after an initial one month trial to determine if authorization may be extended. Patients on the 10mg/kg/day dose whose blood phenylalanine levels have not decreased from baseline after 1 month of treatment should increase to 20mg/kg/day. These patients will be approved for another one month trial at the higher dose. Patients on the 20mg/kg/day dose whose blood phenylalanine levels have not decreased from baseline after 1 month are considered non-responders, and treatment with sapropterin should be discontinued in these patients.
<b>Prior Authorization Group</b>	SAVELLA
<b>Drug Names</b>	SAVELLA, SAVELLA TITRATION PACK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation showing history of widespread pain involving the extremities for 3 months and localized areas of tenderness. Must have chart documentation or claims history showing a trial of gabapentin at a dose of at least 1200mg/day with inadequate response or significant side effects/toxicity despite slow dose titration or have a contraindication to this therapy. Must have chart documentation or claims history showing a trial of a tricyclic antidepressant (e.g. amitriptyline) or muscle relaxant (e.g. cyclobenzaprine) with inadequate response or significant side effects/toxicity or have a contraindication to these therapies. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SCEMBLIX
<b>Drug Names</b>	SCEMBLIX
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SECUADO
<b>Drug Names</b>	SECUADO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have tried and failed 2 atypical antipsychotics.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SEROSTIM
<b>Drug Names</b>	SEROSTIM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have tried and failed 2 other medications used for AIDS wasting (e.g. dronabinol, megestrol). For reauth: must have documentation from prescriber that member has experienced weight stabilization or weight gain.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 84 days. Reauth: 252 dys. Total treatment not to exceed: 336 dys/yr.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SIGNIFOR
<b>Drug Names</b>	SIGNIFOR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of confirmed pituitary source of Cushing's syndrome. Must previously have had pituitary surgery (e.g. transsphenoidal surgery) that was not curative unless not a candidate for surgery. Must have baseline 24-hour urinary free cortisol level. Must have recent (within 6 months) baseline assessments of fasting plasma glucose, liver function tests, electrocardiogram, gallbladder ultrasound, pituitary hormones (e.g. TSH, free T4, growth hormone, IGF-1), and hemoglobin A1C. For reauth: must have documentation from prescriber indicating improvement in condition based on reduction in 24-hour urinary free cortisol level from baseline level as well as signs and symptoms of improvement in the disease (e.g. blood pressure, lipids, weight) and must have documentation that hemoglobin A1C, fasting plasma glucose, liver function tests, gallbladder ultrasound, pituitary hormones, and electrocardiogram have all been reassessed within 3 months of starting pasireotide and at regular intervals thereafter.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SILDENAFIL
<b>Drug Names</b>	SILDENAFIL CITRATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Current use of nitrate product
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have WHO Functional Class II-IV symptoms. For sildenafil suspension: must have chart documentation of the clinical rationale for why sildenafil tablet cannot be used. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	SIROLIMUS
<b>Drug Names</b>	SIROLIMUS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Must have diagnosis of lymphangioleiomyomatosis or prophylaxis of organ rejection. For prophylaxis of organ rejection, must have undergone solid organ transplant and must have at least one of the following: renal dysfunction, coronary allograft vasculopathy following heart transplant, OR trial and failure (defined as intolerance to regimen or inability of regimen to prevent rejection at appropriate therapeutic dosing) of anti-rejection regimen containing at least 2 drugs (including cyclosporine, tacrolimus, azathioprine, mycophenolate mofetil, mycophenolate sodium).
<b>Age Restrictions</b>	Prophylaxis of organ rejection: age 13 years or older. Lymphangioleiomyomatosis: age 18 years or older.
<b>Prescriber Restrictions</b>	Prophylaxis of organ rejection: by or in consultation with a transplant specialist. Lymphangioleiomyomatosis: pulmonologist, hematologist, or oncologist.
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	B vs. D determination will be made prior to clinical criteria being applied.
<b>Prior Authorization Group</b>	SIRTURO
<b>Drug Names</b>	SIRTURO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have previously had inadequate response to at least one first-line TB regimen containing isoniazid and rifampin OR have chart documentation of susceptibility testing of Mycobacterium tuberculosis isolates demonstrating resistance to isoniazid and rifampin. Must be using bedaquiline in combination with at least three other drugs active against pulmonary TB. For reauth: must have documentation from prescriber indicating member's initial response to therapy and clinical rationale for continuation of treatment or for re-treatment AND must have chart documentation of susceptibility testing of Mycobacterium tuberculosis isolates demonstrating continued susceptibility to bedaquiline.
<b>Age Restrictions</b>	Age 5 years or older
<b>Prescriber Restrictions</b>	By or in consultation with an infectious disease specialist or pulmonologist
<b>Coverage Duration</b>	180 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	SKELETAL MUSCLE RELAXANTS
<b>Drug Names</b>	CARISOPRODOL, CHLORZOXAZONE, CYCLOBENZAPRINE HYDROCHLO, METHOCARBAMOL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have attestation from prescriber assessing the risks and benefits of therapy and desire to prescribe a muscle relaxant.
<b>Age Restrictions</b>	Age 65 years or older: criteria apply. Age less than 65 years: criteria do not apply.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SKYRIZI
<b>Drug Names</b>	SKYRIZI, SKYRIZI PEN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of another biologic agent in combination with risankizumab.
<b>Required Medical Information</b>	Negative TB skin test. For psoriatic arthritis (PsA): must have active disease. For plaque psoriasis (PS): must have moderate to severe active dx. For psoriatic arthritis (peripheral disease): must have moderately to severely active psoriatic arthritis AND must have trial of 1 NSAID at target anti-inflammatory dose and of 1 conventional systemic therapy (e.g. methotrexate, cyclosporine, leflunomide, sulfasalazine) with inadequate responses or significant side effects/toxicities or have contraindication to these therapies. For psoriatic arthritis (axial, skin, nail, enthesitis, or dactylitis dominant): must have a trial of 2 NSAIDs at target anti-inflammatory dose with inadequate response or sig. side effects/toxicities unless contraindicated. For plaque psoriasis: must have chronic moderate to severe plaque psoriasis, must have minimum BSA involvement of at least 5% (not required if plaque psoriasis on palms, soles, head/neck, or genitalia), must have trial of 1 topical treatment or phototherapy or photochemotherapy with inadequate response or significant side effects/toxicity or have a contraindication, and must have trial of 1 conventional systemic therapy (e.g. methotrexate, acitretin, cyclosporine) with inadequate response or significant side effects/toxicity or have a contraindication. For Crohns: trial of 1 conventional therapy including corticosteroid or immunosuppressant with inadequate response or sig. side effects/toxicity unless contraindicated. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older.
<b>Prescriber Restrictions</b>	Psoriatic arthritis: rheumatologist, dermatologist. Plaque psoriasis: dermatologist. Crohns disease: gastroenterologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SODIUM PHENYLBUTYRATE
<b>Drug Names</b>	SODIUM PHENYLBUTYRATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation describing how diagnosis was confirmed (e.g. genetic testing results, enzyme assays, ammonia levels, progress notes, etc.). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a hematologist, nephrologist, or physician who specializes in the treatment of inherited metabolic disorders.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SOMAVERT
<b>Drug Names</b>	SOMAVERT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of acromegaly. Must have following baseline labs: elevated serum IGF-1 level for gender/age range (including lab reference range) and elevated growth hormone level defined as GH at least 1ng/mL during oral glucose tolerance test. Must have inadequate response to surgery or radiation therapy or documentation that these therapies are inappropriate. Must have inadequate response to 1 medical therapy (e.g. octreotide, lanreotide) or documentation that these therapies are inappropriate. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with an endocrinologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SPRITAM
<b>Drug Names</b>	SPRITAM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to generic levetiracetam and one other generic antiepileptic drug (such as carbamazepine, oxcarbazepine, or phenytoin).
<b>Age Restrictions</b>	Age 4 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SPRYCEL
<b>Drug Names</b>	SPRYCEL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	STELARA
<b>Drug Names</b>	STELARA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Negative TB skin test. For plaque psoriasis (PS): must have chronic moderate to severe dx, must have minimum BSA of at least 5% (not required if on palms, soles, head/neck, genitalia), and must have had an inadequate response, intolerance, or contraindication to Enbrel or Humira. For Crohn's disease: must have moderately to severely active disease, and must have had an inadequate response, intolerance, or contraindication to Humira. For psoriatic arthritis: must have had an inadequate response, intolerance, or contraindication to Enbrel, Humira or Xeljanz/Xeljanz XR. For ulcerative colitis: must have had an inadequate response, intolerance, or contraindication to Humira or Xeljanz/Xeljanz XR. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Plaque psoriasis, psoriatic arthritis: 6 years or older. Crohn's disease: 18 years or older.
<b>Prescriber Restrictions</b>	Psoriatic arthritis: rheumatologist, dermatologist. Plaque psoriasis: dermatologist. Crohn's disease: gastroenterologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	STIVARGA
<b>Drug Names</b>	STIVARGA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	STRENSIQ
<b>Drug Names</b>	STRENSIQ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have an onset of symptoms prior to age 18. Reauth: must have chart documentation of improvement in the member's condition, including skeletal manifestations.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with an endocrinologist, geneticist, or metabolic disorders specialist
<b>Coverage Duration</b>	Initial: 180 days. Reauth: 365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SUCRAID
<b>Drug Names</b>	SUCRAID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For congenital sucrase-isomaltase deficiency: must have low sucrase activity on duodenal biopsy with other disaccharidases normal on same duodenal biopsy OR must have stool pH less than 6, increase in breath hydrogen of greater than 10ppm when challenged with sucrose after fasting, and negative lactose breath test. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 5 months or older
<b>Prescriber Restrictions</b>	Gastroenterologist, endocrinologist, or metabolic specialist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SUTENT
<b>Drug Names</b>	SUNITINIB MALATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SYLVANT
<b>Drug Names</b>	SYLVANT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. HIV or HHV 8 positive
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation demonstrating a history of (1)lymphadenopathy in greater than one lymph node site and (2)constitutional symptoms such as fever, night sweats, significant weight loss, fatigue, weakness, anorexia, anemia.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	SYMPAZAN
<b>Drug Names</b>	SYMPAZAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of Lennox-Gastaut syndrome. Must have had an inadequate response or intolerance to generic clobazam tablets and 1 other generic antiepileptic drug (e.g. lamotrigine, topiramate, felbamate) and be using as adjunctive therapy to other anti-epileptic drugs.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	SYNAGIS
<b>Drug Names</b>	SYNAGIS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. If under age 12 mo at start of RSV season w/ no other medical dx: must have gestational age (GA) less than 29 wks. If under age 24 mo at start of RSV season during 1st year of life w/ Chronic Lung Disease (CLD) of prematurity: must have GA less than 32 weeks 0 days AND required greater than 21% oxygen (O2) for at least first 28 days of life. If under age 24 mo at start of RSV season during 2nd year of life w/ CLD of prematurity: must have GA less than 32 weeks 0 days AND required greater than 21% O2 for at least first 28 days of life AND have continued to require medical support (chronic corticosteroid therapy, diuretic therapy, supplemental O2) during 6 months before start of 2nd RSV season. If under age 12 mo at start of RSV season w/ heart disease: must have hemodynamically significant Congenital Heart Disease (CHD) (and be on drugs to control heart failure) OR have acyanotic heart disease (and be on drugs to control heart failure and require cardiac surgery) OR have mod-sev pulm HTN OR have cardiac lesions adequately corrected by surgery (and still continue to be on drugs for heart failure). If under age 12 mo at start of RSV season w/ neuromuscular disease or congenital anomaly: must demonstrate that disease/anomaly impairs ability to clear secretions from upper airway b/c of ineffective cough. If under age 24 mo at start of RSV season and profoundly immunocompromised: must have doc of reason (e.g. severe combined immunodeficiency, severe T-cell deficiency, severe AIDS, AML, acute lymphoblastic leukemia, receiving chemotx, received hematopoietic SCT). If under age 24 mo w/ cystic fibrosis (CF): during 1st year of life must have clinical evidence of CLD and/or nutritional compromise OR during 2nd year of life must have manifestation of severe lung disease (prior hospitalization for pulmonary exacerbation in 1st year of life, abnormalities on chest radiography/CT that persist when stable, weight for length is less than 10th %).
<b>Age Restrictions</b>	Less than 12 months or less than 24 months of age at start of RSV season depending on criteria
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Maximum of 5 doses per RSV season.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	TABLOID
<b>Drug Names</b>	TABLOID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TABRECTA
<b>Drug Names</b>	TABRECTA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming MET exon 14 mutation.
<b>Age Restrictions</b>	Age 18 and older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TADALAFIL
<b>Drug Names</b>	ALYQ, TADALAFIL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Current use of nitrate product
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have WHO Functional Class II-IV symptoms. Must have inadequate response or intolerance to sildenafil (Revatio). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	TAFINLAR
<b>Drug Names</b>	TAFINLAR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For unresectable, metastatic, or completely resected melanoma: must have chart documentation of lab result confirming BRAFV600E or BRAFV600K mutation. For metastatic non-small cell lung cancer or locally advanced or metastatic anaplastic thyroid cancer: must have chart documentation of lab result confirming BRAFV600E mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TAGRISSE
<b>Drug Names</b>	TAGRISSE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming epidermal growth factor receptor (EGFR) T790M mutation, exon 19 deletions, or exon 21 L858R mutations.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	TAKHZYRO
<b>Drug Names</b>	TAKHZYRO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of HAE and confirmatory laboratory values on 2 separate instances (copy of laboratory reports required, must include reference ranges). For Type I: low C4 complement level in mg/dL, normal C1q complement component level in mg/dL (C1q complement component level not required for patients under age of 18 or patients whose symptoms began before age 18), and either low C1 esterase inhibitor antigenic level in mg/dL or low C1 esterase inhibitor functional level expressed as a percent. For Type II: low C4 complement level in mg/dL, normal C1q complement component level in mg/dL (C1q complement component level not required for patients under age of 18 or patients whose symptoms began before age 18), and low C1 esterase inhibitor functional level expressed as a percent. For Type III: chart documentation of exclusion of other possible diagnoses and/or causes of angioedema. For reauth, must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 12 years or older
<b>Prescriber Restrictions</b>	By or under the direction of a HAE specialist (defined as an allergist/immunologist who attests to clinical experience in HAE).
<b>Coverage Duration</b>	Initial: 120 days. Reauth: 365 days.
<b>Other Criteria</b>	Must be used as prophylactic therapy for prevention of HAE attacks
<b>Prior Authorization Group</b>	TALTZ
<b>Drug Names</b>	TALTZ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Negative TB skin test. For plaque psoriasis (PS): must have chronic moderate to severe dx, must have minimum BSA of at least 5% (not required if on palms, soles, head/neck, genitalia), and must have had an inadequate response, intolerance, or contraindication to Enbrel or Humira. For psoriatic arthritis: must have had an inadequate response, intolerance, or contraindication to Enbrel, Humira or Xeljanz/Xeljanz XR. For ankylosing spondylitis: must have had an inadequate response, intolerance, or contraindication to Enbrel or Humira. For active non-radiographic axial spondyloarthritis: no drug trials required. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Plaque psoriasis: 6 years or older. Psoriatic arthritis, ankylosing spondylitis, or active non-radiographic axial spondyloarthritis: 18 years or older.
<b>Prescriber Restrictions</b>	Psoriatic arthritis: rheumatologist, dermatologist. Plaque psoriasis: dermatologist. Ankylosing spondylitis: rheumatologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	TALZENNA
<b>Drug Names</b>	TALZENNA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TARGRETIN
<b>Drug Names</b>	BEXAROTENE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, dermatologist, or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TASIGNA
<b>Drug Names</b>	TASIGNA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TAZVERIK
<b>Drug Names</b>	TAZVERIK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 16 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	TEGSEDI
<b>Drug Names</b>	TEGSEDI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Documentation of platelet count greater than 100x10 <sup>9</sup> /L. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist or a provider who specializes in the treatment of inherited disorders.
<b>Coverage Duration</b>	Initial: 6 months. Reauth: 1 year.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TEPMETKO
<b>Drug Names</b>	TEPMETKO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Must be prescribed by or in consultation with an oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TERIPARATIDE
<b>Drug Names</b>	TERIPARATIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure of bisphosphonate therapy or raloxifene unless intolerant or contraindicated. Must have bone mineral density T-score of less than or equal to -2.5 at conventional skeletal sites including the total hip, femoral neck, lumbar spine (post-anterior, not lateral) or radius OR must have history of fracture due to osteoporosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	TETRABENAZINE
<b>Drug Names</b>	TETRABENAZINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Actively suicidal. Uncontrolled depression. Currently using a monoamine oxidase inhibitor or reserpine. Hepatic impairment.
<b>Required Medical Information</b>	Diagnosis. Must have confirmed Huntington's disease either by Huntington Disease Mutation analysis (with laboratory result indicating expanded CAG repeat of greater than or equal to 36 in the huntington gene) or a positive family history of Huntington's Disease with autosomal dominant inheritance pattern. Must have clinical signs of Huntington's Disease to include chart documentation of a clinical work-up showing one or more of the following signs: motor (e.g. finger tapping, rigidity), oculomotor, bulbar (e.g. dysarthria, dysphagia), affective (e.g. depression), cognitive. Must have chart documentation of chorea associated with Huntington's Disease. For doses greater than 50mg/day: must have chart documentation of a trial of 50mg/day dose with inadequate response OR must be CYP2D6 intermediate or extensive metabolizer (as documented through CYP2D6 genotyping results), must provide documentation of slow dose titration with close monitoring of side effects. For reauth: must have documentation from the prescriber indicating improvement in condition and showing monitoring for depression and suicidal ideation. For reauth for doses greater than 50mg/day: must have chart documentation from prescriber showing inadequate efficacy of lower doses and slow titration of dose with close monitoring of side effects.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Maximum dose approved is 100mg/day.
<b>Prior Authorization Group</b>	THALOMID
<b>Drug Names</b>	THALOMID
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist, hematologist, or infectious disease specialist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	THIOLA
<b>Drug Names</b>	TIOPRONIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have severe homozygous cystinuria with urinary cystine level greater than 500mg/day. Must have chart documentation of how diagnosis was confirmed. Must have baseline (within 6 months) urinalysis, complete blood cell count, platelet count, hemoglobin, and liver function tests. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a urologist, nephrologist, or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TIBSOVO
<b>Drug Names</b>	TIBSOVO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	TOLVAPTAN
<b>Drug Names</b>	TOLVAPTAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Anuria, on concomitant therapy with a strong CYP3A inhibitor, underlying liver disease (including cirrhosis), hypovolemic hyponatremia
<b>Required Medical Information</b>	Diagnosis. Must have serum sodium less than 125mEq/L OR symptomatic hyponatremia that resisted correction with 72 hours after consideration of discontinuation of agents known to cause SIADH when clinically feasible (e.g. chlorpropamide, selective serotonin reuptake inhibitors (SSRIs), tricyclic antidepressants, clofibrate, carbamazepine, vincristine, nicotine, narcotics, antipsychotic drugs, ifosfamide, cyclophosphamide, NSAIDs, MDMA, desmopressin, oxytocin, vasopressin). Must have CrCl greater than 10mL/min. Must be initiated and titrated in hospital setting with close serum sodium monitoring. Must be able to sense and appropriately respond to thirst. If SIADH is underlying cause of hyponatremia, must confirm that clinical work-up has been performed to rule out other diagnoses.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Endocrinologist or nephrologist
<b>Coverage Duration</b>	30 days
<b>Other Criteria</b>	Due to risk of liver injury, tolvaptan should not be administered for more than 30 days.
<b>Prior Authorization Group</b>	TOPICAL LIDOCAINE
<b>Drug Names</b>	LIDOCAINE, LIDOCAINE/PRILOCAINE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	If being used as part of a compounded product, all active ingredients in the compounded product must be FDA approved for topical use.

<b>Prior Authorization Group</b>	TOPICAL TACROLIMUS AND PIMECROLIMUS
<b>Drug Names</b>	PIMECROLIMUS, TACROLIMUS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Weakened or compromised immune system
<b>Required Medical Information</b>	Diagnosis. For topical tacrolimus: must have trial and failure of moderate to high potency topical corticosteroid or have a contraindication to this therapy (such as dermatitis on face, genitalia). For topical pimecrolimus: must have trial and failure of moderate to high potency topical corticosteroid or have a contraindication to this therapy (such as dermatitis on face, genitalia) AND must have trial and failure of topical tacrolimus with inadequate response or significant side effect/toxicity or have a contraindication to this therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Pimecrolimus, tacrolimus 0.03%: age 2 years or older. Tacrolimus 0.1%: age 16 years or older.
<b>Prescriber Restrictions</b>	No Prescriber Restrictions.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TRACLEER
<b>Drug Names</b>	BOSENTAN, TRACLEER
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Current use of glyburide or cyclosporine
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have WHO Functional Class II-IV symptoms. For adult patients with WHO Functional Class II and III symptoms: must have previous inadequate response or intolerance to ambrisentan (Letairis). Must have baseline liver function tests (AST, ALT), prior to initiation of therapy. Must have baseline negative pregnancy test prior to initiation of therapy if a female of child-bearing potential. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	TRIENTINE
<b>Drug Names</b>	TRIENTINE HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of how diagnosis was confirmed including at least one of the following: hepatic parenchymal copper content greater than or equal to 250 micrograms per gram dry weight, presence of Kayser-Fleischer Ring in cornea, serum ceruloplasmin level less than 200mg/L, basal 24-hour urinary excretion of copper greater than 100 micrograms (1.6 millimoles), or genetic testing indicating mutation in ATP7B gene. Must have trial of penicillamine (Depen) with an inadequate response or significant side effects/toxicity or must have a contraindication to this therapy. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	By or in consultation with a gastroenterologist, ophthalmologist, or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	Initial: 90 days. Reauth 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TRINTELLIX AND VIIBRYD
<b>Drug Names</b>	TRINTELLIX, VIIBRYD STARTER PACK, VILAZODONE HYDROCHLORIDE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have trial and failure or intolerance to 2 generic antidepressants from SSRI and SNRI classes.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TRUSELTIQ
<b>Drug Names</b>	TRUSELTIQ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	TUKYSA
<b>Drug Names</b>	TUKYSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TURALIO
<b>Drug Names</b>	TURALIO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	TYKERB
<b>Drug Names</b>	LAPATINIB DITOSYLATE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. ECOG Performance Status.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	UPTRAVI
<b>Drug Names</b>	UPTRAVI, UPTRAVI TITRATION PACK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist or pulmonologist. Combination therapy with two or more PAH agents must be prescribed by or in consultation with a pulmonary hypertension specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VALCHLOR
<b>Drug Names</b>	VALCHLOR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	VECAMYL
<b>Drug Names</b>	VECAMYL
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Mild, moderate, and/or labile hypertension. Coronary insufficiency or recent myocardial infarction. Renal insufficiency manifested by rising or elevated BUN level. Uremia. Concurrent use of antibiotics and sulfonamides. Glaucoma. Organic pyloric stenosis. Hypersensitivity to mecamlamine.
<b>Required Medical Information</b>	Diagnosis of moderately severe to severe essential hypertension or uncomplicated malignant hypertension. Must have documented trials of 2 formulary antihypertensive medications from different classes (such as an angiotensin receptor blocker [i.e. irbesartan] and a thiazide diuretic [i.e. hydrochlorothiazide]) with therapeutic failure. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Cardiologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VELTASSA
<b>Drug Names</b>	VELTASSA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have documentation of elevated serum potassium. Must have tried modification of medication regimen to reduce the risk of hyperkalemia if clinically appropriate. Must have trial of, contraindication to, or intolerance to sodium polystyrene sulfonate. For reauth: must have documentation of persistent hyperkalemia and prior reduction in serum potassium levels with Veltassa (patiomer).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	VEMLIDY
<b>Drug Names</b>	VEMLIDY
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Hepatitis B Virus Drug Resistance panel showing resistance to prior tx w/ tenofovir
<b>Required Medical Information</b>	Diagnosis. Must have documentation of results of Hep B Virus Drug Resistance panel if previously received antiviral tx regimen for Hep B. Must have documentation of baseline eval and results for following tests: Hep B viral (HBV) DNA load, hepatitis B e antigen (HBeAg), antibody to hepatitis B e antigen (anti-HBe), hepatitis B surface antigen (HBsAg), antibody to hepatitis surface antigen (anti-HBs), liver biopsy (if available), alanine aminotransferase (ALT) level and assay reference range. For reauth: must have doc from prescriber indicating continued benefit from tx, doc of recent HBV DNA level, chart doc of HBV Drug Resistance panel if mbr has evidence or virologic breakthrough (greater than 10-fold increase in serum HBV DNA from nadir during tx in mbr who had initial virologic response), and doc of HBeAg/Anti-HBe/HBsAg/Anti-HBs (for mbrs with HBeAg positive and for mbrs with HBeAg negative not falling under any other indications).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Infectious disease physician, gastroenterologist, hepatologist, or transplant physician
<b>Coverage Duration</b>	Pregnant mbr: 6 months. All others: 365 days until disease progression or clearance.
<b>Other Criteria</b>	Regimens/requirements based upon AASLD Practice Guidelines for Chronic Hepatitis B. For HBeAg+ chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN OR evidence of moderate/severe inflammation or signif. fibrosis on biopsy) and have HBV DNA level greater than 20,000 IU/mL. For HBeAg- chronic HBV: must meet 1 ALT criterion (ALT greater than or equal to 2xULN, ALT greater than 1xULN w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, ALT less than or equal to ULN w/ ALT increased over time) and 1 HBV DNA criterion (HBV DNA greater than 20,000 IU/mL, HBV DNA greater than 2,000 IU/mL w/ evidence of moderate/severe inflammation or signif. fibrosis on biopsy, HBV DNA less than or equal to 2,000 IU/mL w/ HBV DNA increased over time). For cirrhosis w/ HBV: must have HBV DNA greater than 2,000 IU/mL OR detectable HBV DNA level w/ elevated ALT. For HBV mbr who had liver txfr for HBV or who received solid organ txfr from HBV+ donor: approve regardless of HBV DNA and ALT levels. For HBV carrier who needs immunosuppressive or cytotoxic tx: must be HBsAg+, have planned course of cancer chemotx or immunosuppressive tx. For HBV in mbr currently pregnant to reduce risk of vertical HBV transmission: must be in 3rd trimester of pregnancy and have serum HBV DNA level greater than 200,000 IU/mL or 1,000,000 copies/mL. Reauth for HBeAg+: approve x1 year until all of following are met (loss of HBeAg, undetectable serum HBV DNA, completed 6-12 months of additional tx after appearance of anti-HBe). Reauth for HBeAg-: approve x1 yr until loss of HBsAg. Reauth for cirrhosis, for liver txfr for HBV, or for solid organ txfr from HBV+ donor: long-term tx approvable. Reauth for pregnancy: no reauth provided for same pregnancy. Reauth for HBV carriers receiving immunosuppressive or cytotoxic tx: mbr w/ baseline HBV DNA less than 2,000 IU/mL

should continue x6 months after completion of chemotx or immunosuppressive tx, mbr w/ baseline HBV DNA greater than 2,000 IU/mL should continue until reach therapeutic endpoints for immunocompetent HBV as listed above.

<b>Prior Authorization Group</b>	VENCLEXTA
<b>Drug Names</b>	VENCLEXTA, VENCLEXTA STARTING PACK
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Members who are on concomitant strong CYP3A4 inhibitors during initiation of therapy.
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VENTAVIS
<b>Drug Names</b>	VENTAVIS
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of PAH (WHO Group I) confirmed diagnosis by right heart catheterization. Must have chart documentation of right heart catheterization that indicates the following hemodynamic values: mean pulmonary arterial pressure greater than or equal to 25 mmHg, pulmonary capillary wedge pressure OR left atrial pressure OR left ventricular end-diastolic pressure less than or equal to 15 mmHg, pulmonary vascular resistance greater than 3 Wood units. Must have WHO Functional Class III-IV symptoms. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Pulmonary hypertension specialist, cardiologist, or pulmonologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	B vs. D determination will be made prior to clinical criteria being applied.

<b>Prior Authorization Group</b>	VERSACLOZ
<b>Drug Names</b>	VERSACLOZ
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure of both clozapine tablet AND clozapine orally-disintegrating tablet or have chart documentation of the clinical rationale for why the tablet and orally-disintegrating tablet versions cannot be used.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No prescriber restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VERZENIO
<b>Drug Names</b>	VERZENIO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No exclusion criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VIGABATRIN
<b>Drug Names</b>	VIGABATRIN, VIGADRUNE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must undergo vision testing prior to beginning treatment. For Refractory Complex Partial Seizures: must have inadequate response to 2 anticonvulsant medications (at least 1 medication must be phenytoin or carbamazepine unless intolerant or contraindicated). Must be using in combination with at least 1 other anticonvulsant medication.
<b>Age Restrictions</b>	Seizure: age 2 years or older. Infantile spasms: age 1 month to 2 years.
<b>Prescriber Restrictions</b>	Neurologist or pediatric neurologist.
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	For infantile spasms will not be extended beyond the age of 2 years.

<b>Prior Authorization Group</b>	VITRAKVI
<b>Drug Names</b>	VITRAKVI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Documentation of lab result confirming neurotrophic receptor tyrosine kinase [NTRK] gene fusion.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VIZIMPRO
<b>Drug Names</b>	VIZIMPRO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VONJO
<b>Drug Names</b>	VONJO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of intermediate or high-risk myelofibrosis (includes primary myelofibrosis, post-polycythemia vera myelofibrosis, and post-essential thrombocythemia myelofibrosis). Documentation of baseline platelet count below $50 \times 10^9/L$ prior to initiation of Vonjo.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Hematologist or oncologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	VORICONAZOLE INJECTION
<b>Drug Names</b>	VORICONAZOLE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	VOSEVI
<b>Drug Names</b>	VOSEVI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Decompensated cirrhosis/moderate or severe hepatic impairment (Child Turcotte Pugh class B or C)
<b>Required Medical Information</b>	Diagnosis of chronic Hep C. Doc of prior treatment (tx) for Hep C. Chart doc of lab genotype (GT) result, detectable baseline HCV RNA level (incl. assay date, ref. range), test indicating presence or absence of cirrhosis (e.g. F4 score on liver biopsy from within past 3 years, MRI, ultrasound, CT scan).
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Infectious disease physician, gastroenterologist, hepatologist, HIV specialist, or transplant physician
<b>Coverage Duration</b>	12 weeks
<b>Other Criteria</b>	Regimens/requirements based on AASLD/IDSA Hep C Tx Guidelines found at <a href="https://www.hcvguidelines.org/">https://www.hcvguidelines.org/</a> .
<b>Prior Authorization Group</b>	VOTRIENT
<b>Drug Names</b>	VOTRIENT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	VRAYLAR
<b>Drug Names</b>	VRAYLAR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have a trial and failure or an inadequate response or intolerance to 2 generic antipsychotics (e.g., haloperidol, fluphenazine, chlorpromazine, perphenazine, aripiprazole, olanzapine, quetiapine, risperidone, ziprasidone).
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	WELIREG
<b>Drug Names</b>	WELIREG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XALKORI
<b>Drug Names</b>	XALKORI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming ALK or ROS1 mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	XCOPRI
<b>Drug Names</b>	XCOPRI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Familial short QT syndrome.
<b>Required Medical Information</b>	Diagnosis. Must have had an inadequate response or intolerance to 2 formulary generic antiepileptic drugs (e.g. lamotrigine, topiramate, felbamate)
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a neurologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XELJANZ
<b>Drug Names</b>	XELJANZ, XELJANZ XR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Evidence of infection. Use of biologic disease-modifying antirheumatic drug or potent immunosuppressive agent (e.g. azathioprine, cyclosporine) in combination with tofacitinib. Severe hepatic impairment.
<b>Required Medical Information</b>	Diagnosis. Must have negative tuberculosis skin test. For moderately to severely active RA: Must have trial and failure of methotrexate with inadequate response (if significant side effects, toxicity, or contraindication to methotrexate, must have trial of hydroxychloroquine, leflunomide, or sulfasalazine). Must have lymphocyte count greater than or equal to 500 cells per cubic mm, ANC greater than or equal to 1000 cells per cubic mm, and Hgb level greater than or equal to 9g/dL. For psoriatic arthritis (peripheral disease): must have moderately to severely active psoriatic arthritis AND must have trial of 1 NSAID at target anti-inflammatory dose and of 1 conventional systemic therapy (e.g. methotrexate, cyclosporine, leflunomide, sulfasalazine) with inadequate responses or significant side effects/toxicities or have contraindication to these therapies. For psoriatic arthritis (axial, skin, nail, enthesitis, or dactylitis dominant): must have a trial of 2 NSAIDs at target anti-inflammatory dose with inadequate response or sig. side effects/toxicities unless contraindicated. For reauth: must have documentation from the prescriber indicating stabilization or improvement in condition AND recent lymphocyte count, ANC, Hgb, lipid levels, liver function tests. Lymphocyte count, ANC, Hgb, lipid levels, liver function tests must be completed within 3 months of therapy initiation and at regular intervals thereafter.
<b>Age Restrictions</b>	Age 2 years or older
<b>Prescriber Restrictions</b>	RA: Rheumatologist, Psoriatic Arthritis: Rheumatologist or dermatologist, Ulcerative colitis: gastroenterologist.
<b>Coverage Duration</b>	Initial: 120 days. Reauth: 365 days.
<b>Other Criteria</b>	Not applicable

<b>Prior Authorization Group</b>	XEOMIN
<b>Drug Names</b>	XEOMIN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For blepharospasm: must have previous treatment with onabotulinumtoxinA. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XERMELO
<b>Drug Names</b>	XERMELO
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For diarrhea: must be associated with carcinoid syndrome. Must have had a previous trial of a somatostatin analog (e.g., octreotide) with failure or inadequate control of symptoms. Must be used in combination with a somatostatin analog. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Oncologist, hematologist, endocrinologist, gastroenterologist or palliative care specialist.
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	XGEVA
<b>Drug Names</b>	XGEVA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Concurrent treatment with denosumab (Prolia).
<b>Required Medical Information</b>	Diagnosis of prevention of skeletal-related events in patients with multiple myeloma or bone metastases from solid tumors. For giant cell tumor, must have disease that is unresectable or where surgical resection is likely to result in severe morbidity and member must be skeletally mature if less than 18 years of age. For hypercalcemia of malignancy: must have a trial and failure of IV bisphosphonate therapy (i.e. zoledronic acid 4mg/5mL or 4mg/100mL), with failure defined as an albumin-corrected calcium greater than 12.5mg/dL (3.1 mmol/L) despite recent treatment with an IV bisphosphonate.
<b>Age Restrictions</b>	Prevention of skeletal events, hypercalcemia of malignancy: age 18 years or older. Giant Cell Tumor: age 13 years or older.
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 Days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XIFAXAN
<b>Drug Names</b>	XIFAXAN
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For hepatic encephalopathy: must have trial and failure of lactulose. For diarrhea-predominant irritable bowel syndrome (IBS-D): must have chart documentation of how the diagnosis was confirmed and a trial and failure of loperamide AND an antispasmodic (e.g. dicyclomine) with an inadequate response or significant side effect/toxicity or have a contraindication to these therapies. Reauth for hepatic encephalopathy: must have chart documentation from prescriber indicating improvement in condition. Reauth for IBS-D: must have documentation from prescriber indicating recurrence of IBS-D symptoms after a successful treatment with rifaximin.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Gastroenterologist, hepatologist, or infectious disease specialist
<b>Coverage Duration</b>	Hepatic encephalopathy: 365 Days. IBS-D initial: 14 days. IBS-D reauth: 14 days.
<b>Other Criteria</b>	Criteria only applies to rifaximin 550mg. Criteria does not apply to rifaximin 200mg. For IBS-D: patients who experience a recurrence of symptoms can be retreated up to two times with the same dosage regimen.

<b>Prior Authorization Group</b>	XOLAIR
<b>Drug Names</b>	XOLAIR
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For moderate to severe allergic asthma: must submit patients weight, must have IgE level greater than or equal to 30 IU/mL AND positive skin or RAST test to perennial aeroallergen. Must have trial of combination therapy with an ICS/LABA (inhaled corticosteroid/long-acting beta-agonist, such as Advair, Breo Ellipta, or Symbicort) AND either a LAMA (long-acting muscarinic antagonist, such as Spiriva) or a leukotriene receptor antagonist (such as montelukast) with inadequate response or significant side effects/toxicities or have a contraindication to these therapies within the last year. Must have asthma symptoms that continue to be uncontrolled (uncontrolled defined as hospitalization for asthma within past year, requirement for oral or parenteral corticosteroids to control exacerbations of asthma on 2 occurrences in the past year, or need for daily corticosteroid with inability to taper off). For chronic idiopathic urticaria: must have chart documentation showing 3-month history of urticaria w/ presence of hives, must have trial of one 2nd generation H1 antihistamine (e.g. levocetirizine) and one leukotriene antagonist (e.g. montelukast) with inadequate responses or significant side effects/toxicity unless contraindicated. For nasal polyps: must have trial of at least one nasal corticosteroid (e.g. fluticasone) and must be used as add-on maintenance treatment. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Persistent asthma: 6 years of age or older. Idiopathic urticaria: 12 years of age or older. Nasal polyps: 18 years of age or older.
<b>Prescriber Restrictions</b>	Urticaria: allergist, dermatologist, immunologist. Asthma, Nasal Polyps: no prescriber restrictions.
<b>Coverage Duration</b>	Urticaria, Nasal Polyps: 90 days initial, 365 days reauth. Asthma: 365 days.
<b>Other Criteria</b>	For asthma: must follow recommended dosing guidelines based upon weight and IgE level. For urticaria: dosages above 300mg every 4 weeks is not covered. For nasal polyps: dosages above 600mg every 2 weeks are not covered.
<b>Prior Authorization Group</b>	XOSPATA
<b>Drug Names</b>	XOSPATA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	XPOVIO
<b>Drug Names</b>	XPOVIO, XPOVIO 100 MG ONCE WEEKLY, XPOVIO 40 MG ONCE WEEKLY, XPOVIO 40 MG TWICE WEEKLY, XPOVIO 60 MG ONCE WEEKLY, XPOVIO 60 MG TWICE WEEKLY, XPOVIO 80 MG ONCE WEEKLY, XPOVIO 80 MG TWICE WEEKLY
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XTANDI
<b>Drug Names</b>	XTANDI
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	XYREM
<b>Drug Names</b>	SODIUM OXYBATE, XYREM
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For cataplexy associated with narcolepsy: must have chart documentation and sleep study to confirm diagnosis. For excessive daytime sleepiness associated with narcolepsy: must have polysomnographic evaluation and chart documentation supporting clinical history of narcolepsy AND must have a trial and failure of 2 central nervous stimulants (e.g. modafinil, armodafinil, amphetamine salts, dextroamphetamine, methylphenidate). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 7 years or older
<b>Prescriber Restrictions</b>	Board-certified sleep specialist, pulmonologist, or neurologist
<b>Coverage Duration</b>	Initial: 90 days. Reauth: 365 days.
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ZAVESCA
<b>Drug Names</b>	MIGLUSTAT
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis of mild to moderate Type I Gaucher disease with any of the following: hepatomegaly (defined as liver size greater than or equal to 1.25 times normal), splenomegaly (defined as spleen size greater than 0.2% of body weight), or bone disease (defined as having one of the following: avascular necrosis, Erlenmeyer flask deformity, lytic disease, marrow infiltrations, osteopenia, osteosclerosis, pathological fracture, or radiological evidence of joint deterioration), or bone marrow disease (defined as having anemia or thrombocytopenia). Must not have enzyme replacement therapy as therapeutic option (e.g. allergy/hypersensitivity to ERT, poor venous access, difficulties w/ infusion). For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	By or in consultation with a hematologist or physician who specializes in the treatment of inherited metabolic disorders
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ZEJULA
<b>Drug Names</b>	ZEJULA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	ZELBORAF
<b>Drug Names</b>	ZELBORAF
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. For Erdheim-Chester disease, must have chart documentation of lab result confirming BRAF V600 mutation. For unresectable or metastatic melanoma: must have chart documentation of lab result confirming BRAF V600E mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ZOLINZA
<b>Drug Names</b>	ZOLINZA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ZONTIVITY
<b>Drug Names</b>	ZONTIVITY
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Past history of stroke, transient ischemic attack, or intracranial hemorrhage. Current active pathological bleeding.
<b>Required Medical Information</b>	Past history of myocardial infarction within the past 2 weeks to 12 months or current diagnosis of peripheral artery disease. Must be on concomitant therapy with another antiplatelet agent, such as clopidogrel. Must have documentation of clinical rationale for use of vorapaxar and assessment of member's underlying risk of bleeding to show benefits of vorapaxar would outweigh risk of bleeding. For reauth: must have documentation from prescriber indicating improvement in condition.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	Cardiologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable



<b>Prior Authorization Group</b>	ZORBTIVE
<b>Drug Names</b>	ZORBTIVE
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	Active malignancy
<b>Required Medical Information</b>	Diagnosis of Short Bowel Syndrome (defined as documented malabsorption from small intestines marked by diarrhea, malnutrition, and steatorrhea and that results from resection of the small intestine). Must be receiving adequate nutritional support as determined by prescriber. For reauth: must have documentation from prescriber indicating improvement in condition and clinical rationale for continuation of treatment.
<b>Age Restrictions</b>	Age 18 years or older
<b>Prescriber Restrictions</b>	No Prescriber Restrictions
<b>Coverage Duration</b>	4 weeks
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ZYDELIG
<b>Drug Names</b>	ZYDELIG
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable
<b>Prior Authorization Group</b>	ZYKADIA
<b>Drug Names</b>	ZYKADIA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis. Must have chart documentation of lab result confirming ALK mutation.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable

<b>Prior Authorization Group</b>	ZYTIGA
<b>Drug Names</b>	ABIRATERONE ACETATE, ZYTIGA
<b>PA Indication Indicator</b>	All FDA-approved Indications
<b>Off-label Uses</b>	-
<b>Exclusion Criteria</b>	No Exclusion Criteria
<b>Required Medical Information</b>	Diagnosis.
<b>Age Restrictions</b>	No Age Restrictions
<b>Prescriber Restrictions</b>	Oncologist or hematologist
<b>Coverage Duration</b>	365 days
<b>Other Criteria</b>	Not Applicable