

## **General Precaution for Silver Solutions**

- Never Freeze a silver solution (or allow it to freeze). Do not refrigerate.
- **Never drink directly from a silver solution bottle**. This will contaminate and degrade the solution.
- Avoid extended exposure to direct UV rays (i.e. sunlight) Inside ambient light is safe.
- Always drink colloidal silver on an empty stomach. This is highly preferred.
- Always ensure a silver solution is **kept sealed** and not exposed to airborne contaminants.
- Never place a paper towel, cotton ball or anything else on the opening of a solution to apply the silver. This leaches contaminants back into the solution. **Simply pour out the silver solution to dispense it**. We provide dropper and spray bottle product options.
- Never dispense a silver solution into a potentially contaminated cup or container.

## **Ionic Silver** (External Use only. Can be used in a nebulizer)

If possible, inspect the silver solution before administration for any sign of cloudiness. If an ionic solution become contaminated, it will rapidly form a cloudy solution. You can inspect an ionic silver solution by holding the bottle up to a bright light and checking for the appearance of a cloudy solution.

**Used for:** eyes, ears, lungs and topically.

**Topically** – dosage is not a concern for eyes and ears (suggested dosage is 1-2 drops in either eyes or ear to treat infection). This is an excellent product to use for treating wounds such as cuts, scrapes and abrasions. Also great for cleaning.

## **Colloidal Silver** (External and Internal Use)

Always ensure that the original yellow color of a colloidal solution is still present before use. Never consume a color-shifted or cloudy colloidal silver solution as this is a sure sign of contamination and instability of the colloid. When a yellow colloidal silver solution becomes pink, green, blue, dark grey or clear, this is a sure sign of particle growth (i.e. agglormeration) and is not recommended for administration

## **Use Colloidal Silver:** oral, nasal, lungs, rectally, vaginally.

Colloidal silver needs to be applied differently for each condition. It is excellent to support the immune system and can be taken internally up to 14 days, after which it is suggested to take a break of about a week or so.

Disclaimer: The Information here is entirely for educational purposes only. This information is not intended as a prescription, prognosis, or diagnosis of any diseases or illness, and should not be used as a replacement for any medical treatment you may currently be undergoing. Application of the information provided without the supervision of a licensed medical professional is done so at the individual's own risk.