



- (51) **International Patent Classification:**
A01H 1/04 (2006.01) *C12N 15/82* (2006.01)
- (21) **International Application Number:**
PCT/US2016/022621
- (22) **International Filing Date:**
16 March 2016 (16.03.2016)
- (25) **Filing Language:** English
- (26) **Publication Language:** English
- (30) **Priority Data:**
62/135,261 19 March 2015 (19.03.2015) US
- (71) **Applicant:** PIONEER HI-BRED INTERNATIONAL INC [US/US]; 7100 N.W. 62nd Avenue, P.O. Box 1014, Johnston, Iowa 50131 (US).
- (72) **Inventors:** ELSING, Michael T.; 8407 Sharon Circle, Urbandale, Iowa 50322 (US). GORDON-KAMM, William; 3916 67th Street, Urbandale, Iowa 50322 (US). PERUGINI, Leandro Daniel; 4018 158th Street, Urbandale,

Iowa 50323 (US). YE, Huaxun; 6909 Aubrey Court, Johnston, Iowa 50131 (US).

- (74) **Agent:** STOOP, Johan M.; E.I. du Pont de Nemours and Company, Legal Patent Records Center, Chestnut Run Plaza 721/2340, 974 Centre Road, PO Box 2915, Wilmington, Delaware 19805 (US).


- (81) **Designated States** (*unless otherwise indicated, for every kind of national protection available*): AE, AG, AL, AM, AO, AT, AU, AZ, BA, BB, BG, BH, BN, BR, BW, BY, BZ, CA, CH, CL, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IR, IS, JP, KE, KG, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PA, PE, PG, PH, PL, PT, QA, RO, RS, RU, RW, SA, SC, SD, SE, SG, SK, SL, SM, ST, SV, SY, TH, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

- (84) **Designated States** (*unless otherwise indicated, for every kind of regional protection available*): ARIPO (BW, GH, GM, KE, LR, LS, MW, MZ, NA, RW, SD, SL, ST, SZ,

[Continued on next page]

- (54) **Title:** METHODS AND COMPOSITIONS FOR ACCELERATED TRAIT INTROGRESSION

Figure 1



- (57) **Abstract:** Compositions and methods are provided for the use of pollen-inhibitor genes and/or color marker genes in accelerated trait introgression. Compositions and methods are also provided for introducing a pollen-inhibitor gene and/or a color marker gene in close proximity to a trait locus of interest. Breeding methods and methods for selecting plants comprising a trait locus of interest in close proximity to at least one pollen-inhibitor gene and/or color marker gene are also disclosed. The methods and compositions employ at least one pollen-inhibitor gene and/or color marker gene to provide an effective system for accelerated trait introgression in the genome of a plant.



TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, RU, TJ, TM), European (AL, AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HR, HU, IE, IS, IT, LT, LU, LV, MC, MK, MT, NL, NO, PL, PT, RO, RS, SE, SI, SK, SM, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, KM, ML, MR, NE, SN, TD, TG).

Published:

— *with international search report (Art. 21(3))*

— *with sequence listing part of description (Rule 5.2(a))*

METHODS AND COMPOSITIONS FOR ACCELERATED TRAIT INTROGRESSION

This application claims the benefit of U.S. Provisional Application No. 62/135261, filed March 19, 2015, which is incorporated herein in its entirety by reference.

5

FIELD

The disclosure relates to the field of plant molecular biology. In particular, methods and compositions are provided for introducing and using pollen-inhibitor loci and color marker loci in accelerated trait introgression

REFERENCE TO A SEQUENCE LISTING SUBMITTED AS A TEXT FILE VIA EFS-WEB

10

The official copy of the sequence listing is submitted electronically via EFS-Web as an ASCII formatted sequence listing with a file named 20160223_BB2237WOPCT_SeqLst.txt, created February 23, 2016, and having a size of 387 KB and is filed concurrently with the specification. The sequence listing
15 contained in this ASCII formatted document is part of the specification and is herein incorporated by reference in its entirety.

BACKGROUND

Recombinant DNA technology has made it possible to insert foreign DNA sequences into the genome of an organism, as well as altering endogenous genes
20 of an organism, thus, altering the organism's phenotype. The most commonly used plant transformation methods are *Agrobacterium* infection and biolistic particle bombardment in which transgenes integrate into a plant genome in a random fashion and in an unpredictable copy number.

Site-specific integration techniques, which employ site-specific recombination
25 systems, as well as, other types of recombination technologies, have been used to generate targeted insertions of genes of interest in a variety of organism. Other methods for inserting or modifying a DNA sequence involve homologous DNA recombination by introducing a transgenic DNA sequence flanked by sequences homologous to the genomic target. U.S. Patent No. 5,527,695 describes
30 transforming eukaryotic cells with DNA sequences that are targeted to a

predetermined sequence of the eukaryote's DNA. Transformed cells are identified through use of a selectable marker included as a part of the introduced DNA sequences. While such systems have provided useful techniques for targeted insertion of sequences of interest, there remains a need for methods and
5 compositions which improve these systems and allow for improved breeding methods and compositions and methods useful for accelerated trait introgression.

BRIEF SUMMARY

Compositions and methods are provided for the use of pollen-inhibitor genes and/or color maker genes in accelerated trait introgression. Compositions and
10 methods are also provided for introducing a pollen-inhibitor gene and/or a color marker gene in close proximity to a trait locus of interest. Breeding methods and methods for selecting plants comprising a trait locus of interest in close proximity to at least one pollen-inhibitor gene and / or a color marker gene are also disclosed.

In one embodiment of the disclosure, the method comprises a method for
15 introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising: a) providing a first plant having within a genomic window at least one trait gene of interest integrated into a first target site located proximal to a telomere, wherein said genomic window is about 10 cM in length, wherein said first plant does not comprise a pollen-inhibitor gene; b)
20 breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a pollen-inhibitor gene integrated into a second target site located proximal to both the telomere and the trait gene of interest of (a); and, c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor
25 gene are genetically linked

In another embodiment, the method comprises a method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising: (a) providing a first plant having within a genomic window a first pollen-inhibitor gene integrated into a first target site,
30 wherein said genomic window is about 10 cM in length; (b) breeding to said first plant a second plant having a trait gene of interest integrated into a second target site within said genomic window; (c) selecting a progeny plant from step (b)

comprising said first pollen-inhibitor gene and said trait gene of interest in said genomic window; (d) providing a third plant having a second pollen-inhibitor gene integrated into a third target site within said genomic window; (e) breeding to said third plant a fourth plant, wherein said fourth plant comprises a pollen-inhibitor
5 maintainer (PIM) gene; (f) selecting a progeny plant from step (e) comprising said second pollen-inhibitor gene and said pollen-inhibitor maintainer (PIM) gene; and, (g) cross pollinating the progeny plant of (c) with the progeny plant of (f) and selecting for a progeny plant that comprises said first pollen-inhibitor gene, said trait gene of interest, and said second pollen-inhibitor gene, wherein said first pollen-
10 inhibitor gene, said trait gene of interest, and said second pollen-inhibitor gene are genetically linked. Optionally, the PIM gene is not genetically linked or has been segregated away from said first pollen-inhibitor gene said trait gene of interest and said second pollen-inhibitor gene. The target sites as described herein can be selected from the group consisting of a recombinase target site, a transgenic SSI
15 target site, a single-strand-break-inducing-agent target site, and a double-strand-break-inducing-agent target site, or any one combination thereof. The double-strand-break-inducing-agent target site includes a target site any double strand break inducing agent. For example, but not limiting to an agent selected from the group of a Cas9 endonuclease, a zinc-finger nuclease, a Tal Effector nuclease
20 (TALEN), a meganuclease, and an engineered endonuclease.

In one embodiment, the method comprises a method for introducing a pollen-inhibitor gene and a color marker gene in close proximity to a trait locus of interest in the genome of a plant, said method comprising: (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is
25 about 10 cM in length; (b) introducing into said genomic window of the plant of (a) a color marker gene; (c) breeding to the plant of (b) a second plant, wherein said second plant is a haploid inducer line capable of producing haploid embryos; (d) selecting haploid embryos from the plant of (c) and introducing into said haploid embryos, a pollen-inhibitor gene; and, (e) producing a double haploid plant from the
30 haploid embryo of (d).

In one embodiment, the method comprises a method for introducing two color marker genes in close proximity to a trait locus of interest in the genome of a plant,

said method comprising: (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is about 10 cM in length; (b) introducing into said genomic window of the plant of (a) a color marker gene; (c) breeding to the plant of (b) a second plant, wherein said second plant is a haploid inducer line capable of producing haploid embryos; (d) selecting haploid embryos from the plant of (c) and introducing into said haploid embryos, a second color marker gene; and, (e) producing a double haploid plant from the haploid embryo of for introducing. As described herein, the color marker or the pollen-inhibitor gene can be introduced into any target site of a double-strand-break-inducing-agent.

In another embodiment, the method comprises a method for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising: a) providing a first plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said genomic window is about 5 cM in length and located within 0.1 cM to 10 cM of the telomere, and wherein said first plant does not comprise a pollen-inhibitor gene; (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said second plant does not comprise said first transgenic target site; and, (c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor gene are genetically linked in said genomic window.

In another embodiment, the method comprises a method for accelerated trait introgression in the genome of a plant, the method comprising: (a) providing a first progeny plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere and a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said first transgenic target site and said pollen-inhibitor gene are genetically linked in said

genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere; (b) cross pollinating the first plant of (a) with pollen from a second plant; and (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said pollen-inhibitor gene. The second plant can be
5 an elite inbred line.

In another embodiment, the method comprises a method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising: (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first
10 transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said genomic window is about 5 cM in length; (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site, wherein said second transgenic SSI target site comprises at least one trait gene of interest wherein said second plant does not comprise said
15 first transgenic target site; (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said second transgenic target site genetically linked in said genomic window, (d) providing a third plant having within said genomic window at least a third transgenic SSI target site and a pollen-inhibitor maintainer (PIM), wherein said third transgenic SSI target site comprises a second pollen-
20 inhibitor gene, (e) using the third plant of step (d) to pollinate the plant of step (c) and selecting a progeny plant wherein said first transgenic SSI target site, said second transgenic SSI target site, and said third transgenic SSI target site are genetically linked to each other, and optionally, wherein the PIM gene has been segregated away

In one embodiment, the method comprises a method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising: (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first
25 transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said genomic window is about 5 cM in length; (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site, wherein said second transgenic SSI target site comprises
30

- at least one trait gene of interest wherein said second plant does not comprise said first transgenic target site; (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said second transgenic target site genetically linked in said genomic window; (d) providing a third plant having within said genomic window at least a third transgenic SSI target site, wherein said third transgenic SSI target site comprises a second pollen-inhibitor gene; (e) Breeding to said third plant a fourth plant, wherein said fourth plant comprises a pollen-inhibitor maintainer (PIM) gene; (f) selecting a progeny plant from step (e) comprising said third transgenic target site and pollen-inhibitor maintainer (PIM) gene; and, (g) cross pollinating the progeny plant of (c) with the progeny plant of (f) and selecting for a progeny plant that comprises said first transgenic SSI target site, said second transgenic SSI target site and said third transgenic SSI target site, wherein said first transgenic SSI target site, said second transgenic SSI target site and said third transgenic SSI target site, are genetically linked.
- Compositions and methods are provided for the use of pollen-inhibitor genes and/or color maker genes in accelerated trait introgression.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant, the method comprising: (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one pollen-inhibitor gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere; (b) cross pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said trait of interest and said pollen-inhibitor gene; and, (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant, the method comprising: (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one color marker gene located proximal to both the

telomere and the trait of interest, wherein said trait of interest and said color marker gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere; (b) cross pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said trait of interest site and said color marker gene; and, (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest, a first pollen-inhibitor gene, and a second pollen-inhibitor gene wherein said genomic window is about 5 cM in length, and wherein said trait of interest is flanked by said first and second pollen-inhibitor gene; (b) cross-pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest, a pollen-inhibitor gene and a color marker gene, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene; (b) cross-pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

In one embodiment, the method comprises a method accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest and at least a first color marker gene integrated into a first target site, a second color marker gene integrated into a

second target site for, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene; (b) cross-pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

Also provided are nucleic acid constructs, plants, plant cells, explants, seeds and grain having at least one pollen-inhibitor gene and/or color marker linked to a trait locus of interest. Additional embodiments of the methods and compositions of the present disclosure are shown herein.

BRIEF DESCRIPTION OF THE DRAWINGS AND SEQUENCE LISTINGS

The invention can be more fully understood from the following detailed description and the accompanying drawings and Sequence Listing that form a part of this application.

Figure 1 shows a schematic of a crossing scheme used to establish and perform accelerated introgression for a telomeric trait locus.

Figure 2 shows a schematic of introducing flanking pollen-inhibitor loci around an internal chromosome location containing a trait, and using the triple-linked site to rapidly introgress the trait into other inbreds.

Figure 3 shows a schematic of the use of sequential targeting into haploid embryos to rapidly create linkage between a trait locus and two flanking pollen-inhibitor loci.

Figure 4 shows a schematic of the use of a guide polynucleotide/Cas endonuclease system to integrate a pollen-inhibitor cassette.

Figure 5-A shows a schematic of the expression of dCAS9 fused to repressor peptides along with three guide-RNAs complementary to different sequences in the maize GT1 promoter.

Figure 5-B shows a schematic of the use of dCAS9~RP as the pollen-inhibitor for accelerated trait introgression.

Figure 5-C shows a schematic of the expression cassettes comprising the DCAS9~RP-mediated Pollen-inhibitor locus.

Figure 6 shows a schematic of a pre-established Targeted Accelerated Trait Integration locus.

Figure 7 shows a schematic of a SATI configuration using intein splicing, allowing two-step screening (i.e. over two generations) for breakage of linkage on both sides of an internal genomic locus.

Figure 8 shows a schematic of a SATI configuration using a two component expression system, allowing two step screening (i.e. over two generations) for breakage of linkage on both sides of an internal genomic locus.

Figure 9 A shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP68884.

Figure 9 B shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in of PHP69519.

Figure 9 C shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP70154.

Figure 9 D shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP46070.

Figure 9 E shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP53203.

Figure 9 F shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP71464

Figure 9 G shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP66566

Figure 9 H shows a schematic of the components located between the right border (RB) and left border (LB) of the TDNA in PHP01.

Figure 9 I shows a schematic of the components located between the right border (RB) and left border (LB) of PHP02.

Figure 9 J shows a schematic of the components located between the right border (RB) and left border (LB) of PHP03.

Figure 9 K shows a schematic of the components located between the right border (RB) and left border (LB) of PHP04.

Figure 9 L shows a schematic of the components located between the right border (RB) and left border (LB) of PHP05.

Figure 9 M shows a schematic of the components located between the right border (RB) and left border (LB) of PHP06.

Figure 9 N shows a schematic of the components located between the right border (RB) and left border (LB) of PHP07.

5 Figure 9 O shows a schematic of the components located between the right border (RB) and left border (LB) of PHP08.

Figure 9 P shows a schematic of the components located between the right border (RB) and left border (LB) of PHP09.

10 Figure 9 Q shows a schematic of the components located between the right border (RB) and left border (LB) of PHP10

Figure 9 R shows a schematic of the components located between the right border (RB) and left border (LB) of PHP11.

Figure 10 shows alpha amylase (parts per million) activities of wild type Pioneer Inbred Line 1 (Inbred 1) (negative control) and eight transgenic events (1-8) containing the PG47::AA1 expression cassette.

15 Figure 11 shows the crossing scheme and subsequent PCR analysis that was performed to demonstrate efficacy of the pollen-inhibitor in the Pioneer Inbred Line 1 (PHN46). Transgenic plants (ZMAA1 plants) were used as the pollen donor onto wild-type PHN46 ears. In a population of 1638 F1 plants, the DNA concentration after extraction of genomic DNA from leaf tissue was too low for reliable PCR, but in the remaining 1588 progeny, no transmission of AA1 and NPTII was observed, demonstrating good efficacy of the pollen-inhibitor in terms of preventing fertilization.

25 Figure 12 shows a cartoon depicting the results of high-resolution mapping (using 2 restriction enzyme Genotype-by-Sequencing single nucleotide polymorphism (SNP) analysis) of chromosomal segments in an undesirable BC1F1 progeny plant after using the F1 hybrid to pollinate the recurrent parent Pioneer Inbred Line 2. As evidenced by the white chromosomal segments, over 65% of the genome was converted to Inbred Line 2. However, the majority of chromosome 10 remained Pioneer Inbred Line 1, and thus this was an example of a progeny plant that even though the pollen-inhibitor locus had been lost (while the trait locus

remained), only a small segment of chromosome 10 had converted to Inbred 2 and plants of this type were discarded.

Figure 13 shows a cartoon depicting the results of high-resolution mapping (using 2 restriction enzyme Genotype-by-Sequencing SNP analysis) of chromosomal segments in a desirable BC1F1 progeny plant after using the F1 hybrid to pollinate the recurrent parent Pioneer Inbred Line 2. As evidenced by the white chromosomal segments, over 67% of the genome was converted to Inbred Line 2. In this example, the linkage was broken between the pollen-inhibitor locus and the trait locus (leaving the trait locus behind) while the majority of chromosome 10 converted to inbred line 2, with only small residual segments of Inbred 1 at the telomeres. This type of result was found in the majority of the 28 plants that had lost the pollen-inhibitor locus, while maintaining the trait, representing a rapid (one generation), precise and almost complete introgression of the trait into the new inbred.

Figure 14 shows the results of haplotype analysis focused on the telomeric segment of chromosome 10 that contained the trait locus (at position 150.7 cM, see asterisk). Haplotype results for wild-type Inbred 1 (black bars) are shown in lane 2, and for wild-type Inbred 2 in lane 1. For the 28 progeny analyzed in this manner, all the haplotype markers proximal to the marker that sat on the trait locus (or above the position of the asterisk) were observed to be SNPs found on Chr10 of Inbred 2. The marker at the trait locus and the two SNP calls closest to the telomere (below) were observed to be those of Inbred 1, with three exceptions. Progeny plant 1 contained only the SNP at the trait locus, indicating that a crossover had occurred within 0.7 cM on the proximal side of the trait and within 1.5 cM on the distal side (the position of the next marker). For progeny plants 2 and 15, two closely-spaced crossovers had also occurred at < 0.7 cM and between 1.5 - 4 cM on the proximal and distal side of the trait locus.

Figure 15 shows a schematic of introducing an aleurone-specific anthocyanin locus (a color marker) on one side of an internal chromosome location containing a trait with an aleurone-specific anthocyanin locus (color marker) being introduced on the opposite side, and using the triple-linked site to rapidly introgress the trait into

other inbreds. PAT = Trait Locus, RK = Red Kernel Locus (color marker locus) and PI = Pollen-inhibitor locus.

Figure 16 shows a schematic of introducing a flanking aleurone-specific anthocyanin loci (color marker locus) on both sides of an internal chromosome location containing a trait, and using the triple-linked site to rapidly introgress the trait into other inbreds. PAT = Trait Locus, RK = Red Kernel Locus (color marker locus) and PI = Pollen-inhibitor locus.

Figure 17 shows a schematic of introducing an crown-specific anthocyanin locus on one side of an internal chromosome location containing a trait with an base-specific anthocyanin locus being introduced on the opposite side, and using the triple-linked site to rapidly introgress the trait into other inbreds. PAT = Trait Locus, RC= Red crown (Top of kernel), RP = Red pedicel (bottom of kernel).

The sequence descriptions summarize the Sequence Listing attached hereto. The Sequence Listing contains one letter codes for nucleotide sequence characters and the single and three letter codes for amino acids as defined in the IUPAC-IUB standards described in *Nucleic Acids Research* 13:3021-3030 (1985) and in the *Biochemical Journal* 219(2):345-373 (1984). The symbols and format used for nucleotide and amino acid sequence data comply with the rules set forth in 37 C.F.R. § 1.822.

Table 1. Description of Sequences.

SEQ ID NO:	Description
1	FRT1
2	FRT87
3	Zm-AA1 PRT
4	Zm-AA1 DNA
5	NPTII PRT
6	NPTII DNA
7	35S TERM
8	PINII TERM
9	LTP2 PRO
10	TAGBFP PRT
11	TAGBFP DNA
12	GZ-W64 TERM
13	PG47 PRO
14	BT1 Transit peptide PRT

15	BT1 Transit peptide DNA
16	IN2 TERM
17	BA-BARSTAR PRT
18	BA-BARSTAR DNA
19	UBI1ZM PRO
20	UBI1ZM INTRON
21	DS-RED2 PRT
22	DS-RED2 DNA
23	PMI PRT
24	PMI DNA
25	BA-BARNASE-INT PRT
26	BA-BARNASE-INT DNA
27	35S ENH DNA
28	GAT891G3 PRT
29	GAT891G3 DNA
30	UBIQ3 TERM
31	CZ19B1 TERM
32	OS-ACTIN PRO
33	OS-ACTIN INTRON
34	MO-PAT PRT
35	MO-PAT DNA
36	ZS-YELLOW1 N1 PRT
37	ZS-YELLOW1 N1 DNA
38	TETR PRT
39	TETR DNA
40	TOP DNA (TET OPERATOR)
41	LOX-FAS DNA
42	LOX-2272 DNA
43	PHP70154
44	FRT1-87 from PHP69519: FRT1-NPTII 3XENH-UBI:GAT LTP2:DSRED-FRT87
45	UBI PRO-FLPm-PINII DNA
46	FRT1-87 from PHP68884: FRT1-NPTIENH: LTP2:TAGBFP ENH:PG47:AA1
47	T-DNA FROM PHP66566 UBI:BARSTAR LTP2:DSRED UBI:PMI
48	PHP71464: FRT1-NPTII LTP2:YELLOW PG47:BARNASE-FRT87
49	DPZM03g014880 - ZM-GT1 PRO DNA
50	PHP5096
51	PHP01
52	PHP02
53	PHP03
54	PHP04

55	PHP05
56	RB
57	LB
58	LOXP
59	MO-PMI DNA
60	MO-PMI PRT
61	ALL STOPS
62	SB-SAG12 TERM
63	NP-INTE-C DNA
64	NP-INTE-C PRT
65	NP-INTE-N DNA
66	NP-INTE-N PRT
67	KID-N DNA
68	KID-N PRT
69	KID-C DNA
70	KID-C PRT
71	MO-CYAN DNA
72	MO-CYAN PRT
73	LTP2 PRO-AMCYAN-GZW64A TERM
74	UBI1ZM PRO-UBI1ZM INTRON-AMCYAN-PINII TERM
75	RAB17 PRO-MOCRE-PINII TERM
76	PHP06
77	PG47 PRO-ZM-BT1 TP~ZM-AA1-IN2-1 TERM
78	PG47 PRO-BA-BARNASE-IN2-1 TERM
79	CYP105A1 DNA
80	CYP105A1 PRT
81	CODA DNA
82	CODA PRT
83	ZM-U6 POLIII CHR8 PRO-GT1 GUIDE RNA-ZM-U6 POLIII CHR8 TERM
84	MOCAS9 EXON1-STLS1 INTRON-MOCAS9 EXON2 DNA
85	MOCAS9 PRT
86	ZM-U6 PRO
87	DCAS9~RP DNA
88	DCAS9~RP PRT
89	RB-LOXP-OACTIN PRO & INTRON-FRT1-MOPAT-35S TERM-FRT87-LB
90	PG47 PRO-BT1 TP-ZMAA1-IN2-1 TERM
91	LTP2 PRO-AMCYAN1-PINII TERM
92	UBI1ZM PRO & INTRON-NPTII-PINII TERM
93	AMCYAN1 DNA
94	AMCYAN1 PRT
95	OACTIN PRO & INTRON-MOPAT-35S TERM

96	DCAS9~RP-NLS Fusion
97	moLEXA DNA
98	moLEXA PRT
99	LEXA UAS DNA
100	moCBF1A DNA
101	moCBF1A PRT
102	PG47-moLEXA~moCBF1A-IN2-1-PG47 PRO-MOCODA-PINII-UBI1ZM PRO-NPTII-PINII
103	5xUAS-45 35s PRO::ZMAA1-IN2-1 TERM-PG47-MOCODA-PINII-UBI PRO&INTRON-PMI-PINII
104	ST-LS1 INTRON1
105	CYS4 RECOGNITION SITE
106	CYS4 (MO) DNA
107	CYS4 (MO) PRT
108	GUIDE RNA
109	PG46 PRO-CYS4(MO)-IN2-1 TERM
110	GT1 PRO TARGET1
111	GT1 PRO TARGET2
112	GT1 PRO TARGET3
113	GT1 PRO TARGET4
114	GT1 PRO TARGET5
115	ZM-U6 POLIII CHR8 PRO
116	UBI-PMI-PINII UBI-ZS-YELLOW1N1-GZW64A 35S ENH-PG47 PRO-ZMAA1-IN2-1
117	UBI-FLPm-PINII TERM OSACTIN-PINII UBI-ODP2-PINII
118	FLPm DNA
119	FLPm PRT
120	minimal FRT5 mutant recombination site
121	minimal FRT6 mutant recombination site
122	minimal FRT12 mutant recombination site

DETAILED DESCRIPTION

Many modifications and other embodiments of the disclosures set forth herein will come to mind to one skilled in the art to which this disclosure pertains

5 having the benefit of the teachings presented in the foregoing descriptions and the associated drawings. Therefore, it is to be understood that the disclosure is not to be limited to the specific embodiments disclosed and that modifications and other

embodiments are intended to be included within the scope of the appended claims. Although specific terms are employed herein, they are used in a generic and

10 descriptive sense only and not for purposes of limitation.

Compositions and methods are provided herein for use of a pollen-inhibitor gene, a color marker gene or a pollen-inhibitor and/or color marker expression cassette in the genome of a plant.

As used herein, a "Pollen-Inhibitor gene" refers to a gene that when
5 expressed in pollen or during pollen development, encodes a protein that renders the pollen grain incapable of germinating to produce a pollen tube, or produces a compromised pollen tube incapable of reaching the ovule and fertilizing the egg. Pollen-inhibitor genes also include genes that, when suppressed or silenced, render the pollen grain incapable of germinating to produce a pollen tube, or produces a
10 compromised pollen tube incapable of reaching the ovule and fertilizing the egg.

A pollen-inhibitor gene (pollen inhibitor gene) can be expressed by a pollen-inhibitor expression cassette. A pollen-inhibitor expression cassette refers to a recombinant DNA construct comprising a promoter that stimulates transcription in pollen or during pollen development operably linked to a pollen-inhibitor gene and a
15 3' regulatory sequence such as a terminator.

A pollen-inhibitor locus (pollen inhibitor locus) refers to a genomic location defined by a genetic and physical position, where a pollen-inhibitor gene is located, or into which a pollen-inhibitor expression cassette has integrated. The genomic location defined by a genetic and physical position, into which a pollen-inhibitor
20 expression cassette has integrated is referred to as a "pollen-inhibitor locus".

A non-conditional pollen-inhibitor gene includes a gene that when expressed in pollen produces a protein that will inhibit pollen germination or pollen tube elongation. Examples of genes that can be used as pollen inhibition include, but are not limited to, the maize alpha-amylase gene (Albertsen et al., 1999, US5962769),
25 the barnase gene from *Bacillus amyloliquefaciens* (Mariani et al., 1990, Nature 347:737-741) and the KID gene from *Escherichia coli* (de al Cueva-Mendez et al., 2003, EMBO J. 22:246-251). Other pollen-inhibitor genes are well known in the art (see van Melderren & de Bast, 2009, PLOS Genetics 5:1-6 and Yamaguchi et al., 2001, Ann Rev Genetics 45:61-79 and Leplae et al., 2001, Nuc Acids Res 1-13).

30 A conditional pollen-inhibitor gene includes a gene that when expressed to encode a protein in pollen grains is not inhibitory until the protein cognate substrate is supplied to the pollen. At this point, the non-inhibitory substrate is converted to an

inhibitory molecule. Conditional inhibitory genes include, but are not limited to, the codA gene (Danielsen et al., 1993, Mol. Microbiol. 6:1335-1344), the dhIA gene (Naested et al., Plant J. 18:571-576), the tms2 gene (Sundaresan et al., 1995, genes Dev. 9:1797-1810), or the CYP105A gene (O'Keefe et al., Plant Physiol. 5 105:473-482). When expressed in plants, the encoded gene products are themselves neutral. However, when a non-inhibitory substrate is supplied, the encoded protein converts this compound to an inhibitory compound or inhibitory derivative. The coda encoded protein converts 5-fluorocytosine (5-FC) to cytotoxic 5-fluorouracil (5-FU), the dhIA-encoded protein hydrolyzes haloalkanes such as 1,2 10 dichloroethane to the cytotoxic halogenated alcohol, the tms2-encoded protein converts indole-3 acetamide to the auxin indole-3-acetic acid, and the P450-monooxygenase gene CYP105A encodes a protein that converts the non-herbicidal sulfonylurea R7402 into a potent herbicide.

Promoters useful for expressing pollen-inhibitor genes include included 15 promoters that are expressed after tetrad formation within the maturing pollen grain, the mature pollen grain or during pollen germination, for example, the maize Zm13 promoter (Hamilton et al., 1998. Plant Mol Biol. 38:663-669), the tomato LAT52 promoter (Twell et al., 1990. Development 109:705-713), the Brassica Bp19 promoter (Albani et al., 1991. PMB 16:501-513), the tobacco NTP303 promoter 20 (Weterings et al., 1995. Plant J. 8:55-63), the wheat TaPSG719 promoter (Chen et al., 2010. Mol Biol. Rep. 37:737-744), maize SEQ ID NO:1 (Allen & Lonsdale, 1995. US5412085), the maize pollen-specific promoter described in Fearing et al (1997: Mol Breeding 3:169-176), the tobacco NTPp13 promoter (Yang et al., 2010. Genetika 46:458-463), and promoters of pollen-specific genes described in Khurana 25 et al. (2012, Critical Rev in Plant Science 31:359-390).

Inducible expression can be driven by a promoter that is activated by a specific ligand or by an environmental stimulus. Examples include, but are not limited to, the tetracyclin-responsive repressor system (Gatz and Quail, 1988, PNAS 85:1394-1397), the ethametsulfuron-responsive repressor system (DuPont 30 Patent Applications, McBride et al), the safener-inducible In2 promoter from maize (DeVelder et al., 1997, Plant Cell Physiol 38:568-577) the copper-inducible ACE1 system (McKenzie et al., 1998), the ethanol-inducible AlcA system (Cadick et al.,

1988, *Nat. Biotechnol.* 16:177-180); Runzhi et al., 2005, *Plant Sci.* 169:463-469), the glucocorticoid GVG inducible expression system (Aoyama and Chua, 1997, *Plant J.* 11:605-612), estradiol-inducible expression system (Bruce et al., 2000, *Plant Cell* 12:65-79); Zuo et al., 2000, *Plant J.* 24:265-273), and the
5 methoxyfenozide-inducible VGE system (Koo et al., 2004, *Plant J.* 37:439-448; Padidum et al., 2003, *Curr. Opin. Plant Biol.* 6:87-91)

As described herein, color markers can be useful for screening kernels for the presence of a specific locus. Seed color markers represent an alternative method to the pollen-inhibitor screen described herein, to screen for progeny that
10 have broken the linkage between the trait and the seed-color locus.

A color marker gene can be expressed by a color marker expression cassette. A color marker expression cassette refers to a recombinant DNA construct comprising a promoter operably linked to a gene encoding a color maker. Color maker genes include genes whose expression result in anthocyanin accumulation
15 including ZM-R (X15806, see Perrot and Cone, 1989, *Nucl. Acids Res* 17:8003), ZM-C1 (NCBI Locus NM_001158182, see Alexandrov et al., 2009, *Plant Mol. Boil* 69:179-194) monocot orthologs of the maize R and C1, genes and the fusion of these two genes as CRC (The C1 DNA-Binding domain, the R gene and the C1 activation domain, fused together in that order). Color markers that could be useful
20 also include genes that encode fluorescent proteins such as Am-CYAN1, AcGFP1, ZS-GREEN, ZS-YELLOW1, DS-RED2, DS-RED-EXPRESS (Clontech).

A color marker locus refers to a genomic location defined by a genetic and physical position, where a color marker gene is located, or into which color marker expression cassette has integrated. The genomic location defined by a genetic and
25 physical position, into which a color marker expression cassette has integrated is referred to as a "color marker locus".

Promoters to control expression of color markers in the seed include, but are not limited to, outer endosperm promoters or aleurone promoters such as the barley LTP1 promoter (Skriver et al., 1992, *Plant Mol. Biol.* 18:585-589), the barley LTP2
30 promoter (Kalla et al., 1994, *Plant J.* 6:849-860), the barley GAmyb and High-pl Alpha amylase promoters (Gubler et al., 1995, *Plant Cell* 7:1879-1891), the wheat Early Methionine promoter (Furtado and Henry, 2005, *Plant Biotechnol. J.* 3:421-

434), the rice Chi26 and LTP2 promoters (Hwang et al., 2001, Plant Cell Rep. 20:647-654), the maize BETL1 promoter (Hueros et al., 1999, Plant Physiol. 121:1143-1152), the maize cystatin (CC7) promoter (US patent 8481811, published July 9, 2013), the maize LEG1A promoter (US patent application US20110271405 A1, published November 3, 2011), maize End 2 promoter (US patent application US6903205, published March 4, 2003), and monocot orthologs or paralogs of the above promoters. Provided herein are plants, plant parts, plant cells or seeds having in its genome a genomic window. A genomic window refers to a segment of a chromosome in the genome of a plant that is desirable for producing at least one trait locus, or the segment of a chromosome comprising at least one trait locus that was produced by the methods provided herein.

The genomic window can be about 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or more centimorgans (cM) in length. In one embodiment, the genomic window can be about 1-2 cM, about 1-3 cM, about 1-4 cM, about 1-5 cM, about 1-6 cM, about 1-7 cM, about 1-8 cM, about 1-9 cM, about 1-10 cM, about 2-3 cM, about 2-4 cM, about 2-5 cM, about 2-6 cM, about 2-7 cM, about 2-8 cM, about 2-9 cM, about 2-10 cM, about 3-4 cM, about 3-5 cM, about 3-6 cM, about 3-7 cM, about 3-8 cM, about 3-9 cM, about 3-10 cM, about 4-5 cM, about 4-6 cM, about 4-7 cM, about 4-8 cM, about 4-9 cM, about 4-10 cM, about 5-6 cM, about 5-7 cM, about 5-8 cM, about 5-9 cM, about 5-10 cM, about 6-7 cM, about 6-8 cM, about 6-9 cM, about 6-10 cM, about 7-8 cM, about 7-9 cM, about 7-10 cM, about 8-9 cM, or about 8-10 cM in length.

A "centimorgan" (cM) or "map unit" is the distance between two linked genes, markers, target sites, genomic loci of interest, loci, or any pair thereof, wherein 1% of the products of meiosis are recombinant. Thus, a centimorgan is equivalent to a distance equal to a 1% average recombination frequency between the two linked genes, markers, target sites, loci, genomic loci of interest or any pair thereof.

The genomic window can be located proximal to a telomere of a chromosome or in a non-telomeric (internal) region of a chromosome. The location of the genomic window proximal to a telomere can be about 0.1 to 1 cM, 0.1 to 10 cM, 1-10 cM, 5-15 cM or 20-25 cM distant from telomeric end of the chromosome.

The genomic window can comprise various components. Such components can include, for example, but not limited to, recombination target sites, target sites for site-specific integration (such as, but not limited to, transgenic SSI target sites), single-strand break target sites, double-strand break target sites, genomic loci of interest, native genes, mutated genes, edited genes, trait loci of interest, pollen-inhibitor genes, and polynucleotides of interest. The genomic window can comprise at least 1, 2, 3, 4, 5 or more target sites for a recombinase, a single-strand-break-inducing agent (such as but not limited to a nickase, a Cas endonuclease), a double-strand-break-target site (such as but not limited to a Cas endonuclease, a Zinc finger nuclease, a TALEN, a meganuclease and/or an engineered endonuclease) such that each target site has a different genomic insertion site within the genomic window. In addition, the genomic window can comprise at least 1, 2, 3, 4, 5, 6, 7, 8, 9, 10 or more trait loci of interest each having a different genomic insertion site. By a "different genomic insertion site" is meant that each component of the genomic window (such as for example target sites and trait loci of interest) is inserted into the genome at a different location and as such each component can segregate independently from one another. For example, the genomic window can comprise a combination of target sites and/or trait loci of interest such that each target site or trait loci of interest has a different genomic insertion site within the genomic window.

The components of the genomic windows provided herein have different genomic insertion sites and as such can segregate independently from one another. As used herein, "segregate independently", is used to refer to the genetic separation of any two or more genes, transgenes, native genes, mutated genes, target sites, genomic loci of interest, markers and the like from one another during meiosis. Assays to measure whether two genetic elements segregate independently are known in the art. As such, any two or more genes, transgenes, native genes, mutated genes, target sites, genomic loci of interest, markers and the like within a genomic window provided herein, have genomic insertion sites located at an appropriate distance from one another so that they generally segregate independently at a rate of about 10% or less. Thus, the components of the genomic windows provided herein can segregate independently from one another at a rate of

about 10%, 9%, 8%, 7%, 6%, 5%, 4%, 3%, 2%, 1%, 0.9%, 0.8%, 0.7%, 0.6%, 0.5%, 0.4%, 0.3%, 0.2%, 0.1% or 0.05 %. Alternatively, the components of the genomic windows provided herein can segregate independently from one another at a rate of about 10-0.1%, about 10-0.5%, about 10-1%, about 10-5%, about 9-0.1%, about 9-0.5%, about 9-1%, about 9-5%, about 8-0.1%, about 8-0.5%, about 8-1%, about 8-4%, about 7-0.1%, about 7-0.5%, about 7-1%, about 7-4%, about 6-0.1%, about 6-1%, about 6-0.5%, about 6-3%, about 5-0.1%, about 5-1%, about 5-0.5%, about 4-0.1%, about 4-1%, about 4-0.5%, about 3-0.1%, about 3-1%, about 3-0.5%, about 2-0.1%, about 2-0.5%, about 1-0.1%, about 1-0.5%, or less than 0.1%. For example, if the genomic window comprises a target site and a trait locus of interest that are about 5 cM from each other, the target site and the trait locus trait locus of interest would segregate independently at a rate of about 5%.

As used herein, a “genomic locus of interest” (plural “genomic loci of interest”) comprises a collection of specific polymorphisms that are inherited together. The terms “trait locus” and “trait locus of interest” (plural “trait loci of interest”) are used interchangeably herein and refer to a genomic locus of interest that comprises a trait of interest. A given trait locus of interest can include but is not limited to, a modified or edited native gene, a transgene, an altered double-strand-break target site, a native gene, or a transgenic SSI target site.

As used herein, a “trait” refers to the phenotype conferred from a particular gene or grouping of genes. A trait gene of interest includes any one gene or grouping of genes that encodes a trait. Any desired trait (also referred to as trait of interest) can be introduced into the genome at a given trait locus of interest. Such traits include, but are not limited to, traits conferring insect resistance, disease resistance, herbicide tolerance, male sterility, abiotic stress tolerance, altered phosphorus, altered antioxidants, altered fatty acids, altered essential amino acids, altered carbohydrates, or sequences involved in site-specific recombination. In terms of relative position of two loci on a chromosome, a locus is more “proximal” if it is closer to the centromere (and farther from the telomere) of that chromosome, and a locus is more “distal” if it is closer to the telomere (and farther from the centromere).

The trait locus of interest can include, for example, any modification that confers a trait, such as a transgene or a native trait. The trait locus of interest can also include a native trait or a selectable marker. Selectable markers are described in more detail further herein and include DNA segments that encode products which provide resistance against otherwise toxic compounds. As used herein, a “native trait” refers to a trait found in nature. In another embodiment, the trait locus of interest comprises a transgene.

A given trait locus of interest has its own genomic insertion site within the genomic window. For example, a trait locus of interest and a target site (for a recombinase, a single-strand-break-inducing agent, a double-strand-break-target site, or others) within the genomic window will have different genomic insertion sites within the genome. A given target site can be found within about 10 cM, 9 cM, 8 cM, 7 cM, 6 cM, 5 cM, 4 cM, 3 cM, 2 cM, 1 cM, 0.9 cM, 0.8 cM, 0.7 cM, 0.6 cM, 0.5 cM, 0.4 cM, 0.3 cM, 0.2 cM, 0.1 cM or 0.05 cM from the trait locus of interest such that the target site and trait locus of interest have different genomic insertion sites.

Compositions and methods are provided for introducing a pollen-inhibitor gene and / or a color marker gene in close proximity to a trait locus of interest in the genome of a progeny plant. In one embodiment of the disclosure, the method comprises: a) providing a first plant having within a genomic window at least one trait gene of interest integrated into a first target site located proximal to a telomere, wherein said genomic window is about 10 cM in length, wherein said first plant does not comprise a pollen-inhibitor gene; b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a pollen-inhibitor gene integrated into a second target site located proximal to both the telomere and said trait gene of interest; and, c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor gene are genetically linked.

In one embodiment of the disclosure, the method comprises: (a) providing a first plant having within a genomic window a first pollen-inhibitor gene integrated into a first target site, wherein said genomic window is about 10 cM in length; (b) breeding to said first plant a second plant having a trait gene of interest integrated into a second target site within said genomic window; (c) selecting a progeny plant

from step (b) comprising said first pollen-inhibitor gene and said trait gene of interest in said genomic window; (d) providing a third plant having a second pollen-inhibitor gene integrated into a third target site within said genomic window; (e) breeding to said third plant a fourth plant, wherein said fourth plant comprises a pollen-inhibitor
5 maintainer (PIM) gene within said genomic window; (f) selecting a progeny plant from step (e) comprising said second pollen-inhibitor gene and said pollen-inhibitor maintainer (PIM) gene; and, (g) cross pollinating the progeny plant of (c) with the progeny plant of (f) and selecting for a progeny plant that comprises, genetically linked to each other. Optionally, the PIM gene is segregated away from said first
10 pollen-inhibitor gene, said trait gene of interest, and said second pollen-inhibitor gene.

As used herein, by "target site" is intended a polynucleotide comprising a nucleotide sequence comprising at least one recognition sequence for an agent such as, but not limited to, a recombinase, a single-strand-break-inducing agent
15 (such as but not limited to a nickase or a Cas endonuclease) or a double-strand-break-inducing target site (such as but not limited to a Cas endonuclease, a Zinc finger nuclease, a TALEN, a meganuclease, an engineered endonuclease, or any one combination thereof).

By "transgenic target site" is meant a target site that is non-native in
20 sequence and/or in genomic location to the plant genome. In some embodiments, the transgenic target site can comprise at least 1, 2, 3, 4, 5 or more recombination sites for site-specific recombination (also referred to as transgenic SSI target site). Site-specific recombination system employ various components which are described herein and in U.S. Patent Nos. 6187994, 6262341, 6331661 and 6300545, each of
25 which is herein incorporated by reference.

The terms "transgenic SSI target site", "transgenic target site for site specific integration (SSI)", and "transgenic target site for SSI" are used interchangeably herein and refer to a polynucleotide comprising a nucleotide sequence flanked by at least two recombination sites (See, for example, US Patent Application US2013-
30 0198888 A1, published on August-01-2013, and US Patent Application US2014-0338070 A1, published on November 13, 2014, each of which is herein incorporated by reference. In some embodiments, the recombination sites of the transgenic SSI

target site are dissimilar and non-recombinogenic with respect to one another. One or more intervening sequences may be present between the recombination sites of the transgenic SSI target site. Intervening sequences of particular interest would include linkers, adapters, selectable markers, pollen-inhibitor genes, polynucleotides of interest, promoters and/or other sites that aid in vector construction or analysis. In addition, the recombination sites of the transgenic SSI target site can be located in various positions, including, for example, within intronic sequences, coding sequences, or untranslated regions.

The transgenic SSI target site can comprise 1, 2, 3, 4, 5, 6 or more recombination sites. In one embodiment, the target site comprises a first recombination site and a second recombination site wherein the first and the second recombination site are dissimilar and non-recombinogenic to each other. In a further embodiment, the target site comprises a third recombination site between the first recombination site and the second recombination site. In such embodiments, the first, second and third recombination sites may be dissimilar and non-recombinogenic with respect to one another. Such first, second and third recombination sites are able to recombine with their corresponding or identical recombination site when provided with the appropriate recombinase.

Pollen-inhibitor genes, color marker genes, or trait loci employed in the methods and compositions provided herein can be integrated into recombination sites that are "corresponding" sites or "dissimilar" sites. By "corresponding recombination sites" or a "set of corresponding recombination sites" is intended that the recombination sites have the same or corresponding nucleotide sequence. A set of corresponding recombination sites, in the presence of the appropriate recombinase, will efficiently recombine with one another (i.e., the corresponding recombination sites are recombinogenic). The recombination sites can also be dissimilar. By "dissimilar recombination sites" or a "set of dissimilar recombination sites" is intended that the recombination sites are distinct (i.e., have at least one nucleotide difference). The recombination sites within "a set of dissimilar recombination sites" can be either recombinogenic or non-recombinogenic with respect to one other. By "recombinogenic" is intended that the set of recombination sites are capable of recombining with one another. Thus, suitable sets of

"recombinogenic" recombination sites for use in the methods and compositions provided herein include those sites where the relative excision efficiency of recombination between the recombinogenic sites is above the detectable limit under standard conditions in an excision assay, typically, greater than 2%, 5%, 10%, 20%, 50%, 100%, or greater. By "non-recombinogenic" is intended the set of recombination sites, in the presence of the appropriate recombinase, will not recombine with one another or recombination between the sites is minimal. Thus, suitable "non-recombinogenic" recombination sites for use in the methods and compositions provided herein include those sites that recombine (or excise) with one another at a frequency lower than the detectable limit under standard conditions in an excision assay, typically, lower than 2%, 1.5%, 1%, 0.75%, 0.5%, 0.25%, 0.1%, 0.075, 0.005%, 0.001%.

Each recombination site within the "set of non-recombinogenic sites" is biologically active and therefore can recombine with an identical site. Accordingly, it is recognized that any suitable non-recombinogenic recombination sites may be utilized, including a FRT site or an active variant thereof, a LOX site or active variant thereof, any combination thereof, or any other combination of non-recombinogenic recombination sites known in the art. FRT sites that can be employed in the methods and compositions disclosed herein can be found, for example, in US patent No. 8,586,361 issued on November 19, 2013, herein incorporated by reference.

By "recombination site" is intended a recombination site and active variants thereof. Many recombination systems are known in the art and one of skill will recognize the appropriate recombination site to be used with the recombination system of interest. Any suitable recombination site or set of recombination sites may be utilized herein, including a FRT site, a biologically active variant of a FRT site (i.e. a mutant FRT site), a LOX site, a biologically active variant of a LOX site (i.e. a mutant LOX site), any combination thereof, or any other combination of recombination sites known in the art. Examples of FRT sites include, for example, the wild type FRT site (FRT1) (SEQ ID NO: 1), and various mutant FRT sites, including but not limited to, FRT5 (SEQ ID NO: 120), FRT6 (SEQ ID NO: 121), FRT12 (SEQ ID NO: 122) and FRT87 (SEQ ID NO: 2). See, for example, U.S. Patent No. 6,187,994 issued on January 13, 2001, US patent No. 8,586,361 issued

on November 19, 2013, and US patent application US2013-0198888A1, published on August-01, 2013, each of which are herein incorporated by reference.

Recombination sites from the Cre/Lox site-specific recombination system can also be used. Such recombination sites include, for example, wild type LOX sites and mutant LOX sites. An analysis of the recombination activity of mutant LOX sites is presented in Lee et al. (1998) *Gene* 216:55-65, herein incorporated by reference. Also, see for example, Schlake and Bode (1994) *Biochemistry* 33:12746-12751; Huang et al. (1991) *Nucleic Acids Research* 19:443-448; Sadowski (1995) In *Progress in Nucleic Acid Research and Molecular Biology* Vol. 51, pp. 53-91; Cox (1989) In *Mobile DNA*, Berg and Howe (eds) American Society of Microbiology, Washington D.C., pp. 116-670; Dixon et al. (1995) *Mol. Microbiol.* 18:449-458; Umlauf and Cox (1988) *EMBO* 7:1845-1852; Buchholz et al. (1996) *Nucleic Acids Research* 24:3118-3119; Kilby et al. (1993) *Trends Genet.* 9:413-421; Rossant and Geagy (1995) *Nat. Med.* 1: 592-594; Albert et al. (1995) *The Plant J.* 7:649-659; Bayley et al. (1992) *Plant Mol. Biol.* 18:353-361; Odell et al. (1990) *Mol. Gen. Genet.* 223:369-378; Dale and Ow (1991) *Proc. Natl. Acad. Sci. USA* 88:10558-10562; Qui et al. (1994) *Proc. Natl. Acad. Sci. USA* 91:1706-1710; Stuurman et al. (1996) *Plant Mol. Biol.* 32:901-913; Dale et al. (1990) *Gene* 91:79-85; Albert et al. (1995) *The Plant J.* 7:649-659 and WO 01/00158; all of which are herein incorporated by reference.

Active variants and fragments of recombination sites are also encompassed by the compositions and methods provided herein. Fragments of a recombination site retain the biological activity of the recombination site and hence facilitate a recombination event in the presence of the appropriate recombinase. Thus, fragments of a recombination site may range from at least about 5, 10, 15, 20, 25, 30, 35, 40 nucleotides, and up to the full-length of a recombination site. Active variants can comprise at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the native recombination site, wherein the active variants retain biological activity and hence facilitate a recombination event in the presence of the appropriate recombinase. Assays to measure the biological activity of recombination sites are known in the art. See, for example, Senecoll et al. (1988) *J. Mol. Biol.* 201:406-421; Voziyanov et al.

(2002) Nucleic Acid Research 30:7, U.S. Patent No. 6,187,994, WO/01/00158, and Albert et al. (1995) The Plant Journal 7:649-659.

By "recombinase" is intended a polypeptide that catalyzes site-specific recombination between compatible recombination sites. For reviews of site-specific
5 recombinases, see Sauer (1994) Current Opinion in Biotechnology 5:521-527; and Sadowski (1993) FASEB 7:760-767; the contents of which are incorporated herein by reference. The recombinase can be a naturally occurring recombinase or a biologically active fragment or variant of the recombinase. Recombinases include
10 recombinases from the Integrase and Resolvase families, biologically active variants and fragments thereof, and any other naturally occurring or recombinantly produced enzyme or variant thereof that catalyzes conservative site-specific recombination between specified DNA recombination sites.

The Integrase family of recombinases has over one hundred members and includes, for example, FLP, Cre, Int, and R. For other members of the Integrase
15 family, see for example, Esposito et al. (1997) Nucleic Acid Research 25:3605-3614 and Abremski et al. (1992) Protein Engineering 5:87-91, both of which are herein incorporated by reference. Other recombination systems include, for example, the streptomyces bacteriophage phi C31 (Kuhstoss et al. (1991) J. Mol. Biol. 20:897-908); the SSV1 site-specific recombination system from *Sulfolobus shibatae*
20 (Maskhelishvili et al. (1993) Mol. Gen. Genet. 237:334-342); and a retroviral integrase-based integration system (Tanaka et al. (1998) Gene 17:67-76). Some recombinase do not require cofactors or a supercoiled substrate. Such recombinases include Cre recombinase, FLP recombinase, or active variants or fragments thereof (See for example US patent No. 8,586,361, issued on November
25 19, 2013, which is herein incorporated by reference).

The FLP recombinase is a protein that catalyzes a site-specific reaction that is involved in amplifying the copy number of the two-micron plasmid of *S. cerevisiae* during DNA replication. As used herein, FLP recombinase refers to a recombinase that catalyzes site-specific recombination between two FRT sites. The FLP protein
30 has been cloned and expressed. See, for example, Cox (1993) Proc. Natl. Acad. Sci. U.S.A. 80:4223-4227. The FLP recombinase for use in the methods and with the compositions may be derived from the genus *Saccharomyces*. One can also

synthesize a polynucleotide comprising the recombinase using plant-preferred codons for optimal expression in a plant of interest. A recombinant FLP enzyme encoded by a nucleotide sequence comprising maize preferred codons (FLPm) (SEQ ID NO: 119) that catalyzes site-specific recombination events is known. See, 5 for example, U.S. Patent 5,929,301, herein incorporated by reference. Additional functional variants and fragments of FLP are known. See, for example, Buchholz et al. (1998) Nat. Biotechnol. 16:617-618, Hartung et al. (1998) J. Biol. Chem. 273:22884-22891, Saxena et al. (1997) Biochim Biophys Acta 1340(2):187-204, and Hartley et al. (1980) Nature 286:860-864, all of which are herein incorporated by 10 reference.

The bacteriophage recombinase Cre catalyzes site-specific recombination between two lox sites. The Cre recombinase is known in the art. See, for example, Guo et al. (1997) Nature 389:40-46; Abremski et al. (1984) J. Biol. Chem. 259:1509-1514; Chen et al. (1996) Somat. Cell Mol. Genet. 22:477-488; Shaikh et al. (1977) J. 15 Biol. Chem. 272:5695-5702; and, Buchholz et al. (1998) Nat. Biotechnol. 16:617-618, all of which are herein incorporated by reference. The Cre polynucleotide sequences may also be synthesized using plant-preferred codons. Such sequences (moCre) are described in WO 99/25840, herein incorporated by reference. It is further recognized that a chimeric recombinase can be used in the methods. By 20 "chimeric recombinase" is intended a recombinant fusion protein which is capable of catalyzing site-specific recombination between recombination sites that originate from different recombination systems. That is, if a set of functional recombination sites, characterized as being dissimilar with respect to one another, is utilized in the methods and compositions and comprises a FRT site and a LoxP site, a chimeric 25 FLP/Cre recombinase or active variant or fragment thereof will be needed or, alternatively, both recombinases may be separately provided. Methods for the production and use of such chimeric recombinases or active variants or fragments thereof are described in WO 99/25840, herein incorporated by reference.

As used herein, the terms "double-strand-break target site", "DSB target site", 30 "DSB target sequence", "double-strand-break-inducing-agent target site", and "target site for a double-strand-break-inducing-agent" are used interchangeably and refer to a polynucleotide sequence in the genome of a plant cell (including

chloroplastic and mitochondrial DNA) that comprises a recognition sequence for a double-strand-break-inducing agent at which a double-strand-break is induced in the cell genome by a double-strand-break-inducing-agent.

As used herein, the terms “single-strand-break-inducing-agent target site”, “single-strand-break target site”, “SSB target site”, “SSB target sequence”, and “target site for a single-strand-break-inducing-agent” are used interchangeably and refer to a polynucleotide sequence in the genome of a plant cell (including chloroplastic and mitochondrial DNA) that comprises a recognition sequence for an agent (such as but not limited to a nickase, a nuclease) at which a single-strand-break is induced in the cell genome.

As used herein, the terms “altered double-strand-break target site”, “altered DSB target site”, “aDSB target site”, and “altered target site for a double-strand-break-inducing-agent” are used interchangeably and refer to a DSB target sequence comprising at least one alteration when compared to a non-altered DSB target sequence. “Alterations” can include, for example: (i) replacement of at least one nucleotide, (ii) a deletion of at least one nucleotide, (iii) an insertion of at least one nucleotide, or (iv) any combination of (i)-(iii).

The DSB target site can be an endogenous site in the plant genome, or alternatively, the DSB target site can be heterologous to the plant and thereby not be naturally occurring in the genome, or the DSB target site can be found in a heterologous genomic location compared to where it occurs in nature. As used herein, the term “endogenous DSB target site” refers to an DSB target site that is endogenous or native to the genome of a plant and is located at the endogenous or native position of that DSB target site in the genome of the plant.

The length of the SSB or DSB target site can vary, and includes, for example, DSB target sites that are at least 4, 6, 8, 10, 12, 14, 16, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67, 68, 69, 70 or more nucleotides in length. It is further possible that the DSB target site could be palindromic, that is, the sequence on one strand reads the same in the opposite direction on the complementary strand. The nick/cleavage site could be within the recognition sequence or the nick/cleavage site could be outside of the

recognition sequence. In another variation, the cleavage could occur at nucleotide positions immediately opposite each other to produce a blunt end cut or, in other cases, the incisions could be staggered to produce single-stranded overhangs, also called “sticky ends”, which can be either 5' overhangs, or 3' overhangs.

5 Pollen-inhibitor genes, color marker genes, or trait loci employed in the methods and compositions provided herein can be integrated into a double-strand-break target site by a double-strand-break-inducing-agent.

 A “double-strand-break-inducing agent” (also referred to as “DSB-inducing-agent”) refers to any nuclease which produces a double-strand break in the target
10 sequence. The double-strand break target site can be, but is not limited to a zinc finger endonuclease target site, an engineered endonuclease target site, a meganuclease target site, a TALENs target site and a Cas endonuclease target site.

 Any nuclease that induces a single or double-strand break into a desired target site can be used in the methods and compositions disclosed herein. A
15 naturally-occurring or native endonuclease can be employed so long as the endonuclease induces a single or double-strand break in a desired target site. Alternatively, a modified or engineered endonuclease can be employed. An “engineered endonuclease” refers to an endonuclease that is engineered (modified or derived) from its native form to specifically recognize and induce a single or
20 double-strand break in the desired target site. Thus, an engineered endonuclease can be derived from a native, naturally-occurring endonuclease or it could be artificially created or synthesized. The modification of the endonuclease can be as little as one nucleotide. Producing a single or double-strand break in a target site or other DNA can be referred to herein as “cutting” or “cleaving” the DSB target site or
25 other DNA.

 Active variants and fragments of the SSB or DSB target sites can comprise at least 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the given DSB target site, wherein the active variants retain biological activity and hence are capable of being recognized
30 and cleaved by an DSB-inducing-agent. Assays to measure the double-strand break of a DSB target site by an endonuclease are known in the art and generally measure the ability of an endonuclease to cut the DSB target site.

Endonucleases are enzymes that cleave the phosphodiester bond within a polynucleotide chain, and include restriction endonucleases that cleave DNA at specific sites without damaging the bases. Restriction endonucleases include Type I, Type II, Type III, and Type IV endonucleases, which further include subtypes. In the Type I and Type III systems, both the methylase and restriction activities are contained in a single complex. Restriction enzymes are further described and classified, for example in the REBASE database (Roberts *et al.*, (2003) *Nucleic Acids Res* 31:418-20), Roberts *et al.*, (2003) *Nucleic Acids Res* 31:1805-12, and Belfort *et al.*, (2002) in *Mobile DNA II*, pp. 761-783, Eds. Craigie *et al.*, (ASM Press, Washington, DC).

Endonucleases also include meganucleases, also known as homing endonucleases (HEases), which like restriction endonucleases, bind and cut at a specific DSB target site, however the DSB target sites for meganucleases are typically longer, about 18 bp or more. Meganuclease domains, structure and function are known, see for example, Guhan and Muniyappa (2003) *Crit Rev Biochem Mol Biol* 38:199-248; Lucas *et al.*, (2001) *Nucleic Acids Res* 29:960-9; Jurica and Stoddard, (1999) *Cell Mol Life Sci* 55:1304-26; Stoddard, (2006) *Q Rev Biophys* 38:49-95; and Moure *et al.*, (2002) *Nat Struct Biol* 9:764. In some examples a naturally occurring variant, and/or engineered derivative meganuclease is used. Any meganuclease can be used herein, including, but not limited to, I-SceI, I-SceII, I-SceIII, I-SceIV, I-SceV, I-SceVI, I-SceVII, I-CeuI, I-CeuAIP, I-CreI, I-CrepsbIP, I-CrepsbIIP, I-CrepsbIIIP, I-CrepsbIVP, I-TliI, I-PpoI, PI-PspI, F-SceI, F-SceII, F-SuvI, F-TevI, F-TevII, I-Amal, I-Anil, I-Chul, I-Cmoel, I-Cpal, I-Cpall, I-Csml, I-Cvul, I-CvuAIP, I-Ddil, I-Ddill, I-Dirl, I-Dmol, I-Hmul, I-Hmull, I-HsNIP, I-Llal, I-Msol, I-Naal, I-NanI, I-NclIP, I-NgrIP, I-NitI, I-Njal, I-Nsp236IP, I-PakI, I-PbolP, I-PculP, I-PcuAI, I-PcuVI, I-PgrIP, I-PoblP, I-PorI, I-PorIIP, I-PbpIP, I-SpBetaIP, I-Scal, I-SexIP, I-SneIP, I-SpomI, I-SpomCP, I-SpomIP, I-SpomIIP, I-SqulP, I-Ssp6803I, I-SthPhiJP, I-SthPhiST3P, I-SthPhiSTe3bP, I-TdeIP, I-TevI, I-TevII, I-TevIII, I-UarAP, I-UarHGPAIP, I-UarHGPA13P, I-VinIP, I-ZbiIP, PI-MtuI, PI-MtuHIP, PI-MtuHIIP, PI-Pful, PI-Pfull, PI-Pkol, PI-Pkoll, PI-Rma43812IP, PI-SpBetaIP, PI-SceI, PI-Tful, PI-Tfull, PI-Thyl, PI-TliI, PI-TliII, or any active variants or fragments thereof.

TAL effector nucleases (also referred to as TALENs) can be used to make double-strand breaks at specific target sequences in the genome of a plant or other organism. TAL effector nucleases can be created by fusing a native or engineered transcription activator-like (TAL) effector, or functional part thereof, to the catalytic domain of an endonuclease, such as, for example, *FokI*. The unique, modular TAL effector DNA binding domain allows for the design of proteins with potentially any given DNA recognition specificity. Thus, the DNA binding domains of the TAL effector nucleases can be engineered to recognize specific DNA target sites and thus, used to make double-strand breaks at desired target sequences. See, WO 2010/079430; Morbitzer *et al.* (2010) *PNAS* 10.1073/pnas.1013133107; Scholze & Boch (2010) *Virulence* 1:428-432; Christian *et al.* *Genetics* (2010) 186:757-761; Li *et al.* (2010) *Nuc. Acids Res.* (2010) doi:10.1093/nar/gkq704; and Miller *et al.* (2011) *Nature Biotechnology* 29:143–148; all of which are herein incorporated by reference.

CRISPR (clustered regularly interspaced short palindromic repeats) loci refers to certain genetic loci encoding factors of DNA cleavage systems, for example, used by bacterial and archaeal cells to destroy foreign DNA (Horvath and Barrangou, 2010, *Science* 327:167-170). A CRISPR locus can consist of a CRISPR array, comprising short direct repeats separated by short variable DNA sequences (called 'spacers'), which can be flanked by diverse Cas (CRISPR-associated) genes. Multiple CRISPR-Cas systems have been described including Class 1 systems, with multisubunit effector complexes, and Class 2 systems, with single protein effectors (such as but not limiting to Cas9, Cpf1, C2c1, C2c2, C2c3). (Zetsche *et al.*, 2015, *Cell* 163,1-13; Shmakov *et al.*, 2015, *Molecular Cell* 60,1-13; Makarova *et al.* 2015, *Nature Reviews Microbiology* Vol. 13 :1-15). The type II CRISPR/Cas system from bacteria employs a crRNA (CRISPR RNA) and tracrRNA (trans-activating CRISPR RNA) to guide a Cas9 endonuclease to its DNA target. The crRNA contains a region complementary to one strand of the double strand DNA target and a region that base pairs with the tracrRNA (trans-activating CRISPR RNA) forming a RNA duplex that directs the Cas9 endonuclease to cleave the DNA target. CRISPR systems belong to different classes, with different repeat patterns, sets of genes, and species ranges. The number of CRISPR-associated

genes at a given CRISPR locus can vary between species (Haft et al. , 2005, Computational Biology, PLoS Comput Biol 1(6): e60.

doi:10.1371/journal.pcbi.0010060; Makarova et al. 2015, Nature Reviews Microbiology Vol. 13 :1-15).

5 The term “Cas gene” herein refers to a gene that is generally coupled, associated or close to, or in the vicinity of flanking CRISPR loci. The terms “Cas gene”, “CRISPR-associated (Cas) gene” are used interchangeably herein.

 The term “Cas endonuclease” herein refers to a protein encoded by a Cas gene. A Cas endonuclease herein, when in complex with a suitable polynucleotide
10 component, is capable of recognizing, binding to, and optionally nicking or cleaving all or part of a specific DNA target sequence. A Cas endonuclease described herein comprises one or more nuclease domains. Cas endonucleases of the disclosure includes those having a HNH or HNH-like nuclease domain and / or a RuvC or RuvC-like nuclease domain. A Cas endonuclease of the disclosure includes a Cas9
15 protein, a Cpf1 protein, a C2c1 protein, a C2c2 protein, a C2c3 protein, Cas3, Cas 5, Cas7, Cas8, Cas10, or complexes of these.

 “Cas9” (formerly referred to as Cas5, Csn1, or Csx12) herein refers to a Cas endonuclease of a type II CRISPR system that forms a complex with a crNucleotide and a tracrNucleotide, or with a single guide polynucleotide, for specifically
20 recognizing and cleaving all or part of a DNA target sequence. Cas9 protein comprises a RuvC nuclease domain and an HNH (H-N-H) nuclease domain, each of which can cleave a single DNA strand at a target sequence (the concerted action of both domains leads to DNA double-strand cleavage, whereas activity of one domain leads to a nick).

25 As used herein, the term “guide polynucleotide”, relates to a polynucleotide sequence that can form a complex with a Cas endonuclease (such as but not limited to a Cas9 endonuclease) and enables the Cas endonuclease to recognize and optionally cleave a DNA target site (U.S. Provisional Application No. 62/023239, filed July 11, 2014). The guide polynucleotide can be a single molecule or a double
30 molecule. The guide polynucleotide sequence can be a RNA sequence, a DNA sequence, or a combination thereof (a RNA-DNA combination sequence). Optionally, the guide polynucleotide can comprise at least one

nucleotide, phosphodiester bond or linkage modification such as, but not limited, to Locked Nucleic Acid (LNA), 5-methyl dC, 2,6-Diaminopurine, 2'-Fluoro A, 2'-Fluoro U, 2'-O-Methyl RNA, phosphorothioate bond, linkage to a cholesterol molecule, linkage to a polyethylene glycol molecule, linkage to a spacer 18 (hexaethylene glycol chain) molecule, or 5' to 3' covalent linkage resulting in circularization. A guide polynucleotide that solely comprises of ribonucleic acids is also referred to as a "guide RNA". A guide RNA can include a fusion of two RNA molecules, a crRNA (CRISPR RNA) comprising a variable targeting domain, and a tracrRNA. In one embodiment, the guide RNA comprises a variable targeting domain of 12 to 30 nucleotide sequences and a RNA fragment that can interact with a Cas endonuclease.

The guide polynucleotide can be a double molecule (also referred to as duplex guide polynucleotide) comprising a first nucleotide sequence domain (referred to as Variable Targeting domain or VT domain) that is complementary to a nucleotide sequence in a target DNA and a second nucleotide sequence domain (referred to as Cas endonuclease recognition domain or CER domain) that interacts with a Cas endonuclease polypeptide. The CER domain of the double molecule guide polynucleotide comprises two separate molecules that are hybridized along a region of complementarity. The two separate molecules can be RNA, DNA, and/or RNA-DNA- combination sequences. In some embodiments, the first molecule of the duplex guide polynucleotide comprising a VT domain linked to a CER domain is referred to as "crDNA" (when composed of a contiguous stretch of DNA nucleotides) or "crRNA" (when composed of a contiguous stretch of RNA nucleotides), or "crDNA-RNA" (when composed of a combination of DNA and RNA nucleotides). The crNucleotide can comprise a fragment of the crRNA naturally occurring in Bacteria and Archaea. In one embodiment, the size of the fragment of the crRNA naturally occurring in Bacteria and Archaea that is present in a crNucleotide disclosed herein can range from, but is not limited to, 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20 or more nucleotides. In some embodiments the second molecule of the duplex guide polynucleotide comprising a CER domain is referred to as "tracrRNA" (when composed of a contiguous stretch of RNA nucleotides) or "tracrDNA" (when composed of a contiguous stretch of DNA

nucleotides) or “tracrDNA-RNA” (when composed of a combination of DNA and RNA nucleotides). In one embodiment, the RNA that guides the RNA/ Cas9 endonuclease complex, is a duplexed RNA comprising a duplex crRNA-tracrRNA..

The guide polynucleotide can also be a single molecule comprising a first
5 nucleotide sequence domain (referred to as Variable Targeting domain or VT domain) that is complementary to a nucleotide sequence in a target DNA and a second nucleotide domain (referred to as Cas endonuclease recognition domain or CER domain) that interacts with a Cas endonuclease polypeptide. By “domain” it is meant a contiguous stretch of nucleotides that can be RNA, DNA, and/or RNA-DNA-
10 combination sequence. The VT domain and / or the CER domain of a single guide polynucleotide can comprise a RNA sequence, a DNA sequence, or a RNA-DNA-combination sequence. In some embodiments the single guide polynucleotide comprises a crNucleotide (comprising a VT domain linked to a CER domain) linked to a tracrNucleotide (comprising a CER domain), wherein the linkage is a nucleotide
15 sequence comprising a RNA sequence, a DNA sequence, or a RNA-DNA combination sequence. The single guide polynucleotide being comprised of sequences from the crNucleotide and tracrNucleotide may be referred to as “single guide RNA” (when composed of a contiguous stretch of RNA nucleotides) or “single guide DNA” (when composed of a contiguous stretch of DNA nucleotides) or “single
20 guide RNA-DNA” (when composed of a combination of RNA and DNA nucleotides). In one embodiment of the disclosure, the single guide RNA comprises a crRNA or crRNA fragment and a tracrRNA or tracrRNA fragment of the type II CRISPR/Cas system that can form a complex with a type II Cas endonuclease, wherein said guide RNA/Cas endonuclease complex can direct the
25 Cas endonuclease to a plant genomic target site, enabling the Cas endonuclease to introduce a double strand break into the genomic target site. One aspect of using a single guide polynucleotide versus a duplex guide polynucleotide is that only one expression cassette needs to be made to express the single guide polynucleotide.

The term “variable targeting domain” or “VT domain” is used interchangeably
30 herein and includes a nucleotide sequence that is complementary to one strand (nucleotide sequence) of a double strand DNA target site. The % complementation between the first nucleotide sequence domain (VT domain) and the target

sequence can be at least 50%, 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100%. The
 5 variable target domain can be at least 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29 or 30 nucleotides in length. In some embodiments, the variable targeting domain comprises a contiguous stretch of 12 to 30 nucleotides. The variable targeting domain can be composed of a DNA sequence, a RNA sequence, a modified DNA sequence, a modified RNA sequence, or any
 10 combination thereof.

The term "Cas endonuclease recognition domain" or "CER domain" of a guide polynucleotide is used interchangeably herein and includes a nucleotide sequence (such as a second nucleotide sequence domain of a guide polynucleotide), that interacts with a Cas endonuclease polypeptide. The CER
 15 domain can be composed of a DNA sequence, a RNA sequence, a modified DNA sequence, a modified RNA sequence (see for example modifications described herein), or any combination thereof.

The nucleotide sequence linking the crNucleotide and the tracrNucleotide of a single guide polynucleotide can comprise a RNA sequence, a DNA sequence, or a
 20 RNA-DNA combination sequence. In one embodiment, the nucleotide sequence linking the crNucleotide and the tracrNucleotide of a single guide polynucleotide can be at least 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, 30, 31, 32, 33, 34, 35, 36, 37, 38, 39, 40, 41, 42, 43, 44, 45, 46, 47, 48, 49, 50, 51, 52, 53, 54, 55, 56, 57, 58, 59, 60, 61, 62, 63, 64, 65, 66, 67,
 25 68, 69, 70, 71, 72, 73, 74, 75, 76, 77, 78, 79, 80, 81, 82, 83, 84, 85, 86, 87, 88, 89, 90, 91, 92, 93, 94, 95, 96, 97, 98, 99 or 100 nucleotides in length. In another embodiment, the nucleotide sequence linking the crNucleotide and the tracrNucleotide of a single guide polynucleotide can comprise a loop sequence, such as, but not limiting to a GAAA loop sequence.

30 Nucleotide sequence modification of the guide polynucleotide, VT domain and/or CER domain can be selected from, but not limited to, the group consisting of a 5' cap, a 3' polyadenylated tail, a riboswitch sequence, a stability control

sequence, a sequence that forms a dsRNA duplex, a modification or sequence that targets the guide poly nucleotide to a subcellular location, a modification or sequence that provides for tracking, a modification or sequence that provides a binding site for proteins, a Locked Nucleic Acid (LNA), a 5-methyl dC nucleotide, a 2,6-Diaminopurine nucleotide, a 2'-Fluoro A nucleotide, a 2'-Fluoro U nucleotide; a 2'-O-Methyl RNA nucleotide, a phosphorothioate bond, linkage to a cholesterol molecule, linkage to a polyethylene glycol molecule, linkage to a spacer 18 molecule, a 5' to 3' covalent linkage, or any combination thereof. These modifications can result in at least one additional beneficial feature, wherein the additional beneficial feature is selected from the group of a modified or regulated stability, a subcellular targeting, tracking, a fluorescent label, a binding site for a protein or protein complex, modified binding affinity to complementary target sequence, modified resistance to cellular degradation, and increased cellular permeability.

Polynucleotides of interest are reflective of the commercial markets and interests of those involved in the development of the crop. Crops and markets of interest change, and as developing nations open up world markets, new crops and technologies will emerge also. In addition, as our understanding of agronomic traits and characteristics such as yield and heterosis increase, the choice of genes for gene editing or transformation will change accordingly.

Polynucleotides/polypeptides of interest include, but are not limited to, herbicide-tolerance coding sequences, insecticidal coding sequences, nematocidal coding sequences, antimicrobial coding sequences, antifungal coding sequences, antiviral coding sequences, abiotic and biotic stress tolerance coding sequences, or sequences modifying plant traits such as yield, grain quality, nutrient content, starch quality and quantity, nitrogen fixation and/or utilization, and oil content and/or composition. More specific polynucleotides of interest include, but are not limited to, genes that improve crop yield, genes encoding polypeptides that improve desirability of crops, genes encoding proteins conferring resistance to abiotic stress, such as drought, nitrogen, temperature, salinity, toxic metals or trace elements, or those conferring resistance to toxins such as pesticides and herbicides, or to biotic

stress, such as attacks by fungi, viruses, bacteria, insects, and nematodes, and development of diseases associated with these organisms.

An herbicide resistance protein or a protein resulting from expression of an herbicide resistance-encoding nucleic acid molecule includes proteins that confer
5 upon a cell the ability to tolerate a higher concentration of an herbicide than cells that do not express the protein, or to tolerate a certain concentration of an herbicide for a longer period of time than cells that do not express the protein. Herbicide resistance traits may be introduced into plants by genes coding for resistance to herbicides that act to inhibit the action of acetolactate synthase (ALS), in particular
10 the sulfonylurea-type herbicides, genes coding for resistance to herbicides that act to inhibit the action of glutamine synthase, such as phosphinothricin or basta (e.g., the bar gene), glyphosate (e.g., the EPSP synthase gene and the GAT gene), HPPD inhibitors (e.g., the HPPD gene) or other such genes known in the art. See, for example, US Patent Nos. 7,626,077, 5,310,667, 5,866,775, 6,225,114,
15 6,248,876, 7,169,970, 6,867,293, each of which is herein incorporated by reference.

Agronomically important traits such as oil, starch, and protein content can be genetically altered in addition to using traditional breeding methods. Modifications include increasing content of oleic acid, saturated and unsaturated oils, increasing levels of lysine and sulfur, providing essential amino acids, and also modification of
20 starch. Hordothionin protein modifications are described in U.S. Patent Nos. 5,703,049, 5,885,801, 5,885,802, and 5,990,389, herein incorporated by reference. Another example is lysine and/or sulfur rich seed protein encoded by the soybean 2S albumin described in U.S. Patent No. 5,850,016, and the chymotrypsin inhibitor from barley, described in Williamson *et al.* (1987) *Eur. J. Biochem.* 165:99-106, the
25 disclosures of which are herein incorporated by reference.

Commercial traits can also be encoded on a polynucleotide of interest that could increase for example, starch for ethanol production, or provide expression of proteins. Another important commercial use of transformed plants is the production of polymers and bioplastics such as described in U.S. Patent No. 5,602,321. Genes
30 such as β -Ketothiolase, PHBase (polyhydroxybutyrate synthase), and acetoacetyl-CoA reductase (see Schubert *et al.* (1988) *J. Bacteriol.* 170:5837-5847) facilitate expression of polyhydroxyalkanoates (PHAs).

Derivatives of the coding sequences can be made by site-directed mutagenesis to increase the level of preselected amino acids in the encoded polypeptide. For example, the gene encoding the barley high lysine polypeptide (BHL) is derived from barley chymotrypsin inhibitor, U.S. Application Serial No. 5 08/740,682, filed November 1, 1996, and WO 98/20133, the disclosures of which are herein incorporated by reference. Other agronomically important genes encode latex, Flourey 2, growth factors, seed storage factors, and transcription factors. Polynucleotides that improve crop yield include dwarfing genes, such as Rht1 and Rht2 (Peng *et al.* (1999) *Nature* 400:256-261), and those that increase plant growth, 10 such as ammonium-inducible glutamate dehydrogenase. Polynucleotides that improve desirability of crops include, for example, those that allow plants to have reduced saturated fat content, those that boost the nutritional value of plants, and those that increase grain protein. Polynucleotides that improve salt tolerance are those that increase or allow plant growth in an environment of higher salinity than 15 the native environment of the plant into which the salt-tolerant gene(s) has been introduced.

Polynucleotides/polypeptides that influence amino acid biosynthesis include, for example, anthranilate synthase (AS; EC 4.1.3.27) which catalyzes the first reaction branching from the aromatic amino acid pathway to the biosynthesis of 20 tryptophan in plants, fungi, and bacteria. In plants, the chemical processes for the biosynthesis of tryptophan are compartmentalized in the chloroplast. See, for example, US Pub. 20080050506, herein incorporated by reference. Additional sequences of interest include Chorismate Pyruvate Lyase (CPL) which refers to a gene encoding an enzyme which catalyzes the conversion of chorismate to pyruvate and pHBA. The most well characterized CPL gene has been isolated from *E. coli* 25 and bears the GenBank accession number M96268. See, US Patent No. 7,361,811, herein incorporated by reference.

These polynucleotide sequences of interest may encode proteins involved in providing disease or pest resistance. By "disease resistance" or "pest resistance" is 30 intended that the plants avoid the harmful symptoms that are the outcome of the plant-pathogen interactions. Pest resistance genes may encode resistance to pests that have great yield drag such as rootworm, cutworm, European Corn Borer, and

the like. Disease resistance and insect resistance genes such as lysozymes or cecropins for antibacterial protection, or proteins such as defensins, glucanases or chitinases for antifungal protection, or *Bacillus thuringiensis* endotoxins, protease inhibitors, collagenases, lectins, or glycosidases for controlling nematodes or insects are all examples of useful gene products. Genes encoding disease resistance traits include detoxification genes, such as against fumonisin (U.S. Patent No. 5,792,931); avirulence (avr) and disease resistance (R) genes (Jones *et al.* (1994) *Science* 266:789; Martin *et al.* (1993) *Science* 262:1432; and Mindrinis *et al.* (1994) *Cell* 78:1089); and the like.

Furthermore, it is recognized that the polynucleotide of interest may also comprise antisense sequences complementary to at least a portion of the messenger RNA (mRNA) for a targeted gene sequence of interest. Antisense nucleotides are constructed to hybridize with the corresponding mRNA. Modifications of the antisense sequences may be made as long as the sequences hybridize to and interfere with expression of the corresponding mRNA. In this manner, antisense constructions having 70%, 80%, or 85% sequence identity to the corresponding antisense sequences may be used. Furthermore, portions of the antisense nucleotides may be used to disrupt the expression of the target gene. Generally, sequences of at least 50 nucleotides, 100 nucleotides, 200 nucleotides, or greater may be used.

In addition, the polynucleotide of interest may also be used in the sense orientation to suppress the expression of endogenous genes in plants. Methods for suppressing gene expression in plants using polynucleotides in the sense orientation are known in the art. The methods generally involve transforming plants with a DNA construct comprising a promoter that drives expression in a plant operably linked to at least a portion of a nucleotide sequence that corresponds to the transcript of the endogenous gene. Typically, such a nucleotide sequence has substantial sequence identity to the sequence of the transcript of the endogenous gene, generally greater than about 65% sequence identity, about 85% sequence identity, or greater than about 95% sequence identity. See, U.S. Patent Nos. 5,283,184 and 5,034,323; herein incorporated by reference.

The polynucleotide of interest can also be a phenotypic marker. A phenotypic marker is a screenable or a selectable marker that includes visual markers and selectable markers whether it is a positive or negative selectable marker. Any phenotypic marker can be used. Specifically, a selectable or
5 screenable marker comprises a DNA segment that allows one to identify, or select for or against a molecule or a cell that contains it, often under particular conditions. These markers can encode an activity, such as, but not limited to, production of RNA, peptide, or protein, or can provide a binding site for RNA, peptides, proteins, inorganic and organic compounds or compositions and the like.

10 Examples of selectable markers include, but are not limited to, DNA segments that comprise restriction enzyme sites; DNA segments that encode products which provide resistance against otherwise inhibitory compounds including antibiotics, such as, spectinomycin, ampicillin, kanamycin, TET-repressor, acycline, Basta, neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase
15 (HPT)); DNA segments that encode products which are otherwise lacking in the recipient cell (e.g., tRNA genes, auxotrophic markers); DNA segments that encode products which can be readily identified (e.g., phenotypic markers such as β -galactosidase, GUS; fluorescent proteins such as green fluorescent protein (GFP), cyan (CFP), yellow (YFP), red (RFP), and cell surface proteins); the generation of
20 new primer sites for PCR (e.g., the juxtaposition of two DNA sequence not previously juxtaposed), the inclusion of DNA sequences not acted upon or acted upon by a restriction endonuclease or other DNA modifying enzyme, chemical, etc.; and, the inclusion of a DNA sequences required for a specific modification (e.g., methylation) that allows its identification.

25 Additional selectable markers include genes that confer resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D).

Methods are provided herein for introducing a pollen- inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant.

30 A trait locus of interest can be integrated into a target site by use of a double-strand-break (DSB) inducing agent. The DSB-inducing agent may be provided by any means known in the art. For example, the DSB-inducing agent can

be provided via a polynucleotide encoding the nuclease. Such a polynucleotide encoding a nuclease can be modified to substitute codons having a higher frequency of usage in a plant, as compared to the naturally occurring polynucleotide sequence. The polynucleotide encoding the DSB-inducing agent can be modified to substitute codons having a higher frequency of usage in a maize or soybean plant, as compared to the naturally occurring polynucleotide sequence. A plant having the DSB target site in its genome can also be provided. The DSB-inducing agent may be transiently expressed or the polypeptide itself can be directly provided to the cell. Alternatively, a nucleotide sequence capable of expressing the DSB-inducing agent may be stably integrated into the genome of the plant. In the presence of the corresponding DSB target site and the DSB-inducing agent, a donor DNA comprising the trait of interest can be inserted into the plant's genome. Alternatively, the components of the system (double strand break inducing agent, DSB target site and donor DNA) may be brought together by sexually crossing transformed plants. Thus a sequence encoding the DSB-inducing agent and/or target site (and optionally a donor DNA comprising a trait of interest) can be sexually crossed to one another to allow each component of the system to be present in a single plant. The DSB-inducing agent may be under the control of a constitutive or inducible promoter. Such promoters of interest are discussed in further detail elsewhere herein. Examples of such double-strand- break inducing systems can be guide polynucleotide/Cas endonuclease systems described herein. See also U.S. Patent Application No. 14/463,687, filed August 20, 2014, which is hereby incorporated in its entirety by reference.

As used herein, a "genomic region" is a segment of a chromosome in the genome of a plant cell that is present on either side of a target site or, alternatively, also comprises a portion of the target site. The genomic region can comprise at least 5-10, 5-15, 5-20, 5-25, 5-30, 5-35, 5-40, 5-45, 5- 50, 5-55, 5-60, 5-65, 5- 70, 5- 75, 5-80, 5-85, 5-90, 5-95, 5-100, 5-200, 5-300, 5-400, 5-500, 5-600, 5-700, 5-800, 5-900, 5-1000, 5-1100, 5-1200, 5-1300, 5-1400, 5-1500, 5-1600, 5-1700, 5-1800, 5- 1900, 5-2000, 5-2100, 5-2200, 5-2300, 5-2400, 5-2500, 5-2600, 5-2700, 5-2800, 5- 2900, 5-3000, 5-3100 or more bases such that the genomic region has sufficient

homology to undergo homologous recombination with the corresponding region of homology.

The structural similarity between a given genomic region and the corresponding region of homology found on the donor DNA can be any degree of sequence identity that allows for homologous recombination to occur. For example, the amount of homology or sequence identity shared by the "region of homology" of the donor DNA and the "genomic region" of the plant genome can be at least 50%, 55%, 60%, 65%, 70%, 75%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or 100% sequence identity, such that the sequences undergo homologous recombination.

The region of homology on the donor DNA can have homology to any sequence flanking the target site. While in some embodiments the regions of homology share significant sequence homology to the genomic sequence immediately flanking the target site, it is recognized that the regions of homology can be designed to have sufficient homology to regions that may be further 5' or 3' to the target site. In still other embodiments, the regions of homology can also have homology with a fragment of the target site along with downstream genomic regions. In one embodiment, the first region of homology further comprises a first fragment of the target site and the second region of homology comprises a second fragment of the target site, wherein the first and second fragments are dissimilar.

Homologous recombination includes the exchange of DNA fragments between two DNA molecules at the sites of homology. The frequency of homologous recombination is influenced by a number of factors. Different organisms vary with respect to the amount of homologous recombination and the relative proportion of homologous to non-homologous recombination. Generally, the length of the region of homology affects the frequency of homologous recombination events, the longer the region of homology, the greater the frequency. The length of the homology region needed to observe homologous recombination is also species-variable. In many cases, at least 5 kb of homology has been utilized, but homologous recombination has been observed with as little as 25-50 bp of homology. See, for example, Singer *et al.*, (1982) *Cell* 31:25-33; Shen and Huang, (1986) *Genetics* 112:441-57; Watt *et al.*, (1985) *Proc. Natl. Acad. Sci. USA* 82:4768-

72, Sugawara and Haber, (1992) *Mol Cell Biol* 12:563-75, Rubnitz and Subramani, (1984) *Mol Cell Biol* 4:2253-8; Ayares *et al.*, (1986) *Proc. Natl. Acad. Sci. USA* 83:5199-203; Liskay *et al.*, (1987) *Genetics* 115:161-7.

Once a single or double-strand break is induced in the DNA, the cell's DNA
5 repair mechanism is activated to repair the break. Error-prone DNA repair mechanisms can produce mutations at double-strand break sites. The most common repair mechanism to bring the broken ends together is the nonhomologous end-joining (NHEJ) pathway (Bleuyard *et al.*, (2006) *DNA Repair* 5:1-12). The structural integrity of chromosomes is typically preserved by the repair, but
10 deletions, insertions, or other rearrangements are possible (Siebert and Puchta, (2002) *Plant Cell* 14:1121-31; Pacher *et al.*, (2007) *Genetics* 175:21-9).

Alternatively, the single or double-strand break can be repaired by homologous recombination between homologous DNA sequences. Once the sequence around the double-strand break is altered, for example, by exonuclease
15 activities involved in the maturation of double-strand breaks, gene conversion pathways can restore the original structure if a homologous sequence is available, such as a homologous chromosome in non-dividing somatic cells, or a sister chromatid after DNA replication (Molinier *et al.*, (2004) *Plant Cell* 16:342-52). Ectopic and/or epigenic DNA sequences may also serve as a DNA repair template
20 for homologous recombination (Puchta, (1999) *Genetics* 152:1173-81).

DNA double-strand breaks appear to be an effective factor to stimulate homologous recombination pathways (Puchta *et al.*, (1995) *Plant Mol Biol* 28:281-92; Tzfira and White, (2005) *Trends Biotechnol* 23:567-9; Puchta, (2005) *J Exp Bot* 56:1-14). Using DNA-breaking agents, a two- to nine-fold increase of homologous
25 recombination was observed between artificially constructed homologous DNA repeats in plants (Puchta *et al.*, (1995) *Plant Mol Biol* 28:281-92). In maize protoplasts, experiments with linear DNA molecules demonstrated enhanced homologous recombination between plasmids (Lyznik *et al.*, (1991) *Mol Gen Genet* 230:209-18).

30 Once a double-strand break is introduced in the DSB target site by the DSB inducing agent, the first and second regions of homology of the donor DNA can undergo homologous recombination with their corresponding genomic regions of

homology resulting in exchange of DNA between the donor and the genome. As such, the provided method results in the integration of the donor DNA (comprising for example a trait of interest or a polynucleotide of interest) into the double-strand break in the DSB target site in the plant genome (as described in U.S. Patent
5 Application No. 14/463,687, filed August 20, 2014, which is hereby incorporated in its entirety by reference.

The donor DNA may be introduced by any means known in the art. For example, the donor DNA may be provided transiently to a plant or plant cell by any method known in the art. The donor DNA may be provided by any transformation
10 method known in the art including, for example, Agrobacterium-mediated transformation or biolistic particle bombardment. The donor DNA may be present transiently in the cell or it could be introduced via a viral replicon. In the presence of a DSB inducing agent and the DSB target site, the donor DNA can be inserted into the transformed plant's genome.

15 A trait locus of interest can also be integrated (introduced; inserted) into a target site (located proximal or distal to a telomere) by use of a site-specific integration (SSI) system discussed in further detail elsewhere herein. The site-specific recombination system employs various components which are described in detail below and in U.S. Patent Nos. 6187994, 6262341, 6331661 and 6300545,
20 each of which is herein incorporated by reference. A recombinase is provided that recognizes and implements recombination at the recombination sites of the transgenic SSI target site and the transfer cassette. The recombinase can be provided by any means known in the art and is described in detail elsewhere herein. The coding region of a transfer cassette can encode a recombinase that facilitates
25 recombination between the first and the second recombination sites of the transfer cassette and the transgenic SSI target site, the second and the third recombination sites of the transfer cassette and the transgenic SSI target site, or the first and the third recombination sites of the transfer cassette and the transgenic SSI target site.

Methods for selecting plant cells with integration at the target site, such as
30 selecting for cells expressing a trait of interest, a polynucleotide of interest, or a selectable marker, are known in the art.

As discussed above, various methods can be used to introduce a trait gene of interest and/or pollen-inhibitor genes into the genome of a plant or plant cell, thereby creating a plant having within a genomic window at least one trait locus of interest and/or a pollen-inhibitor gene integrated into a target site.

5 Non-limiting examples of various DNA constructs, transgenic SSI target sites, and transfer cassettes that can be used to insert a polynucleotide of interest into a plant or plant cell are described in PCT/US12/47202 application filed July 18, 2012, incorporated by reference in its entirety herein. In short, once the trait gene of interest has integrated into the target site or once the pollen-inhibitor cassette has
10 integrated into the target site, the appropriate selective agent can be employed to identify the plant cell having the desired DNA construct. Once a target site has been established within the genome, additional target sites may be introduced by incorporating such sites within the nucleotide sequence of the transfer cassette. Thus, once a SSI target site has been established, it is possible to subsequently add
15 or alter sites through recombination or DSB technology. Such methods are described in detail in WO 99/25821, herein incorporated by reference.

Transformation protocols as well as protocols for introducing polypeptides or polynucleotide sequences into plants may vary depending on the type of plant or plant cell, i.e., monocot or dicot, targeted for transformation. By "stably
20 incorporated" or "stably introduced" is intended the introduction of a polynucleotide into the plant such that the nucleotide sequence integrates into the genome of the plant and is capable of being inherited by progeny thereof. Any protocol may be used for the stable incorporation of the DNA constructs or the various components of the pollen-inhibitor system employed herein.

25 Suitable methods of introducing polypeptides and polynucleotides into plant cells include microinjection (Crossway *et al.* (1986) *Biotechniques* 4:320-334), electroporation (Riggs *et al.* (1986) *Proc. Natl. Acad. Sci. USA* 83:5602-5606, *Agrobacterium*-mediated transformation (U.S. Patent No. 5,563,055 and U.S. Patent No. 5,981,840), direct gene transfer (Paszkowski *et al.* (1984) *EMBO J.*
30 3:2717-2722), and ballistic particle acceleration (see, for example, U.S. Patent Nos. 4,945,050; U.S. Patent No. 5,879,918; U.S. Patent No. 5,886,244; and, 5,932,782; Tomes *et al.* (1995) in *Plant Cell, Tissue, and Organ Culture: Fundamental*

Methods, ed. Gamborg and Phillips (Springer-Verlag, Berlin); McCabe *et al.* (1988) *Biotechnology* 6:923-926); and Lec1 transformation (WO 00/28058). Also see Weissinger *et al.* (1988) *Ann. Rev. Genet.* 22:421-477; Sanford *et al.* (1987) *Particulate Science and Technology* 5:27-37 (onion); Christou *et al.* (1988) *Plant*
 5 *Physiol.* 87:671-674 (soybean); McCabe *et al.* (1988) *Bio/Technology* 6:923-926 (soybean); Finer and McMullen (1991) *In Vitro Cell Dev. Biol.* 27P:175-182 (soybean); Singh *et al.* (1998) *Theor. Appl. Genet.* 96:319-324 (soybean); Datta *et al.* (1990) *Biotechnology* 8:736-740 (rice); Klein *et al.* (1988) *Proc. Natl. Acad. Sci. USA* 85:4305-4309 (maize); Klein *et al.* (1988) *Biotechnology* 6:559-563 (maize);
 10 U.S. Patent Nos. 5,240,855; 5,322,783; and, 5,324,646; Klein *et al.* (1988) *Plant Physiol.* 91:440-444 (maize); Fromm *et al.* (1990) *Biotechnology* 8:833-839 (maize); Hooykaas-Van Slogteren *et al.* (1984) *Nature (London)* 311:763-764; U.S. Patent No. 5,736,369 (cereals); Bytebier *et al.* (1987) *Proc. Natl. Acad. Sci. USA* 84:5345-5349 (Liliaceae); De Wet *et al.* (1985) in *The Experimental Manipulation of Ovule*
 15 *Tissues*, ed. Chapman *et al.* (Longman, New York), pp. 197-209 (pollen); Kaeppler *et al.* (1990) *Plant Cell Reports* 9:415-418 and Kaeppler *et al.* (1992) *Theor. Appl. Genet.* 84:560-566 (whisker-mediated transformation); D'Halluin *et al.* (1992) *Plant Cell* 4:1495-1505 (electroporation); Li *et al.* (1993) *Plant Cell Reports* 12:250-255 and Christou and Ford (1995) *Annals of Botany* 75:407-413 (rice); Osjoda *et al.*
 20 (1996) *Nature Biotechnology* 14:745-750 (maize via *Agrobacterium tumefaciens*); all of which are herein incorporated by reference.

In other embodiments, any of the polynucleotides employed herein may be introduced into plants by contacting plants with a virus or viral nucleic acids. Generally, such methods involve incorporating a desired polynucleotide within a
 25 viral DNA or RNA molecule. It is recognized that a sequence employed in the methods or compositions provided herein may be initially synthesized as part of a viral polyprotein, which later may be processed by proteolysis *in vivo* or *in vitro* to produce the desired recombinant protein. Further, it is recognized that promoters employed herein also encompass promoters utilized for transcription by viral RNA
 30 polymerases. Methods for introducing polynucleotides into plants and expressing a protein encoded therein, involving viral DNA or RNA molecules, are known in the art. See, for example, U.S. Patent Nos. 5,889,191, 5,889,190, 5,866,785,

5,589,367, 5,316,931, and Porta *et al.* (1996) *Molecular Biotechnology* 5:209-221; herein incorporated by reference.

The trait gene(s) of interest, color marker genes, and/or pollen-inhibitor gene(s) can be provided to a plant using a variety of transient transformation methods. "Transient transformation" is intended to mean that a polynucleotide is introduced into the host (i.e., a plant) and expressed temporally. Such transient transformation methods include, but are not limited to, the introduction of any of the components of the pollen-inhibitor system or active fragments or variants thereof directly into the plant or the introduction of the transcript into the plant. Such methods include, for example, microinjection or particle bombardment. See, for example, Crossway *et al.* (1986) *Mol Gen. Genet.* 202:179-185; Nomura *et al.* (1986) *Plant Sci.* 44:53-58; Hepler *et al.* (1994) *Proc. Natl. Acad. Sci.* 91: 2176-2180 and Hush *et al.* (1994) *The Journal of Cell Science* 107:775-784, all of which are herein incorporated by reference. Alternatively, the polynucleotide can be transiently transformed into the plant using techniques known in the art. Such techniques include viral vector system and the precipitation of the polynucleotide in a manner that precludes subsequent release of the DNA. Thus, the transcription from the particle-bound DNA can occur, but the frequency with which it is released to become integrated into the genome is greatly reduced. Such methods include the use particles coated with polyethylimine (PEI; Sigma #P3143).

The cells that have been transformed may be grown into plants in accordance with conventional ways. See, for example, McCormick *et al.* (1986) *Plant Cell Reports* 5:81-84. These plants may then be grown, and either pollinated with the same transformed strain or different strains, and the resulting progeny having constitutive expression of the desired phenotypic characteristic identified. Two or more generations may be grown to ensure that expression of the desired phenotypic characteristic is stably maintained and inherited and then seeds harvested to ensure expression of the desired phenotypic characteristic has been achieved. In this manner, transformed seed having the recited DNA construct stably incorporated into their genome is provided.

In specific embodiments, the transgenic SSI target site of the plant cell, plant, plant part and seed further comprises a third recombination site between the first

and the second recombination site, wherein the third recombination site is dissimilar and non-recombinogenic to the first and the second recombination sites. The first, second, and third recombination sites can comprise, for example, FRT1, FRT5, FRT6, FRT12, FRT62 (described in US patent US8318493 issued on November 27, 2012, herein incorporated by reference), or FRT87. Also, provided is a plant cell, plant, or seed wherein the first recombination site is FRT1, the second recombination site is FRT12 and the third recombination site is FRT87.

Plants, plant cells, or seeds having in their genome a genomic window comprising at least one trait locus and at least one pollen-inhibitor gene or a color marker gene provided herein are also encompassed.

Plants described herein include plants that comprise a trait locus that is flanked by a pollen-inhibitor gene on each side; plants that comprise a trait locus that is flanked by a color maker gene on each side, or plants that comprise a trait locus that is flanked by a color marker gene on one side and a pollen-inhibitor gene on the opposite side of the trait locus. The proximity of the pollen-inhibitor gene or color marker gene to the trait locus of interest can be about at least 0.05 cM, 0.1 cM, 0.2 cM, 0.3 cM, 0.4 cM, 0.5 cM, 0.6 cM, 0.7 cM, 0.8 cM, 0.9 cM, 1.0 cM, 1.1 cM, 1.2 cM, 1.3 cM, 1.4 cM, 1.5 cM, 1.6 cM, 1.7 cM, 1.8 cM, 1.9 cM, 2.0 cM, 2.1 cM, 2.2 cM, 2.3 cM, 2.4 cM, 2.5 cM, 2.6 cM, 2.7 cM, 2.8 cM, 2.9 cM, 3.0 cM, 3.1 cM, 3.2 cM, 3.3 cM, 3.4 cM, 3.5 cM, 3.6 cM, 3.7 cM, 3.8 cM, 3.9 cM, 4.0 cM or 5.0 cM.

In one embodiment the composition comprises a plant comprising at least one trait gene of interest, a first recombinant DNA construct comprising a color marker gene, and a second recombinant DNA construct comprising a second color marker gene, wherein said first recombinant DNA construct and said second recombinant DNA construct are genetically linked and flank said trait gene of interest.

In one embodiment the composition comprises a plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a color marker gene, wherein said trait locus of interest and said color marker gene segregate independently from one another at a rate of about 10% to about 0.1%.

In one embodiment the composition comprises a plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a pollen-

inhibitor gene, wherein said trait locus of interest and said a pollen-inhibitor gene segregate independently from one another at a rate of about 10% to about 0.1%.

Compositions as described herein include plants wherein the first pollen-inhibitor gene (or color maker gene), the second pollen-inhibitor gene (or color maker gene) and the trait gene of interest are located within 0.05 cM, 0.1 cM, 0.2 cM, 0.3 cM, 0.4 cM, 0.5 cM, 0.6 cM, 0.7 cM, 0.8 cM, 0.9 cM, 1.0 cM, 1.1 cM, 1.2 cM, 1.3 cM, 1.4 cM, 1.5 cM, 1.6 cM, 1.7 cM, 1.8 cM, 1.9 cM, 2.0 cM, 2.1 cM, 2.2 cM, 2.3 cM, 2.4 cM, 2.5 cM, 2.6 cM, 2.7 cM, 2.8 cM, 2.9 cM, 3.0 cM, 3.1 cM, 3.2 cM, 3.3 cM, 3.4 cM, 3.5 cM, 3.6 cM, 3.7 cM, 3.8 cM, 3.9 cM, 4.0 cM or 5.0 cM of each other.

Compositions as described herein include plants wherein the first pollen-inhibitor gene and the second pollen-inhibitor is selected from the group consisting of barnase, alpha amylase, KID, or any combination thereof. Compositions as described herein include plants wherein the second pollen-inhibitor gene is selected from the group consisting a non-conditional gene, a conditional gene and an inducible gene.

As used herein, the term plant includes plant cells, plant protoplasts, plant cell tissue cultures from which a plant can be regenerated, plant calli, plant clumps, and plant cells that are intact in plants or parts of plants such as embryos, pollen, ovules, seeds, leaves, flowers, branches, fruit, kernels, ears, cobs, husks, stalks, roots, root tips, anthers, and the like. Grain is intended to mean the mature seed produced by commercial growers for purposes other than growing or reproducing the species. Progeny, variants, and mutants of the regenerated plants are also included herein, provided that these parts comprise the recited DNA construct.

A transgenic plant includes, for example, a plant which comprises within its genome a heterologous polynucleotide introduced by a transformation step. The heterologous polynucleotide can be stably integrated within the genome such that the polynucleotide is passed on to successive generations. The heterologous polynucleotide may be integrated into the genome alone or as part of a recombinant DNA construct. A transgenic plant can also comprise more than one heterologous polynucleotide within its genome. Each heterologous polynucleotide may confer a different trait to the transgenic plant. A heterologous polynucleotide can include a sequence that originates from a foreign species, or, if from the same species, can

be substantially modified from its native form. Transgenic can include any cell, cell line, callus, tissue, plant part or plant, the genotype of which has been altered by the presence of heterologous nucleic acid including those transgenics initially so altered as well as those created by sexual crosses or asexual propagation from the initial
5 transgenic. The alterations of the genome (chromosomal or extra-chromosomal) by conventional plant breeding methods, by the genome editing procedure described herein that does not result in an insertion of a foreign polynucleotide, or by naturally occurring events such as random cross-fertilization, non-recombinant viral infection, non-recombinant bacterial transformation, non-recombinant transposition, or
10 spontaneous mutation are not intended to be regarded as transgenic.

In certain embodiments of the disclosure, a fertile plant is a plant that produces viable male and female gametes and is self-fertile. Such a self-fertile plant can produce a progeny plant without the contribution from any other plant of a gamete and the genetic material contained therein. Other embodiments of the
15 disclosure can involve the use of a plant that is not self-fertile because the plant does not produce male gametes, or female gametes, or both, that are viable or otherwise capable of fertilization. As used herein, a "male sterile plant" is a plant that does not produce male gametes that are viable or otherwise capable of fertilization. As used herein, a "female sterile plant" is a plant that does not produce
20 female gametes that are viable or otherwise capable of fertilization. It is recognized that male-sterile and female-sterile plants can be female-fertile and male-fertile, respectively. It is further recognized that a male fertile (but female sterile) plant can produce viable progeny when crossed with a female fertile plant and that a female fertile (but male sterile) plant can produce viable progeny when crossed with a male
25 fertile plant.

In one embodiment, the plant is a soybean or maize plant, wherein the genomic window described herein is not more than 0.1, 0.2, 0.3, 0.4, 0.5, 1, 2, 5, 10 cM in length.

In one embodiment the plant is a plant comprising at least one trait gene of
30 interest, a first recombinant DNA construct comprising a first pollen-inhibitor gene, and a second recombinant DNA construct comprising a second pollen-inhibitor gene, wherein said first recombinant DNA construct and said second recombinant

DNA construct are genetically linked and flank said trait gene of interest. The first pollen-inhibitor gene and the second pollen-inhibitor can be selected from the group consisting of barnase, alpha amylase, KID, coda or CYP105A, or any combination thereof. The second pollen-inhibitor gene can be ,a non-conditional gene , a
 5 conditional gene and an inducible gene.

In one embodiment the plant is a plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a pollen-inhibitor gene, wherein said trait locus of interest and said a pollen-inhibitor gene segregate independently from one another at a rate of about 10% to about 0.1%.

10 In one embodiment, the plant is a soybean or maize plant, wherein the genomic window comprises at least one transgene and at least one pollen-inhibitor gene, , wherein the transgene confers a trait selected from the group consisting of herbicide tolerance, insect resistance, disease resistance, male sterility, site-specific recombination, abiotic stress tolerance, altered phosphorus, altered antioxidants,
 15 altered fatty acids, altered essential amino acids, altered carbohydrates, herbicide tolerance, insect resistance and disease resistance.

The trait gene(s) of interest, color marker gene(s), and/or pollen- inhibitor gene(s) described herein can be of used in any plant species, including, but not limited to, monocots and dicots. Examples of plant species of interest include, but are
 20 not limited to, corn (maize) (*Zea mays*), *Brassica* sp. (e.g., *B. napus*, *B. rapa*, *B. juncea*), particularly those *Brassica* species useful as sources of seed oil, alfalfa (*Medicago sativa*), rice (*Oryza sativa*), rye (*Secale cereale*), sorghum (*Sorghum bicolor*, *Sorghum vulgare*), millet (e.g., pearl millet (*Pennisetum glaucum*), proso millet (*Panicum miliaceum*), foxtail millet (*Setaria italica*), finger millet (*Eleusine coracana*)),
 25 sunflower (*Helianthus annuus*), safflower (*Carthamus tinctorius*), wheat (*Triticum aestivum*), soybean (*Glycine max*), tobacco (*Nicotiana tabacum*), potato (*Solanum tuberosum*), peanuts (*Arachis hypogaea*), cotton (*Gossypium barbadense*, *Gossypium hirsutum*), sweet potato (*Ipomoea batatas*), cassava (*Manihot esculenta*), coffee (*Coffea* spp.), coconut (*Cocos nucifera*), pineapple (*Ananas comosus*), citrus trees
 30 (*Citrus* spp.), cocoa (*Theobroma cacao*), tea (*Camellia sinensis*), banana (*Musa* spp.), avocado (*Persea americana*), fig (*Ficus casica*), guava (*Psidium guajava*), mango (*Mangifera indica*), olive (*Olea europaea*), papaya (*Carica papaya*), cashew

(*Anacardium occidentale*), macadamia (*Macadamia integrifolia*), almond (*Prunus amygdalus*), sugar beets (*Beta vulgaris*), sugarcane (*Saccharum* spp.), oats, barley, vegetables, ornamentals, and conifers.

Vegetables include tomatoes (*Lycopersicon esculentum*), lettuce (e.g., *Lactuca sativa*), green beans (*Phaseolus vulgaris*), lima beans (*Phaseolus limensis*), peas (*Lathyrus* spp.), and members of the genus *Cucumis* such as cucumber (*C. sativus*), cantaloupe (*C. cantalupensis*), and musk melon (*C. melo*). Ornamentals include azalea (*Rhododendron* spp.), hydrangea (*Macrophylla hydrangea*), hibiscus (*Hibiscus rosasanensis*), roses (*Rosa* spp.), tulips (*Tulipa* spp.), daffodils (*Narcissus* spp.),
 10 petunias (*Petunia hybrida*), carnation (*Dianthus caryophyllus*), poinsettia (*Euphorbia pulcherrima*), and chrysanthemum.

Conifers that may be employed include, for example, pines such as loblolly pine (*Pinus taeda*), slash pine (*Pinus elliotii*), ponderosa pine (*Pinus ponderosa*), lodgepole pine (*Pinus contorta*), and Monterey pine (*Pinus radiata*); Douglas-fir (*Pseudotsuga menziesii*); Western hemlock (*Tsuga canadensis*); Sitka spruce (*Picea glauca*);
 15 redwood (*Sequoia sempervirens*); true firs such as silver fir (*Abies amabilis*) and balsam fir (*Abies balsamea*); and cedars such as Western red cedar (*Thuja plicata*) and Alaska yellow-cedar (*Chamaecyparis nootkatensis*). In specific embodiments, plants are crop plants (for example, corn, alfalfa, sunflower, *Brassica*, soybean, cotton, safflower, peanut, sorghum, wheat, millet, tobacco, etc.). In other embodiments, corn
 20 and soybean plants are optimal, and in yet other embodiments corn plants are optimal.

Other plants of interest include grain plants that provide seeds of interest, oil-seed plants, and leguminous plants. Seeds of interest include grain seeds, such as corn, wheat, barley, rice, sorghum, rye, etc. Oil-seed plants include cotton,
 25 soybean, safflower, sunflower, *Brassica*, maize, alfalfa, palm, coconut, etc. Leguminous plants include beans and peas. Beans include guar, locust bean, fenugreek, soybean, garden beans, cowpea, mungbean, lima bean, fava bean, lentils, chickpea, etc.

The terms “polynucleotide,” “polynucleotide sequence,” “nucleic acid
 30 sequence,” and “nucleic acid fragment” are used interchangeably herein. These terms encompass nucleotide sequences and the like. A polynucleotide may be a polymer of RNA or DNA that is single- or double-stranded, that optionally contains

synthetic, non-natural or altered nucleotide bases. A polynucleotide in the form of a polymer of DNA may be comprised of one or more segments of cDNA, genomic DNA, synthetic DNA, or mixtures thereof. The use of the term "polynucleotide" is not intended to limit the present invention to polynucleotides comprising DNA.

5 Those of ordinary skill in the art will recognize that polynucleotides, can comprise ribonucleotides and combinations of ribonucleotides and deoxyribonucleotides. Such deoxyribonucleotides and ribonucleotides include both naturally occurring molecules and synthetic analogues. The polynucleotides provided herein also encompass all forms of sequences including, but not limited to, single-stranded
10 forms, double-stranded forms, hairpins, stem-and-loop structures, and the like.

The compositions provided herein can comprise an isolated or substantially purified polynucleotide. An "isolated" or "purified" polynucleotide is substantially or essentially free from components that normally accompany or interact with the polynucleotide as found in its naturally occurring environment. Thus, an isolated or
15 purified polynucleotide is substantially free of other cellular material, or culture medium when produced by recombinant techniques, or substantially free of chemical precursors or other chemicals when chemically synthesized. Optimally, an "isolated" polynucleotide is free of sequences (optimally protein encoding sequences) that naturally flank the polynucleotide (i.e., sequences located at the 5'
20 and 3' ends of the polynucleotide) in the genomic DNA of the organism from which the polynucleotide is derived. For example, in various embodiments, the isolated polynucleotide can contain less than about 5 kb, 4 kb, 3 kb, 2 kb, 1 kb, 0.5 kb, or 0.1 kb of nucleotide sequence that naturally flank the polynucleotide in genomic DNA of the cell from which the polynucleotide is derived.

25 The terms "recombinant polynucleotide" and "recombinant DNA construct" are used interchangeably herein. A recombinant construct can comprise an artificial or heterologous combination of nucleic acid sequences, e.g., regulatory and coding sequences that are not found together in nature. For example, a transfer cassette can comprise restriction sites and a heterologous polynucleotide of interest. In other
30 embodiments, a recombinant construct may comprise regulatory sequences and coding sequences that are derived from different sources, or regulatory sequences and coding sequences derived from the same source, but arranged in a manner

different than that found in nature. Such a construct may be used by itself or may be used in conjunction with a vector. If a vector is used, then the choice of vector is dependent upon the method that will be used to transform host cells as is well known to those skilled in the art. For example, a plasmid vector can be used. The skilled artisan is well aware of the genetic elements that must be present on the vector in order to successfully transform, select and propagate host cells comprising any of the isolated nucleic acid fragments provided herein. The skilled artisan will also recognize that different independent transformation events will result in different levels and patterns of expression (Jones *et al.*, *EMBO J.* 4:2411-2418 (1985); De Almeida *et al.*, *Mol. Gen. Genetics* 218:78-86 (1989)), and thus that multiple events must be screened in order to obtain lines displaying the desired expression level and pattern. Such screening may be accomplished by Southern analysis of DNA, Northern analysis of mRNA expression, immunoblotting analysis of protein expression, or phenotypic analysis, among others.

The trait gene(s) of interest, color marker genes, and/or pollen-inhibitor gene(s) described herein can be provided in an expression cassette for expression in a plant or other organism or cell type of interest. The cassette can include 5' and 3' regulatory sequences operably linked to a polynucleotide provided herein. "Operably linked" is intended to mean a functional linkage between two or more elements. For example, an operable linkage between a polynucleotide of interest and a regulatory sequence (i.e., a promoter) is a functional link that allows for expression of the polynucleotide of interest. Operably linked elements may be contiguous or non-contiguous. When used to refer to the joining of two protein coding regions, by operably linked is intended that the coding regions are in the same reading frame. The cassette may additionally contain at least one additional gene to be cotransformed into the organism. Alternatively, the additional gene(s) can be provided on multiple expression cassettes. Such an expression cassette is provided with a plurality of restriction sites and/or recombination sites for insertion of a recombinant polynucleotide to be under the transcriptional regulation of the regulatory regions. The expression cassette may additionally contain selectable marker genes.

The expression cassette can include in the 5'-3' direction of transcription, a transcriptional and translational initiation region (i.e., a promoter), a recombinant polynucleotide provided herein, and a transcriptional and translational termination region (i.e., termination region) functional in plants. The regulatory regions (i.e., promoters, transcriptional regulatory regions, and translational termination regions) and/or a polynucleotide provided herein may be native/analogous to the host cell or to each other. Alternatively, the regulatory regions and/or a polynucleotide provided herein may be heterologous to the host cell or to each other. As used herein, "heterologous" in reference to a sequence is a sequence that originates from a foreign species, or, if from the same species, is substantially modified from its native form in composition and/or trait locus by deliberate human intervention. For example, a promoter operably linked to a heterologous polynucleotide is from a species different from the species from which the polynucleotide was derived, or, if from the same/analogous species, one or both are substantially modified from their original form and/or genomic locus, or the promoter is not the native promoter for the operably linked polynucleotide. Alternatively, the regulatory regions and/or a recombinant polynucleotide provided herein may be entirely synthetic.

The termination region may be native with the transcriptional initiation region, may be native with the operably linked recombinant polynucleotide, may be native with the plant host, or may be derived from another source (i.e., foreign or heterologous) to the promoter, the recombinant polynucleotide, the plant host, or any combination thereof. Convenient termination regions are available from the Ti-plasmid of *A. tumefaciens*, such as the octopine synthase and nopaline synthase termination regions. See also Guerineau *et al.* (1991) *Mol. Gen. Genet.* 262:141-144; Proudfoot (1991) *Cell* 64:671-674; Sanfacon *et al.* (1991) *Genes Dev.* 5:141-149; Mogen *et al.* (1990) *Plant Cell* 2:1261-1272; Munroe *et al.* (1990) *Gene* 91:151-158; Ballas *et al.* (1989) *Nucleic Acids Res.* 17:7891-7903; and Joshi *et al.* (1987) *Nucleic Acids Res.* 15:9627-9639.

In preparing the expression cassette, the various DNA fragments may be manipulated, so as to provide for the DNA sequences in the proper orientation. Toward this end, adapters or linkers may be employed to join the DNA fragments or other manipulations may be involved to provide for convenient restriction sites,

removal of superfluous DNA, removal of restriction sites, or the like. For this purpose, *in vitro* mutagenesis, primer repair, restriction, annealing, resubstitutions, e.g., transitions and transversions, may be involved.

A number of promoters can be used in the expression cassettes provided
5 herein. The promoters can be selected based on the desired outcome. It is recognized that different applications can be enhanced by the use of different promoters in the expression cassettes to modulate the timing, location and/or level of expression of the polynucleotide of interest. Such expression constructs may also contain, if desired, a promoter regulatory region (e.g., one conferring inducible,
10 constitutive, environmentally- or developmentally-regulated, or cell- or tissue-specific/selective expression), a transcription initiation start site, a ribosome binding site, an RNA processing signal, a transcription termination site, and/or a polyadenylation signal.

In some embodiments, an expression cassette provided herein can be
15 combined with constitutive, tissue-preferred, or other promoters for expression in plants. Examples of constitutive promoters include the cauliflower mosaic virus (CaMV) 35S transcription initiation region, the 1'- or 2'-promoter derived from T-DNA of *Agrobacterium tumefaciens*, the ubiquitin 1 promoter, the Smas promoter, the cinnamyl alcohol dehydrogenase promoter (U.S. Pat. No. 5,683,439), the Nos
20 promoter, the pEmu promoter, the rubisco promoter, the GRP1-8 promoter and other transcription initiation regions from various plant genes known to those of skill. If low level expression is desired, weak promoter(s) may be used. Weak constitutive promoters include, for example, the core promoter of the Rsyn7 promoter (WO 99/43838 and U.S. Pat. No. 6,072,050), the core 35S CaMV
25 promoter, and the like. Other constitutive promoters include, for example, U.S. Pat. Nos. 5,608,149; 5,608,144; 5,604,121; 5,569,597; 5,466,785; 5,399,680; 5,268,463; and 5,608,142. See also, U.S. Pat. No. 6,177,611, herein incorporated by reference.

Examples of inducible promoters are the Adh1 promoter which is inducible by
30 hypoxia or cold stress, the Hsp70 promoter which is inducible by heat stress, the PPK promoter and the pepcarboxylase promoter which are both inducible by light. Also useful are promoters which are chemically inducible, such as the In2-2

promoter which is safener induced (U.S. Pat. No. 5,364,780), the ERE promoter which is estrogen induced, and the Axig1 promoter which is auxin induced and tapetum specific but also active in callus (PCT US01/22169).

5 Examples of promoters under developmental control include promoters that initiate transcription preferentially in certain tissues, such as leaves, roots, fruit, seeds, or flowers. An exemplary promoter is the anther specific promoter 5126 (U.S. Pat. Nos. 5,689,049 and 5,689,051). Examples of seed-preferred promoters include, but are not limited to, 27 kD gamma zein promoter and waxy promoter, Boronat, A. *et al.* (1986) *Plant Sci.* 47:95-102; Reina, M. *et al.* *Nucl. Acids Res.* 10 18(21):6426; and Kloesgen, R. B. *et al.* (1986) *Mol. Gen. Genet.* 203:237-244. Promoters that express in the embryo, pericarp, and endosperm are disclosed in U.S. Pat. No. 6,225,529 and PCT publication WO 00/12733. The disclosures for each of these are incorporated herein by reference in their entirety.

15 Chemical-regulated promoters can be used to modulate the expression of a gene in a plant through the application of an exogenous chemical regulator. Depending upon the objective, the promoter may be a chemical-inducible promoter, where application of the chemical induces gene expression, or a chemical-repressible promoter, where application of the chemical represses gene expression. Chemical-inducible promoters are known in the art and include, but are not limited 20 to, the maize In2-2 promoter, which is activated by benzenesulfonamide herbicide safeners, the maize GST promoter, which is activated by hydrophobic electrophilic compounds that are used as pre-emergent herbicides, and the tobacco PR-1a promoter, which is activated by salicylic acid. Other chemical-regulated promoters of interest include steroid-responsive promoters (see, for example, the 25 glucocorticoid-inducible promoter in Schena *et al.* (1991) *Proc. Natl. Acad. Sci. USA* 88:10421-10425 and McNellis *et al.* (1998) *Plant J.* 14(2):247-257) and TET repressoracycline-inducible and TET repressoracycline-repressible promoters (see, for example, Gatz *et al.* (1991) *Mol. Gen. Genet.* 227:229-237, and U.S. Pat. Nos. 5,814,618 and 5,789,156), herein incorporated by reference.

30 Tissue-preferred promoters can be utilized to target enhanced expression of a polynucleotide of interest within a particular plant tissue. Tissue-preferred promoters are known in the art. See, for example, Yamamoto *et al.* (1997) *Plant J.*

12(2):255-265; Kawamata *et al.* (1997) *Plant Cell Physiol.* 38(7):792-803; Hansen *et al.* (1997) *Mol. Gen. Genet.* 254(3):337-343; Russell *et al.* (1997) *Transgenic Res.* 6(2):157-168; Rinehart *et al.* (1996) *Plant Physiol.* 112(3):1331-1341; Van Camp *et al.* (1996) *Plant Physiol.* 112(2):525-535; Canevascini *et al.* (1996) *Plant Physiol.* 112(2):513-524; Yamamoto *et al.* (1994) *Plant Cell Physiol.* 35(5):773-778; Lam (1994) *Results Probl. Cell Differ.* 20:181-196; Orozco *et al.* (1993) *Plant Mol Biol.* 23(6):1129-1138; Matsuoka *et al.* (1993) *Proc Natl. Acad. Sci. USA* 90(20):9586-9590; and Guevara-Garcia *et al.* (1993) *Plant J.* 4(3):495-505. Such promoters can be modified, if necessary, for weak expression.

10 Leaf-preferred promoters are known in the art. See, for example, Yamamoto *et al.* (1997) *Plant J.* 12(2):255-265; Kwon *et al.* (1994) *Plant Physiol.* 105:357-67; Yamamoto *et al.* (1994) *Plant Cell Physiol.* 35(5):773-778; Gotor *et al.* (1993) *Plant J.* 3:509-18; Orozco *et al.* (1993) *Plant Mol. Biol.* 23(6):1129-1138; and Matsuoka *et al.* (1993) *Proc. Natl. Acad. Sci. USA* 90(20):9586-9590. In addition, the promoters
15 of cab and rubisco can also be used. See, for example, Simpson *et al.* (1958) *EMBO J* 4:2723-2729 and Timko *et al.* (1988) *Nature* 318:57-58.

Root-preferred promoters are known and can be selected from the many available from the literature or isolated *de novo* from various compatible species. See, for example, Hire *et al.* (1992) *Plant Mol. Biol.* 20(2):207-218 (soybean root-specific glutamine synthetase gene); Keller and Baumgartner (1991) *Plant Cell* 3(10):1051-1061 (root-specific control element in the GRP 1.8 gene of French bean); Sanger *et al.* (1990) *Plant Mol. Biol.* 14(3):433-443 (root-specific promoter of the mannopine synthase (MAS) gene of *Agrobacterium tumefaciens*); and Miao *et al.* (1991) *Plant Cell* 3(1):11-22 (full-length cDNA clone encoding cytosolic glutamine
20 synthetase (GS), which is expressed in roots and root nodules of soybean). See also Bogusz *et al.* (1990) *Plant Cell* 2(7):633-641, where two root-specific promoters isolated from hemoglobin genes from the nitrogen-fixing nonlegume *Parasponia andersonii* and the related non-nitrogen-fixing nonlegume *Trema tomentosa* are described. The promoters of these genes were linked to a β -glucuronidase reporter
25 gene and introduced into both the nonlegume *Nicotiana tabacum* and the legume *Lotus corniculatus*, and in both instances root-specific promoter activity was preserved. Leach and Aoyagi (1991) describe their analysis of the promoters of the
30

highly expressed rolC and rolD root-inducing genes of *Agrobacterium rhizogenes* (see *Plant Science* (Limerick) 79(1):69-76). They concluded that enhancer and tissue-preferred DNA determinants are dissociated in those promoters. Teeri *et al.* (1989) used gene fusion to lacZ to show that the *Agrobacterium* T-DNA gene encoding octopine synthase is especially active in the epidermis of the root tip and that the TR2' gene is root specific in the intact plant and stimulated by wounding in leaf tissue, an especially desirable combination of characteristics for use with an insecticidal or larvicidal gene (see *EMBO J.* 8(2):343-350). The TR1' gene, fused to nptII (neomycin phosphotransferase II) showed similar characteristics. Additional root-preferred promoters include the VfENOD-GRP3 gene promoter (Kuster *et al.* (1995) *Plant Mol. Biol.* 29(4):759-772); and rolB promoter (Capana *et al.* (1994) *Plant Mol. Biol.* 25(4):681-691. See also U.S. Pat. Nos. 5,837,876; 5,750,386; 5,633,363; 5,459,252; 5,401,836; 5,110,732; and 5,023,179. The phaseolin gene (Murai *et al.* (1983) *Science* 23:476-482 and Sengopta-Gopalen *et al.* (1988) *PNAS* 82:3320-3324.

The expression cassette containing the polynucleotides provided herein can also comprise a selectable marker gene for the selection of transformed cells. Selectable marker genes are utilized for the selection of transformed cells or tissues. Marker genes include genes encoding antibiotic resistance, such as those encoding neomycin phosphotransferase II (NEO) and hygromycin phosphotransferase (HPT), as well as genes conferring resistance to herbicidal compounds, such as glufosinate ammonium, bromoxynil, imidazolinones, and 2,4-dichlorophenoxyacetate (2,4-D) and sulfonylureas. Additional selectable markers include phenotypic markers such as beta-galactosidase and fluorescent proteins such as green fluorescent protein (GFP) (Su *et al.* (2004) *Biotechnol. Bioeng.* 85:610-9 and Fetter *et al.* (2004) *Plant Cell* 16:215-28), cyan fluorescent protein (CYP) (Bolte *et al.* (2004) *J. Cell Science* 117:943-54 and Kato *et al.* (2002) *Plant Physiol.* 129:913-42), and yellow fluorescent protein (PhiYFP.TM. from Evrogen; see, Bolte *et al.* (2004) *J. Cell Science* 117:943-54). Such disclosures are herein incorporated by reference. The above list of selectable marker genes is not meant to be limiting. Any selectable marker gene can be used in the compositions presented herein.

Where appropriate, the sequences employed in the methods and compositions (i.e., the polynucleotide of interest, the recombinase, the endonuclease, etc.) may be optimized for increased expression in the transformed plant. That is, the genes can be synthesized using plant-preferred codons for improved expression. See, for example, Campbell and Gowri (1990) *Plant Physiol.* 92:1-11 for a discussion of host-preferred codon usage. Methods are available in the art for synthesizing plant-preferred genes. See, for example, U.S. Patent Nos. 5,380,831, and 5,436,391, and Murray *et al.* (1989) *Nucleic Acids Res.* 17:477-498, herein incorporated by reference.

Fragments and variants of the various components of the DSB-inducing-agent system, such as for example the guide polynucleotide/Cas endonuclease system and the site-specific integration system (transgenic SSI target site, a donor DNA, a transfer cassette, various site-specific recombination sites, site-specific recombinases, polynucleotides of interest or any active variants or fragments thereof) are also encompassed herein. By "fragment" is intended a portion of the polynucleotide or a portion of the amino acid sequence and hence protein encoded thereby. Fragments of a polynucleotide may encode protein fragments that retain the biological activity of the native protein (i.e., a fragment of a recombinase implements a recombination event). As used herein, a "native" polynucleotide or polypeptide comprises a naturally occurring nucleotide sequence or amino acid sequence, respectively. Thus, fragments of a polynucleotide may range from at least about 20 nucleotides, about 50 nucleotides, about 100 nucleotides, and up to the full-length polynucleotide. A fragment of a polynucleotide that encodes a biologically active portion of a protein employed in the methods or compositions will encode at least 15, 25, 30, 50, 100, 150, 200, or 250 contiguous amino acids, or up to the total number of amino acids present in a full-length protein. Alternatively, fragments of a polynucleotide that are useful as a hybridization probe generally do not encode fragment proteins retaining biological activity. Thus, fragments of a nucleotide sequence may range from at least about 10, 20, 30, 40, 50, 60, 70, 80 nucleotides or up to the full length sequence.

A biologically active portion of a polypeptide can be prepared by isolating a portion of one of the polynucleotides encoding the portion of the polypeptide of

interest and expressing the encoded portion of the protein (e.g., by recombinant expression *in vitro*), and assessing the activity of the portion of the polypeptide. For example, polynucleotides that encode fragments of a recombinase polypeptide can comprise nucleotide sequence comprising at least 16, 20, 50, 75, 100, 150, 200, 250, 300, 350, 400, 450, 500, 550, 600, 650, 700, 800, 900, 1,000, 1,100, 1,200, 1,300, or 1,400 nucleotides, or up to the number of nucleotides present in a nucleotide sequence employed in the methods and compositions provided herein.

"Variants" is intended to mean substantially similar sequences. For polynucleotides, a variant comprises a polynucleotide having deletions (i.e., truncations) at the 5' and/or 3' end; deletion and/or addition of one or more nucleotides at one or more internal sites in the native polynucleotide; and/or substitution of one or more nucleotides at one or more sites in the native polynucleotide. For polynucleotides, conservative variants include those sequences that, because of the degeneracy of the genetic code, encode the amino acid sequence of one of the polypeptides employed in the compositions and methods provided herein. Naturally occurring allelic variants such as these, or naturally occurring allelic variants of polynucleotides can be identified with the use of well-known molecular biology techniques, as, for example, with polymerase chain reaction (PCR) and hybridization techniques as outlined below. Variant polynucleotides also include synthetically derived polynucleotides, such as those generated, for example, by using site-directed mutagenesis. Generally, variants of a particular polynucleotide employed in the methods and compositions provided herein will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to that particular polynucleotide as determined by sequence alignment programs and parameters as described elsewhere herein.

Variants of a particular polynucleotide employed in the methods and compositions provided herein (trait gene(s) of interest and/or pollen- inhibitor gene(s), recombinases, nucleases) can also be evaluated by comparison of the percent sequence identity between the polypeptide encoded by a variant polynucleotide and the polypeptide encoded by the reference polynucleotide. Thus, for example, an isolated polynucleotide that encodes a polypeptide with a given

percent sequence identity to the polypeptide are disclosed. Percent sequence identity between any two polypeptides can be calculated using sequence alignment programs and parameters described elsewhere herein. Where any given pair of polynucleotides provided herein is evaluated by comparison of the percent
5 sequence identity shared by the two polypeptides they encode, the percent sequence identity between the two encoded polypeptides is at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity.

"Variant" protein is intended to mean a protein derived from the native protein
10 by deletion (so-called truncation) of one or more amino acids at the N-terminal and/or C-terminal end of the native protein; deletion and/or addition of one or more amino acids at one or more internal sites in the native protein; or substitution of one or more amino acids at one or more sites in the native protein. Variant proteins employed in the methods and compositions provided herein are biologically active,
15 that is they continue to possess the desired biological activity of the native protein. Such variants may result from, for example, genetic polymorphism or from human manipulation. Biologically active variants of a native protein provided herein will have at least about 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or more sequence identity to the
20 amino acid sequence for the native protein as determined by sequence alignment programs and parameters described elsewhere herein. A biologically active variant of a protein provided herein may differ from that protein by as few as 1-15 amino acid residues, as few as 1-10, such as 6-10, as few as 5, as few as 4, 3, 2, or even 1 amino acid residue.

25 Proteins may be altered in various ways including amino acid substitutions, deletions, truncations, and insertions. Methods for such manipulations are generally known in the art. For example, amino acid sequence variants of the recombinase proteins can be prepared by mutations in the DNA. Methods for mutagenesis and nucleotide sequence alterations are well known in the art. See, for example, Kunkel
30 (1985) *Proc. Natl. Acad. Sci. USA* 82:488-492; Kunkel *et al.* (1987) *Methods in Enzymol.* 154:367-382; U.S. Patent No. 4,873,192; Walker and Gaastra, eds. (1983) *Techniques in Molecular Biology* (MacMillan Publishing Company, New York) and

the references cited therein. Guidance as to appropriate amino acid substitutions that do not affect biological activity of the protein of interest may be found in the model of Dayhoff *et al.* (1978) *Atlas of Protein Sequence and Structure* (Natl. Biomed. Res. Found., Washington, D.C.), herein incorporated by reference.

- 5 Conservative substitutions, such as exchanging one amino acid with another having similar properties, may be preferable.

Thus, the polynucleotides used herein can include the naturally occurring sequences, the "native" sequences, as well as mutant forms. Likewise, the proteins used in the methods provided herein encompass both naturally occurring proteins
10 as well as variations and modified forms thereof. Obviously, the mutations that will be made in the polynucleotide encoding the variant polypeptide must not place the sequence out of reading frame and preferably will not create complementary regions that could produce secondary mRNA structure. See, EP Patent Application Publication No. 75,444.

- 15 The deletions, insertions, and substitutions of the protein sequences encompassed herein are not expected to produce radical changes in the characteristics of the protein. However, when it is difficult to predict the exact effect of the substitution, deletion, or insertion in advance of doing so, one skilled in the art will appreciate that the effect will be evaluated by routine screening assays.

- 20 Variant polynucleotides and proteins also encompass sequences and proteins derived from a mutagenic and recombinogenic procedure such as DNA shuffling. With such a procedure, for example, one or more different recombinase coding sequences can be manipulated to create a new recombinase protein possessing the desired properties. In this manner, libraries of recombinant
25 polynucleotides are generated from a population of related sequence polynucleotides comprising sequence regions that have substantial sequence identity and can be homologously recombined *in vitro* or *in vivo*. Strategies for such DNA shuffling are known in the art. See, for example, Stemmer (1994) *Proc. Natl. Acad. Sci. USA* 91:10747-10751; Stemmer (1994) *Nature* 370:389-391; Crameri *et al.* (1997) *Nature Biotech.* 15:436-438; Moore *et al.* (1997) *J. Mol. Biol.* 272:336-347; Zhang *et al.* (1997) *Proc. Natl. Acad. Sci. USA* 94:4504-4509; Crameri *et al.*
30 (1998) *Nature* 391:288-291; and U.S. Patent Nos. 5,605,793 and 5,837,458.

The following terms are used to describe the sequence relationships between two or more nucleic acids or polynucleotides. As used herein, "reference sequence" is a defined sequence used as a basis for sequence comparison. A reference sequence may be a subset or the entirety of a specified sequence; for example, as
5 a segment of a full-length cDNA or gene sequence, or the complete cDNA or gene sequence. As used herein, "comparison window" makes reference to a contiguous and specified segment of a polynucleotide sequence, wherein the polynucleotide sequence in the comparison window may comprise additions or deletions (i.e., gaps) compared to the reference sequence (which does not comprise additions or
10 deletions) for optimal alignment of the two sequences. Generally, the comparison window is at least 20 contiguous nucleotides in length, and optionally can be 30, 40, 50, 100, or longer. Those of skill in the art understand that to avoid a high similarity to a reference sequence due to inclusion of gaps in the polynucleotide sequence a gap penalty is typically introduced and is subtracted from the number of matches.

15 Sequence relationships can be analyzed and described using computer-implemented algorithms. The sequence relationship between two or more polynucleotides, or two or more polypeptides can be determined by determining the best alignment of the sequences, and scoring the matches and the gaps in the alignment, which yields the percent sequence identity, and the percent sequence
20 similarity. Polynucleotide relationships can also be described based on a comparison of the polypeptides each encodes. Many programs and algorithms for the comparison and analysis of sequences are well-known in the art.

"Sequence identity" or "identity" in the context of nucleic acid or polypeptide sequences refers to the nucleic acid bases or amino acid residues in two sequences
25 that are the same when aligned for maximum correspondence over a specified comparison window.

The term "percentage of sequence identity" refers to the value determined by comparing two optimally aligned sequences over a comparison window, wherein the portion of the polynucleotide or polypeptide sequence in the comparison window
30 may comprise additions or deletions (i.e., gaps) as compared to the reference sequence (which does not comprise additions or deletions) for optimal alignment of the two sequences. The percentage is calculated by determining the number of

positions at which the identical nucleic acid base or amino acid residue occurs in both sequences to yield the number of matched positions, dividing the number of matched positions by the total number of positions in the window of comparison and multiplying the results by 100 to yield the percentage of sequence identity. Useful
5 examples of percent sequence identities include, but are not limited to, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95%, or any integer percentage from 50% to 100%. These identities can be determined using any of the programs described herein.

Sequence alignments and percent identity or similarity calculations may be
10 determined using a variety of comparison methods designed to detect homologous sequences including, but not limited to, the MegAlign™ program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Within the context of this application it will be understood that where sequence analysis software is used for analysis, that the results of the analysis will be based on the “default values” of
15 the program referenced, unless otherwise specified. As used herein “default values” will mean any set of values or parameters that originally load with the software when first initialized.

The “Clustal V method of alignment” corresponds to the alignment method labeled Clustal V (described by Higgins and Sharp, (1989) *CABIOS* 5:151-153;
20 Higgins *et al.*, (1992) *Comput Appl Biosci* 8:189-191) and found in the MegAlign™ program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). For multiple alignments, the default values correspond to GAP PENALTY=10 and GAP LENGTH PENALTY=10. Default parameters for pairwise alignments and calculation of percent identity of protein sequences using the Clustal
25 method are KTUPLE=1, GAP PENALTY=3, WINDOW=5 and DIAGONALS SAVED=5. For nucleic acids these parameters are KTUPLE=2, GAP PENALTY=5, WINDOW=4 and DIAGONALS SAVED=4. After alignment of the sequences using the Clustal V program, it is possible to obtain a “percent identity” by viewing the “sequence distances” table in the same program.

30 The “Clustal W method of alignment” corresponds to the alignment method labeled Clustal W (described by Higgins and Sharp, (1989) *CABIOS* 5:151-153; Higgins *et al.*, (1992) *Comput Appl Biosci* 8:189-191) and found in the MegAlign™

v6.1 program of the LASERGENE bioinformatics computing suite (DNASTAR Inc., Madison, WI). Default parameters for multiple alignment (GAP PENALTY=10, GAP LENGTH PENALTY=0.2, Delay Divergen Seqs (%)=30, DNA Transition Weight=0.5, Protein Weight Matrix=Gonnet Series, DNA Weight Matrix=IUB). After alignment of
5 the sequences using the Clustal W program, it is possible to obtain a "percent identity" by viewing the "sequence distances" table in the same program.

"BLAST" is a searching algorithm provided by the National Center for Biotechnology Information (NCBI) used to find regions of similarity between biological sequences. The program compares nucleotide or protein sequences to
10 sequence databases and calculates the statistical significance of matches to identify sequences having sufficient similarity to a query sequence such that the similarity would not be predicted to have occurred randomly. BLAST reports the identified sequences and their local alignment to the query sequence.

It is well understood by one skilled in the art that many levels of sequence
15 identity are useful in identifying polypeptides from other species or modified naturally or synthetically wherein such polypeptides have the same or similar function or activity. Useful examples of percent identities include, but are not limited to, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90% or 95%, or any integer percentage from 50% to 100%. Indeed, any integer amino acid identity from 50% to
20 100% may be useful in describing the present disclosure, such as 51%, 52%, 53%, 54%, 55%, 56%, 57%, 58%, 59%, 60%, 61%, 62%, 63%, 64%, 65%, 66%, 67%, 68%, 69%, 70%, 71%, 72%, 73%, 74%, 75%, 76%, 77%, 78%, 79%, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98% or 99%.

25 Sequence identity/similarity values can also be obtained using GAP Version 10 (GCG, Accelrys, San Diego, CA) using the following parameters: % identity and % similarity for a nucleotide sequence using GAP Weight of 50 and Length Weight of 3, and the nwsgapdna.cmp scoring matrix; % identity and % similarity for an amino acid sequence using GAP Weight of 8 and Length Weight of 2, and the
30 BLOSUM62 scoring matrix (Henikoff and Henikoff (1989) *Proc. Natl. Acad. Sci USA* 89:10915); or any equivalent program thereof. By "equivalent program" is intended any sequence comparison program that, for any two sequences in question,

generates an alignment having identical nucleotide or amino acid residue matches and an identical percent sequence identity when compared to the corresponding alignment generated by GAP Version 10

GAP uses the algorithm of Needleman and Wunsch (1970) *J. Mol. Biol.* 48:443-453, to find the alignment of two complete sequences that maximizes the number of matches and minimizes the number of gaps. GAP considers all possible alignments and gap positions and creates the alignment with the largest number of matched bases and the fewest gaps. It allows for the provision of a gap creation penalty and a gap extension penalty in units of matched bases. GAP must make a profit of gap creation penalty number of matches for each gap it inserts. If a gap extension penalty greater than zero is chosen, GAP must, in addition, make a profit for each gap inserted of the length of the gap times the gap extension penalty. Default gap creation penalty values and gap extension penalty values in Version 10 of the GCG Wisconsin Genetics Software Package for protein sequences are 8 and 2, respectively. For nucleotide sequences the default gap creation penalty is 50 while the default gap extension penalty is 3. The gap creation and gap extension penalties can be expressed as an integer selected from the group of integers consisting of from 0 to 200. Thus, for example, the gap creation and gap extension penalties can be 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, 10, 15, 20, 25, 30, 35, 40, 45, 50, 55, 60, 65 or greater.

GAP presents one member of the family of best alignments. There may be many members of this family, but no other member has a better quality. GAP displays four figures of merit for alignments: Quality, Ratio, Identity, and Similarity. The Quality is the metric maximized in order to align the sequences. Ratio is the quality divided by the number of bases in the shorter segment. Percent Identity is the percent of the symbols that actually match. Percent Similarity is the percent of the symbols that are similar. Symbols that are across from gaps are ignored. A similarity is scored when the scoring matrix value for a pair of symbols is greater than or equal to 0.50, the similarity threshold.

As used herein, "breeding" is the genetic manipulation of living organisms. Plants are bred through techniques that take advantage of the plant's method of pollination. A plant is self-pollinated if pollen from one flower is transferred to the

same or another flower of the same plant. A plant is sib-pollinated when individuals within the same family or line are used for pollination. A plant is cross-pollinated if the pollen comes from a flower on a different plant from a different family or line. In a breeding application, a breeder initially selects and crosses two or more parental plants. As used herein, "crossing" can refer to a simple X by Y cross, or the process of backcrossing, depending on the context.

The introduction of transgenes (transgenic trait of interest) into many major crops is typically performed in a single plant variety that is most amenable to the transformation, tissue culture and regeneration processes. For example, in corn, the readily transformable genotype referred to Hi-II (Armstrong and Green, 1985, 1985, *Planta* 164(2): 207-214) has been used for many years across the industry for genetic transformation. However, the agronomic characteristics of this genotype are not commercially desirable, so once a transgenic trait locus (genomic locus where transgene has integrated into the plant genome) has been created, the transgenic trait locus must be introgressed from Hi-II into commercially relevant inbreds.

Introgression, also known as introgressive hybridization, refers to The movement of a gene (gene flow) from one species into the gene pool of another, which can be accomplished by repeated backcrossing of an interspecific hybrid with one of its parent species.

A transgenic inbred (A) can be crossed to a non-transgenic inbred (B) in order to introgress (or transfer) the transgenic trait into the new germplasm by repeated back-crossing to the recurrent parent (B). The result of the first cross between inbred A and inbred B is the F1 hybrid. The F1 hybrid can then be used as the pollen donor to again cross with the recurrent parent (inbred B) to produce the first backcross generation (the BC1 generation). Successive backcrosses can be performed until the majority of genetic material from the original inbred A has been lost through meiotic recombination and segregation, leaving the transgenic locus in the new inbred background.

Commercial corn seed companies typically work with hundreds of inbreds in different heterotic groups, and thus the transgenic trait locus must be introgressed into numerous inbreds for further efficacy testing – a process that can take up to many successive generations to completely eliminate as much of the undesired

germplasm (such as for example the Hi-II germplasm in maize) as possible. A major impediment in the introgression process is identifying progeny in successive crosses in which meiotic recombination has occurred in close proximity on either side of the transgenic locus, replacing as much of the flanking undesired germplasm chromosome with the new inbred chromosome. To identify progeny where closely-spaced (such as but not limited to less than 0-1 cM, 0-2 cM, 0-3 cM, 0-4 cM, 0-5 cM, 0-6 cM, 0-7 cM, 0-8 cM, 0-9 cM, 0-10 cM, 1-1 cM, 1-2 cM, 1-3 cM, 1-4 cM, 1-5 cM, 1-6 cM, 1-7 cM, 1-8 cM, 1-9 cM, 1-10 cM) meiotic recombination on either side of a transgenic locus has occurred a molecular screening using genetic markers can be used thereby adding more work and cost to each generation of screening. In addition, for certain crop species which naturally self-pollinate such as soybean, making crosses through emasculation and hand-pollination on the scale needed in order to identify low-frequency recombination events is extremely labor intensive.

In order to accelerate the introgression process (also referred to as “Accelerated Trait Introgression” and greatly reduce the labor involved, a (screening) method that selects for recombination in the gametes would be of great benefit. One method of accomplishing such as screen is to position a gamete-specific inhibitor gene (or gamete inhibitor gene) in close proximity to the transgenic locus of interest. One can also position at least one color marker gene in close proximity of a trait locus of interest, in combination with or without a pollen-inhibitor gene as described herein. Described herein are compositions and methods to position a pollen-inhibitor gene and/ or a color marker in close proximity to a trait locus of interest in the genome of a progeny plant. As described herein the trait locus can be flanked by a pollen-inhibitor gene on each side, a color maker gene on each side, or a combination of a color marker gene and a pollen-inhibitor gene. The proximity of the pollen-inhibitor gene or color marker gene to the trait locus of interest can be about at least 0.05 cM, 0.1 cM, 0.2 cM, 0.3 cM, 0.4 cM, 0.5 cM, 0.6 cM, 0.7 cM, 0.8 cM, 0.9 cM, 1.0 cM, 1.1 cM, 1.2 cM, 1.3 cM, 1.4 cM, 1.5 cM, 1.6 cM, 1.7 cM, 1.8 cM, 1.9 cM, 2.0 cM, 2.1 cM, 2.2 cM, 2.3 cM, 2.4 cM, 2.5 cM, 2.6 cM, 2.7 cM, 2.8 cM, 2.9 cM, 3.0 cM, 3.1 cM, 3.2 cM, 3.3 cM, 3.4 cM, 3.5 cM, 3.6 cM, 3.7 cM, 3.8 cM, 3.9 cM , 4.0 cM or 5.0 cM

Compositions and methods are provided herein for the use of pollen-inhibitor genes and/or color maker genes in accelerated trait introgression.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant, the method comprising: (a) providing a first
5 plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one pollen-inhibitor gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere; (b) cross
10 pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said trait of interest and said pollen-inhibitor gene; and, (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.

15 In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant, the method comprising: (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one color marker gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said color marker
20 gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere; (b) cross pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said trait of interest site and said color marker gene; and, (d) optionally, backcrossing the progeny plant of (c) as the pollen
25 donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.

In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest, a first pollen-inhibitor gene,
30 and a second pollen-inhibitor gene wherein said genomic window is about 5 cM in length, and wherein said trait of interest is flanked by said first and second pollen-inhibitor gene; (b) cross-pollinating the first plant of (a) with pollen from a second

plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

5 In one embodiment, the method comprises a method of accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest, a pollen-inhibitor gene and a color marker gene, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;(b)
10 cross-pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

15 In one embodiment, the method comprises a method accelerated trait introgression in the genome of a plant comprising: (a) providing a first plant having within a genomic window at least one trait of interest and at least a first color marker gene integrated into a first target site, a second color marker gene integrated into a second target site for, wherein said genomic window is about 5 cM in length, and
20 wherein trait of interest is flanked by said first and second pollen-inhibitor gene;(b) cross-pollinating the first plant of (a) with pollen from a second plant; and, (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and, (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and
25 selecting progeny plants comprising the trait of interest.

The location of the trait locus can determine how many pollen-inhibitor loci (or color marker loci) are introduced and where the pollen-inhibitor or color marker loci are positioned. If the trait locus is in close proximity (such as but not limited to less than 0-1 cM, 0-2 cM, 0-3 cM, 0-4 cM, 0-5 cM, 0-6 cM, 0-7 cM, 0-8 cM, 0-9 cM, 0-10
30 cM, 1-1 cM, 1-2 cM, 1-3 cM, 1-4 cM, 1-5 cM, 1-6 cM, 1-7 cM, 1-8 cM, 1-9 cM, 1-10 cM) to a telomere (near the end of the chromosome), only a single pollen-inhibitor locus, located just proximal to the trait locus of interest (i.e. 0.5 to 1.0 cM closer to

the centromere) is required. As described in Example 1 and illustrated in Figure 1, expression cassettes comprising a pollen-inhibitor gene (such as, but not limited to, barnase or alpha amylase) can be introduced just proximal to a trait locus near the telomere. By propagating the pollen-inhibitor locus through the female during sexual crosses, linkage between the trait and the AA1-inhibitor can be established. These plants (labeled "A") containing the linked loci can now be crossed to any inbred "B" to create an F1 hybrid. F1 seed are planted and the F1 plants are now used as the pollen source back to the recurrent inbred parent "B". After taking advantage of the herbicide resistance expression cassette within the trait locus to select against any progeny that were derived from wild-type pollen, only progeny in which the linkage has been broken during meiosis will be viable and all viable progeny will contain just the telomeric segment with the trait locus from the original inbred (A) while the majority of the chromosome is all from parent B. In this manner, selection for accelerated introgression of a trait locus from inbred A into the new chromosome from inbred B has been successfully accomplished.

For trait loci in internal (for example, in non-telomeric locations (more toward the centromere of the chromosome), two flanking pollen-inhibitor loci located one on either side of the trait locus can be used (as described in Example 2). As described in Example 2 and illustrated in Figure 2, when the transgenic plants carrying the pollen-inhibitor gene are produced, these two loci must be linked through conventional crossing to the trait locus requiring that at least one of the pollen-inhibitor loci must be able to produce pollen in order to make the 3-way linkage. During this time, either expression of the inhibitor gene must be off or the inhibitor must be inactivated by another protein (a maintainer). After the triple-linked inbred line is crossed to the 100 inbreds of interest (producing 100 F1 populations) the pollen-inhibitor must now be on. Various technical approaches can be used to accomplish this, including the use of a bacterial inhibitor/anti-inhibitor combination (for example barnase/barstar), using an inducible inhibitor, use of a repressor protein (with no chemical induction), use of inteins to switch on the pollen screen, and use of a double haploid step with all of the above strategies. Inhibitor/anti inhibitor systems useful for gamete inhibition include, but are not limited to barnase/barstar, kid/kis and numerous other well-characterized bacterial inhibitors

(Finbarr Hayes, 2003, Science 301:1496-1499; Yamaguchi et al., 2011, Ann. Rev. Gent. 45:61-79). Conditionally-lethal genes such as the codA gene can also be used.

Another alternative for setting up three linked loci without the use of a maintainer, while also obviating the need for conventional breeding methods to establish the linkage, is to use CRISPR-mediated introduction of the pollen-inhibitors directly into the flanking sites in a chromosome that already contains the trait locus. For pollen inhibition, the alpha-amylase gene can be particularly useful, since the breakdown of starch by the expressed protein renders the pollen incapable of forming a pollen tube.

Methods are provided for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant using breeding techniques. For example, a first plant having within a genomic window at least one trait locus of interest integrated into a first target site located proximal to a telomere, wherein said genomic window is about 10 cM in length and located within 10 cM of the telomere, wherein said first plant does not comprise a pollen-inhibitor gene; can be crossed with a second plant, wherein said second plant comprises in said genomic window a pollen-inhibitor gene integrated into a second target site located distal to the telomere, wherein said second plant does not comprise said first target site. A progeny plant is then selected comprising said trait locus of interest and said pollen-inhibitor gene, wherein said trait locus of interest and said pollen-inhibitor gene are genetically linked. Selecting a progeny plant comprising both the trait locus of interest and the pollen-inhibitor gene can be done through various methods. For example, a phenotypic analysis can be performed whereby the activity of the trait of interest or said pollen-inhibitor is detected in the progeny plant. Alternative methods that assay for the presence of said trait locus of interest and said pollen-inhibitor gene which are specific to the said trait locus of interest and said pollen-inhibitor gene include techniques such as PCR, hybridization, Isozyme electrophoresis, Restriction Fragment Length Polymorphisms (RFLPs), Randomly Amplified Polymorphic DNAs (RAPDs), Arbitrarily Primed PCR (AP-PCR), DNA Amplification Fingerprinting (DAF), Sequence Characterized Amplified Regions

(SCARs), Amplified Fragment length Polymorphisms (AFLPs), Simple Sequence Repeats (SSRs), and Single Nucleotide Polymorphisms (SNPs).

The meaning of abbreviations is as follows: "sec" means second(s), "min" means minute(s), "h" means hour(s), "d" means day(s), "μL" means microliter(s),
5 "mL" means milliliter(s), "L" means liter(s), "μM" means micromolar, "mM" means millimolar, "M" means molar, "mmol" means millimole(s), "μmole" means micromole(s), "g" means gram(s), "μg" means microgram(s), "ng" means nanogram(s), "U" means unit(s), "bp" means base pair(s) and "kb" means kilobase(s).

10 Non-limiting examples of compositions and methods disclosed herein are as follows:

1. A method for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:

15 (a) providing a first plant having within a genomic window at least one trait gene of interest integrated into a first target site located proximal to a telomere, wherein said genomic window is about 10 cM in length, wherein said first plant does not comprise a pollen-inhibitor gene;

(b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a pollen-inhibitor gene integrated into
20 a second target site located proximal to both the telomere and the trait gene of interest of (a); and,

(c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor gene are genetically linked.

25 2. A method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:

(a) providing a first plant having within a genomic window a first pollen-inhibitor gene integrated into a first target site, wherein said genomic window is about 10 cM in length;

30 (b) breeding to said first plant a second plant having a trait gene of interest integrated into a second target site within said genomic window;

- (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene and said trait gene of interest in said genomic window;
- (d) providing a third plant having a second pollen-inhibitor gene integrated into a third target site within said genomic window;
- 5 (e) breeding to said third plant a fourth plant, wherein said fourth plant comprises a pollen-inhibitor maintainer (PIM) gene;
- (f) selecting a progeny plant from step (e) comprising said second pollen-inhibitor gene and said pollen-inhibitor maintainer (PIM) gene; and,
- (g) cross pollinating the progeny plant of (c) with the progeny plant of (f) and
- 10 selecting for a progeny plant that comprises said first pollen-inhibitor gene said trait gene of interest and said second pollen-inhibitor gene, wherein said first pollen-inhibitor gene, said trait gene of interest and said second pollen-inhibitor gene are genetically linked.
3. The method of embodiments 1-2, wherein the first target site and the second target site is selected from the group consisting of a recombinase target site, a
- 15 transgenic SSI target site, a single-strand-break-inducing-agent target site, and a double-strand-break-inducing-agent target site, or any one combination thereof.
4. The method of embodiments 1-2, wherein the double-strand-break-inducing-agent target site is a target site for an agent selected from the group of a Cas9 endonuclease, a zinc-finger nuclease, a Tal Effector nuclease (TALEN), a
- 20 meganuclease, and an engineered endonuclease.
5. A method for introducing a pollen-inhibitor gene and a color marker gene in close proximity to a trait locus of interest in the genome of a plant, said method comprising:
- 25 (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is about 10 cM in length;
- (b) introducing into said genomic window of the plant of (a) a color marker gene;
- (c) breeding to the plant of (b) a second plant, wherein said second plant is a
- 30 haploid inducer line capable of producing haploid embryos;
- (d) selecting haploid embryos from the plant of (c) and introducing into said haploid embryos, a pollen-inhibitor gene; and,

- (e) producing a double haploid plant from the haploid embryo of (d).
6. A method for introducing two color marker genes in close proximity to a trait locus of interest in the genome of a plant, said method comprising:
- 5 (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is about 10 cM in length;
- (b) introducing into said genomic window of the plant of (a) a color marker gene;
- (c) breeding to the plant of (b) a second plant, wherein said second plant is a haploid inducer line capable of producing haploid embryos;
- 10 (d) selecting haploid embryos from the plant of (c) and introducing into said haploid embryos, a second color marker gene; and,
- (e) producing a double haploid plant from the haploid embryo of (d).
7. The method of embodiments 5-6, wherein color marker or the pollen-inhibitor gene are introduced into a target site of a double-strand-break-inducing-agent target site.
- 15 8. A method of accelerated trait introgression in the genome of a plant, the method comprising:
- (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one pollen-inhibitor gene
- 20 located proximal to both the telomere and the trait of interest, wherein said trait of interest and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;
- (b) cross pollinating the first plant of (a) with pollen from a second plant; and,
- 25 (c) selecting a progeny plant from step (b) comprising said trait of interest and said pollen-inhibitor gene; and,
- (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.
- 30 9. A method of accelerated trait introgression in the genome of a plant, the method comprising:

- (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one color marker gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said color marker gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;
- (b) cross pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said trait of interest site and said color marker gene; and ,
- (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.
10. A method of accelerated trait introgression in the genome of a plant comprising:
- (a) providing a first plant having within a genomic window at least one trait of interest , a first pollen-inhibitor gene, and a second pollen-inhibitor gene wherein said genomic window is about 5 cM in length, and wherein said trait of interest is flanked by said first and second pollen-inhibitor gene;
- (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,
- (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.
11. A method of accelerated trait introgression in the genome of a plant comprising:
- (a) providing a first plant having within a genomic window at least one trait of interest, a pollen-inhibitor gene and a color marker gene, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;
- (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,
- (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest..

12. A method of accelerated trait introgression in the genome of a plant comprising:
- (a) providing a first plant having within a genomic window at least one trait of interest and at least a first color marker gene integrated into a first target site, a second color marker gene integrated into a second target site for, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;
 - (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
 - (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,
 - (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest..
13. A method for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
- (a) providing a first plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said genomic window is about 5 cM in length and located within 0.1 cM to 100 cM of the telomere, and wherein said first plant does not comprise a pollen-inhibitor gene;
 - (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said second plant does not comprise said first transgenic target site; and,
 - (c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor gene are genetically linked in said genomic window.
14. A method of accelerated trait introgression in the genome of a plant, the method comprising:

- (a) providing a first progeny plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere and a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest,, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said first transgenic target site and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;
- (b) cross pollinating the first plant of (a) with pollen from a second plant (elite inbred); and,
- (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said pollen-inhibitor gene;
15. The method of embodiment 8, further comprising :
- (d) backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait gene of interest.
16. The method of any one of embodiments 13-14, wherein the pollen-inhibitor gene is selected from the group consisting of barnase, alpha amylase and KID.
17. The method of anyone of embodiments 13-14 wherein the trait gene of interest is selected from the group consisting of a selectable marker, a modified or edited native gene, a transgene, an altered double-strand-break target site, a native gene, or a transgenic SSI target site.
18. The method of anyone of embodiments 13-14, wherein the trait gene of interest and the pollen-inhibitor gene are genetically linked and located within 0.5 cM to 1 cM from each other.
19. A method for introducing two pollen- inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
- (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said genomic window is about 5 cM in length;

- (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site, wherein said second transgenic SSI target site comprises at least one trait gene of interest wherein said second plant does not comprise said first transgenic target site;
- (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said second transgenic target site genetically linked in said genomic window;
- (d) providing a third plant having within said genomic window at least a third transgenic SSI target site and a pollen-inhibitor maintainer (PIM), wherein said third transgenic SSI target site comprises a second pollen-inhibitor gene; and,
- (e) using the third plant of step (d) to pollinate the plant of step (c) and selecting a progeny plant wherein said first transgenic SSI target site, said second transgenic SSI target site, and said third transgenic SSI target site are genetically linked to each other.
20. A method of accelerated trait introgression in the genome of a plant comprising :
- (a) providing a first plant having a pollen-inhibitor maintainer gene and a genomic window, wherein said genomic window is about 5 cM in length and wherein said genomic window comprises at least a first transgenic SSI target site, a second transgenic SSI target site and a third transgenic SSI target site, wherein said first transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said second transgenic SSI target site comprises at least one trait gene of interest, wherein said third transgenic SSI target site comprises a second pollen-inhibitor gene, wherein said second transgenic target site is flanked by said first transgenic SSI target site and said third transgenic SSI target site;
- (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said first transgenic target site, said second transgenic target site and said third transgenic target site.

21. The method of embodiment 20, further comprising cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
22. A method of accelerated trait introgression in the genome of a plant comprising:
- 5 (a) providing a first plant having within a genomic window a trait of interest and at least a first pollen-inhibitor gene integrated into a first target site for a Cas9 endonuclease, a second pollen-inhibitor gene integrated into a second target site for a Cas9 endonuclease, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;
- 10 (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene;
23. The method of embodiment 22, further comprising cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
- 15 24. A method for introducing two pollen- inhibitor genes in close proximity to a trait gene of interest in the genome of a progeny plant, said method comprising:
- (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said genomic window is about 5 cM in length;
- 20 (b) introducing into the plant of (a) a first pollen-inhibitor gene, wherein said first pollen-inhibitor gene is genetically linked to said trait gene of interest
- 25 (c) selecting a progeny plant from (b) comprising the pollen-inhibitor gene genetically linked to the trait gene of interest.
- (d) breeding to said progeny plant of (c) a haploid-inducer plant and producing haploid immature embryos .
- 30 (e) introducing into the haploid immature embryos of (d) a second pollen-inhibitor gene linked to said trait gene of interest and said first pollen-inhibitor gene, wherein the trait gene of interest is flanked by the first and second pollen-inhibitor gene; and,

- (f) producing a homozygous plant from the haploid immature embryo of (e);
25. A method of accelerated trait introgression in the genome of a plant comprising
- (a) providing a first plant having a genomic window comprising a trait gene of interest flanked by and genetically linked to a first pollen-inhibitor gene and a second pollen-inhibitor gene, wherein said genomic window is about 5 cM in length;
- (b) cross pollinating the first plant of (a) with pollen from a second plant ;and,
- (c) selecting a progeny plant from step (b) comprising said trait gene of interest flanked by and genetically linked to said first pollen-inhibitor gene and said second pollen-inhibitor gene.
26. The method of embodiment 25 further comprising cross-pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
27. A plant comprising at least one trait gene of interest, a first recombinant DNA construct comprising a first pollen-inhibitor gene, and a second recombinant DNA construct comprising a second pollen-inhibitor gene, wherein said first recombinant DNA construct and said second recombinant DNA construct are genetically linked and flank said trait gene of interest.
28. A plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a pollen-inhibitor gene, wherein said trait locus of interest and said a pollen-inhibitor gene segregate independently from one another at a rate of about 10% to about 0.1%.
29. The plant of embodiment 27, wherein the first pollen-inhibitor gene, the second pollen-inhibitor gene and the trait gene of interest are located within 5 cM of each other.
30. The plant of embodiment 27, wherein the first pollen-inhibitor gene and the second pollen-inhibitor is selected from the group consisting of barnase, alpha amylase, KID, or any combination thereof
31. The plant of embodiment 27, wherein the second pollen-inhibitor gene is selected from the group consisting a non-conditional gene, a conditional gene and an inducible gene.

32. A plant comprising at least one trait gene of interest, a first recombinant DNA construct comprising a color marker gene, and a second recombinant DNA construct comprising a second color marker gene, wherein said first recombinant DNA construct and said second recombinant DNA construct are genetically
5 linked and flank said trait gene of interest.
33. A plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a color marker gene, wherein said trait locus of interest and said color marker gene segregate independently from one another at a rate of about 10% to about 0.1%.
- 10 34. The plant of embodiment 33, wherein the first color marker gene, the second color marker gene and the trait gene of interest are located within 5 cM of each other.

EXAMPLES

In the following Examples, unless otherwise stated, parts and percentages
15 are by weight and degrees are Celsius. It should be understood that these Examples, while indicating embodiments of the disclosure, are given by way of illustration only. From the above discussion and these Examples, one skilled in the art can make various changes and modifications of the disclosure to adapt it to various usages and conditions. Such modifications are also intended to fall within
20 the scope of the appended claims.

EXAMPLE 1

Using a single pollen-inhibitor locus to screen for accelerated trait introgression of a telomeric trait locus.

Illustrated in Figure 1 and described below is an example of how a single
25 pollen- inhibitor gene can be located near a trait of interest, and its use for accelerated trait introgression of a telomeric trait locus of maize.

Figure 1 shows a schematic of a crossing scheme used to establish and perform accelerated introgression for a telomeric trait locus. Step 1: Choose an inbred amenable to transformation, identify accessions that contain two SSI target
30 sites, the first at a position close to the telomere (labeled as "1") and the second SSI site being distal (closer to the centromere, labeled as "2"), the distance between Site 1 and Site 2 being within a close genetic distance, for example 1cM. Step 2: In two

separate transformation experiments, use the SSI transformation method to introduce a trait gene (for example, UBI::GAT:pinII, abbreviated as GAT in the figure, which confers resistance to glyphosate) into Site 1, and a Pollen-inhibitor gene (for example PG47::ZmAA1::pinII, abbreviated as PI in the figure) into Site 2.

- 5 Step 3: Carry pollen from the GAT-containing plant to PI-containing plant and screen the progeny by molecular markers in order to identify plants containing both GAT and PI. Step 4: Carry pollen from the Wild Type inbred plant to the plants containing both GAT and PI to establish the linkage between Site 1 and Site 2. Use molecular markers to screen the progeny in order to identify plants in which GAT
- 10 and PI are genetically linked (1% of the progeny at a genetic distance of 1cM). Step 5: Carry pollen from different Elite inbred (into which you wish to introgress the GAT trait) onto ears of plants containing the linked GAT and PI. Use molecular markers to identify F1 progeny that contain the linked GAT and PI. Step 6: Carry pollen from the F1 progeny containing linked GAT and PI back to the Recurrent Parent (i.e. the
- 15 Elite inbred of step 5) to generate a progeny pool.
 - a. If the linkage between GAT and PI is broken during meiosis then the pollen is viable and GAT can be transmitted to progeny.
 - b. Spray with glyphosate to eliminate any progeny derived from wild-type pollen grains.
 - 20 c. The surviving plants will have GAT gene with a minimal linkage drag from the transformation inbred.

A. Introducing the pollen-inhibitor gene – barnase- in close proximity to a trait locus of interest (GAT, glyphosate resistance trait) in the genome of a maize plant.

- 25 A donor sequence for site specific integration (SSI) (SEQ ID NO: 44) was introduced into a pre-existing transgenic SSI target site containing a first expression cassette comprising a ubiquitin promoter driving a phosphomannose isomerase (PMI) and a pin II terminator, wherein the PMI was preceded by a FRT1 recombination site (UBI PRO::FRT1::PMI::pinII) linked to a second expression
- 30 cassette comprising an actin promoter driving a moPAT selectable marker and a pin II terminator, followed by a FRT87 recombination site (ACTIN PRO:: moPAT::pinII-FRT87) located on chromosome 3 at genetic position 0.9 cM (close to the end of the

short arm of the chromosome), replacing the PMI and moPAT genes with FRT1-NPTII:PINII TERM, 3x 35S ENH:UBI PRO:UBI INTRON:GAT891G3:UBQ3 TERM AND LTP2:GZ-W64A TERM-FRT87 (SEQ ID NO: 44). In a second independent transformation experiment, the pollen-inhibitor cassette PG47::barnase (comprising a PG47 [a maize promoter from a polygalacturonase gene] promoter driving the pollen-inhibitor gene barnase and a pinII terminator; bp 4438 to bp 8039 in PHP70154, SEQ ID NO: 43) was introduced into a pre-existing SSI target site also containing FRT1-PMI + Actin::moPAT-FRT87) located on chromosome 3 at genetic position 3.2 cM, replacing PMI and moPAT with PG47::Barnase.

These two transgenic events were crossed together, creating an F1 generation. Progeny in the F2 generation are then screened for linkage and a progeny plant (maize Pioneer inbred line 1, such as for example PHN46) is selected comprising the linked GAT (glyphosate resistant trait locus of interest) and barnase (pollen-inhibitor gene loci

B. Use of a pollen-inhibitor gene barnase to break linkage with a glyphosate resistance trait (GAT) on chromosome three of maize for accelerated trait introgression.

The maize Pioneer inbred line 1 carrying the linked GAT (glyphosate resistant trait locus of interest) and barnase (pollen-inhibitor gene) described above, can be used as the female in a cross with an (elite) inbred into which GAT will be introgressed. The resultant F1 plants can then be used as pollen-donors back onto the recurrent parent. Pollen that carries both the GAT locus and the pollen-specific barnase expression cassette are non-viable. Progeny of this cross can be screened by exposing the germinating seedlings to glyphosate which eliminates any progeny not carrying the trait locus. In viable progeny that pass this screen (i.e. progeny that contains the GAT trait and is resistant to the herbicide), the linkage between GAT and the pollen-specific barnase inhibitor would have been broken during meiosis. The resultant progeny now carry only the terminal end of the original Pioneer inbred line 1 chromosome 3 carrying the GAT locus, and the remainder of chromosome 3 has been replaced by chromosome 3 from the recurrent parent inbred (Figure 1).

C. Introducing the pollen-inhibitor gene -alpha amylase- in close proximity to a trait locus of interest (GAT, glyphosate resistance trait) in the genome of a maize plant.

A Pioneer inbred Line 1 (PHN46) was identified that comprised a pre-existing transgenic SSI target site located at 152 cM on chromosome 10 containing a first expression cassette comprising a ubiquitin promoter driving a phosphomannose isomerase (PMI) and a pin II terminator, wherein the PMI was preceded by a FRT1 recombination site (UBI PRO::FRT1::PMI::pinII) linked to a second expression cassette comprising an actin promoter driving a moPAT selectable marker and a pin II terminator, followed by a FRT87 recombination site (ACTIN PRO:: moPAT::pinII-FRT87). That Pioneer inbred line 1 was then used as the transformation target for particle-gun-mediated delivery and introduction of a GAT-resistance expression cassette between two dissimilar FLP-recombinase sites (FRT sites FRT1-NPTII::PINII TERM + 3X(35S ENH): UBI1ZM PRO:UBI1ZM INTRON::GAT891G3::UBQ3 TERM + LTP2 PRO::DS-RED2::GZ-W64a TERM-FRT87(SEQ ID NO:44), along with a separate plasmid cassette (comprising a ubiquitin promoter driving a FLP recombinase terminated by a pinII terminator , UBI1ZM PRO:UBI1ZM INTRON:FLPM::PINII TERM (bp 411 to bp 4012 of PHP5096, SEQ ID NO:50), resulting in RMCE and the replacement of PMI & moPAT by GAT in the pre-existing transgenic SSI target site located at 152 cM on chromosome 10. At a position 2 cM proximal (150 cM on chromosome 10), an SSI donor sequence (SEQ ID NO:46) containing FRT1-NPT-II:pinII + 35S ENH:LTP2 PRO::TAGBFP::GZ-W64A TERM + ZM-PG47-PRO::Zm-AA1::IN2-1 TERM-FRT87 (comprising a PG47 promoter driving the Zea mays alpha amylase (Zm-AA1) gene) was also introduced via particle-gun-mediated RMCE in a separate transformation experiment with Pioneer inbred line 1. The expression of alpha amylase in maize plant comprising the expression cassette PG47-PRO::Zm-AA1::pinII inhibits pollen tube growth and these plants must therefore be crossed with plants containing the GAT locus through the female ear in order to establish linkage.

30

D. Use of a pollen-inhibitor gene –alpha amylase- to break linkage with a glyphosate resistance trait (GAT) on chromosome three of maize for accelerated trait introgression.

The maize Pioneer inbred line 1 carrying the linked GAT (glyphosate resistant
5 trait locus of interest) and alpha-amylase (pollen-inhibitor gene) described above,
can be used as the female in a cross with an (elite) inbred into which GAT will be
introgressed. The resultant F1 plants can then be used as pollen-donors back onto
the recurrent parent. Pollen that carries both the GAT locus and the pollen-specific
alpha-amylase expression cassette are non-viable. Progeny of this cross can be
10 screened by exposing the germinating seedlings to glyphosate which eliminates any
progeny not carrying the trait locus. In viable progeny that pass this screen (i.e.
progeny that contains the GAT trait and is resistant to the herbicide), the linkage
between GAT and the pollen-specific alpha-amylase inhibitor would have been
broken during meiosis. The resultant progeny now carry only the terminal end of the
15 original Pioneer inbred line 1 chromosome 10 carrying the GAT locus, and the
remainder of chromosome 10 has been replaced by chromosome 3 from the
recurrent parent inbred (Figure 1).

In a fashion similar to Example 1 A-D described above, a trait locus of
interest located at position of 0.9 cM on chromosome 1 containing the trait
20 expression cassette ACTIN PRO::moPAT::pinII and the PG47 PRO::ZmAA1::pinII
cassette can be introduced into on chromosome 1, at position 3.2 cM of the Pioneer
Proprietary genomic map (PHD 3.2 cM). Establishing linkage (between the pollen-
inhibitor genes such as ZM-AA1 and the trait loci of interest such as GAT891G3)
and using this linked pair to cross to many different commercially important inbreds,
25 can create a population of F1 hybrids. Each of the F1 hybrids can be used as a
pollen donor back onto the same commercially-important inbred and progeny can be
screened for accelerated introgression of the trait into each of the commercially-
important inbreds.

EXAMPLE 2Using two pollen-specific inhibitor loci flanking a trait of interest for accelerated trait introgression of a non-telomeric trait locus.

Illustrated in Figure 2 and described below is an example of how two pollen-specific inhibitor loci can be located such that they flank a trait of interest, and its use for accelerated trait introgression of a non-telomeric trait locus of maize.

Figure 2 shows a schematic of introducing flanking pollen-inhibitor loci around an internal chromosome location containing a trait, and using the triple-linked site to rapidly introgress the trait into other inbreds. Step 1: Choose a transformable inbred that contains three SSI target sites, the first site for introduction of the trait (i.e. GAT at site 2) and two flanking sites (Sites 1 & 3), each a close genetic distance from the trait locus (for example, 1 cM), into which a pollen-inhibitor cassette (PI) will be introduced. In a fourth transformation, Agrobacterium is used to randomly integrate a Pollen-inhibitor maintainer (PIM). Step 2: Use three separate SSI transformation experiments to introduce PI (for example, PG47::bar^{ma}::pinII) into Sites 1 and 3 and GAT (UBI::GAT::pinII) into Site 2. Use Agrobacterium to randomly integrate PIM (for example, UBI::barstar::pinII) into the same inbred background. Identify the PIM transgenic line that expresses the highest level of PIM to use as a maintainer (of male fertility) in crosses performed in steps 3 & 4 below. Step 3: Carry pollen from the GAT-containing plant to a PI-containing plant (Site 1) and screen the progeny using molecular markers to identify plants containing both GAT and PI. Meanwhile, carry pollen from PIM-containing plant to PI-containing plant (Site 3) and screen progeny with molecular markers find plants containing both PIM and PI. Step 4: Carry pollen from plants containing both PIM/PI to plants containing GAT/PI generate progeny having all four genes. During this step, the first linkage between Site 1 (PI) and Site 2 (GAT) has been established. Step 5: Carry pollen from a Wild-type inbred plant to the plants containing all four genes and use molecular markers to screen progeny in order to identify plants that contain Site1(PI), Site2(GAT) and SITE3(PI) with no PIM. Linkage has now been established in these plants for sites 1,2 and 3. Step 6: Carry pollen from a different Elite inbred to Linked WT plants and use molecular markers to screen for F1 progeny that contain the triple-linked PI-GAT-PI. Step 7: Carry pollen from PI-GAT-

PI F1 progeny back to the Recurrent Parent (the elite inbred) to generate a progeny pool. Screen progeny for glyphosate resistance by spraying with the herbicide. The surviving plants will contain the GAT gene with a minimal linkage drag the transformation inbred (i.e. less than 1 cM of flanking chromosome remains from this inbred).

A. Introducing two pollen-inhibitor genes in close proximity to a trait locus of interest (GAT, glyphosate resistance trait) in the genome of a maize plant.

A transgenic trait locus containing the GAT expression locus (GAT, Figure 2) can be introduced using SSI to introduce the expression cassettes between FRT1 and FRT878 of PHP69519 (SEQ ID NO: 44) into chromosome six (6) at a genetic position of 23.7 centimorgans (cM) in the maize Pioneer inbred line 1 (such as for example PHN46). This will be abbreviated in the remainder of this example as the GAT trait locus. Two flanking transgenic SSI target sites at genetic positions 22.7cM and 24.7 cM can be used to set up the recombination screen (i.e. screening for two recombinations in one generation), setting up the screen to function during pollen germination (as illustrated in Figure 2). Before introducing the pollen-inhibitor expression cassette (PI, figure 2), the transgenic line containing the GAT trait locus can also be transformed using *Agrobacterium* with T-DNA to create a randomly-integrated maintainer locus. In a male sterile line, a maintainer locus modulates or antidotes the pollen-inhibitor locus so that the pollen is viable when the maintainer is present and the plant can be used as a pollen donor. For example, if a pollen-expressed barnase is used in the pollen-inhibitor locus, its cognate antidote protein barstar expressed by a similar pollen-specific promoter would be used as the maintainer. When barstar segregates away from barnase, pollen-inhibitor effect is restored. In this example, we use a variation on this theme, with a pollen-specific promoter with TET repressor operator sequences surrounding the TATA box controlling expression of the ZM-AA1 gene that confers pollen sterility, and a pollen-expressed TET repressor gene at another genomic locus to prevent expression of the pollen-inhibitor, thus acting as a maintainer. To create the maintainer germplasm containing TET repressor, a separate *Agrobacterium*-mediated transformation is performed to introduce the T-DNA shown in PHP01 (Figure 9-H,

SEQ ID NO: 51) which introduces the following expression cassettes: RB-35S-Enhancer::PG47 PRO::TET repressor (MOD 1)::SB-SAG12 TERM + UBI1ZM PRO: UBI1ZM INTRON::PMI::pinII-LB (RB= right border, PG47 = PG47 promoter, TETR=TET repressor, which is a maize-optimized gene that encoded the *E. coli* TET repressor protein, pinII terminator, UBI= ubiquitin promoter, PMI = phosphomannanase isomerase, LB= left border of T-DNA) . In this construct the 35S-enhancer element can be used to enhance expression of the PG47-driven TET repressor (TET repressor) gene. The expressed TET repressor is capable of binding to the TET operator (OpT) sequence placed between the PG47 promoter and the ZM-AA1 pollen-inhibitor gene.

Pollen inhibition can be accomplished by using a pollen-specific promoter driving the alpha-amylase gene (AA) which is introduced both at positions 22.7 cM and 24.7 cM on chromosome six using SSI-mediated integration of the seed-specific DS-RED2 and PG47:Top3:ZM-AA1 contained in PHP02 (Figure 9-I, SEQ ID NO: 52) at position 22.7 cM, and the seed-specific YFP and PG47:Top3:ZM-AA1 contained in PHP03 (Figure 9-J, SEQ ID NO: 53) at position 24.7 cM.. By crossing the TET repressor (TETR) germplasm expressing TETRrepressorwith germplasm containing PG47:Top3:ZM-AA1and identifying progeny that now contain both loci, expression of the ZM-AA1 pollen-inhibitor is repressed and these progeny plants can be used as the pollen donor for further crossing, eventually bringing all three transgenic loci (DS-RED/ZM-AA1 at position 22.7, GAT at position 23.7, and YFP/ZM-AA1 at position 24.7, all three forming a linkage group on the same chromosome. Progeny plants confirmed to contain all three linked loci can be used to outcross to another inbred (referred to as the recurrent parent in backcross breeding), beginning the introgression process (see section B below).

B. Use of two pollen-inhibitor genes flanking a trait locus on interest (ZMAA1-GAT-ZMAA1) to break linkage with a glyphosate resistance trait (GAT) for accelerated trait introgression in maize.

Once all three loci are physically linked (ZMAA1—GAT—ZMAA1), this material can then be used to cross with other inbreds (for example, Pioneer Inbred Line 3, Figure 2) to begin the introgression process. The resultant F1 hybrid

contains one copy of chromosome 6 that contains the three linked loci, and the other copy of chromosome 6 with no transgenes (from the recurrent parent). F1 hybrids are screened for the presence of TET repressor using PCR, and individuals in which the TET repressor locus has segregated away are identified for the next

5 cross. For the next cross, the F1 hybrid is used as the male for pollination of ears on wild-type recurrent parent plants. In pollen grains in which no meiotic recombination occurs between the three linked loci, expression of alpha amylase depletes the pollen grain of starch and pollen tube growth is inhibited. In pollen grains in which a single cross-over occurs between GAT and one of the flanking

10 ZMAA1 loci, there is still an active ZMAA1 expression cassette linked to GAT on the side where the cross-over did not occur and the pollen does not germinate. However, in pollen grains in which recombination occurred between GAT and each of the flanking ZZAA1 loci, the pollen grains contain no pollen-inhibitor and can successfully germinate to deliver the sperm cells to the ovule resulting in successful

15 fertilization and embryo development. Progeny seedlings can be screened for GAT by spraying with glyphosate, eliminating any progeny resulting from wild-type pollen. The remaining viable plants now contain the GAT gene and only a small surrounding segment of the original Pioneer inbred line 1 chromosome 6 (i.e. less than 1 cM on either side of GAT) in a Pioneer Inbred Line 3 chromosome 6. To

20 further confirm that the two pollen-inhibitor loci were lost, progeny seed exhibit no red or yellow fluorescence. Using inbred-specific markers provides the final confirmation that less than 1 cM on either side of the GAT locus is from the original transformed inbred and the remainder of chromosome 6 is from the recurrent parent Pioneer Inbred Line 3.

25 Using marker-assisted background selection methods has been widely used in agriculture since the 1990's (for example, see Frisch et al., 1999. Crop Science 39(5):1295-1301). However, the difficulty in introgressing a single trait is actually identifying meiotic recombination events that have occurred close the trait of interest. Our method provides a genetic selection at the gamete level (pollen) that

30 identifies cells and progeny in which this has occurred. Once the chromosome that contains the trait has been substantially converted to the new inbred background,

marker-assisted background selection can be used to more rapidly convert the remaining nine chromosomes.

5 *C. Use of one flanking pollen-inhibitor and a seed color marker on the other flank of an internal trait locus for accelerated trait introgression in maize.*

Figure 15 shows a schematic of identifying an internal trait locus (referred to as PAT, Step 1) and then selfing to produce a homozygous trait locus (Step 2). In Step 3, an aleurone-specific anthocyanin cassette (RK for Red Kernel phenotype, also referred to as a seed color marker) is introduced into a locus on one side of an internal chromosome location containing a trait through targeted integration into the trait-containing chromosome. This line, now containing the linked RK and Trait loci is pollinated by a haploid inducer line (RWS) to produce haploid immature embryos (Step 4), and then the haploid embryos are used for targeted integration of the pollen-inhibitor (PI) cassette (Step 4). The haploid embryos are treated with a doubling agent such as colchicine to produce a double-haploid plant (Step 5). The triple-linked inbred is now ready for rapid introgression of the trait into other inbreds.

In Step 6, pollen is carried from a different Elite inbred (the recurrent parent into which the trait will be introgressed) onto the triple-linked plants (containing RK—Trait—PI) and use molecular markers to screen for F1 progeny that contain the triple-linked RK-GAT-PI. In Step 7, pollen is carried from the RK-Trait-PI F1 progeny back to the Recurrent Parent (the elite inbred) to generate a progeny pool. Kernels are screened for red color. If the kernel is red, it is discarded or set aside for another round of crossing. Yellow kernels were produced by pollen that had lost the pollen-inhibitor through breaking the linkage between the Trait and the PI locus, and had also lost the Red Kernel phenotype by breaking the linkage between the Trait and RK. Yellow kernels are then germinated in herbicide (or germinated and then sprayed with herbicide) to eliminate wild-type progeny, and the surviving seedlings contain the trait locus that has been rapidly introgressed into the trait-carrier chromosome.

30

D. Use of two flanking seed color markers for accelerated trait introgression in maize.

Figure 16 shows a schematic of identifying an internal trait locus (referred to as PAT, Step 1) and then selfing to produce a homozygous trait locus (Step 2). In Step 3, an aleurone-specific anthocyanin cassette (RK for Red Kernel phenotype) is introduced into a locus on one side of an internal chromosome location containing a trait through targeted integration into the trait-containing chromosome. This line, now containing the linked RK and Trait loci is pollinated by a haploid inducer line (RWS) to produce haploid immature embryos (Step 4), and then the haploid embryos are used for targeted integration of the same aleurone-specific anthocyanin cassette (Step 4). The haploid embryos are treated with a doubling agent such as colchicine to produce a double-haploid plant (Step 5). The triple-linked inbred is now ready for rapid introgression of the trait into other inbreds. In Step 6, pollen is carried from a different Elite inbred (the recurrent parent into which the trait will be introgressed) onto the triple-linked plants (containing RK—Trait—RK) and use molecular markers to screen for F1 progeny that contain the triple-linked RK-GAT-RK. In Step 7, pollen is carried from the RK-Trait-RK F1 progeny back to the Recurrent Parent (the elite inbred) to generate a progeny pool. Kernels are screened for red color. If the kernel is red, it is discarded or set aside for another round of crossing. Yellow kernels are produced by pollen that had lost the Red Kernel phenotype by breaking the linkage between the Trait both RK loci. Yellow kernels are then germinated in herbicide (or germinated and then sprayed with herbicide) to eliminate wild-type progeny, and the surviving seedlings contain the trait locus that has been rapidly introgressed into the trait-carrier chromosome.

25

E. Use of one flanking seed color marker expressed in the crown of the kernel and a second seed color marker expressed in the base of the kernel on the other flank of an internal trait locus for accelerated trait introgression in maize.

Figure 17 shows a schematic of identifying an internal trait locus (referred to as PAT, Step 1) and then selfing to produce a homozygous trait locus (Step 2). In Step 3, a crown-specific anthocyanin cassette (RC for Red Crown phenotype) is introduced into a locus on one side of an internal chromosome location containing a

30

trait through targeted integration into the trait-containing chromosome. This line, now containing the linked RC and Trait loci is pollinated by a haploid inducer line (RWS) to produce haploid immature embryos (Step 4), and then the haploid embryos are used for targeted integration of base-specific anthocyanin cassette (referred to in Step 4 as RP for red basal expression in the endosperm or aleurone underneath the pedicel). The haploid embryos are treated with a doubling agent such as colchicine to produce a double-haploid plant (Step 5). The triple-linked inbred is now ready for rapid introgression of the trait into other inbreds.

In Step 6, pollen is carried from a different Elite inbred (the recurrent parent into which the trait will be introgressed) onto the triple-linked plants (containing RK—Trait—RP) and use molecular markers to screen for F1 progeny that contain the triple-linked RK-GAT-RP. In Step 7, pollen is carried from the RK-Trait-RP F1 progeny back to the Recurrent Parent (the elite inbred) to generate a progeny pool. Kernels are screened for red color. If the kernel has a red crown or a red base, or both, it is discarded or set aside for another round of crossing (with the expression pattern being an indication of which side broke linkage). Yellow kernels are produced by pollen that had lost the both the RC and RP phenotypes by breaking the linkage between the Trait both flanking loci. Yellow kernels are then germinated in herbicide (or germinated and then sprayed with herbicide) to eliminate wild-type progeny, and the surviving seedlings contain the trait locus that has been rapidly introgressed into the trait-carrier chromosome.

EXAMPLE 3

Using guide polynucleotide/Cas endonuclease mediated integration to introduce two pollen-specific inhibitor loci flanking a trait locus for accelerated introgression of a non-telomeric trait locus .

Illustrated in Figure 4 and described below is an example of how a guide polynucleotide/Cas endonuclease system can be used to mediate integration to introduce two pollen-specific inhibitor loci flanking a trait locus for accelerated introgression of a non-telomeric trait locus .

Figure 4 shows a schematic of the use of the guide polynucleotide/Cas endonuclease system to integrate a pollen-inhibitor cassette. Two expression cassettes are introduced along with the DNA to be integrated, using an equimolar

amount of the three DNAs to attach to 0.6 uM gold particles for particle bombardment. The first expression cassette contains the UIB1ZM PRO & INTRON::CAS9 and a PINII TERM. The second expression cassette contains the maize U6 PolIII Chr8 promoter driving expression of the guide RNA and a terminator

5 in the form of ZM-U6 PRO::gRNA::PINII TERM. The DNA to be integrated contains two flanking sequences that are homologous to the inbred genomic sequence at the target site (HR1 at the 5' end and HR2 at the 3' end) with the following three expression cassettes in between HR1 and HR2; PG47 PRO::ZM-AA1::PINII TERM + LTP2 PRO::ZS-YELLOW1 N1::PINII AND UIB1ZM PRO & INTRON::PMI::PINII.

10 Upon particle gun-mediated delivery of all three DNA molecules, CAS9 protein is encoded and guide-RNA is transcribed. The guide RNA associates with the CAS9 protein and guides the complex to a genomic sequence (complementary to the guide-RNA) in between the genomic sequences that correspond to the HR1 and HR2 sequences in the DNA to be integrated. CAS9 creates a double-stranded

15 break at the target site, stimulating homologous recombination between the HR sequences and their corresponding genomic sequences, integrating the three expression cassettes containing ZM-AA1, ZS-YELLOW1 N1 and PMI. Transgenic events that have successfully integrated these genes at this specific genomic site are selected by culturing on medium containing mannose and confirmed through

20 yellow fluorescence, PCR and sequencing.

In this example, the trait locus can be a GAT expression cassette (UBI::GAT::pinII) integrated into chromosome 1 at genetic position 53.14 cM. A pollen-inhibitor expression cassette (PG47-PRO::ZM-AA1::pinII) can be introduced at positions 52.56 cM and 54.56 cM on chromosome 1 using guide

25 polynucleotide/Cas endonuclease mediated integration of the a sequence containing three expression cassettes, an expression cassette with PG47 PRO::ZM-AA1::PINII TERM, a cassette with the LTP2 PRO::YFP::PINII TERM (or with DS-RED2 in place of YFP), and a cassette with UIB1ZM PRO: UIB1ZM INTRON::PMI::PINII TERM (Figure 4). The contiguous sequence containing all

30 three expression cassettes is also flanked by maize genomic sequences that are homologous to the genomic sequence to which the guide-RNA will direct CAS9 for targeted double stranded breakage. Using the inbred line that is homozygous for

the GAT expression cassette for transformation, UBI::CAS9::pinII and a guide RNA that directs the CAS9 protein to position 52.56 cM or 54.56 cM can be introduced into immature embryo cells via particle bombardment along with the *Zea mays* alpha amylase (Zm-AA1) expression cassette flanked by genomic regions for homologous recombination from the inbred (HR regions, see Figure 4). The inbred HR sequences can be about 1-2kb sequences that are homologous to the endogenous inbred chromosome sequences flanking the Cas9 recognition sequence (a DNA sequence that is recognized and optionally cleaved by the polynucleotide-guided Cas endonuclease). For position 52.56 cM, the three expression cassettes shown in figure 4 are integrated using HR sequences that correspond to flanking sequences at 52.56 cM. For position 54.56 cM, a similar integration construct is integrated except that DS-RED2 is the fluorescence gene in the first expression cassette (between LTP2 PRO and PINII TERM) and the HR sequences correspond to flanking sequences at this position. Transgenic events containing the targeted integration can be recovered by selecting on medium containing 5 g/l maltose and 12.5 g/l mannose. Transgenic plants can be regenerated and used as the female parent for pollination with RWS homozygous for a LEC1::GFP::pinII cassette. The resulting immature embryos can be screened first for lack of green fluorescence (indicating diploid embryos which are discarded), then for DsRED (containing the ZM-AA1 locus), can be re-transformed using the guide polynucleotide/Cas endonuclease system, but this time integrating the ZM-AA1 expression cassette along with UBI::NPT-II::pinII at PDH 54.56 cM on chromosome 1. After re-transformation, the immature embryos can be exposed to 0.06% colchicine for 8 hours in culture and can be then moved onto kanamycin to select events that have integrated AA1 and NPT-II at position 54.56 cM on chromosome 1. The resultant plants that are regenerated are homozygous for the triple-linked loci (AA—GAT—AA). This material can now be used as a female to bulk up seed. The resultant hemizygous progeny can then be used to cross with other inbreds (for example, Pioneer Inbred Line 3) to begin the introgression process. The resulting F1 plants can then be used as the male crossing once again to the recurrent parent (Pioneer Inbred Line 3). In pollen grains in which no meiotic recombination occurs, expression of alpha amylase depletes the pollen grain of starch and pollen tube

growth is inhibited. In pollen grains in which a single cross-over occurs between GAT and one of the flanking AA loci, there can still be an active AA expression cassette linked to GAT and the pollen does not germinate. However, in pollen grains in which recombination occurred between GAT and each of the flanking AA loci, the pollen grains can successfully germinate and deliver the sperm cells to the ovule resulting in successful fertilization and embryo development. Progeny seedlings can be screened for GAT by spraying with glyphosate, eliminating any progeny resulting from wild-type pollen. The remaining viable plants now contain the GAT gene and only a small surrounding segment of the original Pioneer inbred line 1 chromosome 1 (for example, but not limited to, less than 2 cM on either side of GAT) in an otherwise complete Pioneer Inbred Line 3 chromosome 1.

EXAMPLE 4

Creating a pre-existing Precision Accelerated Trait Integration Site (PATI Site)

Illustrated in Figure 6 and described below is an example of how a pre-existing Precision Accelerated Trait Integration Site (PATI Site) can be created.

A SSI-generated trait locus can be established at a chromosomal location that is confirmed to be agronomically neutral, have no deleterious impacts on expression patterns of surrounding endogenous genes, and that supports good transgene expression levels. For example, as shown in Figure 6, a SSI target site (RTL) can be introduced into position 52 cM on maize Pioneer Inbred Line 3 chromosome 1, using Site-Specific Integration. This event can contain three expression cassettes; UBI1ZM PRO & INTRON:FRT1:NPTII::PINII + UBI1ZM PRO & INTRON::GAT891G3::UBI3 TERM + LTP2::DS-RED2::GZ-W64 TERM-FRT87, with the sequence and features from FRT1 to FRT87 being shown in Figure 9-B, (SEQ ID NO: 44) which were found to support expression of NPTII at consistently high levels across multiple generations. In immature embryos homozygous for the NPTII-containing locus, guide polynucleotide/Cas endonuclease-mediated targeted integration (see also Figure 4) of PG47 PRO::ZM-AA1::pinII + LOXP-UBI PRO & INTRON::PMI::pinII-LOXP + UBI PRO & INTRON::CYAN::pinII can be performed to introduce these expression cassettes at chromosomal position 50 cM on chromosome 1, and in a successive transformation, guide polynucleotide/Cas endonuclease systems can be used again to integrate a third combination of

expression cassettes (PG47 PRO::ZM-AA1::pinII + LOX2272-OS-ACTIN PRO::MOPAT::35S TERM-LOX2272 + UBI PRO & INTRON::ZS-GREEN::pinII) at position 54 cM on chromosome one. Once the final integration event (on 3 mg/l bialaphos) and plants are regenerated, the T0 plants can be crossed to another transgenic line

5 containing RAB17 PRO::MO-CRE::PINII (SEQ ID NO: 75), which expresses CRE recombinase in response to desiccation (i.e. during late embryogenesis) or cold temperatures. In cells containing the triple-linked PATI locus and expression CRE recombinase, both the PMI and MOPAT expression cassettes are excised, leaving PG47 PRO::ZM-AA1::pinII + LOX2272 + UBI PRO & INTRON::CYAN::pinII at both

10 flanking positions in the PATI locus. This germplasm, now containing the three linked loci (trait locus and the two flanking pollen-inhibition loci, Figure 6) can be used for subsequent testing of new trait constructs and for rapid product development. Using either guide polynucleotide/Cas endonuclease systems or RMCE, new trait cassettes can be introduced into the trait locus (replacing NPTII

15 with an herbicide resistance gene for selection along with new traits (or trait-stacks). When efficacious events are identified, the T1 plants can be crossed to new inbreds producing the F1 hybrid for introgression. The F1 can be used as the male to cross back to the same recurrent inbred parent and the BC1-progeny can be screened for the new trait. At this point, the only remnant of Pioneer Inbred Line 3 chromosome

20 1 will be a fragment somewhere between 50 and 54 cM (the exact junctions depending on precisely where the meiotic recombinations occurred) containing the trait, while the remainder of chromosome 1 (from somewhere between about 50-54 centimorgans from the new inbred)

Any proven PATI site with a trait locus linked to a pollen-inhibition locus (for a

25 telomeric trait locus) or to two pollen-inhibition loci (for an internal or non-telomeric trait locus) that has been demonstrated to support appropriate transgene expression and not be deleterious to plant growth and productivity can be used in this fashion, as a pre-established site for transgene introduction and accelerated trait

introgression. Pre-existing Targeted Accelerated Integration Sites such as these

30 can be used to introduce the trait gene, and can be used to introduce either single genes or molecular stacks and then rapidly introgressed these into many new inbreds for testing.

EXAMPLE 5Two flanking pollen-inhibitor intein halves used in two-stage screening for two sequential recombinations.

This experiment utilizes a bacterial endoribonuclease from *Escherichia coli* referred to as KID (Ruiz-Echevarria et al., 1991, Mol. Microbiol. 5:2685-2693. This gene was split into two fragments and then the sequences encoding the amino- (KID-N, SEQ ID NO: 67) and carboxy- (KID-C, SEQ ID NO: 69) fragments were fused to the amino (NP-INTE-N, SEQ ID NO: 65) and the carboxy (NP-INTE-C, SEQ ID NO: 63) intein halves, respectively. These intein halves (from *Nostoc punctiforme*, see Iwai et al., 2006, FEBS Lett. 580:1853-1858) encode two cognate intein, that when expressed in the same cell when fused to two cognate protein fragments peptides (fusing KID-N with NP-INTE-N in that order, and fusing NP-INTE-C with KID-C in that order) will bind with each other and catalyze their own excision, effectively splicing the two protein fragments (i.e. KID-N AND KID-C) together to form a fully functional protein (i.e. KID).

An internal trait locus such as the RMCE locus at position 23.7 cM on chromosome 6 (Figure 7) containing UBI1ZM PRO:: UBI1ZM INTRON::moPAT::PINII (SEQ ID NO: 19, SEQ ID NO: 20, SEQ ID NO: 35, and SEQ ID NO: 8, respectively) can be flanked by PG47 PRO::NP-INTE-C~KID-C::PINII (SEQ ID NOs: 13, 63, 69 and 8, respectively) + LTP2 PRO::MO-CYAN::PINII (SEQ ID NOs: 9, 71 and 8, respectively) at position 23.8 cM and by PG47 PRO::KID-N~NP-INTE-N::PINII (SEQ ID NOs: 13, 67, 65 and 8, respectively) + LTP2 PRO::MO-CYAN::PINII at position 23.6 cM (shown in Figure 7). When both i) the carboxy- half of the KID protein fused to the carboxy-half of the *Nostoc* intein pair, and ii) the amino-half of the intein pair fused to the amino-half of alpha-amylase are expressed in the same pollen grain, the intein pair splices the two halves of the KID protein together resulting in full enzyme function and concomitant pollen inhibition. Seed containing both KID-INTE also fluoresce blue. Pioneer inbred line 1 containing the trait locus linked to both of the KID-intein loci (hemizygous) can be crossed to other inbreds for introgression of the trait, producing F1 populations. The F1 progeny can then be used as the pollen donor back onto the recurrent inbred. In pollen in which a double crossover occurs breaking the linkage on both sides of the

trait locus, the pollen can be viable and produces seed with just the trait locus introgressed into the new inbred chromosome 6. However, if recombination to break linkage occurs on only one side of the trait (removing either the carboxy- or amino-half of KID) the pollen can also be viable leaving behind a non-functional KID-intein half and blue fluorescence. This blue seed can then be grown and used as the pollen donor and the resultant progeny are germinated in the presence of glufosinate (eliminating progeny that came from wild-type pollen). Plants that are herbicide resistant and also produce no blue seed have the UBI::moPAT trait locus that has been introgressed with less than 0.1 cM of Pioneer inbred line 1 chromosome remaining on either side of the locus.

As an alternative to the fluorescent proteins used in the second screening step (to monitor the second recombination and breakage of the linkage), a conditional negative selection marker such as the coda gene, the dhIA gene or the CYP105A gene can be used. When these genes are expressed in plant cells there is no adverse effect until their cognate substrates (5-fluorocytosine, dihaloalkanes or sulfonylurea R4702, respectively) are provided to the plant cell at which point these non-inhibitory substrates are converted to an inhibitor. Using such as marker, fluorescent microscopy is not needed to identify plant cells or tissues in which the genetic linkage has been broken and there is also no need for a maintainer line to render the inhibitor (coda, dhIA or CYP105A) inactive when not needed.

A trait such as resistance to the glufosinate herbicides such as Basta and Liberty (OS-ACTIN PRO & INTRON::MOPAT::35S TERM (SEQ ID NO: 95) can be integrated at Chromosome 6, 23.7 cM in inbred Pioneer inbred line 1 (Figure 7) which can represent the internal trait locus to be introgressed into other inbreds. At position Chr6-23.8 cM the following can be introduced; FRT1-NPTII-PINII TERM + LTP2 PRO-DSRED-GZ-W64A TERM + PG47 PRO-NPINTEC-KIDC-PINII TERM (PHP04, Figure 9-K and SEQ ID NO: 54). At position Chr6-23.6 cM the following can be introduced; FRT1-NPTII-PINII TERM + LTP2 PRO-ZSYELLOW1N1-GZ-W64A TERM + PG47 PRO-KIDN-NPINTEN-PINII TERM (PHP04 in Figure 9-K). Thus, expression of the cassettes at 23.6 cM encode a pollen-specific KID-N (the amino-fragment of the KID protein fused to the *Nostoc punctiforma* DnaE amino-intein half) and a seed-specific yellow fluorescent protein. Expression of the

cassettes at 23.8 cM encode a pollen-specific KID-C (the carboxy-fragment of the KID protein fused to the *Nostoc punctiforme* DnaE carboxy-intein half) and a seed-specific red fluorescent protein. When both intein halves are expressed in the same pollen grain, the intein-halves undergo self-catalyzed cleavage resulting in perfect reconstitution of the full-length, active KID toxin protein, and the pollen are non-viable. However, if linkage on either side of the trait gene occurs between the trait and the intein/fluorescent locus on that side, only one half of the intein-pair is expressed in the pollen grain and the pollen grain is viable. In this manner, breaking the linkage on either side of the trait can readily be determined because the pollen grain is viable and only one color of fluorescence is passed on to the BC1 (1st generation of backcrossing) progeny. These progeny can then be used for a subsequent round of pollination back to the recurrent parent, and BC2 progeny screened for lack of any fluorescence. At this point, successful introgression of the trait into chromosome 6 has been accomplished.

EXAMPLE 6

Rapid creation of two pollen-specific inhibitor loci flanking a trait of interest for accelerated trait introgression.

In this example we describe the use of sequential guide polynucleotide/Cas endonuclease system mediated integration to introduce a first flanking pollen-inhibitor cassette, followed by SSI-mediated targeting along with haploid embryo transformation to produce two flanking pollen-specific inhibitor loci linked to an intervening trait locus (illustrated in Figure 3).

Figure 3 shows a schematic of the use of sequential targeting into haploid embryos to rapidly create linkage between a trait locus and two flanking pollen-inhibitor loci. Step 1: The Pioneer inbred Line 3 containing a 35S::PAT expression cassette at position 53.1cM on Chromosome 1 (conferring the trait of bialophos-resistance) is used as the starting germplasm in this strategy. Step 2: The hemizygous PAT-containing plants are self-pollinated to create homozygous PAT/PAT progeny. Step 3: The CAS/CRISPR system is used to target a PI (PG47::ZmAA1::pinII + UBI::DsRED::pinII + UBI::PmI::pinII) with into 51.56cM and establish the first linkage between PI and GAT. Transgenic events are selected on mannose and hemizygous T0 plants are regenerated. Step 4: T0 plants containing

PI(51.56) and PAT(53.1) are pollinated by a DuPont/Pioneer RWS-derived haploid-inducer line (Big Kahuna) progeny immature embryos are screened by color to discard diploid embryos and harvest DsRED-expressing embryos for re-transformation. The haploid embryos containing PI(52.3) linked to PAT(53.1) are
 5 used for a second round of CAS/CRISPR transformation in order to introduce a second PI (PG47::barnase::pinII + UBI::CYAN::pinII + UBI::NPTII::pinII) at position 54.56cM, establishing linkage between all three transgenic loci. Transgenic events are selected on G418. Step 5: The embryos are treated briefly with colchicine to double the chromosome number and homozygous plants containing the three linked
 10 loci on both copies of chromosome 1 are produced. Step 6: Pollen from the Elite inbred into which PAT is being introgressed (for example, Pioneer Inbred Line 2) is carried to the triple-linked Pioneer Inbred Line 3 ears to produce F1 progeny. Step 7: Pollen from the F1 progeny is used to pollinate the recurrent parent (Pioneer Inbred Line 2) to generate progeny. Progeny are sprayed with bialaphos herbicide
 15 which eliminates progeny derived from wild-type pollen. The surviving plants will have GAT gene with a minimal linkage drag from donor inbred.

The Pioneer inbred Line 3 homozygous for a specific trait locus referred to as TRAIT1 located at 53.1cM (Pioneer Proprietary genomic map) on chromosome 1 can be transformed using particle bombardment of immature embryos to introduce
 20 via guide polynucleotide/Cas endonuclease systems (Figure 4). A donor DNA sequence containing UBI1ZM PRO::UBI1ZM INTRON::PMI::PINII + UBI1ZM PRO::UBI1ZM INTRON::ZS-YELLOW N1::GZ-W64 TERM + 35S ENH:PG47 PRO::BT1 TP::ZM-AA1::IN2-1 TERM (Figure 9-O and SEQ ID NO: 116) can be integrated at position 51.6 cM on chromosome one. Successful targeted integration
 25 events can be selected on medium containing mannose. Plants can be regenerated from these events and grown in a greenhouse. Perfect homologous recombination at position 51.6 cM can be confirmed by PCR reactions specific for the recombination product. When plants are mature and the ears have produced silks, these plants can be pollinated using RWS haploid inducer line (see Rober et al.,
 30 2005, Maydica 50:275-283). In progeny immature embryos, diploid embryos expressing an anthocyanin marker can be discarded. Haploid embryos that fluoresce yellow (indicating that they contain SEQ ID NO: 117 at position 51.6 cM)

can be used for a second round of targeting, using particle gun delivery of UBI::moFLP::pinII, OS-ACTIN::WUS::PIN, UBI PRO & INTRON::ODP2::PINII plus an SSI-donor DNA containing a second pollen-inhibitor cassette, a second selectable marker and a second fluorescent marker (FRT1::NPTII::pinII + 35S

5 ENH:UBI1ZM PRO::UBI1ZM INTRON::DS-RED2::PINII + 35S ENH:PG47 PRO::ZM-AA1::IN2-1 TERM-FRT87 (Figure 9-M and SEQ ID NO: 76) at position 54.56 cM on chromosome one. Successful RMCE into this recombinant trait locus (RTL) can activate NPTII gene expression which permits primary selection of the RMCE events on kanamycin- or G418-containing medium with a secondary

10 selection for red fluorescence, that can target integration into a RTL at 54.56cM on chromosome 1 that can contain the same two dissimilar IFRTb sites (FRT1 and FRT87). Events can be briefly exposed to colchicine early in tissue culture to double the chromosome number. Events can be selected for NPTII expression using the selective agent G418 and can further be confirmed by visually selecting

15 for red fluorescence. Events can be screened using PCR for the correct RMCE into the Pioneer Inbred Line 3 RTL located at 54.56cM on chromosome 1. These events can be regenerated into plants and grown in a greenhouse, and upon further characterization using PCR analysis, can be confirmed to be homozygous for the DsRED locus at 51.56 cM, homozygous for the DP4114 trait locus at 53.1cM and

20 homozygous for the ZS-YELLOW N1 locus at 54.56 cM. These plants can be grown alongside an inbred into which the TRAIT1 will be introgressed, for example Pioneer Inbred Line 2. Pollen from Pioneer Inbred Line 2 can be used to pollinate the PH2HT plants containing the triple-linked loci to produce an F1 hybrid. The F1 seed can then be planted and the resulting F1 plants can be used to pollinate the

25 recurrent parent Pioneer Inbred Line 2. Only pollen grains in which the linkage has been broken will i) be viable successfully fertilizing the wild-type Pioneer Inbred Line 2 egg cells to produce kernels, and ii) produce bialaphos-resistant plants (conferred by the TRAIT1 locus) – resulting in rapid introgression of this trait into the Pioneer Inbred Line 2 chromosome 1.

30

Performing two sequential targeted integrations to introduce two screening markers on either flank of a transgenic trait locus (without a haploid step).

A trait locus located on Chr1-50.5 cM in Pioneer inbred line 1 has been demonstrated to be efficacious and is to be introgressed into many new inbreds for testing. Cas9-mediated targeted integration using immature embryo transformation can be used to introduce a pollen-specific promoter driving expression the pollen-tube inhibitor AA1 at Chr1-49.5 cM, and single-copy targeted-integration T0 plants can be identified using PCR. These plants can then be pollinated using wild-type pollen of Pioneer inbred line 1, and the T1 immature embryos can be isolated for a second round of CAS9-mediated integration of a second AA1 Marker at Chr1-51.5 cM. Using two back-to-back transformations of immature embryos permits rapid creation of the triple-linked AA1—Trait—AA1 that can then be used for rapid introgression.

EXAMPLE 7

Flanking an internal trait locus with a non-conditional pollen-inhibitor gene on one side and a conditional lethal gene on the other, for two-step screening for an introgressed trait locus.

This example describes a method for flanking an internal (non-telomeric) trait locus with a non-conditional pollen-inhibitor gene on one side and a conditional lethal gene on the other. This can be used for traits where breaking the linkage on both sides of the trait simultaneously (using back-crossing for introgression) occurs at too low a frequency (for example, due to very short genetic distances (between the trait locus and the flanking pollen-inhibitor loci on both sides) or due to meiotic recombination-interference),

The two pollen-inhibitor flanking loci can be as follows: one flanking locus can contain a non-conditional pollen-inhibitor such as PG47 PRO::ZM-BT1 TP~ZM-AA1::IN2-1 TERM (SEQ ID NO: 77) or PG47 PRO::BA-BARNASE::PINII TERM (SEQ ID NO: 78), and the opposite flanking locus can contain a conditional pollen-inhibitor expression cassette. Examples of a conditional expression cassette can be PG47 PRO::CODAcodA::PINII TERMpini (Ffor CODA gene, see SEQ ID NO: 81, and for encoded protein see SEQ ID NO: 82). The encoded CODA protein has been shown to be non-inhibitory when expressed in plant cells, until a non-inhibitory

substrate is added (5-fluorocytosine, see Koprek et al., 1999, Plant Journal 6:719-726). Upon addition of the substrate, 5-fluorocytosine is converted by the encoded protein into the inhibitory product 5-fluorouracil (see Koprek et al., 1999, Plant Journal 6:719-726).

5 Expressing a non-conditional pollen-inhibitor such as AA on one side can indicate that this locus can be crossed through the female to establish linkage, while the conditional codA locus does not inhibit pollen in the absence of 5-FC and can be crossed through the male to establish the three-way linkage and can be made homozygous. The Triple-linked plant (AA—GAT—codA) can be crossed to the
10 inbred-of-interest to create the F1. At this juncture, the F1 can be crossed back to the recurrent parent with no 5-FC application. Of the pollen-grains containing the triple-stacked locus, only those that have lost AA will be viable. The progeny can be sprayed with glyphosate to recover the GAT—codA progeny. These can again be crossed back to the recurrent parent but this time the plants can be sprayed with the
15 non-toxic 5-FC immediately before pollen-shed. Of the pollen grains containing codA before meiosis, only those pollen grains in which the linkage was broken will be viable.

This strategy can be used in the following, non-limiting examples:

- 1) when the genetic distances on either side are very small (and thus the
20 frequency of a double cross-over can be very small)
- 2) when recombinational-interference makes it impractical to screen for two simultaneous cross-overs
- 3) in crops where the number of pollen grains is not as great as in maize.

EXAMPLE 8

25 Creating a Cas9 mediated system for pollen inhibition:

In this example we describe how one can create a Cas9 mediated system for pollen inhibition useful for providing an alternative for accomplishing pollen inhibition for use in accelerated trait introgression (see Figure 5a, 5b and 5c).

Figure 5-A shows a diagram of the expression of a modified Cas
30 endonuclease protein fused to repressor peptides along with three guide-RNAs complementary to different sequences in the maize GT1 promoter. The modified Cas9 gene (dCAS9) has been modified to encode two altered amino acids in the

protein, changing aspartic acid at position-10 to alanine and changing histidine at position 840 to alanine. These two amino acid changes inactivate the CAS9 nuclease activity while maintaining the Cas9 DNA-binding function with the mutated version being referred to as dCAS9. The guide-RNAs (labeled A, B and C)

5 associate with the dCAS9 protein fused to ear-motif repressor peptides and guide the fusion-protein to the GT1 promoter, where the repressor motifs block transcription, effectively silencing the gene while the dCAS9~RP fusion is being expressed along with the guide-RNAs.

Figure 5-B shows a diagram of the use of dCAS9~RP as the pollen-inhibitor
10 for accelerated trait introgression. An expression cassette conferring resistance to glufosinate herbicides such as Basta and Liberty, containing RB-LOXP-OS-ACTIN PRO::OS-ACTIN INTRON:: MO-PAT::35S TERM-LOXP-LB (SEQ ID NO: 89) is integrated on Chromosome 2 at genetic position of 5 cM in the inbred Pioneer inbred line 1, while a second DNA containing the dCAS9~RP-mediated pollen-
15 inhibitor locus is integrated using SSI at a distance of one centimorgan proximal (toward the centromere, at 6 cM). When dCAS9~RP and the guide RNAs for the endogenous glycosyltransferase 1 gene (labeled as "G" on chromosome 3 are expressed (see Figure 5-A), the dCAS9~RP/gRNA complexes can bind to their cognate sequences in the GT1 promoter, thereby silencing the gene and resulting in
20 non-viable pollen grains.

Figure 5-C shows a diagram of expression cassettes comprising the DCAS9~RP-mediated Pollen-inhibitor locus. Three expression cassettes can be used; i) the pollen-specific promoter PG47 PRO can be used to drive expression of the fusion protein DCAS9~RP, ii) the maize constitutive promoter ZM-U6 POLIII
25 CHR PRO driving expression of five complexes consisting of a CYS4 recognition site, a GT1-PRO TARGET SEQUENCE (1 – 5) and a guide-RNA sequence (expanded cartoons for target sequence 1 and target sequence 5 are shown, but all five have the same general composition, varying only in the specific target sequences corresponding to five distinct sequences in the GT1 endogenous promoter sequence), and iii) the pollen-specific promoter PG47 PRO driving
30 expression of a maize-optimized CYS4 gene.

There are three components to the Cas9 mediated pollen-inhibitor system:

First, expressing a catalytically inactive double mutant of CAS9 (referred to as dCas9 and containing mutations in the RuvC1 and HNH nuclease domains) which can still associated with a guide-RNA and bind to a specific genomic
 5 sequence but without inducing a double strand break. While it's been demonstrated in *E. coli* that simple binding of dCas9 protein to a promoter can interfere with transcription (see Qi et al., 2013, Cell 152:1173-1183), the transcriptional interference can be enhanced, as described herein, namely by fusing a second component of the system to dCas9. The second component of the system
 10 comprises at least one repressor peptide that can be fused to dCas9. These repressor peptides comprise highly conserved motifs, such as "EAR-motifs" (with consensus signatures such as LxLxL or LxLxPP) that actively repress transcription (see Kagale & Rozwadowski 2011, Epigenetics 6:141-146). By fusing the repressor to dCas9, and expressing the protein behind a pollen-specific promoter, and using
 15 the third component, namely at least one expressed guide –RNA that direct the dCas9~LxLxPP fusion protein to an endogenous promoter (or multiple endogenous promoters) whose encoded protein(s) is(are) required for either pollen development or pollen tube growth, the result will be non-viable pollen.

Examples of genes that can be targeted to produce non-viable pollen;

20 a) A GT1 (glycosyltransferase gene) gene. It has been demonstrated in rice that knocking out the GT1 gene results in non-viable pollen. The maize ortholog of the rice GT1 gene (glycosyltransferase 1) is located on the short arm of chromosome 3 relatively near the centromere. Figures 5-A and 5-B show how dCAS9~RP can be directed to bind the maize GT1 promoter and inhibit pollen
 25 function. This specific example is for a single gene that when transcriptionally-repressed will result in pollen inhibition. The next two examples (described in "b" and "c" below) describe how using a single locus which contains the pollen-specific dCAS9~RP and an expression cassette producing multiple guide RNAs directed at multiple promoters in a gene family ("b") or duplicated genes ("c") can result in non-
 30 viable pollen.

b) It has been documented that in the triple-mutant oas-tLABC (in which the A, B and C family members for the gene encoding O-acetylserine(thiol)lyase are

knocked out, pollen germination does not occur or is so impaired as to be non-functional (see Birke et al., 2013, Plant Physiol 163:959-972). However, trying to create a triple-mutant and have all three mutant isoforms segregate together is impractical. By using a pollen-specific promoter to express *dcas9~LxLxPP* and the
 5 guide-RNA's that will target *dcas9~LxLxPP* to the promoters of *oar-tlA*, *oar-tlB* and *oar-tlC* – pollen tube growth in that pollen grain is blocked, and because these are expressed only in the pollen, there are no whole-plant pleiotropic effects.

c) This system can be used for any gene whose encoded product is essential for pollen viability and function. Another example of maize genes whose down-
 10 regulation will result in non-viable pollen are the maize chalcone synthase genes *Whp* (white pollen) and *C2* which both encode a chalcone synthase protein necessary for pollen viability [Coe et al., 1981. J Heredity 72:318-320; Franken et al., 1991. EMBO J. 10(9):2605-2612].

EXAMPLE 9.

15 Use of a two-component expression system to control pollen-inhibition in the first of a two-step screening process, and a conditional negative marker as the second screen to rapidly identify introgressed trait loci in two generations.

Two component transactivation expression systems have been used for many years in plants, for example, see Schwechheimer et al., 1998, Plant Mol. Biol.
 20 36:195-204) in which a fusion protein consisting of the GAL4 DNA-binding domain and the herpes simplex virus PV16 activation domain is expressed behind a promoter such as CaMV 35S, which binds to upstream activation sequences (UAS) in front of a minimal -45 CaMV promoter and the reporter gene beta-glucuronidase. However, it is also known that the GAL4 UAS is methylated in plants which inhibits
 25 binding (Gälweiler et al., 2000, Plant J. 23:143-157). As an alternative to GAL4~VP16, the use of alternative DNA-binding domains such as LEXA (REF) fused to plant activation domains and the cognate UAS sequence for LEXA make a good alternative for two component expression in plants (Boddepalli et al., US2013/055791). One such plant transcriptional activation domain comes from the
 30 Arabidopsis CBF1A protein (Stockinger et al., 1997, PNAS 94:1035-1040; Wang et al., 2005, Plant Mol. Biol. 58:543-559).

A two-step (two-generations) method comprising a two-component transactivation system to drive expression of a pollen-inhibitor gene (step one), along with a pollen-specific conditional-inhibitor gene (step two) can be used to break linkage on both sides of an internal trait locus in two successive crosses. In the constructs described herein, moLEXA is a maize-optimized gene (SEQ ID NO: 97) which encodes the DNA binding domain of the Escherichia coli LEXA protein (LEXA described in Brent & Ptashne, 1985, Cell 43:729-736; see SEQ ID NO: 98), UAS is the upstream activation sequence to which the LEXA fragment binds (SEQ ID NO: 99), and CBF1A is the polynucleotide sequences encoding the activation domain from the Arabidopsis CBF1A transcription factor (SEQ ID NO: 100).

An RTL exists at Chr1-52cM in Pioneer Inbred Line 3 is used to introduce a trait expression cassette (labeled "GAT" in Figure 8) such as UBI1ZM PRO (SEQ ID NO: 19), UBI1ZM INTRON (SEQ ID NO: 20), GAT891G3 (SEQ ID NO: 26) and the PINII TERM (SEQ ID NO: 8) and events are selected on medium containing glyphosate. In a second transformation, CAS/CRISPR-mediated integration is used (See Example 3) to introduce three expression cassettes at Chr1-50 cM; PG47 PRO::moLEXA~CBF1A::IN2-1 TERM + UBI1ZM PRO:UBI1ZM INTRON::CODA::PINII TERM + UBI1ZM PRO:UBI1ZM INTRON::NPTII::PINII (SEQ ID NO: 102) and events are selected on medium containing G418. In a third transformation, CAS/CRISPR-mediated integration is again used to introduce three expression cassettes at Chr1-54 cM; 5x(UAS):-45 35s PRO::ZM-AA1::PINII TERM + UBI1ZM PRO:UBI1ZM INTRON::CODA::PINII TERM + UBI1ZM PRO:UBI1ZM INTRON::PMI::PINII (SEQ ID NO: 103) and transgenic events are selected on medium containing mannose. The resultant 3-way linkage is shown in Figure 8. Once this linkage has been generated in Pioneer Inbred Line 3, inbreds into which the GAT trait will be introgressed (for example, Pioneer Inbred Line 2) are used as pollen donors onto silks of the PH2HT germplasm containing the 3-way linkage. The resultant F1 hybrid is then used as the pollen donor back onto the recurrent parent (Pioneer Inbred Line 2). Pollen grains containing both the transgenic locus at 50 cM (expressing the LEXA~CBF1A transactivator) and the transgenic locus at 54 cM containing the 5xUAS::ZM-AA1 expression cassette will be inhibited and will not transmit to the next generation. However in pollen grains in which either flanking

transgenic locus (50 or 54 cM) was lost due to meiotic recombination, the pollen will be viable and transmit to the next generation (at a frequency of about 1/50 progeny). The progeny that lost either flanking transgenic locus (and that are still glyphosate resistant) are grown to maturity and again used again as the pollen donor back onto the recurrent parent (back onto Pioneer Inbred Line 2). Progeny seed are germinated in water containing both glyphosate and 5-fluorocytosine and only progeny which contain the glyphosate trait but have lost both flanking loci will be viable. As an alternative to using a conditional negative marker such as coda, a seed specific promoter such as LTP2 PRO can be used to drive expression of a fluorescent protein (i.e. DS-RED2 or ZS-YELLOW1 N1) and progeny seed expressing the fluorescent protein can be discarded before germinating seed in the presence of glyphosate. Seed with no fluorescence that produce glyphosate-resistant seedlings contain the trait without either flanking locus.

EXAMPLE 10

Using a single pollen-inhibitor locus to screen for accelerated trait introgression of a telomeric trait locus.

Illustrated in Figure 1 and described below is an example of how a single pollen-inhibitor gene can be located near a trait of interest, and its use for accelerated trait introgression of a telomeric trait locus of maize.

Figure 1 shows a schematic of a crossing scheme used to establish and perform accelerated introgression for a telomeric trait locus, the trait locus being created through for example, but not limited to, either random integration or through targeted integration (for example, but not limited to, through targeted integration using Site Specific Integration, targeted meganucleases, or CAS/CRISPR technologies). Step 1: Choose an inbred amenable to transformation, and identify accessions that contain a trait locus and a "targeted-integration site", the trait locus being located at a position close to the telomere (labeled as "1") and the targeted-integration site being proximal (closer to the centromere, labeled as "2"), the distance between the target site 1 and Site 2 being within a close genetic distance, for example 1cM. Step 2: In two separate transformation experiments, introduce a trait gene (for example, UBI::GAT:pinII, abbreviated as GAT in the figure, which confers resistance to glyphosate) into the trait locus (Site 1), and using targeted

- integration, introduce a Pollen-inhibitor gene (for example PG47::ZmAA1::pinII, abbreviated as PI in the figure) into Site 2. Step 3: Carry pollen from the GAT-containing plant to PI-containing plant and screen the progeny by molecular markers in order to identify plants containing both GAT and PI. Step 4: Carry pollen from the
- 5 Wild Type inbred plant to the plants containing both GAT and PI to establish the linkage between Site 1 and Site 2. Use molecular markers to screen the progeny in order to identify plants in which GAT and PI are genetically linked (1% of the progeny at a genetic distance of 1cM). Step 5: Carry pollen from different Elite inbreds (into which you wish to introgress the GAT trait) onto ears of plants
- 10 containing the linked GAT and PI. Use molecular markers to identify F1 progeny that contain the linked GAT and PI. Step 6: Carry pollen from the F1 progeny containing linked GAT and PI back to the Recurrent Parent (i.e. the Elite inbred of step 5) to generate a progeny pool.
- a. If the linkage between GAT and PI is broken during meiosis then the
 - 15 pollen is viable and GAT can be transmitted to progeny.
 - b. Spray with glyphosate to eliminate any progeny derived from wild-type pollen grains.
 - c. The surviving plants will contain the GAT locus with minimal linkage drag from the transformation (donor) inbred.

20

A. Introducing the pollen-inhibitor gene - alpha amylase - in close proximity to a trait locus of interest (GAT, glyphosate resistance trait) in the genome of a maize plant.

- A Pioneer Inbred Line1 (PHN46) (the target line) was identified that comprised a pre-existing transgenic SSI target site located at 150.7 cM on
- 25 chromosome 10 containing a first expression cassette comprising a ubiquitin promoter driving a phosphomannose isomerase (PMI) and a pin II terminator, wherein the PMI was preceded by a FRT1 recombination site (UBI PRO::FRT1::PMI::pinII) linked to a second expression cassette comprising an actin promoter driving a moPAT selectable marker and a pin II terminator, followed by a
- 30 FRT87 recombination site (ACTIN PRO:: moPAT::pinII-FRT87). The target line was then used as the transformation target for particle-gun-mediated delivery and introduction of a GAT-resistance expression cassette between two dissimilar FLP-

recombinase sites (FRT sites FRT1-NPTII::PINII TERM + 3X(35S ENH):UBI1ZM
 PRO:UBI1ZM INTRON::GAT891G3::UBQ3 TERM + LTP2 PRO::DS-RED2::GZ-
 W64a TERM-FRT87(SEQ ID NO:44), along with a separate plasmid cassette
 (comprising a ubiquitin promoter driving a FLP recombinase terminated by a pinII
 5 terminator , UBI1ZM PRO:UBI1ZM INTRON:FLPM::PINII TERM (bp 411 to bp 4012
 of PHP5096, SEQ ID NO:50), which resulted in RMCE and the replacement of PMI
 & moPAT by GAT in the pre-existing transgenic SSI target site located at 150.7 cM
 on chromosome 10. At a position 0.5 cM proximal (150 cM on chromosome 10), an
 SSI donor sequence (SEQ ID NO:46) containing FRT1-NPT-II:pinII + 35S
 10 ENH:LTP2 PRO::TAGBFP::GZ-W64A TERM + ZM-PG47-PRO::Zm-AA1::IN2-1
 TERM-FRT87 (comprising a PG47 promoter driving the Zea mays alpha amylase
 (Zm-AA1) gene) was also introduced via particle-gun-mediated RMCE in a separate
 transformation experiment with Pioneer Inbred Line 1.

To confirm that the PG47::AA1::pinII expression cassette targeted to Chr10-
 15 150 cM resulted in pollen tube inhibition, we first verified that AA1 protein was being
 produced and then assessed the efficacy to inhibit pollen tube growth. For the first
 step (assessing expression), protein extracted from segregating pollen produced by
 eight SSI-targeted events (all at this same location) was analyzed by ELISA to
 assess AA1 protein levels. In these assays, wild-type Pioneer Inbred Line 1 pollen
 20 was used as the negative control and no AA1 protein was detected (labeled "Inbred
 1" in Figure 10). However, for the eight transgenic events containing the
 PG47::AA1::pinII, AA1 protein was detected at levels between 66 and 128 PPM
 (labeled 1-8 in Figure 10). The transgenic events that were determined by PCR to
 be clean single-copy "Recombinase-mediated cassette exchange" events were
 25 tested for efficacy of the AA1 expression cassette to inhibit pollen tube growth and
 fertilization. This was assessed by using these AA1 events as the pollinators back
 onto wild-type Pioneer Inbred Line 1 silks and analyzing progeny for the
 transmission of the AA1 locus (Figure 11). In this experiment, 1688 F1 progeny
 plants were analyzed by PCR for the presence or absence of the ZmAA1 and NPTII
 30 genes. Of the 1688 DNA samples analyzed, the DNA concentration was too low to
 perform the NPTII assay in 32 samples, was too low for the ZmAA1 assay in 14
 samples, and was too low for both assays in 4 samples. Excluding samples for

which the DNA concentration was too low to insure a reliable PCR reaction, of the 1588 F1 progeny that provided adequate DNA levels, all were null for AA1 and NPTII (see Figure 11). Thus, the expression of alpha amylase in maize plant comprising the expression cassette PG47-PRO::Zm-AA1::pinII inhibited pollen tube growth as expected (with a low escape rate) and plants from these events were subsequently used as the female in crosses with plants containing the GAT locus in order to establish linkage. The targeted-integration event containing GAT + DsRED (at genetic position Chr10-250.5 cM) was also evaluated for proper Mendelian inheritance and when outcrossed to wild-type Pioneer Inbred Line 1 exhibited the expected 1:1 segregation ratio.

B. Use of a pollen-inhibitor gene alpha amylase (AA1) to break linkage with a glyphosate resistance trait (GAT) on chromosome ten of maize for accelerated trait introgression.

An efficacious event containing AAI + BFP at Chr10-250 cM (referred to as PI in Figure 1) and the event containing GAT + DsRED at Chr10-250.5 (referred to as GAT in Figure 1) were crossed together (Figure 1, Step 2), and progeny were identified using PCR that contained both transgenic loci (Figure 1, Step 3). The progeny plants containing both loci were then pollinated with wild-type Pioneer Inbred Line 1 (such as PHN46) pollen, and the next generation of progeny were again analyzed by PCR for both loci (Figure 1, Step 4); of 1146 plants screened for the AA1 and GAT loci, 5 plants were found to contain both (contained both loci linked on the same chromosome). Based on this data for Pioneer Inbred Line 1 (PHN46), it appeared that the genetic distance between the two loci was 0.4 cM. All the steps up to this point were necessary to first create the two independent targeted-integration events containing either AA1/BFP or GAT/DsRED, and then using conventional breeding methods to link the two loci, which represented the required preparatory work before actually conducting the evaluation as to whether this configuration would aid in accelerating the introgression process.

The next step of the experiment was the first step in testing whether this closely linked pollen-inhibitor locus would facilitate more rapid and more precise introgression into another inbred. First, Pioneer Inbred line 1 plants containing the

linked AA1 and GAT loci were crossed to Pioneer inbred Line 2 (such as inbred line PHH5G, the inbred into which the trait would be introgressed), using Pioneer inbred Line 2 as the pollen donor and producing the first filial generation (more commonly called the F1 hybrid) in the first step of the introgression process (Step 5 in Figure 1). The resultant F1 hybrid plants were then used as the recurrent parent to pollinate wild-type Pioneer Inbred line 2 (PHH5G) (Step 6) to produce BC1F1 progeny (BackCross1 after the filial generation). These BC1F1 progeny were then analyzed for inheritance and introgression of the trait locus into the Pioneer Inbred Line 2 Chr10. This analysis consisted of three steps, each providing a progressively greater level of resolution. First, 4800 BC1F1 progeny were screened by PCR and 28 progeny were found to contain the GAT/DsRED locus at Chr10-150.7 cM with no AA1/BFP locus, providing a first indication that meiotic recombination had occurred during pollen formation between Chr10-150.5 and Chr10-150 cM, producing the type of BC1F1 offspring indicated at the bottom of Step 6 (Figure 1) in which the progeny plant now carried a copy of chromosome 10 in which only the telomeric segment containing the GAT trait was from the original Pioneer Inbred Line 1 donor and the remainder of this chromosome had a high likelihood of being entirely Pioneer Inbred Line 2. In the second analytical step, the frequency of converting the remainder of chromosome ten to Pioneer Inbred Line 2 (i.e. from position ~150.7 cM to 0 cM at the other end of the chromosome) was determined using 2 restriction-enzyme Genotype-by-Sequencing (GbS) on all 28 progeny that had lost the GAT/DsRED locus. Three replicates were run for each progeny plant to validate the consistency of the results, and for all 28 progeny the three replicates corroborated each other. For the 28 progeny, 22 events contained more than 80% of Pioneer Inbred Line 2 Chr10 (except for both telomeric ends where recombination appeared to be more frequent and remnant Pioneer Inbred Line 1 chromosome persisted), and 13 plants contained more than 90% Pioneer Inbred Line 2 chromosome. An examples of GbS results for a progeny plant carrying the GAT/DsRED locus within a small segment of Pioneer Inbred Line 1 chromosome on the end of the long arm of Chr10, followed by a small segment of Pioneer Inbred Line 2 chromosome, and the remainder of Chr10 being Pioneer Inbred Line 1 (Figure 12) represents one of the 6 progeny plants that were discarded due to the high percentage of Pioneer Inbred

Line 1 Chr10 that remained. In contrast, Figure 13 shows the GbS results from one of the desirable plants in which the majority of Chr 10 has been successfully converted to Pioneer Inbred Line 2 in just one generation. Finally, in the third analytical step, haplotype analysis was performed, using nine pairs of diagnostic SNPs between 132 cM and 155.0 cM surrounding the trait locus at 150.7 cM. As seen in Figure 14, for each of the 28 BC1F1 plants analyzed, the Pioneer Inbred Line 1 haplotypes are depicted as black bars and the Pioneer Inbred Line 2 haplotypes are depicted in grey. Genomic DNA from wild-type Pioneer Inbred Line 2 and Pioneer Inbred Line 1 contained only the expected wild-type SNPs for each inbred (grey or black bars, respectively). Of the transgenic BC1F1 plants analyzed, 25 contained Pioneer Inbred Line 1 chromosome at the distal end of the chromosome including the GAT/DsRED trait locus at 150.7 cM, with recombination occurring somewhere between the trait locus and the first proximal haplotype marker at approximately 149.3 cM, at which point the chromosome converts to Pioneer Inbred Line 2. Surprisingly, three plants contained chromosome 10 that was predominantly Pioneer Inbred Line 2 and where a double recombination had occurred on either side flanking the trait locus, with Progeny Number 1 showing evidence of a proximal cross-over within 0.7 cM of the trait and a distal cross-over within 1.5 cM on the other side, and Progeny 2 and 15 in which two crossovers occurred within 0.7 cM on the proximal side and somewhere between 1.5 and 4.0 cM away from the trait on the distal side.

These results demonstrated that using a closely-located pollen-inhibitor locus next to a trait of interest could be effectively used to rapidly (with one generation from the F1 hybrid) screen and identify progeny that had converted the transgene-carrier chromosome (Chr10) to the recurrent parent Pioneer Inbred Line 2.

C. Use of the AA1 pollen-inhibitor locus for accelerated trait introgression of a telomeric trait located at 0.9 cM on the short arm of Chromosome one.

A trait locus on Chr 1 of Pioneer Inbred Line 1 (PHN46) at genetic position 0.9 cM contained both a moPAT (maize-optimized phosphinithricin acetyl transferase) and PMI (phosphomannose isomerase) (illustrated as GAT on Figure 1, step 2). Using Site-Specific Integration, a pollen-inhibitor locus (illustrated as PI on

Figure 1, step 2) was introduced into Pioneer Inbred Line 1 at a genetic position of 2.3 cM on Chr1 (UBI::NPTII::pinII and PG47 PRO::ZM-AA1::pinII) and the two loci were genetically-linked using conventional breeding methods (Figure 1, Steps 3 and 4).

5 Pollen from wild-type Pioneer Inbred Line 2 (PHH5G) was used to pollinate ears of the doubly-linked (Traits + AA1) Pioneer Inbred Line 1 to produce the F1 hybrid (Figure 1, Step 5). The F1 hybrid was then used to pollinate the recurrent parent (RP) Pioneer Inbred Line 2 (Figure 1, Step 6). Of 3587 progeny screened by PCR, 42 contained the trait locus but no longer contained the AA1, again
10 demonstrating that the pollen-inhibitor could be effectively used to rapidly screen for progeny in which Chr1 had undergone a meiotic recombination, potentially creating a predominantly Pioneer Inbred Line 2 Chr1 with <2.3 cM of telomeric Chr1 from Pioneer Inbred Line 1 containing the transgenic trait locus.

Both of the above examples on the long arm of Chr10 and the short arm of
15 Chr1 demonstrated the ability to use pollen-inhibitor loci to rapidly and precisely introgress a transgenic trait locus into a new inbred with a minimum of yield drag (unwanted flanking donor inbred chromosome segments remaining next to the introgressed trait).

EXAMPLE 11.

20 Use of two sequential screenings using a pollen-inhibitor locus on one side of the trait locus and a locus encoding a visible kernel phenotype on the other side, in order to rapidly identify progeny in which the linkage had been broken on both sides of the trait.

An internal trait locus containing at least one trait gene of interest in the
25 Pioneer Inbred Line 1 (such as PHN46) is located at Chr1-51.8 cM. Using Cas9-mediated targeted integration, a pollen tube inhibitor expression cassette (PG47::Zm-AA1::pinII) is positioned at Chr1-50.8 cM, and in a second round of CAS9-mediated targeted integration, a seed-specific color marker (LTP2 PRO::CRC::pinII) is introduced at Chr1-52.8 cM. The seed is bulked up by
30 pollinating with the wild-type Pioneer Inbred Line 1.

To begin the introgression process, ears of the triple-linked transgenic inbred are pollinated with pollen from the various inbreds (for example, Pioneer Inbred Line

2, such as PHH5G) into which trait of interest will be introgressed to produce F1 hybrids. The F1 hybrids are then used as the male to pollinate the recurrent parent (i.e. Pioneer Inbred Line 2) and the BC1F1 seed are examined. Kernels are separated into two pools; the red-kernel phenotype imparted by the anthocyanin fusion-gene CRC and yellow kernels (non-red). Because of the pollen-inhibitor locus that was originally on the distal flank of the trait, any F1 pollen grains containing this locus would not form pollen tubes, and only wild-type pollen grains or transgenic pollen grains that had lost the AA1 locus due to meiotic recombination would produce a pollen tube and hence BC1F1 progeny. Non-red kernels are germinated in the presence of herbicide (which kills all the progeny derived from wild-type pollen) and only the non-red kernels (that had lost the color marker due to meiotic recombination) will survive.

If no surviving progeny are recovered in the first BC1F1 screening, the red BC1F1 seeds can be germinated and used to again pollinate the wild-type recurrent parent (i.e. Pioneer Inbred Line 2). The resultant BC2 seed are again separated into red and non-red kernels and the yellow kernels are germinated in herbicide. Resultant plants that germinate in the presence of herbicide have broken linkage on both sides of the trait and the trait has been introgressed in the RP with < 1.0 cM of remaining Pioneer Inbred Line 1 chromosome on either side of the trait.

EXAMPLE 12

Using a flanking pollen-inhibitor proximal to a telomeric trait along with a meiotically-expressed targeted nuclease between the pollen-inhibitor and the trait locus to simulate targeted recombination and chromosome exchange.

A trait locus in Pioneer Inbred line 1 (such as PHN46) containing a moPAT and PMI at Chr1-0.9 cM is produced, and a T-DNA containing a pollen-inhibitor expression cassette is positioned at Chr1-1.0 cM (0.1 cM proximal to the trait) using Cas9/CRISPR systems. In addition to the pollen-inhibitor expression cassette (PI), the T-DNA contains a 2 kb spacer sequence, an expression cassette containing a meiosis-specific promoter such as the SPO11 PRO in front of Cas9 (CAS9 locus) and an expression cassette driving expression of a guide-RNA that will target a Double-Stranded Break in between the T-DNA and the trait locus.

After establishing these two linked loci (the trait locus and the PI/CAS9 locus) on the same chromosome, pollen from various inbreds (for example Pioneer Inbred Line 1) is used to pollinate the ears of Pioneer Inbred Line 1 comprising the Trait—PI/CAS9 to produce the F1 hybrid. The F1 hybrids are grown to maturity and then
5 used to pollinate the recurrent parent, Pioneer Inbred Line 2. BC1F1 progeny are screened for introgression of the Trait Locus into the Pioneer Inbred Line2Chromosome 1, which normally is predicted to occur in 1/1000 progeny, based on a genetic distance between the two loci of 0.1 cM. However, because of the targeted cutting activity of the CAS9 protein during meiosis, targeted meiotic
10 recombination between the trait and the pollen-inhibitor is stimulated resulting in a higher frequency of progeny that have introgressed the trait into the Pioneer Inbred Line 1 chromosome 1 at very close genetic distance (i.e. < 0.1 cM).

In addition to using Cas9, other double strand break inducing systems can be used in the methods described herein. One can express a fusion between a well-
15 established DNA-binding-domain (DBD) such as GAL4 or LEXA and a nuclease (or meganuclease; abbreviated MN) such as I-SceI, I-CreI, I-DmoI, PI-SceI, PI-PfuI, FokI, One can also use the DBD fused to Spo11 in this type of a scenario. One can include the pollen-inhibitor (or the red-aleurone marker) in the T-DNA followed by a meiosis-specific promoter driving expression of fusion protein (comprising a DNA-
20 binding domain and a meganuclease (or a topoisomerase), a 2-3 kb spacer and then the DNA sequence that is bound by the DNA-binding-domain), which positions the MN at this cleavage site. This provides the pollen-inhibitor (or the seed color marker) to be used as the screening method(s) to identify the progeny in which the linkage has been broken within a specified nearby genetic interval, a meiosis-
25 specific catalyst to create double-strand breaks, and the cutting target site provided by the target sequence of the DNA-binding domain. Targeted double-strand breaks in this region during meiosis will stimulate localized homologous recombination (crossovers) and the pollen and/or seed screening tools will permit rapid identification.

THAT WHAT IS CLAIMED

1. A method for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
 - 5 (a) providing a first plant having within a genomic window at least one trait gene of interest integrated into a first target site located proximal to a telomere, wherein said genomic window is about 10 cM in length, wherein said first plant does not comprise a pollen-inhibitor gene;
 - (b) breeding to said first plant a second plant, wherein said second plant
10 comprises in said genomic window a pollen-inhibitor gene integrated into a second target site located proximal to both the telomere and the trait gene of interest of (a); and,
 - (c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said
15 pollen-inhibitor gene are genetically linked.
2. A method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
 - (a) providing a first plant having within a genomic window a first pollen-
20 inhibitor gene integrated into a first target site, wherein said genomic window is about 10 cM in length;
 - (b) breeding to said first plant a second plant having a trait gene of interest integrated into a second target site within said genomic window;
 - (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor
25 gene and said trait gene of interest in said genomic widow;
 - (d) providing a third plant having a second pollen-inhibitor gene integrated into a third target site within said genomic window;
 - (e) breeding to said third plant a fourth plant, wherein said fourth plant comprises a pollen-inhibitor maintainer (PIM) gene;
 - 30 (f) selecting a progeny plant from step (e) comprising said second pollen-inhibitor gene and said pollen-inhibitor maintainer (PIM) gene; and,
 - (g) cross pollinating the progeny plant of (c) with the progeny plant of (f) and selecting for a progeny plant that comprises said first pollen-inhibitor gene

said trait gene of interest and said second pollen-inhibitor gene, wherein said first pollen-inhibitor gene, said trait gene of interest and said second pollen-inhibitor gene are genetically linked.

- 5 3. The method of claims 1-2, wherein the first target site and the second target site is selected from the group consisting of a recombinase target site, a transgenic SSI target site, a single-strand-break-inducing-agent target site, and a double-strand-break-inducing-agent target site, or any one combination thereof.
- 10 4. The method of claims 1-2, wherein the double-strand-break-inducing-agent target site is a target site for an agent selected from the group of a Cas9 endonuclease, a zinc-finger nuclease, a Tal Effector nuclease (TALEN), a meganuclease, and an engineered endonuclease.
- 15 5. A method for introducing a pollen-inhibitor gene and a color marker gene in close proximity to a trait locus of interest in the genome of a plant, said method comprising:
- (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is about 10 cM in length;
- 20 (b) introducing into said genomic window of the plant of (a) a color marker gene;
- (c) breeding to the plant of (b) a second plant, wherein said second plant is a haploid inducer line capable of producing haploid embryos;
- (d) selecting haploid embryos from the plant of (c) and introducing into said
- 25 haploid embryos, a pollen-inhibitor gene; and,
- (e) producing a double haploid plant from the haploid embryo of (d).
6. A method for introducing two color marker genes in close proximity to a trait locus of interest in the genome of a plant, said method comprising:
- 30 (a) providing a first plant having a trait of interest located within a genomic window, wherein said genomic window is about 10 cM in length;

- (b) introducing into said genomic window of the plant of (a) a color marker gene;
- (c) breeding to the plant of (b) a second plant, wherein said second plant is a haploid inducer line capable of producing haploid embryos;
- 5 (d) selecting haploid embryos from the plant of (c) and introducing into said haploid embryos, a second color marker gene; and,
- (e) producing a double haploid plant from the haploid embryo of (d).
7. The method of claims 5-6, wherein color marker or the pollen-inhibitor gene are
10 introduced into a target site of a double-strand-break-inducing-agent.
8. A method of accelerated trait introgression in the genome of a plant, the method comprising:
- 15 (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one pollen-inhibitor gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;
- 20 (b) cross pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said trait of interest and said pollen-inhibitor gene; and,
- (d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of
25 interest.
9. A method of accelerated trait introgression in the genome of a plant, the method comprising:
- 30 (a) providing a first plant having within a genomic window at least one trait of interest located proximal to a telomere, and at least one color marker gene located proximal to both the telomere and the trait of interest, wherein said trait of interest and said color marker gene are genetically linked in said

genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;

(b) cross pollinating the first plant of (a) with pollen from a second plant; and,

(c) selecting a progeny plant from step (b) comprising said trait of interest site

5 and said color marker gene; and ,

(d) optionally, backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait of interest.

10 10. A method of accelerated trait introgression in the genome of a plant comprising:

(a) providing a first plant having within a genomic window at least one trait of interest , a first pollen-inhibitor gene, and a second pollen-inhibitor gene wherein said genomic window is about 5 cM in length, and wherein said trait of interest is flanked by said first and second pollen-inhibitor gene;

15 (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,

(c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,

(d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

20

11. A method of accelerated trait introgression in the genome of a plant comprising:

(a) providing a first plant having within a genomic window at least one trait of interest, a pollen-inhibitor gene and a color marker gene, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;

25

(b) cross-pollinating the first plant of (a) with pollen from a second plant; and,

(c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,

(d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest.

30

12. A method of accelerated trait introgression in the genome of a plant comprising:
- (a) providing a first plant having within a genomic window at least one trait of interest and at least a first color marker gene integrated into a first target site, a second color marker gene integrated into a second target site for, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;
 - (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
 - (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene; and,
 - (d) optionally, cross pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants comprising the trait of interest..
13. A method for introducing a pollen-inhibitor gene in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
- (a) providing a first plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said genomic window is about 5 cM in length and located within 0.1 cM to 100 cM of the telomere, and wherein said first plant does not comprise a pollen-inhibitor gene;
 - (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said second plant does not comprise said first transgenic target site; and,
 - (c) selecting a progeny plant from step (b) comprising said trait gene of interest and said pollen-inhibitor gene, wherein said trait gene of interest and said pollen-inhibitor gene are genetically linked in said genomic window.
14. A method of accelerated trait introgression in the genome of a plant, the method comprising:

- (a) providing a first progeny plant having within a genomic window at least a first transgenic SSI target site located proximal to a telomere and a second transgenic SSI target site located proximal to both the telomere and the trait gene of interest,, wherein said first transgenic SSI target site comprises at least one trait gene of interest, wherein said second transgenic SSI target site comprises a pollen-inhibitor gene, wherein said first transgenic target site and said pollen-inhibitor gene are genetically linked in said genomic window, wherein said genomic window is about 5 cM in length and located within 10 cM of the telomere;
- (b) cross pollinating the first plant of (a) with pollen from a second plant (elite inbred); and,
- (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said pollen-inhibitor gene;
- 15 15. The method of claim 8, further comprising :
- (e) backcrossing the progeny plant of (c) as the pollen donor onto a recurrent parent plant and selecting progeny plants comprising the trait gene of interest.
- 20 16. The method of any one of claims 13-14, wherein the pollen-inhibitor gene is selected from the group consisting of barnase, alpha amylase and KID.
17. The method of anyone of claims 13-14, wherein the trait gene of interest is selected from the group consisting of a selectable marker, a modified or edited native gene, a transgene, an altered double-strand-break target site, a native gene, or a transgenic SSI target site.
- 25 18. The method of anyone of claims 13-14, wherein the trait gene of interest and the pollen-inhibitor gene are genetically linked and located within 0.5 cM to 1 cM from each other.
- 30

19. A method for introducing two pollen-inhibitor genes in close proximity to a trait locus of interest in the genome of a progeny plant, said method comprising:
- (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said genomic window is about 5 cM in length;
 - (b) breeding to said first plant a second plant, wherein said second plant comprises in said genomic window a second transgenic SSI target site, wherein said second transgenic SSI target site comprises at least one trait gene of interest wherein said second plant does not comprise said first transgenic target site;
 - (c) selecting a progeny plant from step (b) comprising said first transgenic target site and said second transgenic target site genetically linked in said genomic window;
 - (d) providing a third plant having within said genomic window at least a third transgenic SSI target site and a pollen-inhibitor maintainer (PIM), wherein said third transgenic SSI target site comprises a second pollen-inhibitor gene; and,
 - (e) using the third plant of step (d) to pollinate the plant of step (c) and selecting a progeny plant wherein said first transgenic SSI target site, said second transgenic SSI target site, and said third transgenic SSI target site are genetically linked to each other.
20. A method of accelerated trait introgression in the genome of a plant comprising :
- (a) providing a first plant having a pollen-inhibitor maintainer gene and a genomic window, wherein said genomic window is about 5 cM in length and wherein said genomic window comprises at least a first transgenic SSI target site, a second transgenic SSI target site and a third transgenic SSI target site, wherein said first transgenic SSI target site comprises a first pollen-inhibitor gene, wherein said second transgenic SSI target site comprises at least one trait gene of interest, wherein said third transgenic

- SSI target site comprises a second pollen-inhibitor gene, wherein said second transgenic target site is flanked by said first transgenic SSI target site and said third transgenic SSI target site;
- (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- 5 (c) selecting a progeny plant from step (b) comprising said first transgenic target site, said second transgenic target site and said third transgenic target site.
21. The method of claim 20, further comprising cross pollinating the progeny plant
- 10 from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
22. A method of accelerated trait introgression in the genome of a plant comprising:
- (a) providing a first plant having within a genomic window a trait of interest and
- 15 at least a first pollen-inhibitor gene integrated into a first target site for a Cas9 endonuclease, a second pollen-inhibitor gene integrated into a second target site for a Cas9 endonuclease, wherein said genomic window is about 5 cM in length, and wherein trait of interest is flanked by said first and second pollen-inhibitor gene;
- 20 (b) cross-pollinating the first plant of (a) with pollen from a second plant; and,
- (c) selecting a progeny plant from step (b) comprising said first pollen-inhibitor gene, said trait of interest, and said second pollen-inhibitor gene;
23. The method of claim 22, further comprising cross pollinating the progeny plant
- 25 from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
24. A method for introducing two pollen- inhibitor genes in close proximity to a trait gene of interest in the genome of a progeny plant, said method comprising:
- 30 (a) providing a first plant having within a genomic window at least a first transgenic SSI target site, wherein said first transgenic SSI target site

- comprises at least one trait gene of interest, wherein said genomic window is about 5 cM in length;
- (b) introducing into the plant of (a) a first pollen-inhibitor gene, wherein said first pollen-inhibitor gene is genetically linked to said trait gene of interest
- 5 (c) selecting a progeny plant from (b) comprising the pollen-inhibitor gene genetically linked to the trait gene of interest.
- (d) breeding to said progeny plant of (c) a haploid-inducer plant and producing haploid immature embryos .
- (e) introducing into the haploid immature embryos of (d) a second pollen-
- 10 inhibitor gene linked to said trait gene of interest and said first pollen-inhibitor gene, wherein the trait gene of interest is flanked by the first and second pollen-inhibitor gene; and,
- (f) producing a homozygous plant from the haploid immature embryo of (e);
- 15 25. A method of accelerated trait introgression in the genome of a plant comprising
- (a) providing a first plant having a genomic window comprising a trait gene of interest flanked by and genetically linked to a first pollen-inhibitor gene and a second pollen-inhibitor gene, wherein said genomic window is about 5 cM in length;
- 20 (b) cross pollinating the first plant of (a) with pollen from a second plant ;and,
- (c) selecting a progeny plant from step (b) comprising said trait gene of interest flanked by and genetically linked to said first pollen-inhibitor gene and said second pollen-inhibitor gene.
- 25 26. The method of claim 25 further comprising cross-pollinating the progeny plant from step (c) to a recurrent parent plant and selecting progeny plants by exposing germinating seedlings to said selectable marker.
27. A plant comprising at least one trait gene of interest, a first recombinant DNA
- 30 construct comprising a first pollen-inhibitor gene, and a second recombinant DNA construct comprising a second pollen-inhibitor gene, wherein said first

recombinant DNA construct and said second recombinant DNA construct are genetically linked and flank said trait gene of interest.

- 5 28. A plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a pollen-inhibitor gene, wherein said trait locus of interest and said a pollen-inhibitor gene segregate independently from one another at a rate of about 10% to about 0.1%.
- 10 29. The plant of claim 27, wherein the first pollen-inhibitor gene, the second pollen-inhibitor gene and the trait gene of interest are located within 5 cM of each other.
- 15 30. The plant of claim 27, wherein the first pollen-inhibitor gene and the second pollen-inhibitor is selected from the group consisting of barnase, alpha amylase, KID, or any combination thereof
31. The plant of claim 27, wherein the second pollen-inhibitor gene is selected from the group consisting a non-conditional gene, a conditional gene and an inducible gene.
- 20 32. A plant comprising at least one trait gene of interest, a first recombinant DNA construct comprising a color marker gene, and a second recombinant DNA construct comprising a second color marker gene, wherein said first recombinant DNA construct and said second recombinant DNA construct are genetically linked and flank said trait gene of interest.
- 25 33. A plant comprising at least one trait locus of interest and a recombinant DNA construct comprising a color marker gene, wherein said trait locus of interest and said color marker gene segregate independently from one another at a rate of about 10% to about 0.1%.
- 30 34. The plant of claim 32, wherein the first color marker gene, the second color marker gene and the trait gene of interest are located within 5 cM of each other.

Figure 1




Figure 2




Figure 3




Figure 5-A




Figure 5-B




Figure 5-C




Figure 6




Figure 7




Figure 8




Figure 9-A




Figure 9-B




Figure 9-C




Figure 9-D




Figure 9-E




Figure 9-F




Figure 9-G




Figure 9-H




Figure 9-I




Figure 9- J




Figure 9-K




Figure 9-L




Figure 9-M




Figure 9-N




Figure 9-O




Figure 9-P




Figure 9-Q




Figure 9-R

UBI::moFLP::pinII + OS-ACTIN::WUS::PIN + UBI PRO & INTRON::ODP2::PINII

Figure 10




Figure 11




Figure 12




Figure 13




Figure 14




Figure 15




Figure 16





Figure 17



INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/022621

A. CLASSIFICATION OF SUBJECT MATTER
INV. A01H1/04 C12N15/82
ADD.

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)
A01H C12N

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

EP0-Internal, BIOSIS, Sequence Search, EMBASE, WPI Data

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	US 2013/198888 A1 (FALCO CARL [US] ET AL) 1 August 2013 (2013-08-01) cited in the application abstract paragraph [0006] - paragraph [0007] paragraph [0025] paragraph [0058] - paragraph [0059] paragraph [0075] paragraph [0080] - paragraph [0081] figure 1 ----- -/-	1-34



Further documents are listed in the continuation of Box C.



See patent family annex.

* Special categories of cited documents :

"A" document defining the general state of the art which is not considered to be of particular relevance

"E" earlier application or patent but published on or after the international filing date

"L" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)

"O" document referring to an oral disclosure, use, exhibition or other means

"P" document published prior to the international filing date but later than the priority date claimed

"T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention

"X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone

"Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art

"&" document member of the same patent family

Date of the actual completion of the international search

6 May 2016

Date of mailing of the international search report

25/05/2016

Name and mailing address of the ISA/

European Patent Office, P.B. 5818 Patentlaan 2
NL - 2280 HV Rijswijk
Tel. (+31-70) 340-2040,
Fax: (+31-70) 340-3016

Authorized officer

Mundel, Christophe

INTERNATIONAL SEARCH REPORT

International application No
PCT/US2016/022621

C(Continuation). DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	MOHAN S MURALI ET AL: "Identification of quantitative trait loci associated with resistance to foliar diseases in sorghum [Sorghum bicolor (L.) Moench]", EUPHYTICA, vol. 176, no. 2, November 2010 (2010-11), pages 199-211, XP002757386, ISSN: 0014-2336 abstract table 4	33
A	----- US 2015/059010 A1 (CIGAN ANDREW MARK [US] ET AL) 26 February 2015 (2015-02-26) the whole document	1-34
A	----- WO 2013/019411 A1 (DU PONT [US]; PIONEER HI BRED INTERNAT [US]; CIGAN ANDREW MARK [US]; F) 7 February 2013 (2013-02-07) the whole document	1-34
A	----- US 2014/283166 A1 (CHOMET PAUL S [US] ET AL) 18 September 2014 (2014-09-18) the whole document -----	1-34

INTERNATIONAL SEARCH REPORT

International application No.

PCT/US2016/022621

Box No. I Nucleotide and/or amino acid sequence(s) (Continuation of item 1.c of the first sheet)

1. With regard to any nucleotide and/or amino acid sequence disclosed in the international application, the international search was carried out on the basis of a sequence listing:
 - a. ☒ forming part of the international application as filed:
 - ☒ in the form of an Annex C/ST.25 text file.
 - ☐ on paper or in the form of an image file.
 - b. ☐ furnished together with the international application under PCT Rule 13~~ter~~.1(a) for the purposes of international search only in the form of an Annex C/ST.25 text file.
 - c. ☐ furnished subsequent to the international filing date for the purposes of international search only:
 - ☐ in the form of an Annex C/ST.25 text file (Rule 13~~ter~~.1(a)).
 - ☐ on paper or in the form of an image file (Rule 13~~ter~~.1(b) and Administrative Instructions, Section 713).
2. ☐ In addition, in the case that more than one version or copy of a sequence listing has been filed or furnished, the required statements that the information in the subsequent or additional copies is identical to that forming part of the application as filed or does not go beyond the application as filed, as appropriate, were furnished.
3. Additional comments:

INTERNATIONAL SEARCH REPORT

Information on patent family members

International application No

PCT/US2016/022621

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
US 2013198888 A1	01-08-2013	AR 089793 A1 CA 2862844 A1 CN 104245939 A EA 201491431 A1 EP 2807257 A1 US 2013198888 A1 WO 2013112686 A1	17-09-2014 01-08-2013 24-12-2014 30-01-2015 03-12-2014 01-08-2013 01-08-2013
US 2015059010 A1	26-02-2015	AU 2014308896 A1 AU 2014308898 A1 AU 2014308899 A1 AU 2014308900 A1 CA 2922040 A1 CA 2922046 A1 CA 2922089 A1 CA 2922090 A1 US 2015059010 A1 US 2015082478 A1 WO 2015026883 A1 WO 2015026885 A1 WO 2015026886 A1 WO 2015026887 A1	10-03-2016 03-03-2016 10-03-2016 10-03-2016 26-02-2015 26-02-2015 26-02-2015 26-02-2015 26-02-2015 19-03-2015 26-02-2015 26-02-2015 26-02-2015 26-02-2015
WO 2013019411 A1	07-02-2013	CA 2844101 A1 EP 2739739 A1 US 2014338070 A1 WO 2013019411 A1	07-02-2013 11-06-2014 13-11-2014 07-02-2013
US 2014283166 A1	18-09-2014	NONE	