



Howard Johnson's

## CRABGRASS CONTROL PLUS WITH 0.37% PRODIAMINE

FOR SELECTIVE PREEMERGENCE CONTROL OF GRASS & BROADLEAF WEEDS IN LAWNS AND ORNAMENTAL TURF GRASSES (EXCLUDING GOLF COURSE PUTTING GREENS), LANDSCAPE ORNAMENTALS, ESTABLISHED PERENNIALS & WILDFLOWER PLANTINGS. STOPS MANY UGLY WEED GRASSES BEFORE THEY GROW. PREVENTS CRABGRASS, ANNUAL BLUEGRASS, FOXTAIL, & MORE.

ACTIVE INGREDIENT:

Prodiamine: [N',N'-Di-n-propyl-2,4-dinitro-6-(trifluoromethyl)-m-phenylenediamine] . . . . .0.37%

INERT INGREDIENTS: . . . . . 99.63%

TOTAL: . . . . . 100.00%

EPA Reg. No. 32802-66

EPA Est. No. 32802-MO-1 (B)

First letter of bag code shows establishment where manufactured.

Manufactured by:  
Howard Johnson's Enterprises, Inc.  
700 W. Virginia St., No. 222  
Milwaukee, WI 53204

Read the entire "Directions for Use" and "Warranty Statements" before using this product. If terms are not acceptable, return the unopened prod-uct container at once to the seller.

### KEEP OUT OF REACH OF CHILDREN

## CAUTION

### PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS AND DOMESTIC ANIMALS

**CAUTION:** Causes moderate eye injury (irritation). Harmful if inhaled or absorbed through the skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust. Wash thoroughly with soap and water after handling. Prolonged or repeated skin contact with product may cause allergic reactions in some individuals. **ENVIRONMENTAL HAZARDS:** This product has low solubility in water. At the limits of solubility, this product is not toxic to fish. However, at concentrations above the level of water solubility, it may be toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent sites. Do not contaminate water when disposing of equipment washwaters.

First Aid	
If in eyes	<ul style="list-style-type: none"><li>Hold eye open and rinse slowly and gently with water for 15-20 minutes.</li><li>Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
If on skin or clothing	<ul style="list-style-type: none"><li>Take off contaminated clothing.</li><li>Rinse skin immediately with plenty of water for 15-20 minutes.</li><li>Call a poison control center or doctor for treatment advice.</li></ul>
If swallowed	<ul style="list-style-type: none"><li>Call a poison control center or physician immediately for treatment advice.</li><li>Have person sip a glass of water if able to swallow.</li><li>Do not induce vomiting unless told to do so by the poison control center or physician.</li><li>Do not give anything by mouth to an unconscious person.</li></ul>
If inhaled	<ul style="list-style-type: none"><li>Move person to fresh air.</li><li>If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably mouth-to-mouth, if possible.</li><li>Call a poison control center or doctor for further treatment advice.</li></ul>
Hot line number:	
Have the product container or label with you when calling a poison control center or doctor, or going for treatment. For information on this pesticide product (including health concerns, medical emergencies, or pesticide incidents), call the National Pesticide Telecommunications Network at 1-800-858-7378.	

Information regarding the contents and levels of metals in this product is available on the internet at <http://www.wa.gov/agr/>

### GUARANTEED ANALYSIS

Spreader setting to apply at the listed rate (lbs per 1000 sq ft)						
	Cone setting	3.0 lbs	4.0 lbs	4.6 lbs	6.2 lbs	9.3lbs
Accu-Pro 2000	6	L 1/2	M 1/2	N 1/2	O 1/2	Q 1/2
SR2000	6	L 1/2	N	O 1/2	P	R
Lesco 021820	0.25 Closed	D 1/2	E 1/2	F 1/2	G 1/2	I 1/2
Spyker 76-22		4.3	4.5	4.7	4.9	5.5
PSB Bigfoot	A.5	J 1/2	K 1/2	L 1/2	M	N 1/2
Lely HR	II-B	5.25	6	6.5	7	8.5
Vicon 203		15	17	19	21	27

Minimum and Maximum Application Rates (by Turf Grass Species) <sup>1</sup>		
Turf Species	Minimum Rate	Maximum Annual Rate
Bahiagrass, Centipedegrass Bermudagrass <sup>2</sup> , Zoysiagrass Seashore Paspalum St Augustinegrass Tall Fescue (including turf-type)	3.2 pounds per 1000 sq. ft	9.3 pounds per 1000 sq. ft. per year
Buffalograss Kentucky Bluegrass Perennial Ryegrass	3.2 pounds per 1000 sq. ft	6.2 pounds per 1000 sq. ft. per year
Creeping Red Fescue	3.2 pounds per 1000 sq. ft	4.6 pounds per 1000 sq. ft. per year
Creeping Bentgrass	3.2 pounds per 1000 sq. ft	4.0 pounds per 1000 sq. ft. per year

<sup>1</sup> These are the maximum rates per calendar year.  
<sup>2</sup> May be used on newly sprigged or plugged Bermudagrass at rates not to exceed 3.3 lbs / 1000 square feet. Newly sprigged or plugged Bermudagrass stolon rooting may be temporarily retarded. Suppression only of foxtail, goosegrass and rescuegrass due to reduced rates used in sprigging situations.

Use Rate	Months Before Overseeding		
lbs / 1000 sq ft	North	Transition	South
3 lbs	4	4	4
4 lbs	5	4	4
4.6 lbs	6	5	5
4.9 lbs	---	6	6
6.2 lbs	---	7	7
7 lbs	---	---	9
8 lbs	---	---	10
9.3 lbs	---	---	12

SCIENTIFIC NAME	COMMON NAME
Abies spp	Fir species** (Balsam, etc.)
Acer palmatum	Japanese Maple
A. platanoides	Norway Maple
Actinidia chinensis	Kiwi*
Agapanthus africanus	Lily-of-the-Nile (African Lily)
Arctostaphylos densiflora	Vine Hill Manzanita
Arctotheca calendula	Cape Weed
Aucuba japonica	Japanese Aucuba
Berberis gladywensis	Barberry
B. julianae	Wintergreen Barberry
B. mentorensis	Mentor Barberry
B. thunbergii	Japanese Barberry
Buxus microphylla	Japanese Boxwood
Callistemon viminalis	Weeping Bottlebrush
Calluna vulgaris	Scotch Heather
Carpobrotus edulis	Hottentot Fig (Ice Plant)
Cassia artemisoides	Feathery Cassia
Ceanothus rigidus	Wild Lilac
Chamaecyparis pisifera	False Cypress
Cleyera japonica	Cleyera
Citrus spp.	Citrus species*
Cornus florida	Flowering Dogwood
C. stolonifera	American Dogwood
Cortaderia selloana	Pampas Grass
Cotoneaster apiculatus	Cranberry Cotoneaster
C. buxifolius	Cotoneaster
C. dammeri	Bearberry Cotoneaster
C. microphyllus	Rockspray Cotoneaster
C. lataguss spp.	Hawthorne
Cupressus sempervirens	Italian Cypress
Delasperma alba	White Trailing Ice Plant
Delphinium spp.	Larkspur, Blue Elf
Dodonea viscosa	Hop Bush
Elaeagnus pungens	Silverberry
Euonymus fortunei	Wintercreeper
E. japonica	Evergreen Euonymus
E. kiautschovica	Spreading Euonymus
Fatsia japonica	Japanese Aralia
Forsythia intermedia	Border Forsythia
F. viridissima	Greenstem Forsythia
Gardenia jasminoides	Gardenia, Cape-Jasmine

SCIENTIFIC NAME	COMMON NAME
Hedera helix	English Ivy
Hibiscus rosa-sinensis	Chinese Hibiscus
Ilex cornuta	Chinese Holly
I. crenata	Japanese Holly
I. opaca	American Holly
Iris spp.	Iris species**
Jasminum nudiflorum	Winter Jasmine
Juniperus chinensis	Chinese Juniper
J. conferta	Shore Juniper
J. horizontalis	Creeping Juniper
Juglans sp.	Walnut*
Justicia brandegeana	Shrimp Plant
Lagerstromia indica	Crepe Myrtle
Ligustrum amurense	Amur Privet
L. japonicum	Japanese Privet
L. lucidum	Glossy Privet (Waxleaf)
Liriope muscari	Big Blue Lily, Silver Mound
Lonicera japonica	Japanese Honeysuckle
L. tatarica	Tatarian Honeysuckle
Maleophora luteola	Ice Plant
Malus sp.	Crabapple*
Nandina domestica	Heavenly Bamboo
Olea europaea	Olive*
Ophiopogon japonicus	Mondo Grass
Osteospermum truncosum	Trailing African Daisy
Oxydendrum arboreum	Sourwood
Persea americana	Avocado
Photinia fraseri	Frasier's Photinia (Redtip)
Picea spp.	Spruce species**
Pieris japonica	Lily-of-the-Valley Bush
Pinus brutia	Calabrian Pine
P. canariensis	Canary Island Pine
P. halepensis	Aleppo Pine
P. nigra	Austrian Pine
P. radiata	Monterey Pine
P. thunbergiana	Japanese Black Pine
P. strobus	Eastern White Pine
P. sylvestris	Scotch Pine
Pistachio sp.	Pistachio*
Pittosporum rhombifolium	Queensland Pittosporum

**STORAGE AND DISPOSAL:** Do not contaminate water, food, or feed by storage or disposal. **Storage:** Store in original container in a locked storage area out of reach of children and domestic animals. **Pesticide disposal:** Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility. **CONTAINER DISPOSAL:** Completely empty bag into application equipment. Nonrefillable container: Do not reuse or refill this container. Offer for recycling, if available. If not available then dispose of empty bag in a sanitary landfill, or by incineration, or, if allowed by State and local authorities, by burning. If burned, stay out of smoke.

Warranty

The directions for use of this product are believed to be adequate and must be followed carefully, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness, or other unintended consequences may result due to such factors as weather conditions, presence or absence of other materials, or the manner of use or application, all of which are beyond the control of Howard Johnson's Enterprises, Inc the manufacturer, or the seller. To the extent consistent with applicable law, the products sold to you are furnished "as is" by Howard Johnson's Enterprises, Inc. The manufacturer and the seller are subject only to the manufacturer's warranties, if any, which appear on the label of the product sold to you. Except as warranted by this label Howard Johnson's Enterprises, Inc, the manufacturer, or the seller makes no warranties, guarantees, or representations of any kind to the buyer or the user, either express or implied, or by usage of trade, statutory or otherwise, with regard to the product sold or use of the product, including, but not limited to, merchantability, fitness for a particular purpose or use, or eligibility of the product for any particular trade usage. To the extent consistent with applicable law, Buyer's or user's exclusive remedy, and Howard Johnson's Enterprises, Inc, the manufacturer's or the seller's total liability shall be limited to damages not exceeding the cost of the product. No agent or employee of Howard Johnson's Enterprises, Inc, or the seller is authorized to amend the terms of this warranty disclaimer or the product's label or to make a presentation or recommendation different from or inconsistent with the label of this product.

To the extent consistent with applicable law, Howard Johnson's Enterprises, Inc, the manufacturer, or the seller shall not be liable for consequential, special, or indirect damages resulting from the use, handling, application, storage, or disposal of this product or for damages in the nature of penalties, and the buyer and the user waive any right that they may have to such damages.

Do not apply to newly set sod until the following year. Application to turf stressed by drought, low fertility, or pest damage may result in turf injury. Disturbing the herbicide barrier with cultural practices (such as core aeration) may result in reduced weed control. Do not apply to putting greens, or areas where dichondra, colonial bentgrass, velvet bentgrass or annual bluegrass (*Poa annua*) are desirable species. Do not apply through any type of irrigation system. Do not apply aerially. Do not graze or feed livestock forage cut from treated areas.

**RATES OF APPLICATION:** This product is recommended for use on the turf grass species listed in the following table. The maximum amount of this product that may be applied per year is given for each turf grass species. Do not exceed the maximum yearly rate listed:

This product may be applied as a single application or in sequential applications to control weeds germinating throughout the year. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rates recommended for each turf species. The length of time of residual weed control provided by this product is related to the rate applied. Use the higher rates listed for longer periods of weed control.

**WHEN TO OVERSEED AFTER APPLICATION:** This product will inhibit germination of turf seed if overseeded too soon after application. Follow rates and intervals in table for best overseeding/ reseeding results.

**LANDSCAPE ORNAMENTALS** (including established perennials and wildflower plantings): This product may be applied for residual preemergence weed control in ornamentals. Apply at the rate of 3.2 to 9.3 lbs / 1000 square feet in fall and/or spring. Use higher rates of application for longer control periods. Sequential applications may be made so long as the total amount of product applied does not exceed the maximum annual application rate of 9.3 lbs / 1000 square feet per year.

**APPLICATION TIMING AND INFORMATION:** This product may be applied to newly transplanted and established ornamentals a broadcast, over the top or directed application. Around new transplants, delay application to allow soil to settle and water thoroughly before applying. Apply after cuttings form roots and become established. Apply before budding/grafting or after buds/grfts have taken to avoid any inhibition of the tissue union. This product is a preemergence herbicide and will not control emerged weeds. Most effective weed control in ornamentals will be obtained when this product is activated by at least 0.5 inch of rainfall or irrigation, or with shallow (1 to 2 inch) mechanical incorporation, prior to weed seed germination listed below are tolerant to this product. Best weed control is obtained when applied to soil free of clods, weeds, and debris such as leaves. Prior to application, control existing vegetation by hand weeding, cultivation or the use of an appropriate postemergence herbicide.

**TOLERANT ORNAMENTAL SPECIES:** This product will not harm most trees, shrubs, vines and flowers. The species listed below are tolerant to this product. Best results will be obtained when product is uniformly applied to the soil surface; avoid applying to ornamental shoots which may defeat the effort to get uniform distribution on the soil surface. This product may be applied over the top of these species. When plants are under stress (such as heat, drought or frost damage) some cultivars of listed plants may be sensitive to this product.

SCIENTIFIC NAME	COMMON NAME
P. tobira	Japanese Pittosporum
Podocarpus macrophyllus	Japanese Yew
Prunus laurocerasus	English Laurel
Prunus sp.	Almond, Apricot
Nectarine	Peach, Plum and Prune*
Pyracantha coccinea	Firethorn (Scarlet)
P. fortuneana	Firethorn
P. koidzumii	Firethorn
Pyrus sp.	Bradford Pear sp.
Quercus spp.	Oak species
Raphiolepis indica	Indian Hawthorne
Rhododendron	'Coral Bells'
(including Azalea)	'Formosa', 'Hino-crimson', 'PJM', 'Roseum Elegans'
Rosa banksiae	Lady Bank's Rose
Rosmarinus officinalis	Rosemary
Santolina virens	Santolina
Sedum album	Stoncrop
Syzygium paniculatum	Japanese Brushcherry
Taxus cuspidata	Japanese Yew
T. media	Yew
Thuja occidentalis	American Arborvitae
Trachelospermum asiaticum	Star Jasmine
Tsuga canadensis	Canada Hemlock
Viburnum japonicum	Japanese Viburnum
V. odoratissimum	Sweet Viburnum
V. plicatum	Japanese Snowball
V. rotundifolium	Canary Island Viburnum
V. tinus	Laurustinus
V. trilobium	Cranberry Bush
V. wrightii	Leatherleaf Viburnum
Vinca major	Greater Periwinkle, Vinca
Vinca minor	Common Periwinkle,
Vinca	
Vitis sp.	Grape*
Weigela florida	Old Fashioned Weigela
Yucca aloifolia	Spanish Bayonet, Yucca
Y. filamentosa	Adam's Needle, Yucca

\*Do not use on food producing trees, vines or plants. \*\*Not for use on container grown plants.