

# QUALI-PRO

## Imidacloprid 1G

### Nursery & Greenhouse Insecticide

*A systemic insecticide for use on ornamental plants in greenhouses, nurseries, and interior landscapes.*

**ACTIVE INGREDIENT:****% BY WEIGHT**

Imidacloprid: 1-[(6-Chloro-3-pyridinyl)methyl]-N-nitro-2-imidazolidinimine . . . . . 1.0%

**INERT INGREDIENTS:** . . . . . 99.0%

**TOTAL:** . . . . . 100.0%

EPA Reg. No. 53883-227-73220

EPA Est. No. 5905-IA-001

## KEEP OUT OF REACH OF CHILDREN CAUTION

#### FIRST AID

**IF IN EYES:**

- Hold eye open and rinse slowly and gently with water for 15-20 minutes.
- Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye.
- Call a poison control center or doctor for treatment advice.

**IF SWALLOWED:**

- Call a poison control center or doctor immediately for treatment advice.
- Have person sip a glass of water if able to swallow.
- Do not induce vomiting unless told to so by the poison control center or doctor.
- Do not give anything by mouth to an unconscious person.

**IF ON SKIN  
OR CLOTHING:**

- Take off contaminated clothing.
- Rinse skin immediately with plenty of water for 15-20 minutes.
- Call a poison control center or doctor for treatment advice.

#### HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact ProSAR at 1-800-308-5391 for emergency medical treatment information.

#### NOTE TO PHYSICIAN

No specific antidote is available. Treat the patient symptomatically.

See inside label booklet for additional precautionary statements.

**Manufactured for FarmSaver.com, LLC**

**4515 Falls of Neuse Road, Suite 300 • Raleigh, NC 27609**

**Net Weight: 30 lbs.**

**STOP - Read the label before use.**  
**Keep out of reach of children.**

**CAUTION**  
**PRECAUTIONARY STATEMENTS**  
**HAZARDS TO HUMANS AND**  
**DOMESTIC ANIMALS**

**CAUTION.** Harmful if swallowed or absorbed through skin. Avoid contact with skin, eyes, or clothing. Wash thoroughly with soap and water after handling. Keep children or pets off treated area until spray is dry.

**PERSONAL PROTECTIVE EQUIPMENT (PPE)**

**Applicators and other handlers must wear:**

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**ENGINEERING CONTROLS STATEMENTS**

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(4)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

**USER SAFETY RECOMMENDATIONS**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**ENVIRONMENTAL HAZARDS**

This product is highly toxic to aquatic invertebrates. Do not apply directly to water, or to areas where surface water is present or to intertidal areas below the mean high water mark.

Do not contaminate water when disposing of equipment washwaters.

This chemical demonstrates the properties and characteristics associated with chemicals detected in groundwater. The use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in groundwater contamination.

**DIRECTIONS FOR USE**

It is a violation of Federal law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

**AGRICULTURAL USE REQUIREMENTS**

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. Exception: If the product is applied by drenching, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof material such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, polyvinylchloride (PVC) or viton
- Shoes plus socks

## GENERAL INFORMATION

QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide will control insects on ornamental plants grown in nurseries and greenhouses. QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide is a systemic insecticide that is translocated upwards within the plant. For best results, QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide must be applied where the roots of the target plants can absorb the active ingredient. After application, irrigate moderately but thoroughly, being sure to avoid leaching and runoff from the container for at least three irrigations or 10 days whichever is longer.

### Use Precautions:

- Do not graze treated areas or use clippings from treated areas for feed or forage.
- Do not apply to soils which are waterlogged or saturated. This condition will not allow penetration of water into the root zone of the plants.
- Do not over-irrigate or allow excessive runoff to occur following an application of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide.
- Do not apply more than 40 lbs. of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide per acre per year.
- On plants with a production cycle of less than one year, application is not to exceed a frequency of more than once each 16 weeks for a particular plant. On stock plants and woody crops with a production cycle of greater than one year, application may not exceed once a year.

**Food Crops:** Treated areas may be replanted with any crop specified on an imidacloprid label, or with any crop for which a tolerance exists for the active ingredient. For crops not listed on an imidacloprid label, or for crops for which no tolerance for the active ingredient have been established, a 12-month plant-back interval should be observed.

**Insect Resistance:** After repeated use of a particular insecticide, some insects may develop a resistance. It is important that the use of this product conforms to resistance management practices established for the use area. Consult your Cooperative

Extension Service for resistance management strategies and recommended pest management practices for your area.

Do not make a foliar application of any chloronicotynyl insecticide following QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide soil application to the same crop.

**Woody Perennials:** Protection in woody perennials is slower than in herbaceous species and a delay of 2 or more weeks should be expected, with longer delays for larger plants. Because of this, applications to woody perennials should be made well in advance of expected insect activity.

**Incorporation:** Incorporation of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide can be accomplished by cultivation, irrigation, rainfall, mechanical placement, or by using mechanical soil or media mixing equipment.

**Bark Media:** When treating bark media with 30-50% or more bark content, the period of protection with QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide will be shortened.

## APPLICATION TO GRASSY AREAS IN NURSERIES

**Note:** QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide may not be used on commercial sod farms.

QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide will control soil-inhabiting pests in grassy areas in or around nurseries such as under or around field or container grown plants, on roadways or other grassy areas. The need for an application can be based on historical monitoring of the site, previous records or experiences, current season adult trappings or other methods. Applications may be made preceding the egg laying activity of the target pests and high levels of control may be achieved when applications are made preceding or during the egg laying period. For best results, make applications prior to egg hatch of the target pests, followed by sufficient irrigation or rainfall to move the active ingredient through the thatch.

PESTS CONTROLLED	
Annual bluegrass weevil ( <i>Hyperodes</i> spp.) Billbugs ( <i>Sphenophorus</i> spp.) Green June Beetle ( <i>Cotinis nitida</i> ) May or June beetle ( <i>Phyllophaga</i> spp.) Oriental beetle ( <i>Anomala orientalis</i> ) Northern and Southern masked chafers ( <i>Cyclocephala borealis</i> , <i>C. immaculata</i> , and/or <i>C. lurida</i> )	Asiatic garden beetle ( <i>Maladera castanea</i> ) European chafer ( <i>Rhizotrogus majalis</i> ) Japanese beetle ( <i>Popillia japonica</i> ) Mole Crickets ( <i>Scapteriscus</i> spp.) Black turfgrass ataenius ( <i>Ataenius spretulus</i> and <i>Aphodius</i> spp.)
PESTS SUPPRESSED	
Cutworms	Hairy Chinchbugs

### Grassy Area Use Precautions

- Applications must NOT exceed a total of 40 lbs. (0.4 lb. of active ingredient) per area per year.
- Applications should NOT be made when grassy areas are waterlogged or the soil is saturated with water because adequate distribution of the active ingredient cannot be achieved when these conditions exist.
- The treated grassy area must be in such a condition that the rainfall or irrigation will penetrate vertically in the soil profile.

### APPLICATION TO ORNAMENTALS GROWN IN FLATS, BENCHES, BEDS AND CONTAINERS.

QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide will suppress or control the following insect pests when applied to ornamentals grown in flats, benches, beds and containers in greenhouses and nurseries.

Adelgids Armored scale (suppression) Japanese beetles (adults) Leafminers Psyllids Soft scale Whiteflies Leafhoppers (including glassy-winged sharpshooter)	Aphids Fungus gnats (larvae only) <sup>1</sup> Lacebugs Mealybugs Root Weevil Thrips (suppression) <sup>3</sup> Leaf beetles (including Elm and Viburnum leaf beetles) Complex (such as Black Vine Weevil, Apopka Weevil, Citrus Root Weevil <sup>2</sup> )
<sup>1</sup> Fungus gnat larvae in the soil will be controlled by incorporation. QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide will not control adult Fungus gnat. Other foliar insect control is achieved by the uptake of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide from a healthy root system translocating the active ingredient up into the plant. <sup>2</sup> Citrus Root Weevil: For use on non-bearing nursery stock. <sup>3</sup> Thrips suppression on foliage only. Thrips in buds and flowers will not be suppressed.	

1 level teaspoon of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide = 4.1 grams  
3 teaspoons = 1 tablespoons

## HERBACEOUS AND WOODY PERENNIAL SPECIES

Herbaceous Species (one or two plants per pot)				
Container Size (inches)	Application Rates			
	Bulk Application		Topical Application	
	Pots/cu. yd. of mix	Bulk Rate lbs./cu. yard	Topical Rate (level teaspoonful)	Grams/pot
2.0 to 3.0	4500 to 3200	7.0 to 3.5	1/8 to 1/4	0.5 to 1.0
3.5 to 5.0	1600 to 1300	3.9 to 3.0	1/4 to 1/3	1.0 to 1.4
5.5 to 7.0	900 to 400	1.8 to 1.2	1/3 to 1/2	1.4 to 2.0
8.0 to 10.0	200 to 160	1.0 to 0.8	1/2 to 2/3	2.0 to 2.7
11.0 to 12.0	80 to 60	0.8 to 0.5	2/3 to 1.0	2.7 to 4.1
Woody Perennial Species or Herbaceous Species (three or more plants per pot)				
2.0 to 3.0	4500 to 3200	9.0 to 7.0	1/4 to 1/3	1 to 1.3
3.5 to 5.0	1600 to 1300	6.0 to 4.0	1/3 to 1/2	1.3 to 2.0
5.5 to 7.0	900 to 400	2.5 to 2.0	1/2 to 2/3	2.0 to 2.7
8.0 to 10.0	200 to 160	1.5 to 1.0	2/3 to 1.0	2.7 to 4.1
11.0 to 12.0	80 to 60	1.0 to 0.8	up to 1 1/2	4.1 to 6.1
<b>Bulk Rate Application:</b> Mix the required amount of QUALI-PRO IMIDACLOPRID 1G NURSERY & GREENHOUSE Insecticide per cubic yard of medium. Mix thoroughly before filling pots and transplanting. <b>Topical Application:</b> May be made at transplanting, repotting or on established containerized plants. Irrigate moderately after application into the root zone. Do not allow significant leaching and runoff for at least 3 irrigations or 10 days whichever is longer.				

## CONTAINERIZED PLANTS

Container Size (Gallons)	Application Rates			
	Bulk Application		Topical Application	
	Pots/cu. yd. of mix	Bulk Rate lbs./cu. yard	Topical Rate (level teaspoonful)	Grams/pot
1	260 to 230	5.0 to 2.5	1 1/4 to 2 1/2	5 to 10
2	125 to 100	4.4 to 2.2	2 1/2 to 5.0	10 to 20
3	70 to 60	3.3 to 1.7	3 2/3 to 7 1/3	15 to 30
5	50 to 30	2.0 to 1.0	4 1/2 to 9.0	18.5 to 37
For optimum control make applications prior to egg hatch of the target pest. Irrigate moderately after application to move the active ingredient into the root zone.				

## ORNAMENTAL PLANTS GROWN IN FLATS, BENCHES OR BEDS

Application Rate	Application Instructions
15 oz. per 1000 sq. ft.	Apply as a broadcast treatment and incorporate into the soil or media before planting. On established plants, irrigate moderately after application to move the active ingredient into the root zone. Minimize leaching and runoff for at least 3 irrigations or 10 days, whichever is longer.

## FOR CONTROL OF WHITE GRUB LARVAE\* ON ORNAMENTALS IN FIELD AND FOREST NURSERIES

Application Rate	Application Instructions
Use 45 oz. per 1000 sq. ft. of row or 3000 sq. ft.	<p>Vegetation in the area to be treated should be mowed to a height of 3 inches or less prior to application. Mowing to the lowest possible height will ensure greater consistency of control.</p> <p>For grub control in areas of turf, apply as a broadcast application using 45 oz. per 3000 sq. ft.</p> <p>Apply May through July. For optimum control, treatment should be followed by rainfall or irrigation. Apply as a uniform band on either side of row using a band six (6) inches wider than the actual root ball diameter to be dug. Do not allow bands in adjacent rows to overlap.</p>

\* White grub larvae include Japanese beetle, Masked Chafer, Oriental beetle, and Asiatic Garden beetle.

### STORAGE AND DISPOSAL

Do not contaminate water, food or feed by storage or disposal.

**PESTICIDE STORAGE:** Store in a cool, dry place and in such a manner as to prevent cross contamination with other pesticides, fertilizers, food, and feed. Store in original container and out of the reach of children, preferably in a locked storage area.

**PESTICIDE DISPOSAL:** Wastes resulting from the use of the product may be disposed of on site or at an approved waste disposal facility.

### CONTAINER DISPOSAL:

**Paper Bags:** Completely empty bag into application equipment. Then dispose of empty bag in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

**Plastic Container:** Triple rinse (or equivalent). Then offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

### IMPORTANT: READ BEFORE USE

#### LIMITATION OF WARRANTY AND LIABILITY

Read the entire Directions for Use, Conditions of Warranties and Limitations of Liability before using this product. If terms are not acceptable, return the unopened product container at once.

By using this product, user or buyer accepts the following Conditions, Disclaimer of Warranties and Limitations of Liability.

**CONDITIONS:** The directions for use of this product are believed to be adequate and must be followed carefully. However, it is impossible to eliminate all risks associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or the manner of use or application, all of which are beyond the control of FarmSaver.com, LLC. All such risks shall be assumed by the user or buyer.

**DISCLAIMER OF WARRANTIES:** To the extent consistent with applicable law, FarmSaver.com, LLC makes no other warranties, express or implied, of merchantability or of fitness for a particular purpose or otherwise, that extend beyond the statements made on this label. No agent of FarmSaver.com, LLC is authorized to make any warranties beyond those contained herein or to modify the warranties contained herein. To the extent consistent with applicable law, FarmSaver.com, LLC disclaims any liability whatsoever for special, incidental or consequential damages resulting from the use or handling of this product.

**LIMITATIONS OF LIABILITY:** To the extent consistent with applicable law, the exclusive remedy of the user or buyer for any and all losses, injuries or damages resulting from the use or handling of this product, whether in contract, warranty, tort, negligence, strict liability or otherwise, shall not exceed the purchase price paid or at FarmSaver.com, LLC's election, the replacement of product.

Quali-Pro is a trademark of FarmSaver.com, LLC.