

SPECIMEN

GROUP 3 15 HERBICIDE



For use as a preemergence weed control herbicide in ornamentals, landscape and grounds maintenance, and other specified noncrop areas

Active Ingredients:

dimethenamid-P: (S)-2-chloro-N-[(1-methyl-2-methoxy)ethyl]-N-(2,4-dimethyl-thien-3-yl)-acetamide 0.75%

pendimethalin: N-(1-ethylpropyl)-3,4-dimethyl-2, 6-dinitrobenzenamine 1.00%

Other Ingredients: 98.25%

Total: 100.00%

50 lbs contains 0.375 lb of dimethenamid-P and 0.5 lb of pendimethalin.

EPA Reg. No. 7969-273

EPA Est. No.

**KEEP OUT OF REACH OF CHILDREN
CAUTION/PRECAUCIÓN**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand this label, find someone to explain it to you in detail.)

See inside for complete **First Aid, Precautionary Statements, Directions For Use, Conditions of Sale and Warranty,** and state-specific crop and/or use site restrictions.

In case of emergency endangering life or property involving this product, call day or night 1-800-832-HELP (4357).

Net Contents:

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709

 **BASF**
The Chemical Company

FIRST AID	
If swallowed	<ul style="list-style-type: none"> • Call a poison control center or doctor immediately for treatment advice. • DO NOT induce vomiting unless told to do so by a poison control center or doctor. • DO NOT give any liquid to the person. • DO NOT give anything by mouth to an unconscious person.
If in eyes	<ul style="list-style-type: none"> • Hold eyes open and rinse slowly and gently with water for 15 to 20 minutes. • Remove contact lenses, if present, after the first 5 minutes; then continue rinsing eyes. • Call a poison control center for treatment advice.
If on skin or clothing	<ul style="list-style-type: none"> • Take off contaminated clothing. • Rinse skin immediately with plenty of water for 15 to 20 minutes. • Call a poison control center or doctor for treatment advice.
If inhaled	<ul style="list-style-type: none"> • Move person to fresh air. • If person is not breathing, call 911 or an ambulance; then give artificial respiration, preferably by mouth to mouth, if possible. • Call a poison control center or doctor for further treatment advice.
HOT LINE NUMBER	
<p>Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact BASF Corporation for emergency medical treatment information: 1-800-832-HELP (4357).</p>	

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION. Harmful if swallowed. Causes moderate eye irritation. Harmful if absorbed through skin. Avoid contact with skin, eyes or clothing. Avoid breathing dust.

Personal Protective Equipment (PPE)

Some materials that are chemically resistant to this product are listed below. For more options, refer to **Category A** on an EPA chemical-resistance category-selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves made of any waterproof materials, such as barrier laminate, butyl rubber ≥ 14 mils, nitrile rubber ≥ 14 mils, neoprene rubber ≥ 14 mils, polyvinyl chloride ≥ 14 mils, or viton ≥ 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. **DO NOT** reuse them. Follow the manufacturer's instructions for cleaning and maintaining PPE. If no such instructions for washables exist, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to fish. **DO NOT** apply directly to water, to areas where surface water is present, or to intertidal areas below the mean high water mark. Drift and runoff from treated areas may be hazardous to aquatic organisms in adjacent aquatic sites. **DO NOT** contaminate water when disposing of equipment washwaters or rinsate.

Dimethenamid-P has properties that may result in ground-water contamination. Application in areas where soils are permeable or coarse and groundwater is near the surface could result in groundwater contamination. Dimethenamid-P has properties that may result in surface water contamination via dissolved runoff and runoff erosion. Practices should be followed to minimize the potential for dissolved runoff and/or runoff erosion.

Endangered Species Protection

The use of any pesticide in a manner that may kill or otherwise harm an endangered species or adversely modify their habitat is a violation of federal law. To determine whether your county has endangered plant species, consult <http://www.epa.gov/espp/usa-map.htm>. Endangered Species Bulletins may also be obtained from county extension offices or state pesticide agencies. If a bulletin is not

available for your specific area, check with the appropriate local state agency to determine if known populations of endangered plants occur in the area.

To avoid adverse effects on endangered plant species, applicators in ornamentals production must comply with the following mitigation measures where and when endangered plant species are known to occur in proximity of the application site:

If applied by ground, leave untreated buffer zones of 200 feet between treatment area and known endangered plant populations.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling. This labeling must be in the possession of the user at the time of pesticide application.

DO NOT apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your state or tribe, consult the agency responsible for pesticide regulation.

Observe all precautions and limitations in this label and the labels of products used in combination with **FreeHand™ 1.75G herbicide**. The use of **FreeHand 1.75G** not consistent with this label can result in injury to crops, animals, or persons. Keep containers closed to avoid spills and contamination.

Unless otherwise directed in supplemental labeling, all applicable directions, restrictions, precautions and **Conditions of Sale and Warranty** are to be followed.

BASF Corporation does not recommend or authorize the use of this product in manufacturing, processing or preparing custom blends with other products for application to ornamentals.

DO NOT apply **FreeHand 1.75G** aerially or as an aerial application.

DO NOT apply **FreeHand 1.75G** in greenhouses, polyhouses or other fully enclosed greenhouse-type structures.

DO NOT apply **FreeHand 1.75G** to any improved or unimproved maintained turfgrass(es) or lawn(s).

FreeHand 1.75G is not for use by consumers or by persons not under direct professional supervision of a person who is authorized/licensed by the state for this type of application.

DO NOT treat plants grown for food or feed.

DO NOT use treated plants for food or feed.

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL OR CROP INJURY.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT enter or allow worker entry into treated areas during the restricted-entry interval (REI) of **24 hours**.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves made of any waterproof materials, such as barrier laminate, butyl rubber ≥14 mils, nitrile rubber ≥14 mils, neoprene rubber ≥14 mils, polyvinyl chloride ≥14 mils, or viton ≥14 mils
- Shoes plus socks
- Protective eyewear

NONAGRICULTURAL USE REQUIREMENTS

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries, or greenhouses.

DO NOT enter or allow others to enter the treated area until dusts have settled.

PESTICIDE STORAGE AND CONTAINER DISPOSAL

DO NOT contaminate water, food or feed by storage or disposal. Open dumping is prohibited.

Pesticide Storage. DO NOT store above 120° F. Store in original containers and keep closed. Store in a cool, dry place.

Pesticide Disposal. Wastes resulting from the use of this product may be disposed of on-site or at an approved waste disposal facility.

CONTAINER DISPOSAL

(for plastic containers)

Nonrefillable Container. DO NOT reuse or refill this container. After completely emptying container into application equipment, offer for recycling, if available, or reconditioning, if appropriate, or puncture and dispose of in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

(for paper or plastic bags)

Nonrefillable Container. DO NOT reuse or refill this container. After completely emptying container into application equipment, dispose of empty bag in a sanitary landfill, or by incineration, or by other procedures approved by state and local authorities.

IN CASE OF EMERGENCY

In case of large-scale spillage regarding this product, call:

CHEMTREC: 1-800-424-9300
BASF Corporation: 1-800-832-HELP (4357)

In case of medical emergency regarding this product, call:

- Your local doctor for immediate treatment
- Your local poison control center (hospital)
- BASF Corporation: 1-800-832-HELP (4357)

GENERAL INFORMATION

Mode of Action

FreeHand™ 1.75G herbicide combines two active ingredients: dimethenamid-P, a chloroacetamide, and pendimethalin, a dinitroaniline. Combined, they inhibit both plant cell microtubule assembly (cells do not divide and multiply) and cell growth, inhibiting the germinating shoot portion of the susceptible weed seedlings. A small amount of uptake may also occur through the emerging root system of the weed seedling. **FreeHand 1.75G** is **Chemical Group 3** and **Chemical Group 15**, Mode of Action Classification, for broader weed control spectrum.

Use Sites

FreeHand 1.75G may be applied for preemergence control of certain annual grasses, annual broadleaf weeds and sedges as they germinate in:

- Commercial ornamental production nurseries (bed, container, field, liner)
- Landscaped ornamental maintenance areas
- General grounds maintenance in specified noncrop areas

Application Areas

- **Commercial ornamental production nurseries (bed, container, field, liner).** Applications can be made, but are not limited to, plant species listed on this label, such as trees, shrubs, ground covers, perennials, ornamental grasses, bedding plants, conifer and hardwood seedling liner nurseries, tree plantations (including Christmas trees), pulpwood farms and fiber farms.
- **Landscaped ornamental maintenance areas** in and around residential and commercial establishments, multi-family dwellings, military and other institutions, university or college campuses, parks, airports, roadsides, schools, picnic grounds, athletic fields, houses of worship, cemeteries, golf courses, and prairie grass areas, common areas in residential developments, and in non-production areas in commercial nurseries, such as storage areas, vegetation filter strips, windbreaks, shelterbelts or chart paths.

- **General grounds maintenance in the following specified noncrop areas:** parking lots, driveways and roadsides, highway rights-of-way, alleyways, bike and jogging paths, vacant lots, buildings, stone gardens and gravel yards, around statuary or monuments, utility substations, markers/borders and fence lines, and mulch beds. It may be used under asphalt or concrete treatments as part of a site preparation program.

FreeHand 1.75G may be applied as a soft-residual bare-ground treatment in the use sites described in the section on **Application Areas**.

FreeHand 1.75G controls weeds as they germinate but will not control emerged and established weeds. Therefore, areas to be treated should be free of established weeds. For the control of established weeds, **FreeHand 1.75G** may be used in conjunction with herbicides registered for postemergence use. Consult the labels of those herbicides for use rates, timings, and precautions or restrictions.

Unusually cold, excessively wet, or hot and dry conditions that delay germination or extend germination over a long period of time can reduce weed control.

The efficacy of **FreeHand 1.75G** will improve if the application is followed by 1/2 inch of rainfall or its equivalent in sprinkler irrigation. If **FreeHand 1.75G** is not activated by rainfall or irrigation within 30 days, erratic weed control may result.

If weeds should develop prior to activation of the herbicide, shallow cultivate to destroy existing weeds or, where practical, remove by hand. When cultivating for any reason, it should be shallow. **FreeHand 1.75G** may be used before or after applications of herbicides registered for postemergence use (i.e., **Basagran® T/O herbicide**, **Finale® herbicide**, **Roundup® herbicide**, **Segment™ herbicide**) for the control of established weeds. **DO NOT** apply sprays containing **Finale** or **Roundup** over the top of desirable plants. A **FreeHand 1.75G** treatment may be followed by any registered herbicide to control weeds not listed on the **FreeHand 1.75G** label.

Applied according to label directions and under normal growing conditions, **FreeHand 1.75G** will not cause injury to desirable plants. Over-application can result in stand loss, plant injury, or soil residues. Uneven application can decrease weed control or cause plant injury.

Seedling diseases, cold weather, excessive moisture, high soil pH, high soil salt concentration, or drought can weaken seedlings and plants, and increase the possibility of plant damage from **FreeHand 1.75G**.

APPLICATION INFORMATION

Apply with properly calibrated spreader equipment to provide uniform distribution. Check application routinely to determine proper calibration. Avoid overlaps that will increase rates above those stated on this label. Avoid application when winds may cause drift. Avoid unintentional contact of granules with driveways, stone, wood or

other porous surfaces. **FreeHand™ 1.75G herbicide** may cause temporary discoloration of treated surfaces. Sweep surfaces clean immediately to avoid staining. If granules are crushed into surface, rinse thoroughly.

Dry fertilizers. FreeHand 1.75G may be mixed with dry fertilizers. When applied as directed, **FreeHand 1.75G**/dry fertilizer mixtures provide weed control equal to that provided by the same rates of **FreeHand 1.75G** applied alone.

Use Rates

For preemergence control of the listed weed species using broadcast spreader equipment, apply **FreeHand 1.75G** at the following rates.

Table 1. Application Rates for FreeHand 1.75G

Application Rate (lbs product/acre)	Amount (lbs product/1000 sq ft)	Amount (lb product/100 sq ft)
100	2.3	0.23
150	3.4	0.34
200	4.6	0.46

To deliver the proper rate, the applicator must calibrate prior to application of product. Rate settings for a specific type of spreader device must be determined by the applicator. The **FreeHand 1.75G** calibration tray may be used to calibrate the correct application rate.

FreeHand 1.75G may be applied in a single application or in sequential applications. For extended weed control, sequential applications of **FreeHand 1.75G** can be made at a minimum of 8 weeks between applications.

In a single application, **DO NOT** apply more than the equivalent of 200 pounds of **FreeHand 1.75G** per acre.

Application Restrictions and Limitations

- In a single growing season, **DO NOT** apply more than the equivalent of 400 pounds of **FreeHand 1.75G** per acre.
- After applying **FreeHand 1.75G**, wait at least **56 days** before making another application.
- **Stress.** Applications to ornamental plants under stress because of lack of moisture, hail damage, flooding, herbicide injury, mechanical injury, or widely fluctuating temperatures may result in crop injury.
- **DO NOT** contaminate irrigation ditches or water used for domestic purposes.

RECOMMENDED SPECIES

FreeHand 1.75G is safe around and over the top of the established plants listed following. However, not all varieties or strains of the ornamental plants listed have been tested. Refer to the section **APPLICATION INFORMATION** in this label prior to any application of **FreeHand 1.75G**. Unintentional consequences, such as ornamental injury, may result because of certain environmental or growing conditions, manner of use or application. Therefore, before treating a large number of plants, apply to a few plants and observe for plant damage prior to full-scale application.

ORNAMENTALS

FreeHand 1.75G may be used on ornamental plant species not listed on this label. The suitability for such uses should be determined by treating a small number of such plants at the recommended rate. Treated plants should be evaluated 1 to 2 months following treatment for possible injury. **TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE USER ASSUMES RESPONSIBILITY FOR ANY CROP DAMAGE OR OTHER LIABILITY ASSOCIATED WITH APPLICATIONS TO ORNAMENTAL PLANTS NOT LISTED ON THIS LABEL.**

TREES

Common Name	Scientific Name
Arborvitae	<i>Thuja</i> spp.
Crape myrtle	<i>Lagerstroemia indica</i>
Fir, Fraser	<i>Abies fraseri</i>

SHRUBS

Common Name	Scientific Name
Abelia, glossy	<i>Abelia grandiflora</i>
Azalea	<i>Rhododendron</i> spp.
Boxwood, Japanese	<i>Buxus microphylla</i>
Butterfly bush†	<i>Buddleia davidii</i>
Euonymus	<i>Euonymus</i> spp.
Gardenia	<i>Gardenia jasminoides</i>
Hawthorne, Indian	<i>Rhaphiolepis indica</i>
Holly	<i>Ilex</i> spp.
Holly, Chinese	<i>Ilex cornuta</i>
Holly, Japanese	<i>Ilex crenata</i>
Hydrangea†	<i>Hydrangea macrophylla</i>
Juniper	<i>Juniperus</i> spp.
Nandina	<i>Nandina domestica</i>
Rose	<i>Rosa</i> spp.
Spiraea	<i>Spiraea</i> spp.
Viburnum	<i>Viburnum</i> spp.

†Crop injury is possible in certain cultivars of these species; therefore, it is recommended to apply **FreeHand 1.75G** at the lowest use rate and apply only to a small number of plants prior to treating an entire block of plants.

HERBACEOUS PERENNIALS

Common Name	Scientific Name
Daylily [†]	<i>Hemerocallis</i> spp.
Flag [†]	<i>Iris</i> spp.
Hosta	<i>Hosta</i> spp.
Lantana	<i>Lantana</i> spp.
Sage	<i>Salvia</i> spp.
Sage, Ramona	<i>Salvia sylvestris</i>

[†]Crop injury is possible in certain cultivars of these species; therefore, it is recommended to apply **FreeHand™ 1.75G herbicide** at the lowest use rate and apply only to a small number of plants prior to treating an entire block of plants.

BEDDING PLANTS

After planting bedding plants and before **FreeHand 1.75G** is applied, the area to be treated should be watered sufficiently to pack soil around roots.

Common Name	Scientific Name
Periwinkle, rose	<i>Catharanthus roseus</i>
Petunia	<i>Petunia</i> spp.

ORNAMENTAL PRECAUTIONS

- Apply **FreeHand 1.75G** to established plantings.
- **DELAY** applying **FreeHand 1.75G** to seedbeds, transplant beds, or liners until plants have become well rooted.
- For container-grown ornamentals, delay first application of this product to bare-root liners 2 to 4 weeks.
- Care must be taken that soil or planting mixes have settled firmly following transplanting and there are no cracks that would allow direct contact of **FreeHand 1.75G** with roots. Applications where soil and media surfaces are uniformly covered will result in best weed control and ornamental tolerance.
- Plant only those desirable plant species listed on this label into soil treated the previous season with any **FreeHand 1.75G** product or injury may occur.

LANDSCAPE AND GROUNDS MAINTENANCE

FreeHand 1.75G can be incorporated into landscape and grounds maintenance programs to provide extended pre-emergence control of most annual grasses, certain broadleaf weeds, and sedges. Areas to be treated include those areas where gravel, pebble, stones, rock and other porous surfaces are used instead of organic mulches as bedding for landscaping. Areas to be treated should be free of emerged weeds before application. To remove emerged weeds, cultivate **OR** apply a postemergence product labeled for such use.

TREE PLANTATIONS

FreeHand 1.75G can be used for preemergence weed control during site preparation, establishment and/or maintenance of tree plantations (including Christmas trees). **FreeHand 1.75G** may also be used for hardwood and conifer regeneration on Conservation Reserve Program land or similar areas.

FreeHand 1.75G may be applied at planting or to established trees. When making an application at planting, it is important that slit closure be achieved to prevent **FreeHand 1.75G** from directly contacting the tree roots or being washed into the root zone via the open slit, or root stunting may occur. **DO NOT** apply to newly transplanted seedlings until plants have been watered and soil has been thoroughly packed and settled around roots.

WEED SPECIES CONTROLLED

Use **FreeHand 1.75G** for preemergence control of the following weed species listed according to use rate.

Weeds controlled when **FreeHand 1.75G** is applied at 100 lbs/acre (2.3 lbs **FreeHand 1.75G**/1000 sq ft)

Common Name	Scientific Name
Amaranth, Palmer	<i>Amaranthus palmeri</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Carpetweed	<i>Mollugo verticillata</i>
Foxtail, giant	<i>Setaria faberi</i>
Foxtail, green	<i>Setaria viridis</i>
Foxtail, yellow	<i>Setaria glauca</i>
Henbit	<i>Lamium amplexicaule</i>
Lambsquarters, common	<i>Chenopodium album</i>
Pigweed, prostrate	<i>Amaranthus blitoides</i>
Pigweed, redroot	<i>Amaranthus retroflexus</i>
Pigweed, smooth	<i>Amaranthus hybridus</i>
Pigweed, tumble	<i>Amaranthus albus</i>
Pusley, Florida	<i>Richardia scabra</i>

150 lbs/A

In addition to the weeds controlled at 100 lbs **FreeHand™ 1.75G herbicide**/acre application rate, the following weeds will be controlled when **FreeHand 1.75G** is applied at 150 lbs/acre (3.4 lbs **FreeHand 1.75G**/1000 sq ft)

Common Name	Scientific Name
Amaranth, Powell	<i>Amaranthus powellii</i>
Bittercress	<i>Cardamine</i> spp.
Bluegrass, annual	<i>Poa annua</i>
Bluegrass, roughstalk	<i>Poa trivialis</i>
Brome, California	<i>Bromus carinatus</i>
Brome, downy	<i>Bromus tectorum</i>
Chamomile, mayweed	<i>Anthemis cotula</i>
Chickweed, common	<i>Stellaria media</i>
Chickweed, mouseear	<i>Cerastium vulgatum</i>
Crabgrass, large	<i>Digitaria sanguinalis</i>
Crabgrass, smooth	<i>Digitaria ischaemum</i>
Cudweed	<i>Gnaphalium</i> spp.
Cupgrass, Southwestern	<i>Eriochloa gracilis</i>
Doveweed	<i>Murdannia nudiflora</i>
Evening primrose	<i>Oenothera biennis</i>
Flatsedge, rice	<i>Cyperus iria</i>
Goosegrass	<i>Eleusine indica</i>
Johnsongrass (from seed)	<i>Sorghum halepense</i>
Knotweed, prostrate	<i>Polygonum aviculare</i>
Nightshade, black	<i>Solanum nigrum</i>
Nightshade, Eastern black	<i>Solanum ptycanthum</i>
Nightshade, cutleaf	<i>Solanum triflorum</i>
Nightshade, hairy	<i>Solanum sarrachoides</i>
Nutsedge, yellow	<i>Cyperus esculentus</i>
Pearlwort*	<i>Sagina procumbens</i> or <i>S. decumbens</i>
Purslane, common	<i>Portulaca oleracea</i>
Red rice	<i>Oryza sativa</i>
Ryegrass, Italian	<i>Lolium multiflorum</i>
Sandbur, field	<i>Cenchrus incertus</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Signalgrass, broadleaf	<i>Brachiaria platyphylla</i>
Speedwell, corn	<i>Veronica arvensis</i>
Spurges, annual	<i>Chamaesyce</i> spp., <i>Euphorbia</i> spp.
Waterhemp, common	<i>Amaranthus rudis</i>
Waterhemp, tall	<i>Amaranthus tuberculatus</i>
Witchgrass	<i>Panicum capillare</i>

*Not controlled in California.

200 lbs/A

In addition to the weeds controlled at 100 lbs and 150 lbs **FreeHand 1.75G**/acre application rates, the following weeds will be controlled when **FreeHand 1.75G** is applied at 200 lbs/acre (4.6 lbs **FreeHand 1.75G**/1000 sq ft)

Common Name	Scientific Name
Burnweed, American*	<i>Erechtites hieraciifolia</i>
Burweed, lawn	<i>Soliva pterosperma</i>
Clover, hop	<i>Trifolium procumbens</i>
Crowfootgrass	<i>Dactyloctenium aegyptium</i>
Cupgrass, woolly	<i>Eriochloa villosa</i>
Dogfennel*	<i>Eupatorium capillifolium</i>
Eclipta	<i>Eclipta alba</i> <i>E. prostrata</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree	<i>Erodium</i> spp.
Itchgrass	<i>Rottboellia exaltata</i>
Junglerice	<i>Echinochloa colona</i>
Kochia	<i>Kochia scoparia</i>
Kyllinga species*	<i>Kyllinga</i> spp.
Liverwort*	<i>Marchantia polymorpha</i>
Lovegrass (from seed)	<i>Eragrostis</i> spp.
Mulberry weed*	<i>Fatoua villosa</i>
Panicum, browntop	<i>Panicum fasciculatum</i>
Panicum, fall	<i>Panicum dichotomiflorum</i>
Panicum, Texas	<i>Panicum texanum</i>
Pepperweed*	<i>Lepidium virginicum</i>
Phyllanthus, long-stalked*	<i>Phyllanthus tenellus</i>
Puncturevine	<i>Tribulus terrestris</i>
Ragweed, common	<i>Ambrosia artemisiifolia</i>
Rocket, London	<i>Sisymbrium irio</i>
Smartweed, Pennsylvania	<i>Polygonum pennsylvanicum</i>
Sowthistle	<i>Sonchus arvensis</i>
Sprangletop, Mexican	<i>Leptochloa uninervia</i>
Sprangletop, red	<i>Leptochloa filiformis</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Willowherb	<i>Epilobium</i> spp.
Woodsorrel, yellow	<i>Oxalis stricta</i>

*Not controlled in California.

Conditions of Sale and Warranty

The **Directions For Use** of this product reflect the opinion of experts based on field use and tests. The directions are believed to be reliable and must be followed carefully. However, it is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials, or use of the product in a manner inconsistent with its labeling, all of which are beyond the control of BASF CORPORATION ("BASF") or the Seller. To the extent consistent with applicable law, all such risks shall be assumed by the Buyer.

BASF warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes referred to in the **Directions For Use**, subject to the inherent risks, referred to above.

To the extent consistent with applicable law, BASF makes no other express or implied warranty of fitness or merchantability or any other express or implied warranty.

To the extent consistent with applicable law, Buyer's exclusive remedy and BASF's exclusive liability, whether in contract, tort, negligence, strict liability, or otherwise, shall be limited to repayment of the purchase price of the product.

To the extent consistent with applicable law, BASF and the Seller disclaim any liability for consequential, special or indirect damages resulting from the use or handling of this product.

BASF and the Seller offer this product, and the Buyer and User accept it, subject to the foregoing **Conditions of Sale and Warranty** which may be varied only by agreement in writing signed by a duly authorized representative of BASF.

0408

***Basagran** is a registered trademark of BASF.*

***FreeHand** and **Segment** are trademarks of BASF.*

***Finale** is a registered trademark of Bayer.*

***Roundup** is a registered trademark of Monsanto Technology, LLC.*

© 2008 BASF Corporation
All rights reserved.

007969-00273.20080513.**NVA 2008-04-308-0166**
Based on: NVA 2007-04-308-0141

BASF Corporation
26 Davis Drive
Research Triangle Park, NC 27709


The Chemical Company