



Purina Mills

## CORNERSTONE® 22:20 BOV BM

Patent No. 5,571,542

Milk replacer to be fed to herd replacement calves.  
For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves.

### ACTIVE DRUG INGREDIENTS

Lasalocid ..... 30 mg/pound  
(as lasalocid sodium) Equal to 60 g/ton

### GUARANTEED ANALYSIS

Crude Protein, not less than ..... 22.0%  
Crude Fat, not less than ..... 20.0%  
Crude Fiber, not more than ..... 0.15%  
Calcium (Ca), not less than ..... 0.75%  
Calcium (Ca), not more than ..... 1.25%  
Phosphorus (P), not less than ..... 0.70%  
Vitamin A, not less than ..... 30,000 IU/lb  
Vitamin D<sub>3</sub>, not less than ..... 5,000 IU/lb  
Vitamin E, not less than ..... 100 IU/lb

### INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Dried Skimmed Milk, Dried Milk Protein, Animal Fat (preserved with Ethoxyquin), Lecithin, Polysorbate 80, DiCalcium Phosphate, L-Lysine, DL-Methionine, Calcium Carbonate, Brewers Dried Yeast, Dried *Saccharomyces cerevisiae* Fermentation Solubles, Vitamin A Acetate, Vitamin D<sub>3</sub> Supplement, Vitamin E Supplement, Thiamine Mononitrate, Riboflavin, Niacin Supplement, Biotin, Ascorbic Acid, Pyridoxine Hydrochloride, Folic Acid, Vitamin B<sub>12</sub> Supplement, Choline Chloride, Roughage Products, Inulin, Calcium Silicate, Zinc Methionine, Manganese Methionine, Copper Lysine, Iron Amino Acid Complex, Cobalt Sulfate, Ethylenediamine Dihydriodide, Selenium Yeast and Natural & Artificial Flavor.

Important: Follow feeding directions.

**CAUTION:** The safety of lasalocid in unapproved species has not been established. This product is to be fed to calves only.

**WARNING:** A withdrawal period has not been established for lasalocid in pre-ruminating calves.

Distributed by:  
PURINA MILLS LLC  
P.O. BOX 66812  
St. Louis, Missouri 63166-6812

© Registered trademark of Land O'Lakes Purina Feed



8 04273 02769 2

0041974

105Z

010

## 50 LB (22.68 kg) Net Weight

### MIXING AND FEEDING DIRECTIONS

- Always weigh milk replacer powder for accurate mixing.
- Mix milk replacer powder according to the schedule below by adding powder to water as indicated and mix thoroughly.

### FEEDING DIRECTIONS (2 Days to Weaning)

Always offer free choice water.

Feed the following amounts of milk replacer twice daily per calf.

|  | MILK REPLACER<br>POWDER | WATER<br>(110-120° F) |
|--|-------------------------|-----------------------|
| <b>Individual Calf</b>   |                         |                       |
| Recommended  | 12 oz.                  | + 2 quarts            |
| <b>Large Batch</b>   |                         |                       |
| Recommended  | 1.5 lb. (24 oz.)        | + 1 gallon            |
| <b>Large Breeds</b> (100 lb of body weight): Feed 2.3 quarts or fill 2 quart bottle to rim and feed twice daily. |                         |                       |

This product contains 30 milligrams of lasalocid per pound. Feed to provide 45 milligrams of lasalocid per 100 pounds (1 mg/2.2 lbs.) of body weight daily. For example, 1.5 lb of powder contains the proper level of lasalocid for a 100 lb calf.

Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

**IMPORTANT:** Thoroughly mix milk replacer powder in warm water, and feed immediately. When preparing large batches (i.e., anything other than individual calf servings) keep the mixture agitated while dispensing into individual feeding containers to prevent any settling which could result in the feeding of improper levels of lasalocid.

- A characteristic of this product is a slight thickening that occurs after a prolonged period of time. If thickening occurs, remix before feeding.
- Wean the calf when eating a minimum of 2 lbs. of Purina Calf Startena® per day, for three consecutive days (5-7 weeks).
- Feed milk replacer solution at body temperature (100°F).
- Provide free-choice clean water to calves daily, starting on day 2.
- Feed a Purina Calf Startena® ration free-choice starting on the 4th day through the end of the 2nd month of age. Provide fresh starter daily.

**CAUTION:** Store in dry area away from insects and rodents. Do not feed moldy or rancid feed to animals as it may cause illness or death.

QUALITY CONTROLLED BY PURINA RESEARCH  
**DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.**  
**DO NOT FEED TO LACTATING COWS.**