



INSTANT COW'S MATCH® BOV BM MEDICATED DAIRY HERD & BEEF CALF MILK REPLACER

Patent No. 5,571,542
For control of coccidiosis caused by *Eimeria bovis* and *Eimeria*
Zuernii in replacement calves.

ACTIVE DRUG INGREDIENTS

Lasalocid 30 mg/pound
(as lasalocid sodium) Equal to 60 g/ton

GUARANTEED ANALYSIS

Crude Protein, not less than 28.0%
Crude Fat, not less than 20.0%
Crude Fiber, not more than 0.15%
Calcium (Ca), not less than 0.75%
Calcium (Ca), not more than 1.25%
Phosphorus (P), not less than 0.70%
Vitamin A, not less than 20,000 IU/lb
Vitamin D₃, not less than 5,000 IU/lb
Vitamin E, not less than 100 IU/lb

INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey
Product, Dried Skimmed Milk, Dried Milk Protein, Animal Fat
(preserved with Ethoxyquin), Lecithin, Polysorbate 80,
DiCalcium Phosphate, Calcium Carbonate, Dried
Saccharomyces cerevisiae Fermentation Extract, Brewers
Dried Yeast, Dried Saccharomyces, Cerevisiae
Fermentation Solubles, Methionine Supplement, L-Lysine,
Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E
Supplement, Thiamine Mononitrate, Riboflavin, Niacin
Supplement, Calcium Pantothenate, Biotin, Ascorbic Acid,
Pyridoxine Hydrochloride, Folic Acid, Vitamin B₁₂
Supplement, Choline Chloride, Roughage Products,
Calcium Silicate, Zinc Methionine, Manganese Methionine,
Copper Lysine, Iron Amino Acid Complex, Cobalt Sulfate,
Ethylenediamine Dihydrochloride, Selenium Yeast and Natural
& Artificial Flavor.

© Registered trademark of Land O'Lakes, Inc.

**DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL
DO NOT FEED TO LACTATING COWS.**

**"This feed was made in a feed manufacturing facility
that does not handle or store products containing
animal proteins prohibited in ruminant feeds."**

INTENSIFIED CALF FEEDING AND MANAGEMENT PROGRAM

Cow's Match calf milk replacer should be fed along with a spe-
cific calf starter designed for intensified feeding. Calf to calf
variations such as calf weight and health must be con-
sidered in implementing this program. Consult with your
feed specialist for specific feeding and management rec-
ommendations to match with your calf performance
goals and expectations.



1485252-106

Manufactured By:

Land O'Lakes Animal Milk Products Co.

SHOREVIEW, MN 55126-2910

50 LB (22.68 kg) Net Weight

MIXING DIRECTIONS

5 Gallon Mix: While stirring, mix 7.5 lbs. of milk replacer powder
in 3 to 4 gallons of 110-120°F water and then add 110-120°F
water to bring final solution to 5 gallons. Powder should be
weighed for precision and best calf performance.

Individual Calf

Week 1: While stirring, pour 0.94 lbs. (15 oz.) of milk replacer
powder into 1.5 quarts of 110-120°F water. Mix thoroughly and
add 110-120°F water to bring to a total volume of 2.5 quarts of
milk replacer solution.

Week 2 through Week 6: While stirring, pour 1.3 lbs. (21 oz.) of
milk replacer powder into 2.5 quarts of 110-120°F water. Mix thor-
oughly and add 110-120°F water to bring to a total volume of 3.5
quarts of milk replacer solution. When under heat stress, delay
this increase until week 3.

Week 7: While stirring, pour 1.3 lbs. (21 oz.) of milk replacer
powder into 2.5 quarts of 110-120°F water. Mix thoroughly and
add 110-120°F water to bring to a total volume of 3.5 quarts of
milk replacer solution.

FEEDING DIRECTIONS (2 Days to End of Week 7)

ALWAYS offer free choice water.

LARGE BREEDS: (125 - 175 lbs. body weight)

Week 1: Feed 2.5 quarts of milk replacer solution twice daily.

Week 2 through Week 6: Feed 3.5 quarts of milk
replacer solution twice daily. When under heat stress,
delay this increase until week 3.

Week 7: Feed 3.5 quarts of milk replacer solution once
per day. Wean only if calf is eating at least 1.5 lbs. of
starter for 3 consecutive days.

(The above mixing and feeding rates are designed for optimal calf growth from
Cow's Match calf milk replacer.)

This product contains 30 milligrams of lasalocid per pound. Feed to provide
45 milligrams of lasalocid per 100 pounds (1 mg/2.2 lbs.) of body weight daily.
Feed during periods of coccidiosis exposure or when experience
indicates coccidiosis is likely to be a problem.

IMPORTANT: Thoroughly mix milk replacer powder in warm
water, and feed immediately. When preparing large batches (i.e.,
anything other than individual calf servings) keep the mixture agi-
tated while dispensing into individual feeding containers to pre-
vent any settling which could result in the feeding of improper lev-
els of lasalocid.

A characteristic of this product is a slight layering that
occurs after a prolonged period of time. If layering occurs,
re-mix before feeding.

GENERAL RECOMMENDATIONS

1. Feed newborn calves 3 quarts of high quality warm
colostrum via nipple bottle within 1 hour of birth and
repeat 12 hours later (or) feed 4 quarts of high qual-
ity warm colostrum by esophageal feeder within 1
hour of birth and repeat 12 hours later via nipple bot-
tle offering as much as they will consume.
2. Consult with your veterinarian on a dry cow and calf vac-
cination program including but not limited to E. Coli, rota
and corona virus, clostridium and salmonella.
3. Begin feeding milk replacer on day 2 and provide fresh,
clean water continuously on a free-choice basis.
4. Begin feeding a calf starter recommended for intensive
feeding at two days of age and offer continuous-
ly on a free-choice basis.
5. At the start of week 7, transition calves from twice per
day feeding to once per day feeding of Cow's Match.
This will encourage the calf to eat more calf starter
and assures an effective weaning.
6. Wean only if calf is eating at least 1.5 lbs. of starter
for 3 consecutive days.
7. Continue to feed appropriate intensive calf starter up
to 12 weeks of age - no hay fed separately.
8. Begin feeding a calf grower recommended for intensive
feeding at 12 weeks of age at the recommended rate.

CAUTION: The safety of lasalocid in unapproved
species has not been established. This
product is to be fed to calves only.

WARNING: A withdrawal period has not been
established for lasalocid in pre-rumi-
nating calves.

069