



INSTANT MAXI CARE® PLUS BOV BM MEDICATED DAIRY HERD & BEEF CALF MILK REPLACER

For control of coccidiosis caused by *Eimeria bovis* and *Eimeria zuernii* in replacement calves.

ACTIVE DRUG INGREDIENT

Lasalocid (as lasalocid sodium) . . . 36 mg/pound
Equal to 72 g/ton

When mixed and fed according to directions, 45 milligrams of lasalocid will be provided per 100 pounds of body weight daily.

GUARANTEED ANALYSIS

Crude Protein, not less than25.0%
Crude Fat, not less than20.0%
Crude Fiber, not more than0.25%
Calcium (Ca), not less than0.75%
Calcium (Ca), not more than1.25%
Sodium (Na), not more than1.0%
Phosphorus (P), not less than0.70%
Vitamin A, not less than20,000 IU/lb
Vitamin D₃, not less than5,000 IU/lb
Vitamin E, not less than100 IU/lb

INGREDIENTS

Dried Whey, Dried Whey Protein Concentrate, Dried Whey Product, Dried Skimmed Milk, Dried Milk Protein, Soy Isolate, Protein Modified Soy Flour, Animal Plasma, Animal Fat (preserved with Ethoxyquin), L-Lysine, DL-Methionine, Lecithin, Polysorbate 80, DiCalcium Phosphate, Calcium Carbonate, Dried Saccharomyces cerevisiae Fermentation Extract, Brewers Dried Yeast, Dried Saccharomyces cerevisiae Fermentation Solubles, Vitamin A Acetate, Vitamin D₃ Supplement, Vitamin E Supplement, Thiamine Mononitrate, Pyridoxine Hydrochloride, Folic Acid, Vitamin B₁₂ Supplement, Choline Chloride, Calcium Silicate, Zinc Methionine, Manganese Methionine, Copper Lysine, Iron Amino Acid Complex, Cobalt Sulfate, Ethylenediamine Dihydrochloride, Selenium Yeast and Natural & Artificial Flavor.



1520010-106

Manufactured By:

Land O'Lakes Animal Milk Products Co.
SHOREVIEW, MN 55126-2910

50 LB (22.68 kg) Net Weight

MIXING DIRECTIONS

• Always weigh milk replacer powder for accurate mixing.

• Add 10 ounces of milk replacer powder to 1.5 quarts of 110-120° F. water and mix thoroughly. Add 110-120° F. water to bring final solution up to 2 quarts and mix thoroughly. Feed immediately after mixing. Each quart of milk replacer will then contain 11.25 milligrams of lasalocid.

*Add 10 ounces milk replacer powder to provide proper lasalocid (Bovatec) medication level.

FEEDING DIRECTIONS (2 Days to Weaning)

Always offer free choice water.

Feed the following amounts of milk replacer twice daily per calf.

Calf Weight (lbs.)	Milligrams Lasalocid Required Per Calf Per Day	Quarts Fed Per Feeding Twice-A-Day
75	34	1.5
100	45	2.0
125	57	2.5
150	68	3.0

Feed during periods of coccidiosis exposure or when experience indicates coccidiosis is likely to be a problem.

IMPORTANT: Thoroughly mix milk replacer powder in warm water, and feed immediately. When preparing large batches (i.e., anything other than individual calf servings) keep the mixture agitated while dispensing into individual feeding containers to prevent any settling which could result in the feeding of improper levels of lasalocid.

A characteristic of this product is a slight layering that occurs after a prolonged period of time. If layering occurs, re-mix before feeding.

GENERAL RECOMMENDATIONS

1. Feed newborn calves 3 quarts of high quality warm colostrum via nipple bottle within 1 hour of birth and repeat 12 hours later (or) feed 4 quarts of high quality warm colostrum by esophageal feeder within 1 hour of birth and repeat 12 hours later via nipple bottle as much as they will consume.
2. Consult with your veterinarian on a dry cow and calf vaccination program including but not limited to E. Coli, rota and corona virus, clostridium and salmonella.
3. For best mixing, continuously stir with wire whip while adding powder to 110-120° F. water (hot as you can tolerate on your hand). Use correct water temperature to avoid product separation. Feed milk replacer at 100 -105° F.
4. Begin feeding milk replacer on day 2 and provide fresh, clean water along with a high quality, palatable calf starter on a free-choice basis.
5. Observe calves closely during the milk replacer feeding period. Avoid underfeeding, which may result in starvation, or overconsumption which may increase incidence of scours.
6. Continue to feed milk replacer until the calf is consuming a minimum of 1.5 pounds of calf starter per day, which usually occurs at 4-6 weeks of age.

CAUTION: The safety of lasalocid in unapproved species has not been established. This product is to be fed to calves only.

WARNING: A withdrawal period has not been established for lasalocid in pre-ruminating calves.
DO NOT USE IN CALVES TO BE PROCESSED FOR VEAL.
DO NOT FEED TO LACTATING COWS.

"This feed was made in a feed manufacturing facility that does not handle or store products containing animal proteins prohibited in ruminant feeds."

© Registered Trademark of Land O'Lakes, Inc.

039