

IMPORTANT CORRECTION

There is an error in a drug dosage protocol on page 365 of Ignatavicius, Workman, & Rebar: *Medical-Surgical Nursing: Concepts for Interprofessional Collaborative Care*, 9th Edition, ISBNs 978-0-323-44419-4 (single volume), 978-0-323-46158-0 (two-volume set), and 978-0-323-63947-7 (binder-ready version). This issue also affects the following editions: 8 (p. 353), 7 (pp. 389-390), and 6 (pp. 392-393).

Error and Correction Information:

On p. 365, in Chart 20-3 and in the text below, the protocol for epinephrine administration is described as follows:

- Be prepared to administer epinephrine IV (preferred) or IM.
 - Epinephrine 1 : 1000 concentration, 0.3 to 0.5 mL IV push or IM

and

“Epinephrine (1 : 1000) 0.3 to 0.5 mL is the first-line drug for anaphylaxis. It is given IM or IV when symptoms appear (see Chart 20-3).”

This protocol should be described as follows:

- Be prepared to administer epinephrine IM.
 - Epinephrine 1 mg/mL concentration, 0.3 to 0.5 mL IM

and

“Epinephrine 1 mg/mL concentration, 0.3 to 0.5 mL IM is the first-line drug for anaphylaxis It is given when symptoms appear (see Chart 20-3).”

Please make these important changes in your printed copy of the textbook, or be sure to download the latest digital version of the book.

IMPORTANT CORRECTION

There is an error in a drug dosage protocol on page 365 of Ignatavicius, Workman, & Rebar: *Medical-Surgical Nursing: Concepts for Interprofessional Collaborative Care*, 9th Edition, ISBNs 978-0-323-44419-4 (single volume), 978-0-323-46158-0 (two-volume set), and 978-0-323-63947-7 (binder-ready version). This issue also affects the following editions: 8 (p. 353), 7 (pp. 389-390), and 6 (pp. 392-393).

Error and Correction Information:

On p. 365, in Chart 20-3 and in the text below, the protocol for epinephrine administration is described as follows:

- Be prepared to administer epinephrine IV (preferred) or IM.
 - Epinephrine 1 : 1000 concentration, 0.3 to 0.5 mL IV push or IM

and

“Epinephrine (1 : 1000) 0.3 to 0.5 mL is the first-line drug for anaphylaxis. It is given IM or IV when symptoms appear (see Chart 20-3).”

This protocol should be described as follows:

- Be prepared to administer epinephrine IM.
 - Epinephrine 1 mg/mL concentration, 0.3 to 0.5 mL IM

and

“Epinephrine 1 mg/mL concentration, 0.3 to 0.5 mL IM is the first-line drug for anaphylaxis It is given when symptoms appear (see Chart 20-3).”

Please make these important changes in your printed copy of the textbook, or be sure to download the latest digital version of the book.